Nascent Transcription Affected by RNA Polymerase IV in *Zea mays*

Karl F. Erhard Jr.,^{*1,2} Joy-El R. B. Talbot,^{†,*,1,3} Natalie C. Deans,[‡] Allison E. McClish,[§] and Jay B. Hollick^{*,†,3,4}

^{*}Department of Plant and Microbial Biology, University of California, Berkeley, California 94720-3102, [†]Department of Molecular and Cell Biology, University of California, Berkeley, California 94720-3200, [‡]Department of Molecular Genetics, Center for RNA Biology, The Ohio State University, Columbus, Ohio 43210, and [§]Department of Biology, Albion College, Albion, Michigan 49224

ORCID IDs: 0000-0001-5334-3806 (J.R.B.T.); 0000-0001-9749-7647 (N.C.D.)

ABSTRACT All eukaryotes use three DNA-dependent RNA polymerases (RNAPs) to create cellular RNAs from DNA templates. Plants have additional RNAPs related to Pol II, but their evolutionary role(s) remain largely unknown. *Zea mays* (maize) RNA polymerase D1 (RPD1), the largest subunit of RNA polymerase IV (Pol IV), is required for normal plant development, paramutation, transcriptional repression of certain transposable elements (TEs), and transcriptional regulation of specific alleles. Here, we define the nascent transcriptomes of *rpdl* mutant and wild-type (WT) seedlings using global run-on sequencing (GRO-seq) to identify the broader targets of RPD1-based regulation. Comparisons of WT and *rpdl* mutant GRO-seq profiles indicate that Pol IV globally affects transcription at both transcriptional start sites and immediately downstream of polyadenylation addition sites. We found no evidence of divergent transcription from gene promoters as seen in mammalian GRO-seq profiles. Statistical comparisons identify genes and TEs whose transcription is affected by RPD1. Most examples of significant increases in genic antisense transcription appear to be initiated by 3'-proximal long terminal repeat retrotransposons. These results indicate that maize Pol IV specifies Pol II-based transcriptional regulation for specific regions of the maize genome including genes having developmental significance.

KEYWORDS RNA polymerase IV; transcription; gene regulation; transposons; paramutation

EUKARYOTES use at least three DNA-dependent RNA polymerases (RNAPs) to transcribe their genomes into functional RNAs. RNAP Pol II generates messenger RNAs (mRNAs) and noncoding RNAs involved in various RNA-mediated regulatory pathways (reviewed by Sabin *et al.* 2013). Flowering plant genomes encode additional RNAP subunits comprising Pol IV and Pol V, which are central to a small interfering RNA (siRNA)-based silencing pathway primarily targeting repetitive sequences such as transposable elements (TEs) (Matzke and

Mosher 2014; Matzke *et al.* 2015). These additional RNAPs derive from duplications of specific Pol II subunits followed by subfunctionalization during plant evolution (Tucker *et al.* 2011), yet the holoenzyme complexes still share some Pol II subunits (Ream *et al.* 2009; Haag *et al.* 2014).

Zea mays (maize) has distinct largest subunits for Pol IV and V and, unlike *Arabidopsis thaliana*, three second-largest subunits (Erhard *et al.* 2009; Sidorenko *et al.* 2009; Stonaker *et al.* 2009) that in distinct combinations form two Pol IV and three Pol V isoforms (Haag *et al.* 2014). Genetic analyses of *rna polymerase d/e 2a* (*rpdl/e2a*) encoding one of the second-largest subunits (Sidorenko *et al.* 2009; Stonaker *et al.* 2009) together with recent proteomic data showing association of a putative RNA-dependent RNA polymerase (RDR2) with only RPD/E2a-containing isoforms (Haag *et al.* 2014) indicate that maize Pol IV isoforms have diverse functional roles in managing genome homeostasis.

Loss of Pol IV function has different consequences in *Arabidopsis*, *Brassica rapa* (a close relative of *Arabidopsis*), and maize, species in which Pol IV mutants have been identified (Herr *et al.* 2005; Onodera *et al.* 2005; Erhard *et al.* 2009; Huang *et al.* 2013), although mutants in all three

Copyright © 2015 by the Genetics Society of America
doi: 10.1534/genetics.115.174714

Manuscript received September 23, 2014; accepted for publication February 2, 2015;
published Early Online February 4, 2015.

Supporting information is available online at <http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.174714/-DC1>.

Sequencing data have been deposited in the Gene Expression Omnibus database under accession no. GSE54166.

¹These authors contributed equally to this work.

²Present address: Children's Hospital Oakland Research Institute, Oakland, CA 94609.

³Present address: Department of Molecular Genetics, Center for RNA Biology, The Ohio State University, Columbus, OH 43210.

⁴Corresponding author: Department of Molecular Genetics, Center for RNA Biology, The Ohio State University, 500 Aronoff Laboratory, 318 West 12th Ave., Columbus, OH 43210. E-mail: hollick.3@osu.edu

species are defective for siRNA production (Zhang *et al.* 2007; Mosher *et al.* 2008; Erhard *et al.* 2009; Huang *et al.* 2013). *Arabidopsis* NUCLEAR RNA POLYMERASE D1A (*NRPD1A*) mutants are late flowering (Pontier *et al.* 2005) and *B. rapa* *nrpd1a* mutants have no obvious phenotypes (Huang *et al.* 2013), while maize *rna polymerase d1* (*rpdl*) mutants have multiple developmental defects and *trans*-generational degradation in plant quality compared to nonmutant siblings (Parkinson *et al.* 2007; Erhard *et al.* 2009). The disparate impacts of *rpdl*/*NRPD1A* mutations in maize vs. Brassicaceae representatives are potentially related to different genomic TE contents as TE sequences are greatly expanded in maize compared to both *Arabidopsis* (Hale *et al.* 2009) and *B. rapa* (Wang *et al.* 2011). However, recently reported cytosine methylome profiles (Li *et al.* 2014) indicate maize TEs are as equally well methylated in the absence of Pol IV as their *Arabidopsis* counterparts (Stroud *et al.* 2013), predicting that Pol IV-dependent cytosine methylation is not required to maintain TE silencing.

The developmental defects observed in *rpdl* mutants are both distinct and nonheritable (Parkinson *et al.* 2007) and therefore unlikely to be related to TE-derived mutations. These defects also appear unrelated to siRNA-induced silencing because other maize mutants affecting siRNA biogenesis, including *rpdl/e2a*, are developmentally normal (Hale *et al.* 2007; Stonaker *et al.* 2009; Barbour *et al.* 2012). We hypothesize that maize has co-opted RPD1/Pol IV to transcriptionally control specific alleles of genes for which TEs and TE-like repeats act as regulatory elements. Supporting this concept, specific *purple plant1* (*pl1*) alleles having an upstream *doppia* TE fragment are regulated by RPD1 (Erhard *et al.* 2013). As the maize genome is composed of >85% TE-like sequences (Schnable *et al.* 2009), many of which occur within 5 kb of genes (Baucom *et al.* 2009; Gent *et al.* 2013), a large number of alleles using TE-like sequences as regulatory elements is possible. Phylogenomic comparisons between *A. thaliana* and *Arabidopsis lyrata* also support the idea that gene-proximate TEs represent a source of regulatory diversity (Hollister *et al.* 2011).

Maize RPD1 was initially identified as a genetic factor required to maintain transcriptional repression of specific alleles subject to paramutation (Hollick *et al.* 2005)—a process by which meiotically heritable changes in gene regulation are influenced by *trans*-homolog interactions (Brink 1956, 1958; Hollick 2012). Presumably because detailed pedigree analyses are required to recognize instances of paramutation, only a few clear examples involving endogenous alleles have been described (Brink 1956; Coe 1961; Hagemann and Berg 1978; Hollick *et al.* 1995; Sidorenko and Peterson 2001; Pilu *et al.* 2009). Similar behaviors involving transgenes have been noted in both plants and animals (Chandler and Stam 2004; Rassoulzadegan *et al.* 2006; Khaitová *et al.* 2011; Ashe *et al.* 2012; de Vanssay *et al.* 2012; Shirayama *et al.* 2012) although it remains unknown whether these examples are due to mechanistically related processes. One strategy for identifying a broader set of alleles subject to paramutation would be to

start with a list of candidate genes, the transcriptional regulation of which is affected by RPD1 function.

The evolutionary function(s) of Pol IV remains enigmatic. Because Pol IV is required for siRNA-directed cytosine methylation (reviewed by Matzke and Mosher 2014 and Matzke *et al.* 2015), it is expected that the regulation of many alleles might be affected by RPD1/*NRPD1* action although few such alleles have been identified in maize (Hollick *et al.* 2005; Parkinson *et al.* 2007; Erhard *et al.* 2013) and *Arabidopsis* (Matzke *et al.* 2007; Ariel *et al.* 2014). To date there have been no reports of genome-wide effects of RPD1/*NRPD1* on gene regulation in any species although several studies have noted correlations between siRNA profiles and nearby genic mRNA abundance (Hollister *et al.* 2011; Eichten *et al.* 2012; Greaves *et al.* 2012). Template competitions between Pol IV and Pol II have been proposed (Hale *et al.* 2009) to account for RPD1-based transcriptional repression seen at the *Pl1-Rhoades* allele of *pl1* (Hollick *et al.* 2005) and for increases in polyadenylated transcripts of some long terminal repeat (LTR) retrotransposons that specifically accompany loss of RPD1 but not loss of two other siRNA biogenesis factors (Hale *et al.* 2009). These results indicate that RPD1-containing Pol IV complexes directly interfere with Pol II transcription of RPD1-targeted genomic regions.

Here we use global run-on sequencing (GRO-seq) (Core *et al.* 2008) to identify genome-wide targets of Pol IV-based transcriptional regulation. This technique profiles RNAs from RNAPs incorporating a brominated UTP ribonucleotide during a short nuclear run-on reaction (Core *et al.* 2008). Maize Pools IV and V can extend transcripts *in vitro*, but ribonucleotide incorporation is attenuated compared to Pol II (Haag *et al.* 2014). Because *Arabidopsis* Pol IV products are rapidly processed to siRNAs (Li *et al.* 2015), transcription rates of maize Pol IV are relatively slow *in vitro* (Haag *et al.* 2014), and no appreciable maize Pol IV RNAs are detected *in vivo* in short run-on experiments (Erhard *et al.* 2009), most non-ribosomal RNA (rRNA), non-transfer RNA (tRNA) transcription detected by GRO-seq is expected to represent Pol II function. Our results show that loss of Pol IV affects transcription profiles at the 5' and 3' gene ends and at a discrete set of unique TEs and genes, the dysregulation of which may contribute to *rpdl* mutant developmental defects.

Materials and Methods

Genetic stocks

The *rpdl-1* null mutation (originally designated *rmr6-1*) (Erhard *et al.* 2009) was introgressed into the B73 inbred background to ~97% by repeated backcrosses of F₂ *rpdl-1* / *rpdl-1* mutant pollen to a recurrent B73 female parent. Families segregating 1:2:1 for *rpdl-1*/*rpdl-1* mutants, heterozygotes, and homozygous *Rpd1*-B73 individuals were used for nuclei isolations and RNA isolations for reverse transcriptase polymerase chain reaction (RT-PCR) and quantitative real-time PCR (qRT-PCR) analysis.

GRO-seq library preparation

Ten homozygous wild-type (WT; *Rpd1-B73/Rpd1-B73*) and 10 homozygous *rpdl-1* mutant (*rpdl* mutant) siblings were identified with a dCAPs marker for the *rpdl-1* lesion (Erhard *et al.* 2013) and used for nuclei isolations. Nuclei were isolated from whole shoots (roots removed) of 8-day-old seedlings. Seedling tissues and dry ice were pulverized in a blade coffee grinder and transferred to a ceramic mortar with 15 ml of ice-cold isolation buffer (40% glycerol, 250 mM sucrose, 20 mM Tris, pH 7.8, 5 mM MgCl₂, 5 mM KCl, 0.25% Triton X-100, 5 mM β-mercaptoethanol). Pulverized tissue in isolation buffer was ground further with a ceramic pestle and filtered through cheesecloth into a 50-ml conical tube. Grindates were filtered again through 40-μm nylon cell strainers (BD Biosciences, San Jose, CA) into 35-ml centrifuge tubes. Nuclei were centrifuged at 6000 × g for 15 min at 4°, and pellets were washed with 15 ml isolation buffer. Washes were repeated two more times, and pellets were resuspended in 100 μl resuspension buffer (50 mM Tris, pH 8.5, 5 mM MgCl₂, 20% glycerol, 5 mM β-mercaptoethanol). Transcription run-ons were performed as described (Hollick and Gordon 1993) with the following changes: 0.5 mM 5-bromouridine 5'-triphosphate (Sigma) was substituted for UTP and 2 μM cold cytidine triphosphate (CTP) was added in addition to 10 μl of α-³²P-CTP (3000 Ci/mmol, 10 mCi/mL; Perkin-Elmer). RNA isolation was as described (Hollick and Gordon 1993) with the following changes: DNase I and Proteinase K digestions were performed for 1 hr each at 37° and 42°, respectively; one acid phenol/chloroform extraction was performed to isolate RNA. High-throughput sequencing libraries were prepared from *in vitro*-labeled RNA as described (Core *et al.* 2008) using agarose bead-conjugated α-bromodeoxyuridine antibody (Santa Cruz Biotechnology).

GRO-seq library processing

Fifty nucleotide (nt) single-end raw reads (54,318,135 for WT library and 54,873,783 for *rpdl* mutant library) were generated on the Illumina HiSeqII platform (Vincent J. Coates Genome Sequencing Laboratory, University of California at Berkeley). Based on FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) Phred score analysis, 10 nt were trimmed from the relatively lower quality 3' end of all reads from both libraries using the FASTQ/A Trimmer script (http://hannonlab.cshl.edu/fastx_toolkit/). The Cutadapt program (Martin 2011) was used to trim adapter sequences and any additional low-quality bases (option -q 10) from the 3' end of all reads; reads shorter than 20 nt after adapter trimming (2,253,657 and 2,488,839, respectively) were excluded (option -m 20). The Fastx Quality Filter (http://hannonlab.cshl.edu/fastx_toolkit/) removed reads with <97% of their bases (option -p 97) above the Phred quality of 10 (option -q 10); 654,703 WT reads and 667,412 *rpdl* mutant reads were removed. Finally, 7546 and 5443 sequencing artifacts were removed from the WT and *rpdl* mutant libraries, respectively, using the Fastx Artifacts Filter (<http://hannonlab.cshl.edu/>

[fastx_toolkit/](http://hannonlab.cshl.edu/fastx_toolkit/)). The resulting libraries consisted of 51,402,229 and 51,712,089 high-quality WT and *rpdl* mutant reads, respectively, with an average length of 32 nt, and these were used for subsequent mapping, computational, and statistical analyses.

Computational analyses of GRO-seq libraries

Alignments to genomic features: For filtering and downstream comparisons, high-quality GRO-seq reads were mapped using Bowtie alignment software (version 0.12.7; Langmead *et al.* 2009) to annotated maize sequence features (see [Supporting Information, Table S1](#), for file types and origins): rRNAs and tRNAs (kindly provided by Blake Meyers, University of Delaware), maize B73 AGPv2 filtered gene set (FGS), Maize Transposable Element Consortium (MTEC) consensus sequences, and maize pseudochromosomes 1–10 representing the sequenced B73 reference genome (AGP version 2, build 5b). Bowtie indices were built from the above annotated maize feature sequences using the bowtie-build command with default parameter settings. Reads that failed to match to the rRNA/tRNA indices with up to two mismatches (option -v 2) comprise the filtered non-rRNA/non-tRNA alignments (41,169,885 and 42,498,964 for WT and *rpdl* mutant libraries, respectively), which were aligned to the maize pseudo-chromosomes 1–10 allowing two mismatches (option -v 2) and to match only once in the genome (option -m 1). These alignments define uniquely mapping reads (12,239,069 and 11,739,444 for WT and *rpdl* mutant libraries, respectively), which were used for the meta-gene and differential expression analyses described below.

Distribution of reads over genomic features: To measure the relative contribution of introns and exons in WT and *rpdl* mutant GRO-seq libraries, we first isolated the subset of 26,987 (68%) maize FGS models with no predicted alternatively spliced transcript isoforms, which we define as single-transcript genes. As a control, we determined the contribution of introns and exons to all single-transcript genes using intron and exon chromosomal coordinates contained in the FGS position annotation file ([Table S1](#)). We compared the control distribution to percentages of uniquely mapping GRO-seq reads overlapping introns or exons within single transcript genes, which were identified using the intersectBed tool, part of the BEDTools suite (Quinlan and Hall 2010), with the following parameters: -f 1 -u -wa. Only reads contained entirely within a gene, exon, or intron were reported (no untranslated regions or exon/intron boundaries are reported by this method). Percentages of introns and exons were calculated as a proportion of unique reads contained within single-transcript genes.

For direct Bowtie alignments of filtered non-rRNA/non-tRNA reads to TE and FGS sequence indices, we allowed two mismatches (option -v 2) and reads to map more than once to the respective set of sequences, but counted each multi-mapping read only once (default option -k 1 to report only one random alignment per read for TEs and option --best to report the best FGS alignment). Analyses of differentially

transcribed TE superfamilies compared numbers of reads aligned directly to MTEC consensus TE sequences (with the same alignment parameters) normalized by total mappable reads of each respective library. For a control alignment to TE sequences, we generated a random sampling of genomic sequences using the Sherman program (<http://www.bioinformatics.babraham.ac.uk/projects/sherman/>). Specifically, a set of 51,402,229 (the number of high-quality reads in the WT GRO-seq library; option -n 51,402,229) random 32-mers (the average length of high-quality GRO-seq reads for both the WT and *rpd1* mutant libraries, weighted for abundances: option -l 32) was generated from the maize genome, inhibiting *in silico* bisulfite conversion with option -cr 0. The resulting 32-mers were aligned directly to TE and FGS sequences using identical Bowtie parameters as GRO-seq read alignments.

To determine overlaps of uniquely and repetitively mapping reads to genomic features (genes, TEs, and intergenic regions), we created alignment tables from the raw Bowtie alignment files (SAM formatted). For each read, alignment characteristics (unmappable, maps uniquely, or maps repetitively) were extracted from the unique alignment of non-rRNA/non-tRNA reads to the B73 genome. These were compared to mapping sense/mapping antisense/not mapping values for the same reads to indices built from the FGS sequences, MTEC TE sequences, or custom sequences extending 5 kb upstream (FGS -5 kb) or downstream (FGS +5 kb) of the original FGS sequences. In all cases, two mismatches were allowed for each reference; for the FGS-only alignment, the best alignment was reported (option -best); for the rest, a random alignment was reported (default option -k 1). The resulting dataset was used to collapse reads into different groups; for example, uniquely mapping TE reads within 5 kb of gene starts would map only once to the B73 genome and map to the MTEC and FGS -5 kb indices, but not to the FGS or FGS +5 kb indices. Only the original B73 genome alignment determined uniquely vs. repetitively mapping. Therefore, when a uniquely mapping read maps within 5 kb of an annotated transcription start site (TSS) and within 5 kb of a 3' gene end, the read is likely between two genes that are <10 kb apart.

Metagene and heatmap profiles of gene boundaries: Uniquely mapping reads overlapping TSSs and 3' gene ends were tallied and binned using a metagene analysis pipeline of custom Python scripts (https://github.com/HollickLab/metagene_analysis). Using *metagene_count.py*, the 5' read ends (option --count_method start) were tallied against the positions of TSSs (option --feature_count start) and 3' ends (option --feature_count end) of FGS models (Table S1, FGS positions) >1 kb in length (31,794 of 39,656 total models). The tallies extended ± 5 or ± 1 kb by changing the padding option (--padding) to 5000 or 1000, respectively; in both cases counting was strand-specific relative to the feature orientation by default. For the 5-kb tallies, only the first 1 kb of genic windows at each gene were kept by excluding windows with starting positions > +1000 from the TSS or < -1000 from the 3' end in R. Tallies from *metagene_count.py* were

strand-specifically binned (option --separate_groups) with *metagene_bin.py* in either 10-nt nonoverlapping windows (--window_size 10 --step_size 10) for detailed metagene plots or 50-nt nonoverlapping windows (--window_size 50 --step_size 50) for the heatmap plots. The resulting count tables were imported to R (version 2.15.2; <http://www.r-project.org/>) for normalization, statistical testing, and plotting.

To view the normalized coverage (reads per million uniquely mapped), a heatmap-like plot was made using *image*, a base R command, to plot coverage (z-axis) on a color scale at each position along the gene model (x-axis) for groups of 60 genes (y-axis). Gene models were ordered by their maximum contribution to a 10-nt window's total abundance (Maximum Sum Contribution) for all heatmap and metagene plots. At each window, a gene's sum contribution represents its influence on the total coverage via the calculation: gene's coverage at the window/total coverage of all genes at the window. The Maximum Sum Contribution was also used to exclude the upper and lower 5% of genes from the metagene plots described below. Heatmap plots were binned into 50-nt nonoverlapping windows and neighboring gene models after sorting by Maximum Sum Contribution were averaged in groups of 60 genes.

To summarize the average behavior of GRO-seq coverage over the FGS models, we created metagene plots summarizing the coverage of each gene using either the total (sum) or the mean coverage at each 10-nt window. Welch's two-sample *t*-tests and 95% confidence intervals were calculated for each 10-nt window across all inner 90% quantile (by Maximum Sum Contribution) gene models. To identify the windows with statistically significant coverage differences $\pm$ RPD1, the individual Welch's *t*-test results were corrected for multiple sampling using the Holm-Bonferroni method ($\alpha = 0.05$) across all 800 windows comprising the regions around the TSSs and 3' ends on both sense and antisense strands. Final plots used base R commands to plot mean or sum abundance as lines (or bars) and 95% confidence intervals as polygons and to highlight statistically significant windows with horizontal line segments.

Correlations between WT and *rpd1* mutant libraries: To determine the correspondence between WT and *rpd1* mutant libraries, the number of uniquely mapping reads per kilobase per million uniquely mapped reads were tallied across various regions. Near-genic analysis of the 31,794 genes analyzed by the metagene profiles divided the region around each gene model into constant 1 kb upstream of TSS, 1 kb downstream of TSS, 1 kb upstream of gene end, and 1 kb downstream of gene end regions. The internal portion (>1 kb away from each gene boundary) of those genes >2 kb in length (23,050 of 31,794) was used to represent the interior gene region. All counts used the 5'-most base of each GRO-seq read. For each region, zero values were artificially set to 1/10 of the lowest nonzero value in either data set; this allowed both the inclusion of zero vs. non-zero data and had minimal (linear regression) to no (Spearman's rank correlation) effect on the summary statistics. Resulting normalized coverages were

log₁₀-transformed for plotting, fitting a linear regression “Data fit” line, and calculating the Spearman’s rank correlation coefficient, all of which were performed in R (version 3.0.2).

siRNA analyses: National Center for Biotechnology Information (NCBI)-sourced profiles of maize 16- to 35-nt RNAs were pooled and overlapped with the gene models used in the GRO-seq metagene analysis. Because no profiles represented 8-day-old B73 seedling shoots, we pooled WT B73 siRNAs from seedling (day 3) root tips (SRR218319: Gent *et al.* 2012), seedling (day 11) shoot apices (SRR488770 and SRR488774: Barber *et al.* 2012), ovule-enriched unfertilized cob (at silk emergence) (SRR408793: Gent *et al.* 2013), and developing ear (SRR1583943 and SRR1583944: Gent *et al.* 2014). Pooled siRNAs from B73 *rdr2* mutant developing ears (SRR1583941 and SRR1583942: Gent *et al.* 2014) represent 24-nt RNA deficiency that should mimic Pol IV loss as RDR2 is required with Pol IV for 24-nt RNA biogenesis (Matzke and Mosher 2014; Matzke *et al.* 2015).

All raw sequences were downloaded from the NCBI Sequence Read Archive and processed to a high-quality set that all had trimmable 3’ adapters and neither low-quality (Phred scores <30) nor ambiguous bases. The resulting high-quality reads (11,163,623 for *rdr2* mutant and 74,787,717 for WT) were filtered against rRNA/tRNA sequences and aligned to the maize B73 v2 genome with 551,194 (*rdr2* mutant) and 11,466,496 (WT) reads mapping uniquely. Uniquely mapping 24-nt reads (154,509 or 28% for *rdr2* mutant and 8,611,691 or 75% for WT) were subjected to metagene analysis as described above, using all size classes for library normalization in reads per million uniquely mapped per region length. For consistency, the order of genes in the heatmaps followed the same order (by Maximum Sum Contribution to total GRO-seq coverage) as the GRO-seq heatmaps.

Definition of alternative transcription start sites: As an alternative TSS definition, we used the 5’-end sequence of full-length complementary DNAs (fcdDNAs) from predominantly 7-day-old seedling tissues (ZM_BFc set: http://www.ncbi.nlm.nih.gov/nucest/?term=ZM_BFc; see Soderlund *et al.* 2009). Positions for the fcdDNA 5’ ends were defined by perfectly (0 mismatches) and uniquely (only 1 B73 alignment) aligning the first 50 nt of the cDNA sequence to the maize B73 reference genome (version 2, build 5b) with Bowtie (version 0.12.7). Identical TSS positions were collapsed. Metagene counting, binning, and plotting of total normalized GRO-seq read abundance were performed ± 1 kb from all of the fcdDNA-defined TSSs in 10-nt nonoverlapping windows.

Identification of differentially transcribed genes and TEs: For differential transcription analysis of genes (gene body only) and TEs with defined genomic coordinates (positionally defined TEs), the DESeq package (Anders and Huber 2010) was used with uniquely aligning reads and the maize FGS GFF3 and MTEC TE GFF3 (Table S1) annotation files.

The counts table for DESeq used tallies of raw reads overlapping features in sense and antisense orientations (-s and -S options, respectively) generated by intersectBed (additional options -c -wa; Quinlan and Hall 2010). The sense and antisense counts tables were processed with the following DESeq parameters: fit = “local”; method = “blind”; and sharingMode = “fit-only”. The Benjamini–Hochberg correction (Benjamini and Hochberg 1995) for multiple sampling adjusted the *P*-values adjusted (padj) for False Discovery Rate (FDR) control, and a 10% FDR threshold was applied to the padj values (Anders and Huber 2010). The list of features passing the DESeq thresholds were further curated and trimmed as described below. Sequence polymorphisms between the introgressed haplotype containing the *rpm1-1* mutation and the homologous B73 chromosome 1 region likely contribute to differences in the abundances of reads aligning to this region between the WT and *rpm1* mutant libraries. While the size of the *rpm1-1*-containing haplotype is unknown, we estimated it to be at most 20 Mb. Therefore, we excluded 12 TEs and 26 genes located within 20 Mb on either side of the *rpm1* locus from the respective lists of features classified as having decreased transcription in *rpm1* mutants. We manually curated genes with increased or decreased sense transcription based on visual inspection for GRO-seq read coverage consistent with the transcription unit defined by the gene annotation; coverage localized to only a portion of the gene model were tagged as unlikely to be related to that gene’s expression (nonbold entries in Table S3). Genes with increased antisense transcription were visually inspected, and TEs <2 kb beyond the genic 3’ end were tallied. We determined whether positionally defined, differentially transcribed TEs were located within annotated genic regions or <5 kb from their 5’ or 3’ ends using the closestBed tool (Quinlan and Hall 2010) with the -d parameter, which in this case reports the distance of the closest gene to each TE analyzed.

Expression analysis

RT-PCR expression analysis: Three homozygous *Rpm1-B73* (WT) and three homozygous *rpm1-1* mutant 8-day-old seedlings from accessions described in *Genetic stocks*, were identified by genotyping as described above and used for RNA isolations. Seedling tissues (whole shoots, as described above) were pulverized separately in ceramic mortars in 1 ml of Trizol reagent (Invitrogen), and RNAs were isolated following the manufacturer’s protocol. cDNA synthesis using oligo(dT) primers (New England Biolabs) and Superscript II (Invitrogen) followed manufacturer’s protocols using 1 μ g RNA as templates for reverse transcription reactions. cDNAs were amplified using gene-specific primers (Table S2) designed from sequences corresponding to the predicted coding regions of *ocl2* (AC235534.1_FG007), GRMZM2G043242 (ATPase-domain containing protein), and GRMZM2G161658 (*Epoxide hydrolase2*), as well as primers matching a control gene *alanine aminotransferase* (*aat*) as described (Woodhouse *et al.* 2006). RT-PCR products were sized on a 2% agarose gel and stained

with ethidium bromide for visualization and quantified with ImageJ software (<http://rsbweb.nih.gov/ij/>) to normalize to *aat* products. For each gene-specific primer pair, PCR amplicons were gel-extracted (Qiaquick Gel Extraction kit) and Sanger-sequenced (University of California at Berkeley Sequencing Facility) to verify that correctly spliced products were being amplified (sequences available upon request).

qRT-PCR expression analysis: Seedling tissues (whole shoots, as described above) were harvested from 8-day-old sibling homozygous *Rpd1-B73* (WT) and *rpdl-1* mutant plants, flash-frozen in liquid nitrogen, and stored at -80° for subsequent RNA extraction. A total of nine seedlings for each genotype were prepared in pools of three seedlings each. These pools of three were pulverized with dry ice in a coffee grinder and subsequently ground in a mortar and pestle with 5 ml of ice-cold isolation buffer (40% glycerol, 250 mM sucrose, 20 mM Tris, pH 7.8, 5 mM $MgCl_2$, 5 mM KCl, 0.25% Triton X-100, 5 mM β -mercaptoethanol). RNAs were then extracted from 1 ml of the grindate with 1 ml of TRIzol reagent (Invitrogen) following the manufacturer's protocol. An aliquot (2 μ g) of total RNA was DNaseI (Roche) treated for 20 min at 37° and heat-inactivated (75° for 10 min) in 5 mM EDTA. The Tetro cDNA synthesis kit (Bioline, Taunton, MA) was used for the first-strand cDNA synthesis, followed by RNA degradation with an RNase A/TI/H cocktail. qRT-PCR reactions each used cDNA from 20 ng of starting total RNA in a $1\times$ SYBR Sensimix (Bioline) reaction with 0.25 μ M of each primer (Table S2). Each biological sample had three technical replicates for both the *aat* and *ocl2* primer sets. The qRT-PCR conditions used 40 cycles with 30 sec at 60° annealing and 45-sec extension steps, the melt curve ramped from 60° to 95° in 20 min. The technical (three per template) and biological (three pools of three seedlings each) replicates were combined to calculate the average *ocl2* abundance relative to *aat* as $2^{(aat\ Ct - ocl2\ Ct)}$.

Results

Global run-on sequencing profiles nascent transcription in maize seedlings

Additional Pol II-derived RNAPs in multicellular plants (Luo and Hall 2007) may affect transcription dynamics across the genome, particularly as all of these RNAPs share accessory subunits with Pol II (Ream *et al.* 2009; Tucker *et al.* 2011; Haag *et al.* 2014). Pol II transcription of both LTR retrotransposons (Hale *et al.* 2009) and certain *pl1* alleles (Erhard *et al.* 2013) is increased in *rpdl* mutants. However, dysregulation of these specific loci cannot explain all the developmental phenotypes observed in *rpdl* mutants (Parkinson *et al.* 2007; Erhard *et al.* 2009). We used GRO-seq to view nascent transcription profiles $\pm$ RPD1 to identify both particular (haplotype specific) and general (locus independent) effects of Pol IV loss on maize genome transcription.

GRO-seq libraries were prepared using sibling *rpdl* mutant and nonmutant (WT) seedlings with each library representing

nuclei from 10 separate individuals (see *Materials and Methods*). Sequencing reads from these libraries were mapped to the B73 reference genome (Schnable *et al.* 2009) (Figure 1A). Transcripts from all five classes of maize RNAPs (Pols I–V) could be represented in the WT GRO-seq library, whereas only those requiring Pol IV function (including Pol IV transcripts) would be absent in the *rpdl* mutant library. To focus predominantly on Pol II, IV, and V transcription, we removed most Pol I and III products by excluding rRNA- and tRNA-aligning reads from subsequent analysis (see *Materials and Methods*). Genomic non-rRNA/non-tRNA reads separated into repetitively aligning and uniquely aligning groups show similar distributions $\pm$ RPD1 (Figure 1A). To evaluate whether the libraries are enriched for nascent transcripts as opposed to spliced mRNAs, we compared the exonic/intronic distributions of genic reads (Figure S1). Both repetitively and uniquely mapping reads include intronic sequences (~ 90 and $\sim 40\%$, respectively), confirming the enriched representation of nascent, unspliced transcripts.

Because genes and TEs are predicted to be differentially affected by Pol IV loss, we categorized each read as having possibly originated from a TE, a gene (including annotated UTRs and introns), both, or neither (intergenic). Most uniquely mapping reads originate from genic or intergenic loci having little overlap with TEs (Figure 1B, Unique). Repetitively mapping reads align to TE-like sequences (Figure 1B, Repetitive), although these TE-like reads usually align to genes as well (Figure 1B, purple boxes). Approximately 96% of repetitive reads aligning to TE sequences could originate from genic transcripts. To ensure that this enrichment is not biased by our categorical analysis, we repeated the analysis on a set of *in silico* sequences randomly generated from the maize genome (see *Materials and Methods*; Figure S2) and found 97.3% of the TE-aligning, *in silico*-generated 32mers also aligned to genes. These results indicate that the majority of TE sequences represented in nascent transcription profiles are likely found within introns and untranslated regions of gene-derived Pol II RNAs.

Exclusively genic reads in both libraries (61 and 59.7% of all mappable WT and *rpdl* mutant reads, respectively) are highly enriched compared to the prevalence of genic sequences in the maize genome (8%) (Figure S3), indicating that these GRO-seq profiles represent largely genic transcription. Because GRO-seq reads provide strand-specific information, we could also identify a sense-oriented strand bias among all genic reads, particularly the uniquely mapping reads that originate from the mapped locus (Figure 1C). Even reads representing TEs embedded in host genes appear biased toward the sense strand: 95% of uniquely aligning TE-like reads (5937 and 5641 reads in WT and *rpdl* mutant libraries) are sense-oriented relative to their host genes. In all comparisons, the read distributions remain similar between WT and *rpdl* mutant libraries, consistent with prior run-on transcription results showing that Pol IV contributes $<5\%$ of the transcribed nuclear RNA pool (Erhard *et al.* 2009) and the recent finding that Pol IV transcripts are rapidly processed to siRNAs (Li *et al.* 2015). These results indicate that loss of RPD1 does

Figure 1 GRO-seq reads are similarly distributed in WT and *rpd1* mutant libraries. (A) Percentages of WT and *rpd1* mutant GRO-seq reads that are unmappable, map to rRNA/tRNA sequences, map repetitively (>1 alignment), or map uniquely to the B73 reference genome. (B) Distribution of repetitively and uniquely mapped reads [reads per million mapped (RPM)] from A to annotated genes (blue), TEs (red), both (purple), or neither (intergenic; yellow). (C) Strandedness of the best alignment to gene models of all potentially genic reads (those that align with genes only or with both genes and TEs; blue and purple regions from B, respectively).

not generally shift transcription away from genes and toward TEs, suggesting that Pol IV has a more precise mechanism for regulating transcription of its targets.

Maize seedling transcription is focused on genic regions

Mappable GRO-seq reads not aligning to either gene or TE features represented the “intergenic” category. These reads could have several origins including unannotated genes and TEs or noncoding DNA sources for siRNAs. However, we find that most nongenic non-TE reads represent pretermination transcription downstream of currently annotated polyadenylation addition sites (PAS) (Figure 2). Nascent transcription profiles in *Homo sapiens* (Core *et al.* 2008), *Drosophila melanogaster* (Core *et al.* 2012), and *Caenorhabditis elegans* (Kruesi *et al.* 2013) also identify transcription beyond the PAS and, in some cases, antisense transcription upstream of the TSS. To test whether nascent transcription is enriched near maize genes, we queried nongenic read alignments at regions 5 kb upstream of annotated TSSs and 5 kb downstream of annotated 3' ends. Both intergenic and TE-only reads consist largely (>80%) of sequences that can align to within 5 kb of a gene model, representing potential genic reads (Figure 2A). The one exception is uniquely mapping TE-only reads, where only half of the reads align within 5 kb of genes. These intergenic and

TE-only reads >5 kb from the nearest gene comprise ~5% of the maize seedling non-rRNA/non-tRNA transcriptome (see *Materials and Methods*). At our current depth of sequence coverage we cannot confidently determine if transcription of those loci far from genes is influenced by RPD1 loss. However, our datasets are enriched over genes (~65% of uniquely mapping reads in both WT and *rpd1* mutant datasets) with an average sense-strand coverage of 173 (*rpd1* mutant) to 187 (WT) raw reads per gene; we therefore continued our focus predominantly on near-genic regions, which are enriched in our dataset.

In support of the idea that near-genic reads in the maize GRO-seq profiles represent Pol II pretermination extensions of genic transcription units, uniquely mapping near-genic reads align predominantly in the downstream 5 kb (up to 84 and 87% of WT and *rpd1* mutant near-genic reads, respectively; Figure 2B). The overlap of uniquely mapping reads between the upstream 5 kb and downstream 5 kb (22% in both genotypes) likely represents reads aligning between genes separated by <10 kb. Repetitively mapping near-genic reads have a more even distribution between upstream and downstream 5 kb, which could represent an artifact of alignments to nonorigin loci. With uniquely mapping reads, where alignments likely correspond to the originating locus, near-genic reads are predominantly sense-stranded (Figure 2C) in accord with the hypothesis that they represent continuation of genic transcription.

We next used pile-ups of uniquely mapped WT GRO-seq reads across genic loci to profile the typical maize genic transcription unit at higher resolution. Combining all sense and antisense profiles (Figure S4) into a metagene composite (Figure 2D) confirms the sense-strand enrichment downstream of currently annotated gene models. Additionally, the composite profile indicates that most pretermination transcription occurring 3' of PASs concludes within 1–1.5 kb of currently annotated gene ends. This result agrees with estimates from metazoan profiles (Core *et al.* 2008, 2012; Kruesi *et al.* 2013). Beyond presumptive genic transcription termination points and upstream of TSSs there is remarkably little evidence of transcription. Although our results do not distinguish RNAs produced from different RNAPs, the nature of the read enrichment at genes (starting near the TSS and extending beyond the 3' PAS) and the sense-strand bias supports the prediction that most of these reads derive from gene-associated nascent Pol II RNAs.

Promoter-proximal transcription in maize is distinct from that seen in metazoans

The composite metagene profile (Figure 2D) highlighted unexpected features of typical maize transcription initiation. The beginning of composite genic transcription is marked with a prominent narrow peak of GRO-seq reads nearly coincident with the TSS (Figure 2D). Sense-oriented peaks located ~50 bp downstream of TSSs are identified by GRO-seq profiles in humans (Core *et al.* 2008) and *Drosophila* (Core *et al.* 2012) and under stress conditions in *C. elegans* (Kruesi *et al.* 2013; Maxwell *et al.* 2014). These peaks are cited as

Figure 2 Nongenic GRO-seq reads are enriched near genes. (A) Percentage of nongenic/non-TE (intergenic) or TE-only reads that align near genes (within 5 kb; dark gray) or >5 kb (light gray) from genes. (B) Nongenic reads within 5 kb of genes found exclusively upstream (Up, left circle), downstream (Down, right circle), or at both ends (intersection) of a nearby gene. (C) Distribution of uniquely mapping near-genic reads [reads per million uniquely mapped (RPMUM)] by strand orientation relative to the nearby gene model. (D) Metagene profile of uniquely mapping WT GRO-seq reads summed over 10-nt windows ± 5 kb from FGS models.

evidence of promoter-proximal Pol II pausing, a transcriptional regulatory mechanism first described at the *hsp70* gene in *Drosophila* (Rougvie and Lis 1988). It is unclear whether the more upstream maize TSS-proximal peak (Figure 2D) represents Pol II pausing. Over 1/4 of maize gene models used in the original metagene profile (Figure 2D) begin with the triplet ATG sequence (27%), compared to only 2% when triplet sequences are sampled 5 kb upstream of the gene models (an approximation of ATG enrichment genome-wide) (Figure S5). This finding indicates that translation, rather than transcription, initiation sites define many current annotations of maize gene start positions. To test if an alternative gene start definition would shift the TSS-proximal peak, we first defined a set of genomic TSS annotations using maize seedling and young leaf full-length cDNA sequences (see *Materials and Methods*; Soderlund *et al.* 2009). A similar metagene analysis using this validated set of TSSs places the sense-oriented GRO-seq peak upstream of the TSS (Figure S6). This result indicates that the peak positions relative to TSSs are dependent on annotation methods but likely do not represent canonical Pol II pausing as described in metazoans.

Nascent transcription profiling (Core *et al.* 2008) and short RNA cDNA libraries (Seila *et al.* 2008) identified evidence of divergent antisense transcription peaking at ~ -250 bp at mammalian promoters. Similar to *Drosophila* GRO-seq profiles (Core *et al.* 2012), our metagene profile has no evidence of such a broad antisense peak (Figure 2D). The only peak at ~ -250 bp is due to antisense-biased GRO-seq coverage of

a single gene (Figure S7), which likely represents a transcription unit unrelated to the annotated gene model. Together, these two characteristics of near-promoter transcription—a TSS-proximal peak and lack of divergent transcription—distinguish the maize seedling transcriptional environment at promoters from that of currently profiled metazoans.

Pol IV affects nascent transcription at gene boundaries

Additional Pol II-derived plant RNAPs (Pols IV and V) represent a key distinction between the transcriptional landscape of multicellular plants and other eukaryotes. To evaluate Pol IV effects on transcriptional activity at functionally important sites surrounding genes, such as promoters and transcription termination sites, we generated and compared metagene profiles displaying the mean GRO-seq read abundance across a composite of annotated gene edges and their flanking genomic regions (Figure 3A). To exclude outliers, we sorted gene models based on their maximum read count contribution to the metagene plot and included only the inner 90% (~ 28.6 thousand) of gene models in the metagene profiles (see *Materials and Methods*).

Profile comparisons reveal changes in transcription at gene boundaries $\pm$ RPD1, while transcription of genic and upstream regions remain unaffected (Figure 3A and Figure S6). Near TSSs, the *rpdl* mutant library has significantly lower read coverage in both strand orientations (Figure 3A, Welch's *t*-tests in 10-nt windows, corrected for multiple sampling by the Holm–Bonferroni method at $\alpha = 0.05$). Heatmap

Figure 3 Pol IV loss alters global transcription profiles at gene boundaries. (A) WT and *rpd1* mutant mean GRO-seq read coverage (black and purple lines, respectively) of 90% of the maize genes within 1 kb of gene start (TSS) or 3' end (End). Gray and purple shading represent 95% confidence intervals; red horizontal bars highlight 10-nt nonoverlapping windows that significantly differ between libraries (Welch's *t*-test by window, corrected to $\alpha = 0.05$ with the Holm–Bonferroni method for multiple sampling). (B) Variation in coverage between WT and *rpd1* mutant libraries for all FGS genes. Fold change was calculated from the average coverage (reads per million uniquely mapped) of 60 neighboring genes when sorted by their maximum sum contribution. Fifty-nucleotide windows with zero coverage in either library are plotted in white. The gold bar highlights the inner 90% of genes used in A.

representations of read abundance differences across individual gene boundaries indicate that the trends observed in the metagene summary apply to most genes (Figure 3B). Elsewhere, the WT and *rpd1* mutant GRO-seq profiles are remarkably similar, particularly for regions having high coverage (sense strand over gene bodies and 1 kb downstream of gene ends) (Figure S8). The sense-strand gene body coverage between libraries has Spearman's rank correlation coefficients (ρ) of 0.971, 0.963, and 0.977 (first 1 kb, middle, and last 1 kb, respectively), which approximate the $\rho = 0.967$ observed

between biological replicates in the original GRO-seq analysis (Core *et al.* 2008). Together, the metagene summary, fold-change heatmap, and Spearman's correlations highlight that RPD1 has no general impact over the coding region of genes. More striking, the pretermination region beyond the PAS has significantly increased read coverage in the absence of RPD1. Downstream of the PAS there is evidently increased transcription for most genes relative to upstream of the PAS, and this 3' transcription is even more pronounced in *rpd1* mutants (Figure 3B). Upstream of currently annotated gene models,

the fold differences in read abundances are variable as expected for regions of relatively slow or underrepresented transcription; fold changes in read abundances representing antisense-oriented transcription (Figure S9) show similarly variable patterns. Although window sizes used for generating fold-change heatmap data are larger (50 vs. 10 nt) than in the metagene profiles, most of these larger windows directly above TSSs still show negative fold changes, indicating increased transcription in WT vs. *rpdl* mutant samples is a common feature of many genes (Figure 3B).

The RPD1-dependent changes in the GRO-seq profiles observed near gene boundaries implicate Pol IV activity near genes. Because Pol IV is required to generate 24-nt RNAs, presumably through downstream processing of Pol IV transcripts by RNA interference-like machinery (Li *et al.* 2015; Matzke and Mosher 2014; Matzke *et al.* 2015), we looked for 24-nt read (24mer) evidence supporting Pol IV action nearby recognized Pol II transcription units. Previous genome-wide profiling of maize 24mers identified enrichment 1.5 kb upstream of genes (Gent *et al.* 2014). To profile 24mers within 1 kb of gene boundaries, we pooled ~75 million B73 16- to 35-nt RNA reads from published datasets for metagene analysis. To increase effective sequencing depth, we pooled four distinct datasets: 3-day-old seedling root (Gent *et al.* 2012), unfertilized cobs (Gent *et al.* 2013, 2014), and 11-day-old seedling shoot apices (Barber *et al.* 2012), all representing the B73 inbred background, and subjected them to similar coverage analysis near genes (see *Materials and Methods*). Most 24mers represent repetitive features so their originating loci cannot be determined. We therefore limited analysis to uniquely mapping 24mers (8.6 million). These 24mers are enriched both upstream and downstream of genes (Figure S10A and B), and because they are uniquely mapping, they are presumably generated from Pol IV transcription occurring in regions immediately flanking genes. Biogenesis of 24-nt RNAs also requires RDR2 to create a double-stranded RNA from Pol IV transcripts (Li *et al.* 2015; Matzke *et al.* 2015). In *Arabidopsis*, RDR2 functions only in physical association with Pol IV (Haag *et al.* 2012). While the maize RDR2 ortholog (Alleman *et al.* 2006) has not been tested for a similar requirement, it physically associates with Pol IV (Haag *et al.* 2014) and is required for 24-nt RNA biogenesis (Nobuta *et al.* 2008). We therefore used an existing siRNA dataset from an *rdr2* mutant (Gent *et al.* 2014) as a proxy for identifying RPD1-dependent 24-nt RNAs. Although this comparative analysis is more limited (11.1 million total reads, ~551,000 unique 16-35mers, and ~154,000 unique 24mers), the 24mer coverage across all genes showed no enrichment flanking gene boundaries (Figure S10A, C, and D), indicating that the near-genic 24-nt RNAs are RDR2-dependent. These 24mer analyses support the presence of Pol IV immediately upstream and downstream of Pol II genes. Together, this analysis of maize *rpdl* mutants at gene boundaries identifies previously unknown interactions by which Pol IV affects Pol II transcription at discrete positions near genic regions. In addition, these results identify both conserved and novel features of transcription in higher plants vs. metazoans.

Transcription of specific genes is affected by RPD1

RPD1 is required for restriction of *silkless1* gene expression from apical inflorescences, thus ensuring proper male flower development (Parkinson *et al.* 2007), although how RPD1 regulates developmentally important genes is unknown. We employed a computational method (DESeq; Anders and Huber 2010) that assigns statistical significance to regions annotated as genes over- or underrepresented in uniquely mapping GRO-seq reads from either WT or *rpdl* mutant libraries. This approach robustly controls for false positives (type I errors) across a dynamic range of coverage levels and, by pooling information from similarly represented loci, can estimate the required mean and variance values for each locus even when the total sequencing depth and/or number of biological replicates is small (Anders and Huber 2010). Using this method, we could treat the WT and *rpdl* mutant libraries as effective biological replicates (see *Materials and Methods*), assuming that at most loci there was no differential transcription, an assumption supported by the trends observed in the global analysis (Figure 1), the metagene profiling (Figure 3), and the Spearman's rank correlation coefficients (Figure S8). As a result, we have higher confidence in our identified differentially transcribed loci than if we treated each library individually, although the method likely underestimates the total number of differentially transcribed loci.

Applying this stringent statistical method identified a total of 209 annotated genes whose seedling-stage transcription across the entire gene body (annotated UTRs, introns, and exons) is significantly increased or decreased by loss of RPD1. We excluded a cluster of 26 gene models having significantly reduced GRO-seq representation in the *rpdl* mutant profiles found within 20 Mb of the *rpdl-1* introgressed haplotype as these were likely identified because of an inability of some polymorphic *rpdl* mutant reads to align to B73 sequences in this interval. The remaining 183 genes represent potential direct or indirect targets of RPD1/Pol IV regulation (Figure 4A and Table S3). To determine whether or not genic TEs were related to RPD1-affected transcription, we re-annotated the 183 genes and compared their TE content with 200 randomly selected genes (Table S3 and Table S4). This analysis indicates that RPD1-regulated genes have an average TE content: 64% of the differentially transcribed genes vs. 66% of the randomly selected genes.

Visual inspection of the differentially transcribed genes having sense-strand changes (148 in total) using genome browser displays identified two classes: those with consistent GRO-seq read coverage across the entire gene model and those having more biased or localized coverage. In total, 32 of 46 (70%) having increased transcription (Table 1) and 96 of 102 (94%) with decreased transcription (boldface entries in Table S3) matched current gene annotations and are likely related to RPD1-dependent effects on genic transcription. The remaining 20 genes identified as differentially expressed by DESeq analysis have changes in GRO read coverage localized to subgenic regions, sometimes overlapping TEs or the

beginning of the gene with persistent coverage extending from the 3' end of an upstream gene.

Three examples of differentially transcribed genes were subsequently examined and validated at mRNA levels. Results of oligo(dT)-primed RT-PCR analyses using similar biological materials confirmed increased sense-oriented mRNA levels of all three genes tested from the set identified by computational analysis of WT and *rpm1* mutant GRO-seq reads (Figure S11). As an example, the predicted fold changes $\pm$ RPD1 of *outer cell layer 2* (*ocl2*) nascent transcripts and mRNAs are $\sim$ 7.9- and 3-fold increases, respectively. Additional qRT-PCR results estimate a 10-fold increase in *ocl2* mRNAs in the absence of RPD1 (Figure S12). These results validate the GRO-seq technique and DESeq analyses in discovering genes whose Pol II transcription is affected by RPD1.

We expected to detect primarily increased sense-specific transcription by comparing WT and *rpm1* mutant GRO-seq profiles because the *Pl1-Rhoades* allele is transcriptionally repressed by RPD1 (Hollick *et al.* 2005). However, the largest fraction ($\sim$ 0.56) of genes affected by loss of RPD1 has lower sense-oriented transcription in *rpm1* mutants (Figure 4B and Table S3). One possible explanation for this result is that genomic features transcriptionally repressed in an *rpm1* mutant background represent indirect effects. Potentially related to this idea, the presumed maize ortholog of the *Arabidopsis* *REPRESSOR OF SILENCING 1* (*ROS1*) gene, which encodes a DNA glycosylase that facilitates demethylation of cytosine residues (Morales-Ruiz *et al.* 2006), is transcriptionally repressed in *rpm1* mutants (2.9-fold decrease) although this differential representation does not pass our statistical cutoff after adjustment for multiple testing by our DESeq test (raw *P*-value: 0.002, adjusted *P*-value: 0.2). Maize *ros1* RNA levels are also reduced in meristems of *rdr2* mutants (Jia *et al.* 2009), although the mechanism by which any genes are repressed in the absence of RPD1 or Pol IV-dependent siRNAs remains unknown.

Among differentially transcribed gene models in *rpm1* mutants (Table S3), we identified several candidates whose dysregulation could result in developmental abnormalities and potentially contribute to *rpm1* mutant phenotypes (Parkinson *et al.* 2007). The candidate showing the second greatest increased transcription is *ocl2* (Figure 4C), a member of the plant-specific homeodomain leucine zipper IV (HD-ZIP IV) family of transcription factors predicted to have leaf epidermis-related functions in maize (Javelle *et al.* 2011). Mature *rpm1-2* mutant plants often exhibit problems maintaining proper leaf polarity (adaxialized leaf sectors) (Parkinson *et al.* 2007). Interestingly, *ocl2* is not normally expressed in epidermal or mesophyll cells (Javelle *et al.* 2011), which comprise the majority of cells used for GRO-seq library generation, indicating that this gene is transcribed outside its normal expression domain in *rpm1* mutants.

Genome browser visualization of GRO-seq reads uniquely mapping to the genomic region containing *ocl2* (Figure 4C) highlights coincident transcription profiles of a proximate TE and this RPD1-regulated gene. A fragment of an LTR

retrotransposon of the *Gypsy* class (RLG) assigned to the *ubid* family located $\sim$ 1.3 kb upstream of the *ocl2* gene is also transcribed in *rpm1* mutants, but in antisense orientation with respect to the *ocl2*-coding region (Figure 4C). These results identify the *ubid* fragment upstream of *ocl2* as a putative controlling element for this allele, with the absence of RPD1 corresponding with transcriptional increases of both the *ubid* fragment and the adjacent gene.

We also identified 36 genes transcriptionally altered in the antisense orientation in *rpm1* mutants (Figure 4B). Pol IV is implicated in the production of an antisense precursor transcript and corresponding 24-nt siRNAs, homologous to the 3' end of the *Arabidopsis* gene *FLOWERING LOCUS C* (*FLC*) (Swiezewski *et al.* 2007), which encodes an epigenetically regulated MADS Box factor important for vernalization and the regulation of flowering time (Dennis and Peacock 2007). However, only four genes (Table S3) show decreased antisense transcription in the *rpm1* mutant, indicating that Pol IV-dependent antisense transcription of genes is unlikely a primary mechanism of its action on a genome-wide scale. Many more genes (32 total) were recognized having increased transcription in antisense orientation (Figure 4B), yet only one of these (GRMZM2G045560; a gene model encoding a WRKY DNA-binding domain-containing protein) had a significant (DESeq method of Anders and Huber 2010) increase in sense transcription as well. Most of these examples appear to represent transcription of noncoding RNAs initiated 3' of the annotated genes (Figure 4D). Additionally, visual inspections indicate that most (22 of 32, 69%) of these transcription units begin at downstream TEs (Figure 4D). Counting MTEC TE annotations within 2 kb of the 3' ends of these 32 genes identifies a large number of LTR retrotransposons immediately downstream (Figure 4E). These results support previous findings showing that Pol IV loss allows increased transcription of certain LTR retrotransposons (Hale *et al.* 2009), and they highlight how such promiscuous Pol II transcription could affect gene regulation (Kashkush and Khasdan 2007). Genes encoding a histidine kinase receptor for cytokinin, an important phytohormone, and a homolog of an *Arabidopsis* HD-ZIP factor ATHB-4 also have elevated antisense transcription profiles in the absence of RPD1 (Table S3), indicating the potential for a biologically significant role for this novel mechanism of RPD1 gene regulation in maize.

Transcription of TE families and specific TEs is affected by RPD1

While overall TE transcription appears modest (Figure 1B and Figure 2A), we compared GRO-seq read representations among specific TE families and at individual TE loci to determine if certain types are preferentially transcribed. Direct alignments of total WT and *rpm1* mutant GRO-seq reads (unique and non-unique) to consensus sequences of annotated maize TE classes and major superfamilies allowed us to compare transcription of these features to their relative abundance in the maize genome (Figure 5A) (Schnable *et al.* 2009). Although the RLG class of LTR TEs is the most abundant TE superfamily in the maize

Figure 4 Specific alleles are susceptible to Pol IV-induced changes in gene expression. (A) Across FGS gene bodies, the log₂ fold change (*rpd1* mutant/WT) of uniquely mapping read coverage vs. total coverage (average of WT and *rpd1* mutant reads). Triangles represent genes with infinite fold change due to zero coverage from WT (top) or *rpd1* mutant (bottom) uniquely mapping reads. Of the 39,656 FGS gene bodies analyzed, those with zero coverage in both WT and *rpd1* mutant datasets (7783 and 9667 for sense and antisense strand transcription, respectively) were excluded from the plots. Purple dots represent genes with significantly (by the DESeq statistical method of Anders and Huber 2010; see *Materials and Methods*) increased or decreased GRO-seq read representation in *rpd1* mutants. Teal dots represent genes within 20 Mb of the *rpd1* locus whose decreased transcription in *rpd1* mutants may reflect alignment artifacts (see *Materials and Methods*) and are excluded from subsequent analysis. (B) Distribution of categories (by direction of the change and strand) among transcriptionally altered genes in *rpd1* mutants. (C) Genome browser view of WT (black peaks) and *rpd1* mutant (green peaks) GRO-seq reads [normalized to reads per million uniquely mapped (RPMUM)] in sense (S) and antisense (AS) orientation over the *ocl2*-coding region and ~3 kb of flanking genomic sequences on chromosome 10. Gray-shaded area highlights the *ubid* TE fragment 5' of *ocl2* having

Table 1 Curated genes with increased sense-oriented transcription in *rpm1* mutants

Gene	Annotation	Fold change (<i>rpm1</i> /WT)	P-value ^a
GRMZM2G303010	NBS-LRR disease resistance protein	9.113	7.07E-05
AC235534.1_FG007	<i>ocl2</i> (HD-ZIP IV)	7.932	2.65E-09
GRMZM2G161658	Epoxide hydrolase 2-like	7.893	3.01E-22
GRMZM2G132763	Putative LRR receptor-like protein kinase	6.681	2.27E-02
GRMZM2G062716	Defense-related protein (type 1 glutamine amidotransferase domain)	5.867	2.49E-02
GRMZM2G047105	Hypothetical, unknown protein	5.867	2.49E-02
GRMZM2G088413	Hypothetical, unknown protein	5.098	1.47E-02
GRMZM5G830269	Hypothetical, unknown protein	4.639	4.71E-02
GRMZM2G333140	Hypothetical, unknown protein	4.490	6.73E-02
GRMZM2G045155	B12D protein	4.243	2.55E-02
GRMZM2G147724	Phosphotidic acid phosphatase	4.023	1.51E-04
GRMZM2G043242	Putative ATP-binding, ATPase-like domain-containing protein	3.963	9.42E-08
GRMZM2G147399	Early nodulin 93	3.897	2.49E-02
GRMZM2G028677	Putative cytochrome P450 superfamily protein	3.824	7.34E-02
GRMZM2G009080	Hypothetical, unknown protein	3.542	7.05E-03
GRMZM2G131421	Early nodulin 93	3.421	8.64E-04
GRMZM2G174449	Hypothetical, unknown protein	3.329	1.26E-02
AC197705.4_FG001	Pyruvate decarboxylase isozyme 1	3.206	1.26E-02
GRMZM2G045560	WRKY DNA-binding domain-containing protein	3.102	1.89E-02
GRMZM2G300965	Respiratory burst oxidase-like protein B	2.897	3.37E-05
GRMZM2G053503	Ethylene-responsive factor-like protein (ERF1)	2.548	2.63E-02
GRMZM2G087063	Hypothetical, unknown protein, DUF 2930	2.444	2.54E-02
GRMZM2G051683	Anthocyanidin 5,3-O-glucosyltransferase	2.415	1.48E-03
GRMZM2G145213	14-3-3-like protein	2.406	1.27E-03
GRMZM2G024996	Pseudogene, transposon relic, upregulated	2.385	1.21E-03
GRMZM5G814164	Peroxisome biogenesis protein 3-2-like	2.360	2.80E-03
GRMZM2G168747	Nrat1 aluminum transporter 1	2.182	2.49E-02
GRMZM2G031827	Splicing factor U2af 38-kDa subunit	2.174	2.65E-02
GRMZM2G392791	Epoxide hydrolase 2-like	2.107	4.14E-02
GRMZM2G083538	Amino-acid-binding protein (ACR5)	2.058	2.54E-02
GRMZM2G021369	Putative AP2/EREBP transcription factor	1.986	2.95E-02
GRMZM2G013448	1-Aminocyclopropane-1-carboxylate oxidase	1.981	3.24E-02

^a P-values were adjusted by the Benjamini–Hochberg method for multiple testing (Benjamini and Hochberg 1995) as part of the DESeq analysis (Anders and Huber 2010).

genome (Figure 5A) (Schnable *et al.* 2009), it is underrepresented in both GRO-seq profiles as compared to the LTR Unknown class (RLX) and to type II DNA TEs (Figure 5B).

To determine if distinct TE groups were differentially represented in nascent transcriptomes, we compared normalized abundances of WT and *rpm1* mutant reads mapping to annotated TE superfamilies (Figure 5C). This comparison identified several TE superfamilies with elevated transcription in the *rpm1* mutant background (Figure 5C), including *Gypsy*, *Mariner*, *Helitron*, and *CACTA* noncoding elements. The majority of TE-derived GRO-seq reads cannot be mapped uniquely to specific genomic coordinates, limiting the detection of individual *rpm1*-affected TE loci to those TEs harboring significant sequence polymorphisms with respect to their family members. We thus employed the same method (DESeq; Anders and Huber 2010) used with genes to identify genomic regions annotated as TEs (Baucom *et al.* 2009; Schnable *et al.* 2009) having statistically significant differences in uniquely mapping GRO-seq read coverage. This method identifies 63 individual TEs (Figure 5D and Table

S5) representing several different superfamilies (Figure 5E) whose transcription is either increased (Figure 5F) or decreased in *rpm1* mutants in either sense or antisense directions. Only 28 of these unique TEs are not within or nearby (± 5 kb) genic regions, and some identify larger TE regions affected by RPD1 (Figure 5F and Table S5). These results agree with previous analyses (Hale *et al.* 2009) indicating that RPD1 prohibits mRNA accumulation of certain LTR retrotransposons by interfering with normal Pol II transcription and RNA processing.

Discussion

Our GRO-seq profiling of WT and *rpm1* mutant maize seedlings represents the first genome-wide nascent transcription analysis in plants and of a Pol IV mutant. The GRO-seq technique facilitates future studies of RNAP dynamics relevant to basic mechanisms of gene control in higher plants. Our results identify Pol IV effects on transcription at most gene boundaries, indicating that distinctions between higher

increased transcription in *rpm1* mutants. (D) Gene browser view of *GRMZM2G171408* showing increased antisense transcription in *rpm1* mutants. Sense (S) and antisense (AS) transcription occur in distinct units of GRO-seq coverage in both WT (black peaks) and *rpm1* mutant (green peaks) libraries. (E) Distribution of downstream features within 2 kb by type. Type I TEs are subdivided into LINE-like elements (RIX) and *Copia* (RLC), *Gypsy* (RLG), and Unknown (RLX) classes of LTR TEs. Color coding in E applies to TEs in browser views. Arrows indicate orientation of gene and TE features.

Figure 5 Pol IV loss affects both entire TE families and individual elements. (A) Distribution of TE categories within the B73 genome (Schnable *et al.* 2009). (B) Distribution of total (unique and repetitive) WT and *rpd1* mutant GRO-seq reads within the different TE categories shown in A. (C) Log₂ ratios (*rpd1* mutant/WT) of GRO-seq reads, normalized to total mappable reads, mapping to annotated TE superfamilies. (D) Log₂ fold change (*rpd1* mutant/WT) of uniquely mapping reads in sense and antisense orientation to genomic regions annotated as TEs vs. total coverage (averages of WT and *rpd1* mutant reads) to those regions. Triangles represent TEs with infinite fold change due to zero coverage from WT (top) or *rpd1* mutant (bottom) uniquely mapping reads. Of the 1,612,638 TE annotations analyzed, those with zero coverage in both WT and *rpd1* mutant datasets (1,392,382 and 1,399,008 for sense and antisense strand transcription, respectively) were excluded from the plots. Purple dots represent TEs with significantly (by the DESeq statistical method of Anders and Huber 2010; see *Materials and Methods*) increased or decreased GRO-seq read representation in *rpd1* mutants; orange stars or triangles represent those differentially transcribed TEs farther than 5 kb from the nearest FGS gene. Teal dots represent TEs within 20 Mb of the *rpd1* locus whose decreased transcription in *rpd1* mutants may reflect alignment artifacts (see *Materials and Methods*) and are excluded from subsequent analysis. (E) Distribution of transcriptionally altered unique TEs among TE categories shown in A. (F) Genome browser view of WT (black peaks) and *rpd1* mutant (green peaks) GRO-seq reads [normalized to reads per million uniquely mapped (RPMUM)] in sense (S) and antisense (AS) orientation

plant and metazoan transcription may be partly related to the plant-specific expansion of RNAP diversity. We also identified specific TE and genic alleles that show significant changes in nascent transcription $\pm$ RPD1, a dataset that should prove useful for better understanding the role(s) of Pol IV function in TE silencing, paramutation, and maize development.

The GRO-seq method captures snapshots of active transcription, which can identify entire transcription units from initiation to termination, helping to identify alternative TSSs and cryptic transcripts. Through comparisons of GRO-seq and RNA-seq profiles, it should be possible to identify the extent to which regulation of gene expression occurs at the level of post-transcriptional RNA stability. As GRO-seq reveals aspects of transcriptional regulation absent from the mature mRNA, nascent transcriptome profiles in metazoans, plants, and fungi will continue to define and distinguish RNAP functions across eukaryotes.

General Pol IV effects on genic transcription

Similar to metazoans, maize transcription extends beyond the PAS with termination occurring within $\sim$ 1–1.5 kb downstream. However, because of its additional RNAPs, plants may have alternative mechanisms to terminate Pol II transcription. At maize 3' gene ends, we speculate that Pol IV plays a role in attenuating aberrant readthrough transcription by Pol II into neighboring genes or TEs. This model is supported by the enrichment of RDR2-dependent 24-nt RNAs immediately downstream of PASs. Another possibility is that the kinetics of cotranscriptional mRNA splicing and/or polyadenylation are affected by Pol IV, perhaps related to the sharing of specific holoenzyme subunits (Haag *et al.* 2014). Together, the GRO-seq profiles and *rpdl* mutant analysis indicate that Pol II termination in maize is unique relative to metazoans.

Maize Pol IV also affects transcription at 5' gene boundaries. Our results show that *rpdl* mutants have decreased transcription at most gene TSSs, identifying a previously unknown role for Pol IV at Pol II initiation sites. Enrichment of 24-nt RNAs immediately (this article) and further upstream (on average 1.5 kb in maize; Gent *et al.* 2014) of genes supports the presence of Pol IV at genic promoters. Because Pol IV is predicted to engage transcription bubble-like DNA templates (Haag *et al.* 2012) and appears to initiate at AT-rich and nucleosome-depleted regions (Li *et al.* 2015), perhaps Pol IV holoenzymes synthesize RNA, either abortively or productively, at loci undergoing Pol II transcription initiation. Such behaviors could account for the relatively higher abundance of both sense and antisense 5' GRO-seq reads found in WT although this idea seems inconsistent with the observed patterns of 24mer vs. GRO-seq enrichment upstream of genes (Figure 3A and Figure S10). These discordant distributions indicate that the decrease in GRO-seq coverage near the TSS

is an indirect effect of Pol IV loss affecting transcription from another RNAP. It remains formally possible that Pol V contributes to this TSS-proximate transcription as *Arabidopsis* Pol V associates with TE-proximal promoters and more transiently with other promoters (Zhong *et al.* 2012). Alternatively, the decrease in promoter proximal GRO-seq read coverage could be due to titration of Pol II to other initiation sites in the absence of Pol IV (Hale *et al.* 2009), such as to the LTR TEs downstream of genes showing increased antisense transcription in *rpdl* mutants (Figure 4D).

The maize TSS-proximal peak of GRO-seq reads appears distinct from the promoter-proximal peak associated with canonical Pol II pausing in metazoans. Whether Pol II pausing, as described in humans and *Drosophila* (Core *et al.* 2008; Core *et al.* 2012), regulates transcription elongation of some maize genes remains unknown, although we observe no strong evidence for Pol II pausing in our datasets. Our experimental design focused on capturing a broad view of nascent transcription $\pm$ RPD1, and as such, we chose seedling tissue in which $>90\%$ of the genes produce detectable mRNAs via ultradeep sequencing (Martin *et al.* 2014). Seedlings grown under laboratory conditions may have no need for Pol II elongation regulation; *C. elegans* tends to show evidence of Pol II pausing only under stress (Kruesi *et al.* 2013; Maxwell *et al.* 2014). Additionally, we omitted Sarkosyl from the run-on reactions not knowing how this detergent might affect Pol IV function. This omission may also prevent detection of promoter-proximal Pol II pausing peaks as Sarkosyl can dissociate inhibitory factors holding Pol II at a canonical paused gene, *hsp70*, in *Drosophila* (Rougvie and Lis 1988). However, it should be noted that GRO-seq profiles in *Drosophila* indicate that the pausing peak, although greatly diminished, can still be detected in the absence of Sarkosyl (Core *et al.* 2012). While the nature of the maize TSS-proximal peak remains unclear, there is still a significant difference in transcription behavior at these regions $\pm$ RPD1.

A third characteristic identified in metazoan GRO-seq profiles is divergent transcription, which may be a by-product of previous Pol II initiations at the same promoter and/or a mechanism to maintain the nucleosome-free region (reviewed by Seila *et al.* 2009). These divergent transcripts may have roles as either regulatory scaffolds or sources of small RNAs (Core *et al.* 2008; Seila *et al.* 2008). Divergent transcription is prevalent at mammalian genes, but less so at *C. elegans* and *Drosophila* promoters, perhaps related to the directional specificity of favored promoter sequences (Core *et al.* 2012; Kruesi *et al.* 2013). We found no evidence of divergent transcription at maize promoters, potentially placing them in a similar category as *Drosophila*, which has a median 32-fold bias for sense-oriented transcription at promoters (Kruesi *et al.* 2013). This finding is curious because evidence in *Arabidopsis* indicates

over a 15-kb interval on chromosome 3 containing an RLX_milt type I element with increased transcription in *rpdl* mutants. Only the element outlined in black has significantly altered GRO-seq read representation in *rpdl* mutants based on the statistical threshold used (Anders and Huber 2010; see Materials and Methods).

that Pol V can be recruited to Pol II promoters (Zhong *et al.* 2012), and profiles of uniquely mapping 24-nt RNAs (Figure S10) place Pol IV near genes. It may be that plant RNAPs provide divergent transcription to help maintain nucleosome-free regions and that our GRO-seq assay conditions do not detect such nascent transcripts.

If Pol IV and/or Pol V are present immediately upstream of genes, then why is there little evidence of nascent transcription upstream of genic units? The Pol IV and V catalytic cores differ from that of Pol II, leading to relatively slow elongation rates and insensitivity to the Pol II inhibitor α -amanitin in both *Arabidopsis* (Haag *et al.* 2012) and maize (Haag *et al.* 2014). These differences may also affect their sensitivity to the limiting CTP present in the GRO-seq run-on reaction that affects Pol II NTP incorporation rates (Core *et al.* 2008) or their ability to incorporate the brominated UTP analog. Because only uniquely mapping reads were analyzed at gene boundaries, repetitive Pol IV- and/or Pol V-derived reads would be excluded. Additionally, any nascent Pol IV RNAs cotranscriptionally processed into siRNAs (Haag *et al.* 2012; Li *et al.* 2015) would not have been incorporated into the GRO-seq libraries. However, we are still able to identify effects of RPD1 loss on Pol II transcription (Table 1; Figure 4C; Figure S11; and Figure S12). Within limits of the GRO-seq assay, our results indicate that Pol II behaviors are generally affected by Pol IV, and this defines fundamental differences in genic transcription between metazoans and higher plants.

Pol IV affects gene regulation

McClintock referred to TEs as controlling elements because of their potential to affect gene regulation (McClintock 1951). TEs are transcriptionally repressed by Pol IV action either through direct competitions with Pol II (Hale *et al.* 2009) or through chromatin modifications dictated by Pol IV small RNAs (Matzke and Mosher 2014; Matzke *et al.* 2015); thus it is not surprising that increasing evidence points to Pol IV as a general source of epigenetic variation affecting gene regulation (Parkinson *et al.* 2007; Hollister *et al.* 2011; Eichten *et al.* 2012; Gent *et al.* 2012; Greaves *et al.* 2012; Erhard *et al.* 2013). Here we found Pol IV responsible for transcriptional control of both TEs and genes, consistent with a role of TEs as regulatory elements for specific alleles.

In accord with prior results (Erhard *et al.* 2009; Hale *et al.* 2009), loss of RPD1 results primarily in increased TE transcription, and both type I and type II TEs are among those affected. We identified only 28 unique TEs whose transcription was affected by RPD1 that were farther than 5 kb of annotated genes (Table S5). While specific repetitive TE classes are differentially affected, our results indicate that the majority of the genome-wide nongenic TEs are not transcribed at the seedling stage of development even in the absence of RPD1. Our analyses, however, likely underestimate the number of transcribed TEs because our sequencing depth, particularly in nongenic regions, is insufficient to detect low-abundance transcripts (Martin *et al.* 2014). Additionally, TE-like reads aligning to the B73 genome representing

unannotated TEs, TEs highly divergent from the Maize TE Consortium canonical set, or chimeras from multiple insertion events may have been misclassified as intergenic reads. Our results contrast with RNA-seq data from *rdt2* mutant meristems (Jia *et al.* 2009) showing significant increases in TE RNAs in the absence of this siRNA biogenesis factor. Assuming that TE RNA levels accurately reflect transcription rates, this difference in experimental results indicates that the mechanisms of TE repression among meristematic and differentiated cell types are distinct. Consistent with the limited cytosine methylation changes seen in the absence of maize RPD1 (Parkinson *et al.* 2007; Erhard *et al.* 2013; Li *et al.* 2014), Pol IV plays a potentially redundant role in repressing most TE transcription in whole seedlings although a fraction appear to be directly controlled by Pol IV action(s). The genomic and/or molecular features that distinguish these two general classes remain to be identified.

In addition to TEs, ~0.5% of all B73 alleles are transcriptionally responsive to RPD1, although loss of RPD1 can result in either increased or decreased transcription and in some cases in antisense orientation. It seems plausible that inappropriate antisense gene transcription could interfere with normal cotranscriptional RNA-processing steps or that sense-antisense RNA pairs could create double-stranded substrates for endonucleases, leading to post-transcriptional degradation. Pol II/Pol IV competitions for gene and TE promoters as previously proposed (Hale *et al.* 2009) and here exemplified by the B73 *ocl2* haplotype profiles (Figure 4C) remain a viable hypothesis to explain RPD1-specific effects. Such competitions could also account for the increased antisense transcription of many genes in the absence of RPD1 where the antisense transcription unit is contiguous with a downstream TE (Figure 4, D and E). It will be important to characterize the make-up of nuclear transcription “factories” in plants to see if and how Pol II and Pol IV potentially compete for specific genomic templates. It will also be necessary to compare whole-genome transcription profiles of mutants deficient for downstream components required for RNA-directed DNA methylation to identify genes whose regulation is associated with modulations of cytosine methylation. Independent of specific mechanisms, our genome-wide analyses indicate that Pol IV plays a significant regulatory role for specific alleles in the grasses. Given that there are multiple functional Pol IV isoforms defined by alternative second largest subunits in the grasses (Sidorenko *et al.* 2009; Stonaker *et al.* 2009; Haag *et al.* 2014; Sloan *et al.* 2014), and that haplotype diversity in maize is largely based on radically different intergenic TE compositions (Wang and Dooner 2006), the potential for regulatory diversity controlled by alternate RNAPs is immense in the maize pangenome.

Transcriptional control affecting paramutation

One feature of the *ocl2* haplotype, transcription of an upstream TE affected by Pol IV, is shared among *pl1* alleles affected by Pol IV loss. Several *pl1* alleles, including *Pl1-Rhoades*, have an upstream type II CACTA-like TE fragment

belonging to the *doppia* subfamily, which defines kernel-specific expression after conditioning in an *rpd1* mutant background (Erhard *et al.* 2013). This *Pl1-Rhoades doppia* fragment may be necessary, but is insufficient, for facilitating paramutations (Erhard *et al.* 2013). A related *doppia* fragment serves as a kernel-specific promoter and is partly required for paramutation occurring at *R-r:standard* haplotypes (Kermicle 1996; Walker 1998). At both *Pl1-Rhoades* and *R-r:standard*, other *doppia*-independent features are clearly important for mediating the *trans*-homolog interactions characteristic of paramutation (Kermicle 1996; Erhard *et al.* 2013). At the *B1-Intense* haplotype, paramutation interactions require a distal 5' transcriptional enhancer composed of direct repeats (Stam *et al.* 2002) that loop to the *b1* promoter region (Louwers *et al.* 2009), but there is currently no evidence supporting a functional role of specific TE sequences. In the four most studied examples of paramutation occurring in maize, transcription and/or transcriptional enhancers are functionally implicated in the mechanism required for establishing meiotically heritable repression associated with paramutation (Kermicle 1996; Sidorenko and Peterson 2001; Stam *et al.* 2002; Gross and Hollick 2007). In the absence of RPD1, high levels of gene expression are restored at *R-r:standard*, *Pl1-Rhoades*, and *B1-Intense* haplotypes previously repressed by paramutation (Hollick *et al.* 2005). At *Pl1-Rhoades*, loss of RPD1 results in increased *pl1* transcription, and this is often associated with meiotically heritable reversions of *Pl1-Rhoades* to a stable and highly expressed nonparamutant state (Hollick *et al.* 2005), resulting in strong plant pigmentation. We purposely excluded *Pl1-Rhoades* from materials used for the GRO-seq libraries reported here to avoid changes in transcription related to light perception that might be affected by seedling pigmentation. However, now having a list of RPD1-regulated alleles, we can use pedigree analyses to test whether these alleles also exhibit paramutation-like properties. Characterizing nascent transcription in different Pol IV and siRNA mutant backgrounds across *Pl1-Rhoades* and other haplotypes subject to paramutation promises to uncover important features of the underlying mechanism.

Our findings indicate that RPD1 uses mechanistically diverse actions, some of which may be independent of its catalytic action within the Pol IV holoenzyme, to regulate alleles in different genomic contexts. The identification of alleles affected by RPD1 loss now presents the opportunity to identify specific haplotype structures that have co-opted direct Pol IV action for their regulation. Given that maize Pol IV defines both mitotically and meiotically heritable patterns of gene regulation (Parkinson *et al.* 2007; Erhard *et al.* 2013), alterations of its function by developmental, environmental, or genealogical sources might lead to both ontogenetic and phylogenetic changes.

Acknowledgments

We thank Leighton Core for consultations regarding the GRO-seq protocol; Blake Meyers for kindly providing maize rRNA and tRNA sequences; and Janelle Gabriel, Brian

Giacopelli, Tzuu Fen Lee, Reza Hammond, Blake Meyers, and Keith Slotkin for comments and discussion. Illumina sequencing was supported by the Vincent J. Coates Genome Sequencing Laboratory, University of California at Berkeley. This work was supported by the National Science Foundation (MCB-0920623). The views expressed are solely those of the authors and are not endorsed by the sponsors of this work.

Literature Cited

- Alleman, M., L. Sidorenko, K. McGinnis, V. Seshadri, J. E. Dorweiler *et al.*, 2006 An RNA-dependent RNA polymerase is required for paramutation in maize. *Nature* 442: 295–298.
- Anders, S., and W. Huber, 2010 Differential expression analysis for sequence count data. *Genome Biol.* 11: R106.
- Ariel, F., T. Jegu, D. Latrasse, N. Romero-Barrios, A. Christ *et al.*, 2014 Noncoding transcription by alternative RNA polymerases dynamically regulates an auxin-driven chromatin loop. *Mol. Cell* 55: 383–396.
- Ashe, A., A. Sapetschnig, E.-M. Weick, J. Mitchell, M. P. Bagijn *et al.*, 2012 piRNAs can trigger a multigenerational epigenetic memory in the germline of *C. elegans*. *Cell* 150: 88–99.
- Barber, W. T., W. Zhang, H. Win, K. K. Varala, J. E. Dorweiler *et al.*, 2012 Repeat associated small RNAs vary among parents and following hybridization in maize. *Proc. Natl. Acad. Sci. USA* 109: 10444–10449.
- Barbour, J. R., I. T. Liao, J. L. Stonaker, J. P. Lim, C. C. Lee *et al.*, 2012 required to maintain repression2 is a novel protein that facilitates locus-specific paramutation in maize. *Plant Cell* 24: 1761–1775.
- Baucom, R. S., J. C. Estill, C. Chaparro, N. Upshaw, A. Jogi *et al.*, 2009 Exceptional diversity, non-random distribution, and rapid evolution of retroelements in the B73 maize genome. *PLoS Genet.* 5: e1000732.
- Benjamini, Y., and Y. Hochberg, 1995 Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. B* 57: 289–300.
- Brink, R. A., 1956 A genetic change associated with the *R* locus in maize which is directed and potentially reversible. *Genetics* 41: 872–889.
- Brink, R. A., 1958 Paramutation at the *R* locus in maize. *Cold Spring Harb. Symp. Quant. Biol.* 23: 379–391.
- Chandler, V. L., and M. Stam, 2004 Chromatin conversations: mechanisms and implications of paramutation. *Nat. Rev. Genet.* 5: 532–544.
- Coe, Jr., E. H., 1961 A test for somatic mutation in the origination of conversion-type inheritance at the *B* locus in maize. *Genetics* 46: 707–710.
- Core, L. J., J. J. Waterfall, and J. T. Lis, 2008 Nascent RNA sequencing reveals widespread pausing and divergent initiation at human promoters. *Science* 322: 1845–1848.
- Core, L. J., J. J. Waterfall, D. A. Gilchrist, D. C. Fargo, H. Kwak *et al.*, 2012 Defining the status of RNA polymerase at promoters. *Cell Reports* 2: 1025–1035.
- Dennis, E. S., and W. J. Peacock, 2007 Epigenetic regulation of flowering. *Curr. Opin. Plant Biol.* 10: 520–527.
- de Vanssay, A., A.-L. Bougé, A. Boivin, C. Hermant, L. Teyssset *et al.*, 2012 Paramutation in *Drosophila* linked to emergence of a piRNA-producing locus. *Nature* 490: 112–115.
- Eichten, S. R., N. A. Ellis, I. Makarevitch, C.-T. Yeh, J. I. Gent *et al.*, 2012 Spreading of heterochromatin is limited to specific families of maize retrotransposons. *PLoS Genet.* 8: e1003127.
- Erhard, Jr., K. F., J. L. Stonaker, S. E. Parkinson, J. P. Lim, C. J. Hale *et al.*, 2009 RNA polymerase IV functions in paramutation in *Zea mays*. *Science* 323: 1201–1205.

- Erhard, Jr., K. F., S. E. Parkinson, S. M. Gross, J. R. Barbour, J. P. Lim *et al.*, 2013 Maize RNA polymerase IV defines *trans*-generational epigenetic variation. *Plant Cell* 25: 808–819.
- Gent, J. I., Y. Dong, J. Jiang, and R. K. Dawe, 2012 Strong epigenetic similarity between maize centromeric and pericentromeric regions at the level of small RNAs, DNA methylation and H3 chromatin modifications. *Nucleic Acids Res.* 40: 1550–1560.
- Gent, J. I., N. A. Ellis, L. Guo, A. E. Harkess, Y. Yao *et al.*, 2013 CHH islands: *de novo* DNA methylation in near-gene chromatin regulation in maize. *Genome Res.* 23: 628–637.
- Gent, J. I., T. F. Madzima, R. Bader, M. R. Kent, X. Zhang *et al.*, 2014 Accessible DNA and relative depletion of H3K9me2 at maize loci undergoing RNA-directed DNA methylation. *Plant Cell* 26: 4903–4917.
- Greaves, I. K., M. Groszmann, H. Ying, J. M. Taylor, W. J. Peacock *et al.*, 2012 *Trans*-chromosomal methylation in *Arabidopsis* hybrids. *Proc. Natl. Acad. Sci. USA* 109: 3570–3575.
- Gross, S. M., and J. B. Hollick, 2007 Multiple *trans*-sensing interactions affect meiotically heritable epigenetic states at the maize *pl1* locus. *Genetics* 176: 829–839.
- Haag, J. R., T. S. Ream, M. Marasco, C. D. Nicora, A. D. Norbeck *et al.*, 2012 *In vitro* transcription activities of Pol IV, Pol V, and RDR2 reveal coupling of Pol IV and RDR2 for dsRNA synthesis in plant RNA silencing. *Mol. Cell* 48: 811–818.
- Haag, J. R., B. Brower-Toland, E. K. Krieger, L. Sidorenko, C. D. Nicora *et al.*, 2014 Functional diversification of maize RNA polymerase IV and V subtypes via alternative catalytic subunits. *Cell Reports* 9: 378–390.
- Hagemann, R., and W. Berg, 1978 Paramutation at the *sulfurea* locus of *Lycopersicon esculentum* Mill.: VII. Determination of the time of occurrence of paramutation by the quantitative evaluation of the variegation. *Theor. Appl. Genet.* 53: 113–123.
- Hale, C. J., J. L. Stonaker, S. M. Gross, and J. B. Hollick, 2007 A novel Snf2 protein maintains *trans*-generational regulatory states established by paramutation in maize. *PLoS Biol.* 5: e275.
- Hale, C. J., K. F. Erhard, Jr., D. Lisch, and J. B. Hollick, 2009 Production and processing of siRNA precursor transcripts from the highly repetitive maize genome. *PLoS Genet.* 5: e1000598.
- Herr, A. J., M. B. Jensen, T. Dalmay, and D. C. Baulcombe, 2005 RNA polymerase IV directs silencing of endogenous DNA. *Science* 308: 118–120.
- Hollick, J. B., 2012 Paramutation: a *trans*-homolog interaction affecting heritable gene regulation. *Curr. Opin. Plant Biol.* 15: 536–543.
- Hollick, J. B., and M. P. Gordon, 1993 A poplar tree proteinase inhibitor-like gene promoter is responsive to wounding in transgenic tobacco. *Plant Mol. Biol.* 22: 561–572.
- Hollick, J. B., G. I. Patterson, E. H. Coe, Jr., K. C. Cone, and V. L. Chandler, 1995 Allelic interactions heritably alter the activity of a metastable maize *pl* allele. *Genetics* 141: 709–719.
- Hollick, J. B., J. L. Kermicle, and S. E. Parkinson, 2005 *Rmr6* maintains meiotic inheritance of paramutant states in *Zea mays*. *Genetics* 171: 725–740.
- Hollister, J. D., L. M. Smith, Y.-L. Guo, F. Ott, D. Weigel *et al.*, 2011 Transposable elements and small RNAs contribute to gene expression divergence between *Arabidopsis thaliana* and *Arabidopsis lyrata*. *Proc. Natl. Acad. Sci. USA* 108: 2322–2327.
- Huang, Y., T. Kendall, and R. A. Mosher, 2013 Pol IV-dependent siRNA production is reduced in *Brassica rapa*. *Biology* 2: 1210–1223.
- Javelle, M., C. Klein-Cosson, V. Vernoud, V. Boltz, C. Maher *et al.*, 2011 Genome-wide characterization of the HD-ZIP IV transcription factor family in maize: preferential expression in the epidermis. *Plant Physiol.* 157: 790–803.
- Jia, Y., D. R. Lisch, K. Ohtsu, M. J. Scanlon, D. Nettleton *et al.*, 2009 Loss of RNA-dependent RNA polymerase 2 (RDR2) function causes widespread and unexpected changes in the expression of transposons, genes, and 24-nt small RNAs. *PLoS Genet.* 5: e1000737.
- Kashkush, K., and V. Khasdan, 2007 Large-scale survey of cytosine methylation of retrotransposons and the impact of readout transcription from long terminal repeats on expression of adjacent rice genes. *Genetics* 177: 1975–1985.
- Kermicle, J. L., 1996 Epigenetic silencing and activation of a maize *r* gene, pp. 267–287 in *Epigenetic Mechanisms of Gene Regulation*, edited by V. E. A. Russo, R. A. Martienssen, and A. D. Riggs. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Khaitová, L. C., M. Fojtová, K. Křížová, and J. Lunerová, J. Fulneček *et al.*, 2011 Paramutation of tobacco transgenes by small RNA-mediated transcriptional gene silencing. *Epigenetics* 6: 650–660.
- Kruesi, W. S., L. J. Core, C. T. Waters, J. T. Lis, and B. J. Meyer, 2013 Condensin controls recruitment of RNA polymerase II to achieve nematode X-chromosome dosage compensation. *ELife* 2: e00808.
- Langmead, B., C. Trapnell, M. Pop, and S. L. Salzberg, 2009 Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10: R25.
- Li, Q., S. R. Eichten, P. J. Hermanson, V. M. Zaunbrecher, J. Song *et al.*, 2014 Genetic perturbation of the maize methylome. *Plant Cell* 26: 4602–4616.
- Li, S., L. E. Vandivier, B. Tu, L. Gao, S. Y. Won *et al.*, 2015 Detection of Pol IV/RDR2-dependent transcripts at the genomic scale in *Arabidopsis* reveals features and regulation of siRNA biogenesis. *Genome Res.* 25: 235–245.
- Louwers, M., R. Bader, M. Haring, R. van Driel, W. de Laat *et al.*, 2009 Tissue- and expression level-specific chromatin looping at maize *b1* epialleles. *Plant Cell* 21: 832–842.
- Luo, J., and B. D. Hall, 2007 A multistep process gave rise to RNA polymerase IV of land plants. *J. Mol. Evol.* 64: 101–112.
- Martin, J. A., N. V. Johnson, S. M. Gross, J. Schnable, X. Meng *et al.*, 2014 A near complete snapshot of the *Zea mays* seedling transcriptome revealed from ultra-deep sequencing. *Sci. Rep.* 4: 4519.
- Martin, M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17: 10–12.
- Matzke, M. A., and R. A. Mosher, 2014 RNA-directed DNA methylation: an epigenetic pathway of increasing complexity. *Nat. Rev. Genet.* 15: 394–408.
- Matzke, M. A., T. Kanno, B. Huettel, L. Daxinger, and A. J. M. Matzke, 2007 Targets of RNA-directed DNA methylation. *Curr. Opin. Plant Biol.* 10: 512–519.
- Matzke, M. A., T. Kanno, and A. J. M. Matzke, 2015 RNA-directed DNA methylation: the evolution of a complex epigenetic pathway in flowering plants. *Ann. Rev. Plant Biol.* 66: 9.1–9.25.
- Maxwell, C. S., W. S. Kruesi, L. J. Core, N. Kurhanewicz, C. T. Waters *et al.*, 2014 Pol II docking and pausing at growth and stress genes in *C. elegans*. *Cell Reports* 6: 455–466.
- McClintock, B., 1951 Chromosome organization and genic expression. *Cold Spring Harb. Symp. Quant. Biol.* 16: 13–47.
- Morales-Ruiz, T., A. P. Ortega-Galisteo, M. I. Ponferrada-Marín, M. I. Martínez-Macías, R. R. Ariza *et al.*, 2006 *DEMETER* and *REPRESSOR OF SILENCING 1* encode 5-methylcytosine DNA glycosylases. *Proc. Natl. Acad. Sci. USA* 103: 6853–6858.
- Mosher, R. A., F. Schwach, D. Studholme, and D. C. Baulcombe, 2008 PolIVb influences RNA-directed DNA methylation independently of its role in siRNA biogenesis. *Proc. Natl. Acad. Sci. USA* 105: 3145–3150.
- Nobuta, K., C. Lu, R. Shrivastava, M. Pillay, E. De Paoli *et al.*, 2008 Distinct size distribution of endogenous siRNAs in

- maize: evidence from deep sequencing in the *mop1-1* mutant. *Proc. Natl. Acad. Sci. USA* 105: 14958–14963.
- Onodera, Y., J. R. Haag, T. Ream, P. Costa Nunes, O. Pontes *et al.*, 2005 Plant nuclear RNA polymerase IV mediates siRNA and DNA methylation-dependent heterochromatin formation. *Cell* 120: 613–622.
- Parkinson, S. E., S. M. Gross, and J. B. Hollick, 2007 Maize sex determination and abaxial leaf fates are canalized by a factor that maintains repressed epigenetic states. *Dev. Biol.* 308: 462–473.
- Pilu, R., D. Panzeri, E. Cassani, F. Cerino Badone, M. Landoni *et al.*, 2009 A paramutation phenomenon is involved in the genetics of maize *low phytic acid1-241* (*lpa1-241*) trait. *Heredity* 102: 236–245.
- Pontier, D., G. Yahubyan, D. Vega, A. Bulski, J. Saez-Vasquez *et al.*, 2005 Reinforcement of silencing at transposons and highly repeated sequences requires the concerted action of two distinct RNA polymerases IV in *Arabidopsis*. *Genes Dev.* 19: 2030–2040.
- Quinlan, A. R., and I. M. Hall, 2010 BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 26: 841–842.
- Rassoulzadegan, M., V. Grandjean, P. Gounon, S. Vincent, I. Gillot *et al.*, 2006 RNA-mediated non-Mendelian inheritance of an epigenetic change in the mouse. *Nature* 441: 469–474.
- Ream, T. S., J. R. Haag, A. T. Wierzbicki, C. D. Nicora, A. D. Norbeck *et al.*, 2009 Subunit compositions of the RNA-silencing enzymes Pol IV and Pol V reveal their origins as specialized forms of RNA polymerase II. *Mol. Cell* 33: 192–203.
- Rougvie, A. E., and J. T. Lis, 1988 The RNA polymerase II molecule at the 5' end of the uninduced *hsp70* gene of *D. melanogaster* is transcriptionally engaged. *Cell* 54: 795–804.
- Sabin, L. R., M. J. Delás, and G. J. Hannon, 2013 Dogma derailed: the many influences of RNA on the genome. *Mol. Cell* 49: 783–794.
- Schnable, P. S., D. Ware, R. S. Fulton, J. C. Stein, F. Wei *et al.*, 2009 The B73 maize genome: complexity, diversity, and dynamics. *Science* 326: 1112–1115.
- Seila, A. C., J. M. Calabrese, S. S. Levine, G. W. Yeo, P. B. Rahl *et al.*, 2008 Divergent transcription from active promoters. *Science* 322: 1849–1851.
- Seila, A., L. J. Core, J. T. Lis, and P. A. Sharp, 2009 Divergent transcription: a new feature of active promoters. *Cell Cycle* 8: 2557–2564.
- Shirayama, M., M. Seth, H.-C. Lee, W. Gu, T. Ishidate *et al.*, 2012 piRNAs initiate an epigenetic memory of non-self RNA in the *C. elegans* germline. *Cell* 150: 65–77.
- Sidorenko, L. V., and T. Peterson, 2001 Transgene-induced silencing identifies sequences involved in the establishment of paramutation of the maize *p1* gene. *Plant Cell* 13: 319–335.
- Sidorenko, L., J. E. Dorweiler, A. M. Cigan, M. Arteaga-Vazquez, M. Vyas *et al.*, 2009 A dominant mutation in *mediator of paramutation2*, one of three second-largest subunits of a plant-specific RNA polymerase, disrupts multiple siRNA silencing processes. *PLoS Genet.* 5: e1000725.
- Sloan, A. E., L. Sidorenko, and K. M. McGinnis, 2014 Diverse gene silencing mechanisms with distinct requirements for RNA polymerase subunits in *Zea mays*. *Genetics* 198: 1031–1042.
- Soderlund, C., A. Descour, D. Kudrna, M. Bomhoff, L. Boyd *et al.*, 2009 Sequencing, mapping, and analysis of 27,455 maize full-length cDNAs. *PLoS Genet.* 5: e1000740.
- Stam, M., C. Bebele, J. E. Dorweiler, and V. L. Chandler, 2002 Differential chromatin structure within a tandem array 100 kb upstream of the maize *b1* locus is associated with paramutation. *Genes Dev.* 16: 1906–1918.
- Stonaker, J. L., J. P. Lim, K. F. Erhard, Jr., and J. B. Hollick, 2009 Diversity of Pol IV function is defined by mutations at the maize *rmr7* locus. *PLoS Genet.* 5: e1000706.
- Stroud, H., M. V. C. Greenberg, S. Feng, Y. V. Bernatavichute, and S. E. Jacobsen, 2013 Comprehensive analysis of silencing mutants reveals complex regulation of the *Arabidopsis* methylome. *Cell* 152: 352–364.
- Swiezewski, S., P. Crevillen, F. Liu, J. R. Ecker, A. Jerzmanowski *et al.*, 2007 Small RNA-mediated chromatin silencing directed to the 3' region of the *Arabidopsis* gene encoding the developmental regulator, FLC. *Proc. Natl. Acad. Sci. USA* 104: 3633–3638.
- Tucker, S. L., J. Reece, T. S. Ream, and C. S. Pikaard, 2010 Evolutionary history of plant multisubunit RNA polymerases IV and V: subunit origins via genome-wide and segmental gene duplications, retrotransposition, and lineage-specific subfunctionalization. *Cold Spring Harb. Symp. Quant. Biol.* 75: 285–297.
- Walker, E. L., 1998 Paramutation of the *r1* locus of maize is associated with increased cytosine methylation. *Genetics* 148: 1973–1981.
- Wang, Q., and H. K. Dooner, 2006 Remarkable variation in maize genome structure inferred from haplotype diversity at the *bz* locus. *Proc. Natl. Acad. Sci. USA* 103: 17644–17649.
- Wang, X., H. Wang, J. Wang, R. Sun, J. Wu *et al.*, 2011 The genome of the mesopolyploid crop species *Brassica rapa*. *Nat. Genet.* 43: 1035–1039.
- Woodhouse, M. R., M. Freeling, and D. Lisch, 2006 Initiation, establishment, and maintenance of heritable *MuDR* transposon silencing in maize are mediated by distinct factors. *PLoS Biol.* 4: e339.
- Zhang, X., I. R. Henderson, C. Lu, P. J. Green, and S. E. Jacobsen, 2007 Role of RNA polymerase IV in plant small RNA metabolism. *Proc. Natl. Acad. Sci. USA* 104: 4536–4541.
- Zhong, X., C. J. Hale, J. A. Law, L. M. Johnson, S. Feng *et al.*, 2012 DDR complex facilitates global association of RNA polymerase V to promoters and evolutionarily young transposons. *Nat. Struct. Mol. Biol.* 19: 870–875.

Communicating editor: E. U. Selker

GENETICS

Supporting Information

<http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.174714/-/DC1>

Nascent Transcription Affected by RNA Polymerase IV in *Zea mays*

Karl F. Erhard Jr., Joy-El R. B. Talbot, Natalie C. Deans, Allison E. McClish, and Jay B. Hollick

SUPPORTING INFORMATION

SI FIGURES

Figure S1 GRO-seq reads align to both exonic and intronic sequences. Percentages of exonic and intronic sequence in all annotated single transcript genes (grey, repeated for unique vs. non-unique sets) and in WT (blue) and *rpd1* mutant (green) GRO-seq reads mapping to single transcript genes.

Figure S2 Most TE-like sequences can align to genes. Numbers and overlap of 32-mers randomly generated from the maize genome that align to annotated maize TE and gene sequences.

Figure S3 Genic reads are highly enriched in GRO-seq mappable reads. Distribution of genic vs. non-genic nucleotides in the maize genome (Whole Genome) and distribution of all non-rRNA / non-tRNA B73 mappable GRO-seq reads to genic vs. non-genic sequences.

Figure S4 Distribution of uniquely mapping WT GRO-seq reads around genes. Reads per million uniquely mapped (RPMUM) WT coverage in 50 nt windows +/- 5 kb from gene boundaries in both sense (top plot) and antisense (bottom plot) orientations. Each row represents the average coverage of 60 neighboring genes sorted by maximum sum contribution.

Figure S5 FGS gene annotations start with ATG more frequently than other loci. The first three bases of all FGS gene models were tallied to determine any bias for the ATG set indicative of translation start codons. As a control for general three base set distribution, the first three bases 5 kb upstream of the FGS start were also tallied. In addition to the ATG set, the top four ranking three base sets are individually highlighted.

Figure S6 Alternative TSS definition has little effect on coverage profiles. TSS sites were defined by the 5' end of full-length cDNAs (flcDNAs) from a library of predominantly 7 day-old seedling tissue (Soderlund *et al.* 2009). Metagene analysis was performed across all flcDNA defined TSSs in 10 nt non-overlapping windows with the total coverage in reads per million uniquely mapped (RPMUM) displayed. Pair-wise Welch's *t*-tests evaluated the difference between WT and *rpd1* mutant read abundance at each window, and a Holm-Bonferoni correction ($\alpha = 0.05$) was applied to identify the windows that were significantly different (red horizontal bar and pink shading).

Figure S7 Strong antisense coverage upstream of GRMZM2G061206. Genome browser view of GRMZM2G061206 showing sense (S) and antisense (AS) WT (black) and *rpd1* mutant (green) reads. Annotated genes are in blue with blue lines indicating introns, thin boxes UTRs and thick boxes CDS, and the arrow indicates the TSS. Annotated Type II TE is in gold with an arrow indicating its orientation.

Figure S8 Correlation between wild-type (WT) and *rpd1* mutant uniquely mapping GRO-seq coverage at genes. Near-genic regions were divided into five sections: 1 kb upstream, the first 1 kb after the TSS, a variable internal region, the last 1 kb before the gene end, and 1 kb downstream. GRO-seq coverage is plotted for both sense and antisense directions (relative to the gene model) in reads per kb per million uniquely mapped. To include genes with zero read abundance on the logarithmic scale, they were forced to 1/10th the minimum non-zero abundance of both datasets for that plot. Data fit lines represent linear regression models of the $\log_{10}$ transformed WT and *rpd1* mutant abundances. Rho (ρ) values represent the Spearman's rank correlation coefficient for each region. Total number of genes in each plot is 31,794 except for the interior of the gene where the total gene length had to be greater than 2 kb; 23,050 genes in total.

Figure S9 Fold change between WT and *rpd1* mutant antisense GRO-seq read coverage. Variation in antisense coverage between WT and *rpd1* mutant libraries for all FGS genes is plotted. Each horizontal line represents the average coverage of 60 neighboring genes when sorted by their maximum sum contribution (see *Materials and Methods*). Fifty nucleotide windows with zero coverage in either library are plotted in white. The gold bar highlights the inner 90% of genes used in Figure 3A.

Figure S10 Coverage of uniquely mapping 24mers near gene boundaries. Small RNA datasets representing B73 (WT) or *rdr2* mutants introgressed into B73 were pooled from publically available datasets, processed, aligned, and uniquely mapping 24mers were tallied near gene boundaries as with the GRO-seq datasets (see *Materials and Methods*). **(A)** Mean coverage for sense (left) and antisense (right) strands within 1 kb of gene boundaries for the same 90% of genes analyzed in the GRO-seq metagene profiles (see Figure 3A). 95% confidence intervals are shaded in grey and pale purple. **(B - C)** Mean coverage of uniquely mapping 24mers in groups of 60 neighboring genes, when sorted as in the GRO-seq analysis (see Figure 3B) in WT **(B)** and *rdr2* mutant **(C)** datasets. **(D)** Log₂ fold-change of *rdr2* mutant mean coverage **(C)** over WT **(B)**. Only blocks (50 nt window over 60 genes) with coverage in both datasets are analyzed--any blocks with zero coverage in either or both datasets are left white. Gold vertical bars highlight the genes that were included in the metagene plots in **(A)**. TSS: transcription start site; end: 3' annotated end of gene; WT: wild-type; RPMUM: reads per million uniquely mapped.

Figure S11 RT-PCR analysis mirrors increased transcription of three genes in the absence of RPD1. Primers specific to each gene and a control gene *alanine aminotransferase (aat)* (see *Materials and Methods*) were used to amplify identical quantities of oligo dT-primed cDNA templates generated from three genotyped homozygous *rpd1-1* mutants and three homozygous wild-type siblings. Amplifications using control samples prepared with no reverse transcriptase were performed and used to subtract background intensity from amplicon intensity signals. Error bars are ± 1 s.e.m.

Figure S12 qRT-PCR analysis mirrors increased transcription of *ocl2* in the absence of RPD1. Three biological replicates each of three 8 day-old seedling shoots were analyzed for *ocl2* and *aat* (control) RNA levels by qRT-PCR with three technical replicates each. The relative RNA abundance represents $2^{(aat\ Ct - ocl2\ Ct)}$ where mean and confidence intervals of ± 1 s.e.m. were calculated on the ΔCt values prior to converting to fold change.

SI TABLES

Table S1 Sources of genomic feature annotations

Feature Type	File Format	Original Location
rRNA/tRNA sequences	FASTA	Dr. Blake Meyers, University of Delaware
FGS sequences	FASTA	release-5b/filtered-set/ZmB73_5b_FGS_genes.fasta.gz ^a
FGS positions	GFF3	release-5b/filtered-set/ZmB73_5b_FGS.gff.gz ^a
MTEC consensus sequences	FASTA	release-5b/repeats/repeat-ls.tar.gz ^a
MTEC consensus positions	GFF3	release-5b/repeats/ZmB73_5a_MTEC_repeats.gff.gz ^a
Maize AGPv2 build 5b B73 genome pseudo-chromosomes	FASTA	release-5b/assembly/ZmB73_RefGen_v2.tar.gz ^a

^a Originally these files were downloaded from ftp://<ftp.maizesequence.org/pub/gramene/maizesequence.org/> base directory. Now you can access these same files from this base directory: ftp://<ftp.gramene.org/pub/gramene/maizesequence.org/>.

Abbreviations: Filtered Gene Set (FGS) and Maize TE Consortium (MTEC)

Table S2 Primers used for RT-PCR and qRT-PCR analysis

Gene Model	Exons	Amplicon	Primer Sequence (5' to 3')	
	Amplified	Size (bp) ^a	Forward	Reverse
<i>ocl2</i> (AC235534.1_FG007)	6 & 7	176 bp	TGTTTCATTGATGGACTGC	AGCGCATAAGAGGCTGGTAG
	8 & 9	125 bp	ATCCTCAGCAATGGTGGCCCTAT	AATCAAGATGCTGTTCAGGTGGGAG
GRMZM2G043242	11 – 14	243 bp	TATGCATGTGGCAGCTAAGG	AATCGTGCTTCGAGTCTTGC
GRMZM2G161658	2 – 4	341 bp	CAGGAATTTGGTGTGCTGA	CCCGGAGCGTTGTATGTAAT
<i>aat</i> (GRMZM2G088064)	11 – 13	283 bp	ATGGGGTATGGCGAGGAT ^b	TTGCACGACGAGCTAAAGACT ^b
	12 & 13	122 bp	CAATATCACTGGTCAAATCCTTGCGA	TTGCACGACGAGCTAAAGACT ^b

^a Shorter product sizes were used for the qRT-PCR reaction, while longer products were used for the RT-PCR experiment.

^b From Woodhouse *et al.* 2006.

Table S3 Genes differentially transcribed in *rpdl* mutants

Increased Transcription - Sense Orientation ^a					
Gene ID	Annotation	TE / Repeat Content ^b	Genomic Position ^c	Log ₂ Fold Change (<i>rpdl</i> / WT) ^d	p-Value ^e
GRMZM5G832375	Pyruvate decarboxylase (<i>pdh3</i>)	none	chr1:45510990-45511749	2.527	3.37E-05
GRMZM2G087063	Hypothetical, unknown protein, DUF 2930	exon and intron	chr1:80620045-80622805	1.290	2.54E-02
GRMZM2G143082	Hypothetical, unknown protein	none	chr1:86260377-86261631	3.652	2.28E-09
GRMZM2G147724	Phosphotidic acid phosphatase	exon (3' UTR)	chr1:147013445-147016840	2.008	1.51E-04
GRMZM2G081151	Beta-amryin synthase	introns	chr1:160287318-160457160	1.407	4.98E-03
GRMZM2G303010	NBS-LRR disease resistance protein	none	chr1:173142735-173144264	3.188	7.07E-05
GRMZM2G392791	Epoxide hydrolase 2-like	exon	chr1:179988453-179990792	1.075	4.14E-02
GRMZM2G132763	Putative leucine-rich repeat receptor-like protein kinase	exon	chr1:187587612-187591058	2.740	2.27E-02
GRMZM2G430455	Ribosomal protein S4	exon (3' UTR)	chr1:188855847-188859402	---	4.69E-05
GRMZM2G161233	NAD-dependent epimerase/dehydratase family protein	none	chr1:192730986-192732596	2.527	4.83E-03
GRMZM2G083538	Amino acid binding protein (ACR5)	exon	chr1:194674440-194677737	1.041	2.54E-02
GRMZM2G088413	Hypothetical, unknown protein	introns	chr1:254419042-254422294	2.350	1.47E-02
GRMZM2G143142	ASF/SF2-like pre-mRNA splicing factor srp30	exon (3' UTR) and intron	chr1:273032625-273036627	1.725	3.18E-03

GRMZM2G161658	Epoxide hydrolase 2-like	none	chr3:36217073-36226741	2.981	3.01E-22
GRMZM2G081363	Hypothetical, unknown protein	exon	chr3:127968485-127969840	2.367	2.49E-02
GRMZM5G866269	Hypothetical, unknown protein	exon	chr3:151450891-151451823	4.021	7.96E-05
GRMZM2G051683	Anthocyanidin 5,3-O-glucosyltransferase	none	chr3:205694156-205695856	1.272	1.48E-03
GRMZM2G145213	14-3-3-like protein	intron	chr4:5962321-5967839	1.266	1.27E-03
GRMZM2G174449	Hypothetical, unknown protein	none	chr4:25070680-25072205	1.735	1.26E-02
GRMZM2G043242	Putative ATP-binding, ATPase-like domain-containing protein	introns+	chr4:218998953-219053414	1.987	9.42E-08
GRMZM2G062716	Defense-related protein (Type 1 glutamine amidotransferase domain)	none	chr4:234873379-234874456	2.553	2.49E-02
GRMZM2G333140	Hypothetical, unknown protein	3' UTR	chr5:19298125-19301282	2.167	6.73E-02
GRMZM2G098438	Hypothetical, unknown protein	exon and intron	chr5:70050931-70063152	1.748	4.19E-06
GRMZM2G168747	Nrat1 aluminum transporter 1	none	chr5:74612262-74615918	1.125	2.49E-02
GRMZM2G028677	Putative cytochrome P450 superfamily protein	none	chr5:159912586-159914501	1.935	7.34E-02
GRMZM2G021369	Putative AP2/EREBP transcription factor	exons	chr5:210273706-210275023	0.990	2.95E-02
GRMZM5G830269	Hypothetical, unknown protein	none	chr6:11200705-11201863	2.214	4.71E-02
GRMZM2G009080	Hypothetical, unknown protein	exons	chr6:84050523-84051872	1.825	7.05E-03
GRMZM2G543070	Hypothetical, unknown protein	none	chr6:90044328-90045333	1.159	2.92E-02
GRMZM2G110742	Putative thioredoxin superfamily protein with	exons (T02	chr7:57059839-57064670	3.652	5.47E-07

	Squamosa-promoter binding (SBP) domain	transcript only)			
GRMZM5G814164	Peroxisome biogenesis protein 3-2-like	exon (5' UTR) and intron	chr7:85874277-85881002	1.239	2.80E-03
GRMZM2G045155	B12D protein	none	chr7:164234830-164235658	2.085	2.55E-02
GRMZM2G146004	Protein induced upon tuberization	exon	chr8:5186351-5187295	1.392	7.56E-05
GRMZM2G053503	Ethylene-responsive factor-like protein (ERF1)	none	chr8:35561736-35562916	1.349	2.63E-02
GRMZM2G031827	Splicing factor U2af 38 kDa subunit	5' UTR, exon and intron	chr8:70136197-70144793	1.120	2.65E-02
AC197705.4_FG001	Pyruvate decarboxylase isozyme 1	exon	chr8:118165588-118167724	1.681	1.26E-02
GRMZM2G013448	1-aminocyclopropane-1-carboxylate oxidase	none	chr8:138861203-138862890	0.986	3.24E-02
GRMZM2G045560	WRKY DNA-binding domain-containing protein	intron	chr8:168823637-168828313	1.633	1.89E-02
GRMZM2G147399	Early nodulin 93	none	chr9:20665757-20666996	1.962	2.49E-02
GRMZM2G131421	Early nodulin 93	none	chr9:20693835-20695096	1.774	8.64E-04
GRMZM2G046528	Hypothetical, unknown protein	exons (3' UTR)	chr9:25386483-25390940	1.487	2.15E-02
GRMZM2G047105	Hypothetical, unknown protein	exon, 3' UTR and intron	chr9:43122694-43125766	2.553	2.49E-02
GRMZM2G024996	Pseudogene, transposon relic - upregulated	exon	chr9:134749957-134751048	1.254	1.21E-03
GRMZM2G300965	Respiratory burst oxidase-like protein B	exon	chr10:24649645-24653324	1.535	3.37E-05

GRMZM2G070575	Cycloartenol synthase	introns	chr10:66168601-66179083	2.974	1.35E-04
AC235534.1_FG007	ocl2 (HD-ZIP IV)	exons and intron	chr10:136932960-136939534	2.988	2.65E-09

Increased Transcription - Antisense Orientation^a

Gene ID	Annotation	TE / Repeat Content ^b	Genomic Position ^c	Log2 Fold Change (<i>rpdl</i> / WT) ^d	p-Value ^e
GRMZM2G385021	Hypothetical, unknown protein	none	chr1:45510990-45511749	2.459	4.11E-03
GRMZM2G043141	Hypothetical, DUF 1336 Domain-containing protein	exon	chr1:75147439-75151689	2.082	4.63E-02
GRMZM2G130034	Putative ureidoglycolate hydrolase	exons, introns and 3' UTR	chr1:167851430-167863929	3.263	1.62E-02
GRMZM2G443308	Hypothetical, unknown protein	none	chr1:177766328-177767115	3.259	3.39E-03
GRMZM2G421579	Putative MYB DNA binding protein	exons and introns	chr1:193768325-193775666	4.161	4.19E-08
GRMZM2G131683	Electron transporter, glutaredoxin-related protein	exon (3' UTR)	chr1:230186289-230191174	2.628	1.75E-05
GRMZM2G131715	Hypothetical, unknown protein	exon (3' UTR)	chr1:230198773-230199704	2.000	3.27E-02
GRMZM2G107540	Histone H2A	none	chr1:270053078-270054144	3.162	1.05E-05
GRMZM5G891855	Hypothetical, unknown protein	intron	chr2:224015038-224047387	1.836	5.44E-02
GRMZM5G837822	Hevamine-A - chitinase (plant defense)	none	chr3:176570295-176571270	2.708	4.82E-03
GRMZM2G430936	Acidic endochitinase-like	none	chr3:176597396-176598753	2.692	1.01E-02
GRMZM2G052365	Subtilase-like protease-like	intron	chr3:193637409-193648778	3.737	1.29E-02

GRMZM2G126239	Homeobox-leucine zipper protein ATHB-4	exon	chr3:214857327-214861719	2.177	6.86E-03
GRMZM2G094900	Type I inositol-1,4,5-triphosphate 5-phosphatase CVP2-like protein	exon	chr5:14650208-14653205	3.848	6.05E-07
GRMZM2G472346	IQ calmodulin-binding motif family protein	exons and intron	chr5:67476538-67482854	3.790	8.89E-10
GRMZM2G074634	Cysteine-type peptidase	none	chr5:84101452-84105175	1.967	3.76E-02
GRMZM2G171408	U-box domain-containing protein 10 or armadillo/beta-catenin-like repeat-containing protein	exons	chr7:115182036-115185167	3.788	2.55E-08
AC225205.3_FG003	Phosphatidylinositol 3-and 4-kinase family like protein	none	chr7:130092349-130094061	2.841	6.18E-04
GRMZM5G824938	Phosphatidylinositol 3-and 4-kinase family like protein	none	chr7:130097614-130101719	3.396	4.78E-09
GRMZM2G181378	E3 ubiquitin-protein ligase UPL5-like	exon and introns	chr7:132186216-132203350	2.350	2.39E-02
GRMZM2G082365	H/ACA ribonucleoprotein complex subunit 4-like protein and putative kinetochore protein	exon	chr7:168403976-168405995	2.213	4.82E-03
GRMZM2G181266	3-hydroxyacyl-CoA dehydratase PASTICCINO 2A-like	exon (5' UTR) and introns	chr8:14597519-14604732	2.932	4.19E-08
GRMZM2G009894	60S ribosomal protein L29	none	chr8:36503543-36505405	2.649	2.58E-05
GRMZM2G396248	Cytochrome P450	none	chr8:166900775-166902970	2.294	2.95E-02
GRMZM2G396246	Hypothetical, unknown protein	exon (3' UTR)	chr8:166909761-166911646	2.615	7.47E-03

GRMZM2G510796	Hypothetical, unknown protein	none	chr8:168817914-168818496	3.193	4.82E-03
GRMZM2G045560	WRKY DNA-binding domain-containing protein	intron	chr8:168823637-168828313	3.019	4.19E-08
GRMZM2G149105	Hypothetical, unknown protein	none	chr8:174306169-174306927	2.761	2.76E-02
GRMZM5G872568	Hypothetical, unknown protein	intron+	chr9:139536440-139543922	3.939	4.11E-03
GRMZM5G884151	Hypothetical, unknown protein	exon (3' UTR)	chr9:142015409-142016063	2.619	2.68E-03
GRMZM2G180258	Hypothetical, unknown protein	exon	chr10:2137242-2141058	2.434	1.82E-04
GRMZM2G084473	Histidine kinase cytokinin receptor	exon	chr10:146545112-146548756	3.075	1.07E-02

Decreased Transcription - Sense Orientation^a

Gene ID	Annotation	TE / Repeat Content ^b	Genomic Position ^c	Log ₂ Fold Change (<i>rpdl</i> / WT) ^d	p-Value ^e
GRMZM2G475380	Gibberellin 20 oxidase	intron	chr1:4767327-4772060	-1.563	4.57E-05
GRMZM2G496991	Hypothetical, unknown protein, DUF581 superfamily	none	chr1:17624743-17625481	-1.511	9.17E-03
GRMZM2G164974	3-ketoacyl-CoA synthase 6-like	none	chr1:27506254-27508489	-1.126	2.49E-02
GRMZM2G089812	Nuclear transcription factor Y subunit C-1 isoform	exon	chr1:34594746-34596756	-1.553	7.77E-02
GRMZM2G054332	ATPase, coupled to transmembrane movement of substance	none	chr1:43054782-43057900	-1.032	7.08E-02
GRMZM2G038973	Glycosyl hydrolase family 38 protein (alpha-mannosidase)	introns+	chr1:117114680-117142745	-2.264	9.47E-12
GRMZM2G337242	Glycosyl hydrolase family 38 protein (alpha-mannosidase)	introns+	chr1:117155139-117165079	-1.268	2.65E-02
GRMZM5G841743	Hypothetical, unknown protein	none	chr1:123015378-123016357	-2.192	4.72E-02

GRMZM2G022804	Hypothetical, unknown protein	intron and exon (3' UTR)	chr1:123254749-123256442	-2.498	1.31E-02
GRMZM2G067929	Hypothetical, unknown protein	none	chr1:124817067-124821052	-2.685	8.54E-06
GRMZM2G081977	Hypothetical, unknown protein, DUF828 superfamily	exons and introns	chr1:149708692-149715244	-1.534	2.27E-02
GRMZM2G093038	Hypothetical, unknown protein	exon and intron (3' UTR)	chr1:156872706-156873991	-2.232	4.96E-02
GRMZM2G429128	DNA-binding storekeeper protein-related	exon (5' UTR)	chr1:157173981-157177692	-2.686	1.97E-07
GRMZM2G331902	Probable polyamine transporter At3g19553-like	exon	chr1:160808686-160810744	-1.608	7.40E-02
GRMZM2G460861	SAUR33-auxin-responsive SAUR family member	none	chr1:209180557-209181568	-1.297	3.98E-02
GRMZM2G153184	Chlorophyll a-b binding protein 4; PSI light-harvesting complex type 4 protein	exon (3' UTR)	chr1:213652786-213654502	-1.423	6.54E-05
AC217050.4_FG007	Terpene synthase 7	introns	chr1:214886200-214892746	-1.830	6.38E-05
AC217910.3_FG004	Hypothetical, unknown protein	none	chr1:217279821-217280603	-1.536	2.69E-02
GRMZM2G143883	Long-chain-alcohol oxidase FAO1-like	exon and intron	chr1:232400407-232405547	-0.914	7.29E-02
GRMZM2G115646	Glutamate-ammonia ligase-like protein	exon and introns	chr1:235645071-235668787	-1.195	2.24E-03
GRMZM2G060742	Citrate transporter family protein	intron	chr1:235973873-235982962	-0.923	8.66E-02
GRMZM5G858417	Nucleobase-ascorbate	none	chr1:262997924-263003009	-1.001	3.68E-02

	transporter LPE1				
GRMZM2G396114	Putative POX domain/homeobox DNA-binding domain family protein	5' UTR, exons and intron	chr1:273091163-273101408	-0.928	9.76E-02
GRMZM2G010034	Hypothetical, unknown protein	none	chr1:275382091-275384621	-1.398	4.14E-02
GRMZM2G089282	Hypothetical, unknown protein	none	chr2:9587226-9588772	-1.313	2.63E-02
GRMZM2G093526	Putative ent-kaurene synthase B	intron	chr2:10568344-10572400	-1.036	7.29E-02
GRMZM2G000166	Putative metal ion-binding protein	none	chr2:13712123-13713766	-1.491	3.88E-03
GRMZM2G168474	Cis-zeatin O-glucosyltransferase 1 (cisZOG1)	exon and intron and 5' UTR	chr2:18460858-18463365	-1.976	4.35E-03
GRMZM2G337113	Glyceraldehyde-3-phosphate dehydrogenase	exon	chr2:42257261-42259297	-1.094	3.93E-02
GRMZM2G134264	Hypothetical protein with Agglutinin superfamily domain	intron	chr2:102136620-102139245	-1.609	1.43E-06
GRMZM2G090028	Hypothetical, unknown protein	5' UTR	chr2:102279931-102283830	-1.387	9.17E-03
GRMZM2G056500	Hypothetical protein with Agglutinin superfamily domain	intron	chr2:102750321-102752520	-1.552	2.41E-05
GRMZM2G090675	Cystathionin beta synthase protein	introns	chr2:102897151-102905974	-0.946	5.84E-02
GRMZM2G152189	Hypothetical, unknown protein	exon (3' UTR)	chr2:209536404-209537584	-1.852	6.76E-02
GRMZM2G009188	11-beta-hydroxysteroid	introns	chr2:222270518-222276029	-1.062	1.80E-02

	dehydrogenase 1B-like	and exon (3' UTR)			
GRMZM2G085019	NADP-dependent malic enzyme, chloroplastic precursor	exons and intron	chr3:7275169-7280519	-1.516	1.34E-05
GRMZM2G339866	Hypothetical, unknown protein DUF3339 superfamily	none	chr3:32466336-32467145	-1.629	1.66E-03
GRMZM2G072298	Glycerol-3-phosphate acyltransferase 1	exon and intron	chr3:35353116-35357015	-3.086	1.26E-07
GRMZM2G062531	Methylsterol monooxygenase 1-1-like	none	chr3:101735928-101737972	-4.456	6.76E-02
GRMZM2G138258	Hypothetical, unknown protein	exon and intron	chr3:147397873-147401506	-1.711	1.65E-06
GRMZM5G882078	Putative ankyrin-kinase	exon	chr3:168959356-168962676	-1.128	2.54E-02
GRMZM2G077333	PSII 22 kDa protein	intron	chr3:174825201-174827787	-1.067	2.01E-02
GRMZM2G132169	L-ascorbate oxidase	none	chr3:183644921-183647436	-1.420	1.26E-02
GRMZM2G121878	Carbonic anhydrase	intron	chr3:215466579-215473931	-1.951	2.55E-09
GRMZM2G162200	Ribulose biphosphate carboxylase/oxygenase activase, chloroplast Precursor	exon (3' UTR)	chr4:693736-696087	-1.082	6.43E-02
GRMZM2G143373	Cis-2, 3-dihydrobiphenyl-2,3-diol dehydrogenase	exon and intron	chr4:17332223-17339190	-1.117	1.62E-02
GRMZM2G337387	Hypothetical, unknown protein, DUF247 superfamily	intron	chr4:30678789-30681127	-2.793	1.34E-05
GRMZM2G374302	Arginine decarboxylase	intron	chr4:144835678-144840927	-2.279	2.28E-09
GRMZM2G038243	Transferase family protein	exon and intron	chr4:151478038-151481556	-1.566	1.34E-05
GRMZM2G163925	Phosphoribosylanthranilate transferase	none	chr4:153318407-153320975	-1.564	1.26E-02

GRMZM2G017268	Putative Myb DNA-binding domain superfamily protein	none	chr4:197071819-197073073	-1.530	8.66E-02
GRMZM2G027479	Thionin-like protein (plant defense peptide)	intron	chr4:204275345-204276126	-1.561	3.64E-03
GRMZM2G155253	Fructose-bisphosphate aldolase	exon	chr4:204599067-204601218	-2.323	1.35E-04
GRMZM2G401040	ATP synthase F1, delta subunit family protein	none	chr5:8334393-8335549	-1.055	2.88E-02
GRMZM2G455869	MYB transcription factor in SANT Superfamily	intron	chr5:10172933-10174782	-1.807	4.72E-02
GRMZM2G377341	Acetyl-CoA carboxylase 1-like	exon (3' UTR) and intron	chr5:38567212-38578529	-0.919	9.76E-02
GRMZM2G042758	Hypothetical protein with similarity to GDSL-motif lipase/hydrolase-like protein (rice and others) and anther-specific proline-rich protein APG	exon (3' UTR) and intron	chr5:48384381-48390703	-1.113	1.03E-02
GRMZM2G020320	Glycerol-3-phosphate acyltransferase 8	none	chr5:69114970-69116904	-1.968	1.21E-03
GRMZM2G147172	Putative F-box family protein	none	chr5:136581652-136583153	-1.537	1.21E-03
GRMZM2G169458	Fatty aldehyde dehydrogenase 1	intron	chr5:190228137-190232283	-1.133	9.68E-03
GRMZM2G398807	Cortical cell-delineating protein	none	chr5:192613610-192614489	-1.157	8.04E-02
GRMZM2G162529	Phosphoribulokinase	none	chr5:199592727-199595057	-1.501	2.63E-02
GRMZM2G466309	Hypothetical, unknown protein	none	chr5:202304835-202305876	-1.258	8.66E-02
GRMZM2G103101	Chlorophyll a-b binding	none	chr5:209873678-209875630	-1.041	4.38E-02

	protein 4; PSI light-harvesting complex type 4 protein				
GRMZM2G052422	1-aminocyclopropane-1-carboxylate oxidase 1 (Acc oxidase)	none	chr5:210776511-210779081	-1.710	1.97E-07
GRMZM2G318662	Hypothetical, unknown protein	exon (3' UTR)	chr5:213264683-213268374	-0.916	7.16E-02
GRMZM2G358540	VAMP protein SEC22	none	chr6:655472-656792	-1.048	7.08E-02
GRMZM2G461716	Putative chloroplast nucleoid DNA-binding protein cnd41	exon (3' UTR)	chr6:61601766-61603753	-1.826	3.63E-02
GRMZM2G142345	E3 ubiquitin-protein ligase XBAT33 and ankyrin repeat-containing protein	exons and intron	chr6:63430823-63436487	-1.568	3.68E-02
GRMZM2G169321	Adiponectin receptor 1	exon	chr6:96919201-96922279	-0.998	4.05E-02
GRMZM2G137421	NRT1/ PTR FAMILY 4.6-like (putative nitrate transporter)	5' UTR, introns, exon and 3' UTR	chr6:103602046-103609491	-1.396	2.49E-02
GRMZM2G030583	Monoterpene synthase	introns	chr6:107307539-107312261	-1.036	4.25E-02
GRMZM2G095404	Peroxidase 11-like	none	chr6:118788607-118790352	-1.356	2.49E-02
GRMZM2G162755	Anthocyanidin 3-O-glucosyltransferase	none	chr6:119876153-119878032	-1.396	2.49E-02
GRMZM2G088501	Transporter-like protein	exon (3' UTR) and introns	chr6:133868502-133877776	-1.482	4.14E-02
GRMZM2G059637	Glycerol-3-phosphate acyltransferase 5	exon and intron	chr6:151462209-151465411	-1.495	9.68E-02
GRMZM2G012397	photosystem I reaction center6	none	chr7:5129273-5130176	-1.077	1.97E-02
GRMZM2G129246	Glycolate oxidase	exon (3'	chr7:5523491-5527292	-1.202	2.99E-03

		UTR)			
GRMZM2G046750	Lipid binding protein	exon and exonic 3'UTR	chr7:9201072-9202357	-1.412	5.30E-02
GRMZM2G177424	Flavonoid 7-O- methyltransferase	intron	chr7:44803178-44815872	-2.003	1.29E-04
GRMZM2G058310	Beta-amylase	none	chr7:155357370-155360570	-1.207	1.96E-02
GRMZM2G072280	Chlorophyll a-b binding protein; PSI antenna protein	exon and intron	chr7:160255313-160258286	-1.199	2.85E-03
GRMZM2G003930	Putative zinc finger, C3HC4 type (RING finger) domain containing protein	none	chr8:10287956-10291623	-0.930	9.78E-02
GRMZM2G097141	Hypothetical, unknown protein	exon (3' UTR)	chr8:38487905-38488708	-1.006	2.62E-02
GRMZM2G028570	Solute carrier family 2, facilitated glucose transporter member 8	exon and introns	chr8:71714905-71721778	-1.804	8.25E-03
GRMZM2G028134	Metal ion binding protein	none	chr8:141293857-141294830	-1.763	2.27E-02
GRMZM5G833406	IAA-amino acid hydrolase ILR1- like	exon and intron	chr8:154028733-154034508	-1.109	3.68E-02
GRMZM2G082376	Putative RING zinc-finger domain superfamily protein	exon	chr8:162182482-162183632	-1.811	1.96E-02
GRMZM2G034360	Omega-hydroxypalmitate O- feruloyl transferase-like	exons	chr8:166481614-166484307	-0.977	7.72E-02
GRMZM2G465046	GDSL esterase/lipase At1g74460-like	none	chr9:4308985-4310333	-1.485	9.68E-02
GRMZM2G083841	Phosphoenolpyruvate carboxylase 1 (PEPCase 1)	none	chr9:61296279-61301686	-1.713	2.37E-07
GRMZM2G039586	GATA transcription factor 20	none	chr9:80170947-80173134	-0.938	9.03E-02

GRMZM2G107886	Zinc finger protein CONSTANS-LIKE 16	none	chr9:106201175-106203143	-1.394	1.35E-04
GRMZM2G012140	Putative glycosyl hydrolase family 17 protein	exon and intron	chr9:126718179-126723107	-1.156	9.11E-03
GRMZM2G022686	Thioredoxin family protein	exon and 3' UTR	chr9:136175680-136180522	-1.518	7.34E-02
GRMZM2G181546	Hypothetical, unknown protein	none	chr10:2402129-2403240	-2.026	1.08E-04
GRMZM2G046284	Fructose-bisphosphate aldolase	none	chr10:10194502-10196450	-1.070	1.38E-02
GRMZM2G079490	Aspartic proteinase nepenthesin-1	exon	chr10:120671025-120672700	-1.763	6.95E-02
GRMZM2G031660	Beta-glucosidase	intron	chr10:130301051-130304967	-1.888	7.96E-05
GRMZM2G043813	NAC domain-containing protein 21/22	intron	chr10:130620219-130622989	-2.465	7.06E-03
GRMZM2G150248	Lysine-specific histone demethylase 1	intron	chr10:147593063-147597356	-1.196	2.73E-03
GRMZM2G092427	Chlorophyll a/b binding apoprotein CP24 precursor	exon	chr10:147890562-147891878	-1.237	2.66E-02

Decreased Transcription - Antisense Orientation^a

Gene ID	Annotation	TE / Repeat Content ^b	Genomic Position ^c	Log ₂ Fold Change (<i>rpdl</i> / WT) ^d	p-Value ^e
GRMZM2G061988	Putative pyridoxamine 5-phosphate oxidase	introns	chr1:140039425-140056837	-1.933	2.22E-02
GRMZM5G819500	Hypothetical, unknown protein	exon	chr1:148364112-148364951	-2.948	2.44E-02
GRMZM2G023033	Hypothetical, unknown protein	intron	chr2:33036267-33037791	-3.858	5.91E-02
AC197146.3_FG002	Putative Myb DNA-binding domain superfamily protein	none	chr10:2466640-2467603	-2.841	9.33E-02

^a Bold entries highlight sense oriented differential transcription that corresponds to the gene-model defined transcription unit as identified by visual inspection.

^b TE content within genes is defined as a positive hit in Repeatmasker (<http://www.girinst.org/censor/help.html#MAP-SIM>) with 0.7 similarity or greater to an annotated *Panicoideae* TE.

^c Genomic position is based on the B73 maize reference genome, version 2, build 5b.

^d Fold change is Log_2 (*rpd1* mutant / WT); for lines where no fold change is indicated, either WT or *rpd1* mutant counts were zero.

^e *p*-values have been adjusted to account for multiple testing with the Benjamini-Hochberg method (Benjamini and Hochberg, 1995) by the DESeq analysis (Anders and Huber, 2010).

Table S4 Repeatmasker analysis of 200 maize pre-mRNAs

Gene ID	TE / Repeat Content ^a	Chromosome	Start Position	End Position
GRMZM2G030567	yes	1	3737629	3742546
GRMZM2G703487		1	11358814	11359340
GRMZM2G015571	yes	1	19901211	19911946
GRMZM2G043470	yes	1	30297523	30302047
GRMZM2G145275	yes	1	41217232	41221207
GRMZM2G043976		1	41441330	41442027
GRMZM2G048821	yes	1	79235489	79241264
GRMZM2G031222		1	83540947	83541696
GRMZM2G071157	yes	1	88011829	88014372
GRMZM5G893936		1	91551111	91552314
AC201966.3_FG001		1	129729964	129730554
GRMZM2G158911		1	146822410	146823472
AC212878.4_FG001		1	172638114	172638272
GRMZM5G865967		1	172878324	172880857
GRMZM2G300788		1	176053059	176054692
GRMZM2G083725	yes	1	194657095	194659202
GRMZM2G017045		1	201625281	201626321
GRMZM2G043819	yes	1	211879253	211896231
GRMZM2G449133		1	227240671	227242553
GRMZM2G346455	yes	1	228245258	228256942
GRMZM2G131815	yes	1	230206756	230209825
GRMZM2G050914	yes	1	272870479	272874905
GRMZM2G033219	yes	2	4834198	4840631
GRMZM2G032977	yes	2	4847231	4850042
GRMZM5G876898	yes	2	8906080	8909067
GRMZM2G045398		2	15140829	15141833
GRMZM2G178681	yes	2	27871624	27892346
GRMZM2G094792	yes	2	42618631	42623189
GRMZM2G362803		2	54204566	54206256
GRMZM2G068590	yes	2	60874151	60875855
GRMZM2G005114	yes	2	78001561	78003312
GRMZM2G000959	yes	2	133189502	133191105
AC211347.4_FG006	yes	2	150091407	150092649
GRMZM5G804382		2	153140326	153142545
GRMZM2G083977	yes	2	161758262	161759154
GRMZM2G074567	yes	2	170296225	170299574

GRMZM2G414252	yes	2	184966846	184968211
GRMZM2G002131	yes	2	186183204	186187268
GRMZM2G329047	yes	2	189225428	189226150
GRMZM2G022921	yes	2	193408724	193413807
GRMZM2G049613	yes	2	197221948	197226964
GRMZM5G814007	yes	2	199822895	199825927
GRMZM2G001247	yes	2	216665547	216666787
GRMZM2G527387	yes	2	223510475	223522044
GRMZM2G099862	yes	2	225159420	225162831
GRMZM2G355760	yes	2	225449048	225451754
GRMZM2G135013		2	232876230	232879140
GRMZM5G848876		3	6589628	6591631
GRMZM2G101646		3	16031024	16032042
GRMZM2G463577		3	16570818	16572010
GRMZM2G112598	yes	3	29619773	29622817
GRMZM2G070264		3	32245135	32250561
GRMZM2G067511	yes	3	38428236	38432826
GRMZM2G440605	yes	3	38633283	38638691
GRMZM5G856092		3	81178553	81178945
GRMZM2G145886	yes	3	83717797	83740902
GRMZM2G339674		3	88185969	88187004
GRMZM2G439884		3	128878836	128881697
GRMZM2G170689	yes	3	152451136	152453723
GRMZM2G146697	yes	3	161846601	161856015
GRMZM2G014720	yes	3	182896768	182900200
GRMZM2G137965	yes	3	204071388	204078435
GRMZM2G371628	yes	3	220693706	220701719
GRMZM2G481362	yes	3	224937881	224941022
GRMZM2G002646		3	225083458	225086011
GRMZM5G803334		4	909819	910536
GRMZM2G106336		4	2649672	2651390
GRMZM2G398057	yes	4	4172535	4175902
GRMZM5G833660	yes	4	5260193	5262527
GRMZM2G007289	yes	4	19819570	19820771
GRMZM2G389944		4	23187939	23189854
GRMZM2G147726	yes	4	25786864	25797883
GRMZM2G702403		4	34832264	34833245
GRMZM2G105901	yes	4	42150834	42152527
GRMZM2G172365	yes	4	59997271	60007705

GRMZM2G131699	yes	4	88514893	88534830
GRMZM2G456756	yes	4	123603966	123604463
AC208559.3_FG003	yes	4	125108205	125110893
GRMZM2G416142	yes	4	127696347	127700700
GRMZM2G159691	yes	4	129532798	129535825
AC194897.4_FG003		4	129620391	129621465
GRMZM2G454581	yes	4	144648371	144651235
GRMZM2G319786	yes	4	151713456	151714591
GRMZM5G881641		4	161408409	161409593
GRMZM2G141036		4	163962237	163964018
GRMZM2G579820	yes	4	166308893	166309390
GRMZM2G025989	yes	4	183596885	183597900
GRMZM2G121905	yes	4	184351460	184378527
GRMZM2G076087	yes	4	186841262	186849745
GRMZM2G332495		4	198072097	198077474
AC191412.3_FG005	yes	4	198870615	198910724
AC202991.3_FG003	yes	4	199556534	199557670
GRMZM2G104419	yes	4	234720860	234727340
GRMZM2G069603	yes	4	239072144	239075476
GRMZM2G025173	yes	5	1439592	1440702
GRMZM2G112548		5	5326925	5328393
GRMZM2G150448		5	10897428	10900964
GRMZM2G079343		5	27594407	27597152
GRMZM5G835721		5	37702606	37703715
GRMZM2G000753	yes	5	42657556	42661577
GRMZM2G423518	yes	5	63156102	63158567
GRMZM2G102845	yes	5	78381834	78389884
GRMZM2G022181	yes	5	87035070	87036937
GRMZM2G396411	yes	5	127067801	127074652
GRMZM2G350284	yes	5	142136328	142137035
GRMZM2G169734	yes	5	147107709	147109875
GRMZM2G002240	yes	5	148226699	148228232
GRMZM2G115480	yes	5	168899560	168908436
GRMZM2G495613	yes	5	180003014	180005743
GRMZM2G170049	yes	5	188916348	188920991
GRMZM2G391487	yes	5	212684729	212687940
GRMZM2G058289	yes	5	215052090	215055981
AC208348.3_FG012	yes	5	216695478	216697052
GRMZM2G093035		6	14133929	14134987

GRMZM2G172980		6	14972850	14974387
GRMZM2G076868	yes	6	26668196	26671075
GRMZM5G839223	yes	6	71677776	71680373
GRMZM2G032537	yes	6	72810648	72815513
GRMZM2G113624	yes	6	78188618	78191058
GRMZM2G306028	yes	6	86113653	86118400
GRMZM2G154845	yes	6	86175191	86181011
GRMZM2G168449	yes	6	86543469	86552114
GRMZM5G862355	yes	6	97113476	97115733
GRMZM2G331283		6	104963322	104965389
GRMZM2G343636	yes	6	106906026	106912138
GRMZM2G167733		6	108857961	108860755
AC186647.3_FG001		6	110030367	110030822
GRMZM2G137593	yes	6	111665045	111676475
GRMZM2G319537		6	113280504	113280779
AC206185.3_FG002	yes	6	116345195	116346376
GRMZM2G063754		6	123748479	123751779
GRMZM2G177026	yes	6	125268897	125272586
GRMZM2G002656	yes	6	128823409	128833681
GRMZM2G301096	yes	6	136648745	136651477
GRMZM2G468535	yes	6	153252044	153255958
GRMZM2G032699	yes	7	35007319	35011752
GRMZM2G164862		7	59685157	59687044
GRMZM2G479906	yes	7	92078951	92089092
GRMZM2G036916	yes	7	103506101	103511006
GRMZM2G055809		7	105062000	105063230
GRMZM2G077655		7	133563863	133566646
GRMZM2G091586	yes	7	143515014	143521045
GRMZM2G436511	yes	7	145724785	145728515
GRMZM2G077140		7	146973952	146974990
GRMZM2G322361		7	147005038	147007540
GRMZM2G700262	yes	7	150654899	150655528
GRMZM2G161611	yes	7	150659401	150662789
GRMZM2G105982	yes	7	172366098	172367449
GRMZM2G333087	yes	7	176525487	176535559
GRMZM2G095366		8	24868167	24869609
GRMZM2G451464	yes	8	29828392	29829280
GRMZM2G086897		8	34485084	34486021
GRMZM2G164129		8	73479234	73479957

GRMZM2G375504	yes	8	95715355	95726032
GRMZM2G456564	yes	8	99417809	99422005
AC196082.3_FG001		8	99954430	99955248
GRMZM2G356162		8	100479139	100479599
GRMZM2G404237	yes	8	108495927	108500185
GRMZM2G090563	yes	8	117149558	117151380
AC197705.4_FG011	yes	8	118078273	118090836
GRMZM2G700655	yes	8	128542778	128548144
GRMZM2G002427	yes	8	129414617	129417699
GRMZM2G088196	yes	8	136659605	136668296
GRMZM2G064336	yes	8	148514901	148519447
GRMZM2G060541		8	149891773	149893456
GRMZM2G700730		8	156336396	156336629
GRMZM2G438704		8	156469002	156471588
GRMZM2G109472	yes	8	158113092	158123850
GRMZM2G068392	yes	8	163509234	163516799
GRMZM2G042218		8	166580075	166581471
GRMZM2G130119		8	175537743	175539495
GRMZM5G820423	yes	9	4881573	4882112
GRMZM2G142057		9	8956846	8957884
GRMZM2G703370		9	11440061	11440240
GRMZM2G023636	yes	9	14563084	14585427
GRMZM2G027958	yes	9	37538687	37540567
GRMZM2G019859		9	40772038	40773556
GRMZM2G024221	yes	9	43231158	43232834
GRMZM5G833635		9	72844239	72844562
GRMZM2G161472	yes	9	78621553	78625626
GRMZM2G111711	yes	9	103075648	103077219
GRMZM2G028254	yes	9	107315294	107317421
AC216730.2_FG005	yes	9	112209899	112210708
GRMZM2G013671	yes	9	116268809	116270671
GRMZM2G046729	yes	9	146083613	146087184
GRMZM2G028432		10	9537710	9539765
GRMZM2G001195		10	12546764	12548674
GRMZM2G012879	yes	10	16959440	16961424
GRMZM2G140737	yes	10	17832709	17837521
AC188722.3_FG002		10	34730619	34730912
GRMZM2G360615	yes	10	39949992	39964688
AC188036.3_FG003		10	82151668	82152585

GRMZM2G029609	yes	10	107805061	107806206
GRMZM2G103725		10	110745904	110751262
GRMZM2G135378		10	115741757	115744192
GRMZM5G887803	yes	10	118989952	118990398
GRMZM2G132875		10	120494047	120495742
GRMZM2G169642	yes	10	135493446	135495893
GRMZM2G086136		10	145174954	145177446
GRMZM2G406074	yes	10	148849233	148855901

^aTE content within pre-mRNAs is defined as a positive hit in Repeatmasker (<http://www.girinst.org/censor/help.html#MAP-SIM>) with 0.7 similarity or greater to an annotated *Panicoideae* TE. 131 of these 200 genes have TE content by this definition.

Table S5 TEs differentially transcribed in *rpdl* mutants

Increased Transcription – Sense Orientation ^a			
TE ID	Genomic Position ^b	Log ₂ Fold Change (<i>rpdl</i> / WT) ^c	p-Value ^d
RLG_prem1_AC201801_7095	chr1:168882596-168883075	5.129	3.66E-02
ZM_hAT_noncoding_120	chr1:196249905-196250023	5.392	8.19E-03
RLX_uwum_AC177933_415	chr2:107508182-107508735	3.395	8.78E-08
RLG_yfages_AC197085_5383	chr3:36221587-36223382	2.896	6.81E-06
RLC_ufonah_AC194064_3852	chr3:36223389-36224416	2.807	1.42E-02
RLG_fege_AC205532_8884	chr3:36224420-36224976	3.459	3.03E-05
RLC_ufonah_AC194064_3852	chr3:36224716-36225104	3.301	1.19E-02
RLX_milt_AC211742_11402	chr3:122962737-122968593	2.739	3.03E-05
RLC_machiavelli_AC200490_6883	chr3:160664585-160664712	2.842	3.78E-05
RLC_machiavelli_AC200490_6883	chr3:160664720-160664907	2.911	4.38E-05
RLX_fageri_AC204875_8470	chr5:70052242-70052969	1.767	8.27E-02
ZM_hAT_8	chr5:84105173-84105372	3.433	3.68E-02
DTM_Zm00460_consensus	chr5:98657883-98658114	2.317	3.39E-04
RLX_name_AC197689_5725	chr7:10972520-10972677	---	4.97E-04
RLG_neha_AC215285_13046	chr8:143766280-143766943	4.954	2.79E-04
RLX_naseup_AC196428_5108	chr9:69638177-69640432	2.984	1.08E-02
RLX_milt_AC209648_10275	chr9:148227395-148228110	1.905	1.41E-02
RLX_baso_AC192251_3423	chr10:23192733-23192950	2.154	6.11E-02
RLC_ji_AC193479_3665	chr10:23192806-23193005	2.191	4.97E-02
RLC_gudyeg_AC206942_9404	chr10:105781195-105781407	5.322	1.09E-02
RLC_gudyeg_AC206942_9404	chr10:105781195-105782219	5.322	1.09E-02

RLC_gudyeg_AC206942_9404	chr10:105781357-105784535	5.322	1.09E-02
RIX_ogepy_AC198740-0	chr10:142977228-142979196	4.492	1.25E-02

Increased Transcription – Antisense Orientation^a

TE ID	Genomic Position ^b	Log ₂ Fold Change (<i>rpdl</i> / WT) ^c	p-Value ^d
RLX_vora_AC206187_9112	chr1:92817640-92817751	6.755	6.12E-09
RLG_flip_AC208040_9765	chr1:118536328-118537641	---	1.14E-03
RLC_giepum_AC211155_11010	chr1:180819578-180819782	3.248	3.17E-02
RLG_xilon-diguus_AC195486_4629	chr2:158762498-158763219	4.138	9.23E-05
RLG_xilon-diguus_AC195486_4629	chr2:158762921-158764622	3.524	5.06E-02
ZM_CACTA_noncoding_2	chr3:7523757-7524489	2.081	1.67E-03
ZM_CACTA_noncoding_2	chr3:7523878-7524489	2.081	1.67E-03
ZM_CACTA_noncoding_1	chr3:7523973-7524489	2.081	1.67E-03
ZM_CACTA_noncoding_7	chr3:7524044-7524580	2.075	1.68E-03
ZM_hAT_noncoding_12	chr3:36220145-36220665	3.019	2.25E-04
RLG_nihep_AC194441_4115	chr3:65920084-65921482	5.322	6.98E-03
RLC_ji_AC190978_2799	chr3:65921484-65921523	---	6.98E-03
RLC_ji_AC215728_13156	chr3:65921492-65922891	4.209	5.06E-02
ZM_CACTA_34	chr3:91408073-91408137	---	1.22E-04
RLX_hani_AC186285_1359	chr3:157967848-157968598	2.443	7.82E-02
RLX_avahi_AC191363_3084	chr4:63005326-63007571	3.113	3.98E-05
ZM_CACTA_89	chr5:123935141-123936168	---	2.31E-02
ZM_CACTA_noncoding_2	chr5:127956983-127957675	2.198	3.02E-03
ZM_CACTA_noncoding_25	chr6:12704019-12704523	2.183	2.66E-02
ZM_CACTA_noncoding_2	chr7:17711484-17712080	3.379	3.79E-02

RLX_vegu_AC190718_85	chr7:115190804-115191388	---	3.02E-03
ZM_CACTA_16	chr7:133871697-133872651	2.000	9.77E-02
RLX_ewib_AC207533_9599	chr8:138863636-138864837	2.550	9.48E-03
RLG_dagaf_AC208646_9966	chr9:69644374-69648016	4.524	2.25E-04
ZM_CACTA_noncoding_2	chr9:79123736-79124344	2.126	1.45E-02
ZM_CACTA_noncoding_10	chr9:115947886-115948286	2.553	4.47E-02
ZM_CACTA_noncoding_2	chr9:115948005-115948494	2.709	2.66E-02
ZM_CACTA_noncoding_7	chr9:115948057-115948584	2.603	5.55E-02
ZM_CACTA_noncoding_1	chr9:151268135-151268651	2.441	2.99E-03
ZM_CACTA_noncoding_7	chr9:151268206-151268741	2.482	2.35E-03
ZM_CACTA_16	chr10:66170238-66171076	4.392	7.51E-05
Decreased Transcription – Sense Orientation^a			
TE ID	Genomic Position ^b	Log ₂ Fold Change (<i>rp1</i> / WT) ^c	p-Value ^d
RLG_flip_AC203163_7675	chr2:64555076-64557648	-5.170	2.76E-02
ZM_Stowaway_30	chr2:102138521-102138680	-1.983	8.69E-02
RLX_yraj_AC205486_8834	chr10:132170975-132171029	---	2.45E-02
Decreased Transcription – Antisense Orientation^a			
TE ID	Genomic Position ^b	Log ₂ Fold Change (<i>rp1</i> / WT) ^c	p-Value ^d
RLX_nuhan_AC206272_9161	chr1:32985086-32985358	-3.970	1.67E-02
RLX_votaed_AC215881_13209	chr1:117129922-117132946	-2.728	7.36E-03
RLX_ubat_AC212211_11652	chr1:117135865-117137824	-3.087	8.15E-04
RLC_labe_AC203760_7937	chr1:140040803-140042013	-3.841	3.91E-02
RLX_votaed_AC215881_13209	chr1:140040939-140041818	-5.087	3.17E-02

RLX_buire_AC194484_4158	chr5:30166464-30166598	-4.129	8.72E-02
--------------------------------	-------------------------------	---------------	-----------------

^a Bold entries are TEs not located within genes or 5 kb of either side (see *Materials and Methods*).

^b Genomic position is based on the B73 maize reference genome, version 2, build 5b.

^c Fold change is Log_2 (*rpd1* mutant / WT); for lines where no fold change is indicated, either WT or *rpd1* mutant counts were zero.