Published in final edited form as:

Addiction. 2009 April ; 104(4): 669–675. doi:10.1111/j.1360-0443.2009.02508.x.

How reaction to cigarette packet health warnings influence quitting: Findings from the ITC Four Country survey

Ron Borland¹, Hua-Hie Yong¹, Nick Wilson², Geoffrey T. Fong^{3,4}, David Hammond³, K. Michael Cummings⁵, Warwick Hosking⁶, and Ann McNeill⁷

¹The Cancer Council Victoria, Melbourne, Australia ²Otago University, Wellington, New Zealand

³University of Waterloo, Waterloo, Canada ⁴Ontario Institute for Cancer Research, Toronto, Canada ⁵Roswell Park Cancer Institute, Buffalo, USA ⁶Victoria University, Melbourne, Australia

⁷University of Nottingham, Nottingham, UK

Abstract

Objectives—To examine prospectively the impact of health warnings on quitting activity.

Design—Five waves (2002–2006) of a cohort survey where reactions to health warnings at one survey wave are used to predict cessation activity at the next wave, controlling for country (proxy for warning differences) and other factors. These analyses were replicated on four wave-to-wave transitions.

Setting and participants—Smokers from Australia, Canada, the UK and USA. Samples were Waves 1–2: n=6525; Waves 2–3: n=5257; Waves 3–4: n=4439; and Waves 4–5: n=3993.

Measures—Warning salience, cognitive responses (thoughts of harm and of quitting), forgoing of cigarettes and avoidance of warnings were examined as predictors of quit attempts, and of quitting success among those who tried (1 month sustained abstinence), replicated across four wave-to-wave transitions.

Results—All four responses to warnings were independently predictive of quitting activity in bivariate analyses. In multivariate analyses, forgoing cigarettes and cognitive responses to the warnings both prospectively predicted making quit attempts in all replications. However, avoiding warnings did not consistently add predictive value, and there was no consistent pattern for warning salience. There were no interactions by country. Some, but not all, of the effects were mediated by quitting intentions. There were no consistent effects on quit success.

Conclusions—This study adds to the evidence that forgoing cigarettes as a result of noticing warnings and quit-related cognitive reactions to warnings are consistent prospective predictors of making quit attempts. This work strengthens the evidence base for governments to go beyond the

Please send all correspondence to: Ron Borland, PhD, Nigel Gray Distinguished Fellow in Cancer Prevention, VicHealth Centre for Tobacco Control, The Cancer Council Victoria, 1 Rathdowne Street, Carlton, VIC 3053, Australia. Ron.Borland@cancervic.org.au, Ph 61-3-9635 5185.

Ethics clearance: The study protocol was approved by the institutional review boards or research ethics boards of the University of Waterloo (Canada), Roswell Park Cancer Institute (United States), University of Strathclyde (UK), University of Stirling (UK), The Open University (UK), and The Cancer Council Victoria (Australia).

Competing interests: The authors declare that they have no competing interests.

FCTC to mandate health warnings on tobacco products that stimulate the highest possible levels of these reactions.

Keywords

Health warnings; graphic warnings; tobacco product labelling; consumer information; tobacco health information

Introduction

There is an increasing body of evidence that larger and graphic health warnings on cigarette packs are noticed more often and increase knowledge about health effects of smoking better than relatively small text-only warnings as found in many countries around the world (e.g., China, United States, Malaysia) [1–3]. The impact of enhanced pack warnings on altering smoking behaviour are less well studied. One study of the transition to larger text-based warnings in Australia in 1995 found that that forgoing cigarettes as a result of noticing the health warnings was a prospective predictor of subsequent quitting activity, but that reported frequency of noticing the warnings was not [4]. This finding shows that immediate reactions to warnings can predict longer term effects. Another study, using a composite measure of depth of cognitive processing of warnings, found this to be a prospective predictor of quit attempts [5]. However, there have been no other predictive studies.

Evaluating the effects of interventions, such as health warnings, that are likely to have only small effects on outcomes such as quitting is difficult, especially when there is limited capacity to control or model other potential determinants of the outcomes [6]. It is unlikely that we could directly demonstrate by-country differences in quitting activity due to health warnings, so we have adopted a less direct approach [6]. We test a mediational model by exploring the relationship between reactions to warnings and subsequent quitting activity. We have already demonstrated that new, stronger warnings led to increased reactions in both the UK [7] and Australia [1], so the next step is to show whether these reactions prospectively predict quitting.

Reactions to warnings could also be associated with reductions in desired outcomes as well as facilitating them. Some researchers have argued that strong warnings and various public health campaigns may inadvertently create psychological reactance that could potentially inhibit desirable behaviour change [8–9], including quitting activity [10]. The possibility of a negative effect of reactance was considered by Hammond et al [3], who found no evidence that avoidance of health warnings reducing quitting [3]. In other areas of health, reactance effects have been found for immediate intentions to change behaviour but not longer term behaviour. For example, emotionally charged messages to promote sun protection behaviour [11] inhibited intentions shortly after being read, but were predictive of increased reported behaviour change at follow-up some months later. Similarly, Jessop & Wade [12] found evidence of greater willingness to binge drink among some study participants in response to a binge drinking and mortality message but no increase in binge drinking over the following week. Therefore, the evidence for reactive effects of interventions inhibiting the behaviour the intervention is designed to motivate is weak and may be short term and/or restricted to

intentions rather than behaviour. However, while it remains a possibility, it is important to have population-based research to rule out reactance effects to cigarette packet health warnings, especially since the enhancement of pack warnings is a key provision of the World Health Organization's Framework Convention on Tobacco Control [13].

In this paper we use data from five survey waves of the International Tobacco Control Policy Research Survey (ITC) study to examine how responses to enhanced pack warnings impact subsequent smoking cessation. The ITC 4-Country study began in 2002 and includes annual surveys of cohorts of smokers in Australia, Canada, UK and USA. The ITC study provides a vehicle for studying the impact of varying forms of health warnings and of changes in warnings within countries. Warnings range from large graphic Canadian warnings to small text-based US warnings that are on the side of the pack rather than on the main faces as is the case in the other three countries. The UK increased the size and contrast of its warnings in early 2003 between waves 1 and 2 of the survey and Australia moved from moderately large text-based warnings to larger graphic warnings in 2006, between waves 4 and 5.

Methods

Sample

Details of the samples used are in Table 1. As detailed in this Table, some respondents contributed data to multiple pairs.

Design

Prospective prediction of reactions to health warnings on quit smoking attempts, and among those who tried, quit success (defined as staying quit for at least 1 month), with outcomes measured at the next year's wave of the survey.

A full description of the ITC Project conceptual framework and methods have been published elsewhere [14–15]. Further details surrounding the ITC Project methods relating to questions on health warnings have also been recently been published [7]. Critical to this paper is that cohorts are followed annually in four countries, USA, UK, Canada and Australia, with similar sets of questions asked each year.

Measures

Salience of health warnings was assessed by combining two questions (alpha = 0.67 – 0.81, waves 1–5): how often over the preceding month respondents had noticed the warnings, and read or looked closely at them, both on 5-point scales: never to very often. Cognitive responses were assessed by combining three items (alpha= 0.78 – 0.80, waves 2–5): the extent in the last month to which the warnings both made the respondent think about the health risks of smoking, and made them more likely to quit smoking (4-point scales: Not at all to A lot), both only asked from wave 2; and in a separate part of the survey, the extent to which “warning labels on cigarette packages” motivated them to think about quitting in the last six months, with response options: Not at all, Somewhat, and Very much. There were two different types of behavioural response: a single item measure of frequency (if ever) of

forgoing cigarettes as a result of the warnings (coded ever vs never): and questions about four kinds of avoiding warnings (cover-up, keep out of sight, use cigarette case, or avoid particular labels) from which a binary variable, no avoidance vs. any avoidance, was computed.

Potential moderator variables assessed were the demographic variables: country of origin, sex, and age (18–24, 25–39, 40–54, 55+ years). Household income and highest level of education attained were assessed where these were equated across the four countries and coded into low, medium and high. In addition we controlled for reported cigarettes per day (using a square root transform to improve normality), and intention to quit (assessed on a 4-point scale: planning in the next month, considering in next 6 months, considering beyond 6 months, and no interest).

Outcomes, assessed at the next wave of the survey, were reporting having made a quit attempt of at least 24 hours since the previous wave, and among those, whether the quit attempt lasted for at least one month as an indicator of short-term success. Smokers quit for less than 1 month at follow-up were excluded from these analyses.

Analysis

Logistic regression analyses were used to predict making quit attempts between waves and, separately, among those who tried, success of attempts. We conducted four wave-to-wave replications of the associations, although only three replications were possible for the cognitive reactions, which were not assessed at Wave 1. Analyses controlled for demographic factors, cigarettes per day, and country. We also checked for interactions with country, particularly looking at the UK for the Wave1–2 analyses and Australia for the Wave 4–5 analyses, as in these two cases, new stronger warnings were introduced between the surveys, so some of the quitting may have been stimulated by this, and thus the prior wave reactions might not fully reflect subsequent impact. As no interactions with country were found for the focal cases or any others, these interaction terms were removed from the final models reported. In secondary analyses, we tested to see whether the effects on quit attempts were mediated by intention at the wave in which the reactions were reported.

Results

Table 2 presents the range of intercorrelations between the four measures of reactions to pack warnings for the waves in which they were collected. All of these were positive and the pattern and strength of associations observed were similar for all survey waves.

Table 3 presents data on percent making quit attempts between waves and percent still quit among those who did. The varying inter-survey intervals is responsible for some of the wave to wave variability. While there was relatively more quitting activity in Australia in the wave where the warning labels was strengthened, there is no such pattern for the UK, or any association for short-term success.

There were significant positive bivariate associations between all four of the warning label reactions and making a quit attempt by the next survey wave (see Table 4), but no consistent pattern of associations for predicting short-term success.

Table 5 presents the summary results for independent prediction of making quit attempts for the reactions for each of the four wave transitions studied (i.e., independent of other reactions). These analyses also controlled for demographic factors and cigarette consumption (Model A), and then additionally controlled for intentions to quit (Model B). It can be seen that both cognitive responses and forgoing cigarettes were consistent predictors of making a quit attempt across all wave-to-wave transitions where they were measured. Both remained significant predictors after controlling for quit intentions, although the size of the effects were reduced, especially for cognitive responses, suggesting partial mediation. Neither warning salience, nor warning avoidance had a consistent independent relationship. Salience had a significant positive relationship in one wave transition, but a significant negative one in another, with nothing significant after controlling for intentions. Avoidance of the product warning was a significant independent predictor for the Wave 1–Wave 2 transition, but was not significant in the other wave transitions, although in all cases the odds ratio was positive.

Table 6 shows the multivariate relationships for the four types of reactions and the success of quit attempts among those who tried. There were no consistent effects.

Discussion

This study demonstrates that some reactions to warning labels have consistent and independent predictive power for the making of subsequent quit attempts, but this does not appear to extend to predicting success of those attempts. This finding for a predictive effect is consistent with most of the previous literature and, at least in the case of making quit attempts, is consistent with theorised effects. The stronger the warnings, the greater they stimulate cognitive and behavioural reactions [1, 7], and this is translated into the stimulation of quit attempts in some cases. The finding that the effects may be partly independent of intentions, suggests that reacting to warnings has the potential to trigger spontaneous quit attempts, something that is quite common [16], as well as to enhance intentions.

We have generally ignored the isolated significant effects, especially when the results across different survey waves were not consistently in the same direction. This means we make no claims for any clear effects on smoking cessation outcomes; ie, preventing relapse. It should come as no surprise that reactions to health warnings when smoking do not predict outcomes once quit, as the ex-smoker no-longer will have cigarette packs with them, so their potential effects on the success of quit attempts are likely to be diluted. Further, there is no reason to believe that reactions to packs while smoking would be strongly enough associated with reactions after quitting for effects to persist.

In this paper we found no consistent effect of warning avoidance on subsequent behaviour, but the trends were for a positive effect, and the bivariate effects were always positive and

significant. This is further evidence against the position that cigarette packet warning avoidance is an indicator of psychological reactance that might inhibit desired behaviour change (e.g., quitting), as postulated by some theorists [10], indeed any effects, albeit indirect, are positive.

We found no evidence of interactions with country in the analyses. This was generally expected as the by-country differences in strength of warnings are directly measured by the strengths of reactions to them. The two instances where this might not be so were where new stronger warnings were introduced between survey waves (Waves 1–2 for the UK, and Waves 4–5 for Australia). We are unclear of what implications there are for failure to find interactions in these cases. Overall, the findings are consistent with the reactions (most notably cognitive responses and forgoing cigarettes) having a causal impact on subsequent quit attempts, however, it remains possible that some unmeasured third variable is responsible for both. The only possibility we can think of is something like overall motivation to quit. However, we had a measure of quit intentions and at least part of the effects were independent of it, so it seems unlikely that motivation can explain the findings, although it is possible that a more comprehensive measure of it might. Further, if motivation was the unmeasured third party, we would expect interactions with country (as a proxy for warning strength) as the variance in warnings should weaken the relationship between the reactions to the warnings and the unmeasured motivation. In the absence of a plausible alternative mechanism for the findings we conclude that a causal role for the reactions to the warnings is likely.

Regardless of their causal role, it is clear that the measures of cognitive responses and forgoing cigarettes are useful indicators of the potential effectiveness of pack health warnings. These measures can now be used as intermediate indicators of warning label effectiveness, which should be useful in pre-testing of potential new warning content and form.

Replicating the associations several times is a major strength of this study as it provides an indication of the stability of the associations. Further, the specificity of the effects demonstrates that the predictive relationships are as expected, stronger for the reactions that are conceptually closest to making quit attempts, such as being triggered to think about quitting or to forgo smoking a cigarette.

Implications for policy makers

These data provide the strongest evidence to date that health warnings stimulate reactions that are predictable antecedents of quitting attempts. While the effects are partly mediated through intentions to quit, there is clear evidence that some of the effect is independent of intentions (at least as measured in this study). Based on the findings from this study, and the absence of any credible alternative hypothesis to explain the findings, we conclude that the stronger the warnings, the greater the reactions, and thus the greater the quitting activity they evoke. That said, the total impact on smoking prevalence is too small to accurately estimate. Health policy makers should be seeking to implement strong, large graphic warnings to maximise their effectiveness in achieving their goal of warning people off continuing to

smoke. This strengthens the case for the strongest possible implementation of the Framework Convention on Tobacco Control (FCTC) requirements on health warnings.

Given the growing international interest in improving health warnings on tobacco products [17] there is also a need for more research on the optimal design of such warnings. In particular, there is a need to strengthen the evidence base in non-English speaking countries, in countries with low literacy levels, and for the impact of warnings on youth experimenters and new smokers. The measures validated here as indicators of subsequent cessation could be used by developers of warnings to test their likely impact.

Acknowledgements

This research was funded by grants from the National Cancer Institute of the United States (R01 CA 100362), the Roswell Park Transdisciplinary Tobacco Use Research Center (P50 CA111236), Robert Wood Johnson Foundation (045734), Canadian Institutes of Health Research (57897 and 79551), National Health and Medical Research Council of Australia (265903 and 450110), Cancer Research UK (C312/A3726), and Canadian Tobacco Control Research Initiative (014578), with additional support from the Centre for Behavioural Research and Program Evaluation, National Cancer Institute of Canada/ Canadian Cancer Society. None of the sponsors played any direct role in the design and conduct of the study; the collection, management, analysis, and interpretation of the data; or the preparation, review, and approval of the manuscript. RB had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. NW, HHY, WH assisted in statistical analyses and the other authors all helped to design the study and write the manuscript.

References

1. Borland R, Wilson N, Fong GT, Hammond D, Cummings KM, Yong HH, Hosking W, Hastings G, McNeill A. Impact of graphic and text warnings on cigarette packs: Findings from four countries over five years. (in review).
2. Hammond D, Fong GT, McNeill A, Borland R, Cummings KM. Effectiveness of cigarette warning labels in informing smokers about the risks of smoking: findings from the International Tobacco Control (ITC) Four Country Survey. *Tob Control*. 2006; 15(Suppl 3):iii19–iii25. [PubMed: 16754942]
3. Hammond D, Fong GT, McDonald PW, Brown KS, Cameron R. Graphic Canadian cigarette warning labels and adverse outcomes: evidence from Canadian smokers. *Am J Public Health*. 2004; 94:1442–1445. [PubMed: 15284057]
4. Borland R. Tobacco health warnings and smoking-related cognitions and behaviours. *Addiction*. 1997; 92:1427–1435. [PubMed: 9519486]
5. Hammond D, Fong GT, McDonald PW, Cameron R, Brown KS. Impact of the graphic Canadian warning labels on adult smoking behaviour. *Tob Control*. 2003; 12:391–395. [PubMed: 14660774]
6. IARC. Methods for Evaluating Tobacco Control Policies. Lyon, France: International Agency for Research on Cancer; IARC Handbooks of Cancer Prevention: Tobacco Control. Volume 12. (in press).
7. Hammond D, Fong GT, Borland R, Cummings KM, McNeill A, Driezen P. Text and graphic warnings on cigarette packages: findings from the international tobacco control four country study. *Am J Prev Med*. 2007; 32:202–209. [PubMed: 17296472]
8. Ruiter R, Abraham C, Kok G. Scary warnings and rational precautions: A review of the psychology of fear appeals. *Psychol Health*. 2001; 16:613.7.
9. Campo S, Cameron KA. Differential effects of exposure to social norms campaigns: a cause for concern. *Health Commun*. 2006; 19:209–219. [PubMed: 16719724]
10. Ruiter R, Kok G. Saying is not (always) doing: cigarette warning labels are useless [letter]. *Eur J Public Health*. 2005; 15:329.
11. Buller DB, Borland R, Burgoon M. Impact of behavioural intention on effectiveness of message features: Evidence from the family sun safety project. *Human Communication Research*. 1998; 24(3):433–453. 16.

12. Jessop DC, Wade J. Fear appeals and binge drinking: A terror management theory perspective. *Br J Health Psychol.* 2008 (E-publication, 21 January).
13. World Health Organization. Final text of WHO Framework Convention on Tobacco Control. Geneva: World Health Organization; 2003. <http://www.who.int/tobacco/en/>.
14. Fong GT, Cummings KM, Borland R, Hastings G, Hyland A, Giovino G, Hammond D, Thompson ME. The conceptual framework of the International Tobacco Control (ITC) Policy Evaluation Project. *Tob Control.* 2006; 15(Suppl 3):iii3–iii11. [PubMed: 16754944]
15. Thompson ME, Fong GT, Hammond D, Boudreau C, Driezen P, Hyland A, Borland R, Cummings KM, Hastings G, Siahpush M, Mackintosh A, Laux F. The methodology of the International Tobacco Control Policy Evaluation (Four-Country) Survey (ITCPES). *Tob Control.* 2006; 15(suppl III):iii12–iii18. [PubMed: 16754941]
16. West R, Sohal T. "Catastrophic" pathways to smoking cessation: findings from national survey. *BMJ.* 2006; 332:458–460. [PubMed: 16443610]
17. Cunningham, R. Package Warnings: Overview of International Developments. Toronto: Canadian Cancer Society; 2007. http://www.smoke-free.ca/warnings/WarningsResearch/Release_WarningLabels_20070320.pdf.

Table 1

Respondent characteristics for the four countries and four wave-to-wave transitions

	Wave 1 to Wave 2 N = 6525	Wave 2 to Wave 3 N = 5257	Wave 3 to Wave 4 N = 4439	Wave 4 to Wave 5 N = 3993
Age (Mean, SD)	42.7 (14.2)	44.1 (14.0)	44.5 (13.5)	45.3 (13.6)
Cigarettes per day (Mean, SD)	17.1 (10.3)	16.9 (10.0)	16.5 (9.5)	16.9 (9.7)
Female %	55.8	56.2	57.9	58.1
Wave of recruitment				
Wave 1 recruits %	100.0	79.6	56.8	43.6
Wave 2 recruits %	–	20.4	12.4	9.4
Wave 3 recruits %	–	–	30.8	20.3
Wave 4 recruits %	–	–	–	26.7
Country				
Canada %	25.0	26.7	26.3	26.8
USA %	19.9	21.0	18.2	20.3
UK %	27.3	25.3	28.2	27.0
Australia %	27.8	27.0	27.3	25.8

SD – Standard deviation

Table 2

Ranges of inter-correlations of the four measures of reactions to warning labels across waves

	1.	2.	3.
1. Salience:	–		
2. Cognitive reactions:	.48 – .49	–	
3. Forgoing:	.27 – .30	.43 – .46	–
4. Avoidance:	.15 – .19	.24 – .29	.19 – .24

Notes: All correlations are significant at the 0.01 level (two – tailed)

Table 3
Percentages making quit attempts, between waves and of those the percentage still quit at the time of the survey

Period	Inter-survey interval (days)	Outcome	Canada %	USA %	UK %	Australia %	Average %
Wave 1 – 2	203	% Quit Attempt	43.9	36.5	31.4	34.3	36.3
		% Still Quit	45.2	39.5	53.6	43.7	45.5
Wave 2 – 3	388	% Quit Attempt	43.4	39.0	36.0	43.6	40.7
		% Still Quit	55.2	49.3	63.0	52.6	55.2
Wave 3 – 4	458	% Quit Attempt	39.8	39.9	41.5	42.7	41.1
		% Still Quit	57.3	50.9	69.7	58.9	59.7
Wave 4 – 5	361	% Quit Attempt	37.5	36.1	35.2	43.9	38.3
		% Still Quit	56.9	56.4	68.0	59.0	60.0

Table 4

Bivariate associations between mean levels of reactions to warnings and making quit attempts by the next survey wave.

		Warning Salience		Cognitive Reactions		Forgoing Warnings		Warning Avoidance	
		Quit Attempts		Quit Attempts		Quit Attempts		Quit Attempts	
		No	Yes	No	Yes	No	Yes	No	Yes
Wave 1 – 2	M	2.73	2.97**	—	—	0.09	0.18**	0.14	0.22**
	SE	0.19	0.03	—	—	0.00	0.01	0.01	0.01
Wave 2 – 3	M	2.95	3.07**	1.70	1.98**	0.09	0.18**	0.15	0.21**
	SE	0.02	0.03	0.01	0.02	0.01	0.01	0.01	0.01
Wave 3 – 4	M	2.83	3.00**	1.69	1.93**	0.09	0.16**	0.15	0.19*
	SE	0.02	0.03	0.01	0.02	0.00	0.01	0.01	0.01
Wave 4 – 5	M	2.81	2.96**	1.68	1.93**	0.08	0.16**	0.12	0.15*
	SE	0.02	0.03	0.01	0.02	0.01	0.01	0.01	0.01

Note: M, mean, SE, standard error,

* p < .01

** p < .001

Table 5

Predictors of having reported quit attempts at surveys in waves 2 to 5

<i>Unweighted</i>	Wave 2 OR (95% CI)	Wave 3 OR (95% CI)	Wave 4 OR (95% CI)	Wave 5 OR (95% CI)
Model A				
Salience	1.07 (1.02 – 1.12) **	0.94 (0.89 – 0.99) *	0.96 (0.91 – 1.02)	0.96 (0.91 – 1.02)
Cognitive reactions	#	1.57 (1.43 – 1.72) ***	1.44 (1.31 – 1.58) ***	1.46 (1.32 – 1.61) ***
Forgoing	1.69 (1.43 – 1.99) ***	1.30 (1.08 – 1.57) **	1.43 (1.18 – 1.74) ***	1.40 (1.14 – 1.72) **
Avoidance	1.32 (1.14 – 1.52) ***	1.15 (0.98 – 1.34)	1.04 (0.89 – 1.22)	1.04 (0.86 – 1.24)
Model B (adjusted for Quit intention)				
Salience	1.02 (0.98 – 1.07)	0.96 (0.90 – 1.02)	0.97 (0.91 – 1.02)	0.97 (0.91 – 1.03)
Cognitive reactions	#	1.28 (1.16 – 1.41) ***	1.14 (1.03 – 1.26) *	1.16 (1.04 – 1.29) **
Forgoing	1.38 (1.16 – 1.65) ***	1.23 (1.01 – 1.49) *	1.42 (1.16 – 1.74) **	1.30 (1.05 – 1.62) *
Avoidance	1.17 (1.01 – 1.37) *	1.09 (0.93 – 1.29)	1.03 (0.87 – 1.22)	1.03 (0.85 – 1.25)

Notes: OR, odds ratio; CI, confidence interval; All odds ratios were adjusted for age, sex, income, education, country and daily cigarette consumption;

*
p<0.05;

**
p<0.01;

p<0.001;

Relevant questions not asked in the Wave 1 survey.

Table 6

Quit success (for at least 1 month) among smokers who had made attempts for 4 wave-to-wave transitions

	Wave 1–2 OR (95% CI)	Wave 2–3 OR (95% CI)	Wave3– 4 OR (95% CI)	Wave 4–5 OR (95% CI)
<i>Unweighted</i>				
Salience	0.94 (0.87 – 1.02)	0.94 (0.86 – 1.04)	1.01 (0.92 – 1.10)	1.00 (0.90 – 1.12)
Cognitive reactions	#	1.02 (0.87 – 1.19)	1.17 (1.00 – 1.37)*	1.02 (0.86 – 1.22)
Forgoing	0.99 (0.76 – 1.30)	1.04 (0.77 – 1.40)	1.02 (0.75 – 1.39)	0.66 (0.48 – 0.93)*
Avoidance	1.02 (0.80 – 1.30)	0.98 (0.76 – 1.26)	0.77 (0.59 – 0.99)*	1.18 (0.85 – 1.64)

Notes: OR, odds ratio; CI, confidence interval; All odds ratios were adjusted for age, sex, income, education, country and daily cigarette consumption;

*
p<0.05,

questions not asked in the predictor wave.