Published in final edited form as:

Curr Psychiatry Rep. 2014 September ; 16(9): 462. doi:10.1007/s11920-014-0462-z.

Meta-Analytic Review of Psychological Interventions for Children Survivors of Natural and Man-Made Disasters

Elana Newman,

University of Tulsa Institute of Trauma, Adversity, and Injustice, Department of Psychology, 800 South Tucker Drive, Tulsa, OK 74103, USA

Betty Pfefferbaum,

Department of Psychiatry and Behavioral Sciences, College of Medicine, Terrorism and Disaster Center, University of Oklahoma Health Sciences Center, PO Box 26901, Williams Pavilion 3470, Oklahoma City, OK 73126-0901, USA

Namik Kirlic,

University of Tulsa Institute of Trauma, Adversity, and Injustice, Department of Psychology, 800 South Tucker Drive, Tulsa, OK 74103, USA

Robert Tett,

University of Tulsa Institute of Trauma, Adversity, and Injustice, Department of Psychology, 800 South Tucker Drive, Tulsa, OK 74103, USA

Summer Nelson, and

University of Tulsa Institute of Trauma, Adversity, and Injustice, Department of Psychology, 800 South Tucker Drive, Tulsa, OK 74103, USA

Brandi Liles

University of Tulsa Institute of Trauma, Adversity, and Injustice, Department of Psychology, 800 South Tucker Drive, Tulsa, OK 74103, USA

Elana Newman: elana-newman@utulsa.edu; Betty Pfefferbaum: Betty-pfefferbaum@ouhsc.edu; Namik Kirlic: Namik-kirlic@utulsa.edu; Robert Tett: Robert-tett@utulsa.edu; Summer Nelson: Summer-nelson@utulsa.edu; Brandi Liles: Brandi-liles@utulsa.edu

© Springer Science+Business Media New York 2014

Correspondence to: Elana Newman, elana-newman@utulsa.edu.

Conflict of Interest None of the authors of this manuscript report any potential or actual conflicts of interest.

Compliance with Ethics Guidelines

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by any of the authors.

This article is part of the Topical Collection on *Disaster Psychiatry*

Disclosures This work was funded in part by the National Institute of Mental Health, the National Institute of Nursing Research, and the Substance Abuse and Mental Health Services Administration (5 R25 MH070569) which established the Child and Family Disaster Research Training and Education Program in the Terrorism and Disaster Center (TDC) at the University of Oklahoma Health Sciences Center. TDC is a partner in the National Child Traumatic Stress Network and is funded in part by the Substance Abuse and Mental Health Services Administration (1 U79 SM57278).

Points of view in this document are those of the authors and do not necessarily represent the official position of the National Child Traumatic Stress Network; the National Institute of Mental Health; the National Institute of Nursing Research; the Substance Abuse and Mental Health Services Administration; the University of Oklahoma Health Sciences Center; the University of Tulsa; or the U.S. Department of Veterans Affairs.

Abstract

Although many post-disaster interventions for children and adolescent survivors of disaster and terrorism have been created, little is known about the effectiveness of such interventions. Therefore, this meta-analysis assessed PTSD outcomes among children and adolescent survivors of natural and man-made disasters receiving psychological interventions. Aggregating results from 24 studies (total $N=2630$) indicates that children and adolescents receiving psychological intervention fared significantly better than those in control or waitlist groups with respect to PTSD symptoms. Moderator effects were also observed for intervention package, treatment modality (group vs. individual), providers' level of training, intervention setting, parental involvement, participant age, length of treatment, intervention delivery timing, and methodological rigor. Findings are discussed in detail with suggestions for practice and future research.

Keywords

Child disaster; Interventions; Meta-analysis; Terrorism; Posttraumatic stress disorder; Acute stress disorder; Natural disaster; Treatment efficacy; Psychosocial treatment

Introduction

Children and adolescents surviving large scale traumatic events, such as disasters, accidents, and terrorism present with a wide range of psychological problems, including posttraumatic stress disorder (PTSD) [1, 2]. Meta-analyses of child interventions focused on PTSD [3•, 4] have revealed promising results, but the lack of specific focus upon the efficacy and effectiveness of child disaster mental health interventions leaves a major public health concern. Recently, several qualitative reviews have examined child disaster mental health interventions [5, 6, 7••, 8••, 9], describing specific trends and gaps in the field.

To extend this work, this meta-analysis was conducted to examine effect sizes of interventions upon PTSD and explore the moderating effects of factors raised in descriptive reviews [5, 6, 7••, 8••, 9]. Our specific aims were to: (1) examine outcomes in children receiving treatment relative to control and waitlist groups; (2) evaluate treatment outcomes yielded by specific intervention packages, and (3) examine the moderating role of intervention packages, treatment modality (individual vs. group), providers' level of training, intervention setting, parental involvement, participant age, length of therapy, intervention delivery timing, and the methodological rigor of the study based on Nathan and Gorman [10] criteria. These moderators have been the subject of critical focus and debate in the field. Broadly, we seek to delineate treatment-improving factors, guiding recommendations for psychological practices in post-disaster environments for children and adolescents presenting with trauma-related symptoms.

Method

Study Selection

Research studies assessing outcomes of interventions for children exposed to mass trauma were identified by searching the following bibliographic databases: Ovid, Medline,

PsycInfo, EBM Reviews, EMBASE, ERIC, and PubMed. The initial search was conducted in 2011 and updated in the winter of 2012. Articles were identified by the following terms: treatment(s), intervention(s), child(ren), adolescent(s), disaster(s), terrorist event(s), terrorism, teen, youth, hurricane, tsunami, disaster intervention, posttraumatic stress disorder, treatment outcomes, grief, drug therapy, early intervention, trauma, and family therapy. Titles and abstracts identified in the searches were reviewed to select material for potential review.

All manuscripts published in English providing useable statistical outcome data were included. Initially we reviewed all studies addressing ongoing political conflict, war, and single incidents such as accidents. However, we excluded articles on ongoing war and terrorism because continuous exposure to terrorism and war likely requires different interventions that address both past and current concerns that are different from past war and terrorism. For similar reasons, we also excluded treatments aimed at single accident events and interpersonal violence.

Of the 74 articles identified in the initial search, 18 were classified as war or chronic terrorism studies and were thus excluded. Additionally, four case studies [11–14] and one mixed sample study [15] were dropped. Seven articles focused on accidents or sudden loss of a loved one were excluded [16–22]. Finally, 19 were excluded due to a lack of outcome data on PTSD, missing appropriate numerical outcome data required to calculate effect sizes for analysis (i.e., means, standard deviations), and/or if consisted of treatment description only [23–41], thus yielding 25 studies. Finally, one study was excluded as an outlier (*Cohen's d* > 3.5) [42]. Seven studies [43–49] offered results for two independent samples (see below). All told, the final sample includes 31 main data points, from 24 distinct studies.

Coding Procedure

Rater pairs coded each useable article on a variety of dimensions, including sample demographics (mean age, % male), trauma exposure, outcome measures, use of control group (placebo, waitlist, non-treatment), total sample size at each time point, days since trauma (i.e., intervention delivery timing), number of treatment sessions, session length, follow up since treatment, and means and standard deviations for each outcome measure of each group and at each time point. Studies were further coded for the presence or absence of blinded assessment, random assignment, use of treatment manuals, clear inclusion and exclusion criteria, treatment adherence, and psychometric properties of the outcome measure (reliability and validity), thereby allowing the classification of each study into Types 1, 2, or 3, according to the Nathan and Gorman [10] levels of methodological rigor. A Type 1 study represents the highest level of rigor, applicable to a treatment study that had comparison groups, random assignment, blinded assessment, clear inclusion and exclusion criteria, state-of-the-art diagnostic measures, adequate power, and clearly described methods [10]. Type 2 studies are missing one or two of the Type 1 criteria, but are otherwise sound. Type 3 studies lack comparison groups, present only pre-post data, are open treatment studies obtaining pilot data, or have obtained outcome information retrospectively. Type 4, 5, and 6 studies were not identified in the literature review. Following the evaluation of individual studies, the research team met to sort articles by intervention package, as follows: Strict Cognitive

Behavioral Therapy (CBT), CBT with Grief Interventions, Eclectic with and without CBT components, Eye Movement Desensitization and Reprocessing (EMDR), Exposure, Relaxation, Psychological First Aid, and Psychological Debriefing/Crisis Intervention. All coding disagreements and inconsistencies were resolved by consensus and consultation.

For values where ranges were given, the midpoint was used. Baseline values were used as pre-treatment measures, while the last follow-up data collection values were used as post-treatment (outcome) values. When effect sizes were calculated for intervention groups relative to controls, the last time point when data were collected for both groups was used. In cases where waitlist control groups were used for comparison, post-intervention data of the intervention group was compared to pre-intervention data of the waitlist control group. Finally, studies comparing active treatments side by side or using a modified treatment as a comparison were separated and treated as individual sources. Fifteen different measures assessed PTSD symptoms (see Table 1).

Coding procedures were refined, and coders trained through pilot coding of five articles outside the useable set but from related domains (e.g., adult trauma and child maltreatment). Coding was refined until coders reached an average agreement of $r=.80$. Median kappa and median percent agreement after coding 18 articles was .80 and 88.9 %, respectively (means, .80 and 88.6). Due to staff changes, the final seven articles added to the study were coded by only one of the raters in consultation with the lead author. All discrepancies were resolved in consultation with the team.

Effect size Calculation

Within group pre-post treatment effect sizes were calculated for each study using Cohen's [67] $d=(M_1-M_2)/\sigma_p$, where M_1 =pre-treatment mean, M_2 =post-treatment mean, and σ_p =pooled standard deviation. The same formula was used to calculate between-group outcome effect sizes, where M_1 =intervention group mean, M_2 =control group mean, and σ_p =pooled standard deviation. Small, medium, and large effect sizes were defined as $d=.20$, $d=.50$, and $d=.80$, respectively [67].

Meta-analysis Procedures

Mean effect sizes and cross-study moderators were assessed using Hunter and Schmidt's [68] "bare bones" meta-analysis procedure, which averages observed effect sizes across studies without correcting for systematic effects of measurement unreliability and range restriction. The "bare bones" approach is recommended when reliabilities and information needed to determine differential range restriction are too infrequently reported to permit the associated corrections, as was judged the case here. The procedure yields two key outputs, per analysis (i.e., overall and by moderator subgroup): (1) the sample N -weighted mean d closely approximates the value of d that would obtain with all participants from all studies combined into a single sample; and (2) $SD(d')$ is the standard deviation of d with variability due to sampling error removed¹. To the degree $SD(d')>0$, the mean weighted d may be considered the mean of multiple uncorrected δ s, warranting consideration of possible moderators. Of further relevance to moderators, $SD(d')$ is used to calculate an 80 % credibility interval [$\text{mean } d \pm 1.28 * SD(d')$], identifying the points at which 10 % of

uncorrected δ s (represented by the given sample of input d s) are expected to fall above and 10 % to fall below. Where $SD(d')=0$, the interval has zero width, and N -weighted mean d is taken to estimate a single uncorrected δ . Success rates, or relative improvement, were assessed using the binomial effect size display (BESD) [69].

Due to the relatively low power conferred by the small number of available studies, we relied on procedures offered by Hunter and Schmidt [68, p. 293–294], based on logic underlying analysis of variance, when testing for moderators. Specifically, moderation is suggested to the degree that (a) mean d varies between subgroup and (b) the k -weighted average $Var(d)$ across subsets is less than that based on the combined (i.e., overall) sample. The Hunter and Schmidt [68] method has been found to yield the most accurate estimate of the moderating effect magnitude, as well as superior regarding accuracy of Type I error rates [70].

We also used a more traditional method of testing moderator effects, comparing subgroups based on the Q statistic [71]. Specifically, between-group heterogeneity (Q_{between}) is tested as χ^2 with $df=N_{\text{groups}} - 1$. For moderators with three or more levels (i.e., intervention package and methodological rigor), an additional conservative omnibus moderator comparison approach was used [71]. This approach resembles the Bonferroni correction used in ANOVA, where a series of comparisons between all possible two-group combinations is made, and the Q_{between} for each of the two groups is examined using a corrected significance level based on the total number of comparisons.

Analyses

Three sets of analyses were conducted: (1) intervention outcomes comparing treatment relative to control groups for PTSD ($k=14$); (2) pre-versus post-treatment intervention data without control groups for PTSD ($k=30$), including 13 of the 14 from the analysis above; and (3) potential moderating effects of intervention package, treatment modality (individual vs. group), providers' level of training, intervention setting, parental involvement, participant age, length of therapy, intervention delivery timing, and the methodological rigor of the study. Moderator subgroups were derived from the 30 pre/post studies, permitting direct comparison of mean subgroup $Var(d)$ to overall $Var(d)$. Intervention packages represented by only one study (i.e., Psychological First Aid, Relaxation, CBT with Grief components, Psychological Debriefing, and Spiritual-Hypnosis Assisted Treatment) were excluded altogether from this particular moderation analysis. To allow more robust comparisons for the five remaining packages with $k \geq 2$, we re-ran the overall analysis without the five $k=1$ intervention packages ($k=25$). Similarly, five input samples did not specify the setting in which the treatment was carried out, and were thus excluded from the moderation analysis examining the effect of setting. We re-ran the analysis without these five studies ($k=25$), to allow more robust comparisons for the remaining samples with

¹Mean d fully corrected for unreliability and range restriction permits interpretation as δ (or mean δ), and $Var(d)$ is further refined by removal of variance attributable to those artifacts. As we do not correct for unreliability or range restriction, mean d denotes the mean uncorrected δ and we use d' to represent our consideration of sampling error only in calculating residual variance [i.e., $Var(d')=Var(d) - Var(e)$].

clearly specified setting. Across all moderator analyses, we report meta-analytic results for $k=2$ cases, but interpret the outcomes with caution.

Results

The 31 input samples had a combined N of 2630 participants (Table 1). Averaging across samples, the mean participant's age was 10.9 years ($SD=2.3$), and 46 % of participants were male. All samples had been exposed to traumatic events, with median intervention starting 365 days after exposure. Interventions averaged 8.4 ($SD=4.9$) hours of therapy across 8.1 ($SD=4.6$) sessions.

Symptom Outcomes

Meta-analytic results, presented in Table 2, suggest that treatment group outcomes are better than control and waitlist group outcomes on PTSD measures, showing a medium effect size (Cohen's $d=.74$, $SD=.59$, 80 % $CI=-.02$ to 1.49). With respect to binomial success rates, participants receiving some form of psychological intervention averaged 66 % more improvement over that of control and waitlist group participants on PTSD measures. The weighted mean effect size for the reduction in PTSD symptoms as a result of psychological intervention (pre- vs. post-treatment) was large ($d=1.13$, $SD=.69$, 80 % $CI=.25$ to 2.02), yielding 74 % relative improvement.

Moderator Analyses

Moderator analyses examined whether the effects of psychological intervention on PTSD outcomes varied as a result of intervention package, treatment modality (individual vs. group), providers' level of training, intervention setting, parental involvement, participant age, length of therapy, intervention delivery timing, and methodological rigor.

Comparisons by intervention package (focusing on the five intervention packages with $k=2$) yielded different mean effect sizes, accompanied by a moderation [$mean\ subgroup\ Var(d)=.42$, $Var(d)_{overall}=.57$; $\chi^2(4, K=25)=149.64$, $p<.001$]. Large treatment effect sizes are evident for *EMDR* ($d=2.15$), followed by *Exposure* ($d=1.56$), *Strict CBT* ($d=1.25$), *Eclectic with CBT* ($d=1.07$), and *Eclectic with no CBT* ($d=.56$). Using the Omnibus moderator comparison approach corrected for multiple comparisons at $p=.005$ level, the $Q_{between}$ analysis revealed that *EMDR* significantly differed from *Eclectic with CBT* [$\chi^2(1, K=12)=63.48$, $p<.001$], *Strict CBT* [$\chi^2(1, K=11)=42.11$, $p<.001$], *Exposure* [$\chi^2(1, K=4)=9.55$, $p<.005$], and *Eclectic with no CBT* interventions [$\chi^2(1, K=4)=115.04$, $p<.001$]. *Exposure* significantly differed from *Eclectic with no CBT* [$\chi^2(1, K=4)=43.39$, $p<.001$], *Eclectic with CBT* [$\chi^2(1, K=12)=12.35$, $p<.001$], but not *Strict CBT* [$\chi^2(1, K=11)=4.84$, $p=.03$] interventions. *Strict CBT* significantly differed from *Eclectic with CBT* [$\chi^2(1, K=19)=10.57$, $p<.005$], and *Eclectic with no CBT* interventions [$\chi^2(1, K=10)=71.55$, $p<.001$]. Finally, *Eclectic with CBT* and *Eclectic with no CBT* interventions differed significantly [$\chi^2(1, K=12)=46.87$, $p<.001$].

The analysis further yielded higher mean effect sizes for children receiving individual treatment ($d=1.38$) relative to those in group therapy ($d=1.04$), accompanied by a moderation [$mean\ subgroup\ Var(d)=.43$, $Var(d)_{overall}=.54$; $\chi^2(1, K=30)=55.38$, $p<.001$].

Interventions carried out by mental health professionals yielded larger effect sizes ($d=1.19$) than those carried out by teachers and other school personal ($d=.93$) for PTSD outcomes [*mean subgroup* $Var(d)=.47$, $Var(d)_{overall}=.54$; $\chi^2(1, K=30)=26.93$, $p<.001$].

Outcomes varied according to the setting in which interventions were carried out [*mean subgroup* $Var(d)=.53$, $Var(d)_{overall}=.56$; $\chi^2(2, K=25)=31.72$, $p<.001$], such that the largest effect sizes were observed for interventions conducted in refugee camps ($d=1.47$), followed by interventions at health or mental health sites ($d=1.31$), and finally, schools ($d=1.11$). Using omnibus moderator comparison approach corrected for multiple comparisons at $p=.017$ level, the $Q_{between}$ analysis revealed that interventions conducted in refugee camps predicted better outcomes than interventions in schools [$\chi^2(1, K=20)=28.25$, $p<.001$], but did not significantly differ from those conducted in mental health settings [$\chi^2(1, K=7)=2.62$, $p=.11$]. Gains were significantly greater for interventions conducted in health or mental health settings than in those conducted in schools [$\chi^2(1, K=23)=6.34$, $p=.012$]. Notably, only two studies in our sample were conducted in refugee camps; the results should be interpreted with this in mind.

Interventions involving parents yielded better outcomes than those not involving parents [$d=1.30$ vs. $d=1.04$; *mean subgroup* $Var(d)=.52$, $Var(d)_{overall}=.54$; $\chi^2(1, K=30)=37.08$, $p<.001$].

Outcomes varied across different age groups [*mean subgroup* $Var(d)=.50$, $Var(d)_{overall}=.54$; $\chi^2(2, K=30)=71.26$, $p<.001$]. Using omnibus moderator comparison approach corrected for multiple comparisons at $p=.017$ level, the $Q_{between}$ analysis revealed that children in the 10–11.6 age group ($d=1.30$) exhibited greater treatment gains than children in the 5.3–9.83 age group ($d=.91$) [$\chi^2(1, K=22)=68.43$, $p<.001$], but not children in the 12–16 age group ($d=1.19$) [$\chi^2(1, K=22)=4.29$, $p=.04$]. Additionally, children in the 12–16 age group exhibited greater treatment gains than children in the 5.3–9.83 age group [$\chi^2(1, K=16)=28.11$, $p<.001$].

Outcomes further varied as a result of number of hours spent in therapy [*mean subgroup* $Var(d)=.53$, $Var(d)_{overall}=.54$; $\chi^2(2, K=30)=36.95$, $p<.001$]. Using omnibus moderator comparison approach corrected for multiple comparisons at $p=.017$ level, the $Q_{between}$ analysis revealed that spending 0.5–4.5 ($d=1.18$) and 6–9 ($d=1.25$) hours in therapy did not significantly differ [$\chi^2(1, K=20)=2.12$, $p=.15$]. Both significantly predicted better outcomes than spending 10–18 ($d=.95$) hours in therapy [$\chi^2(1, K=20)=19.36$, $p<.001$ and $\chi^2(1, K=20)=34.37$, $p<.001$, respectively].

Interventions delivered within four months following trauma exposure yielded largest effect sizes ($d=1.47$), followed by interventions delivered between 39 and 69 months ($d=1.20$) and interventions delivered between 4.5 and 12 months following trauma exposure ($d=.76$), supported by a moderation [*mean subgroup* $Var(d)=.48$, $Var(d)_{overall}=.54$; $\chi^2(2, K=30)=185.08$, $p<.001$]. Using the Omnibus moderator comparison approach corrected for multiple comparisons at $p=.017$ level, the $Q_{between}$ analysis revealed that interventions delivered within the first four months significantly differed from interventions delivered between 4.5 and 12 [$\chi^2(1, K=17)=178.60$, $p<.001$] months and interventions delivered

between 39 and 69 months following trauma exposure [$\chi^2 (1, K=23)=27.76, p<.001$]. Finally, interventions delivered between 4.5 and 12 months differed significantly from those delivered between 39 and 69 months following trauma exposure [$\chi^2 (1, K=20)=78.29, p<.001$].

With respect to methodological rigor, the largest effect sizes were observed in Type 2 studies ($d=1.22$), followed by Type 1 (highest rigor) ($d=1.20$), and Type 3 ($d=.93$) studies. A moderating effect of methodological rigor was supported by Q_{between} [$\chi^2 (2, K=30)=39.48, p<.001$], but not a $\text{Var}(d)$ comparison [$\text{mean subgroup Var}(d)=.54, \text{Var}(d)_{\text{overall}}=.54$]. Using the Omnibus moderator comparison approach corrected for multiple comparisons at $p=.017$ level, the Q_{between} analysis revealed that Type 1 studies significantly differed from Type 3 [$\chi^2 (1, K=18)=22.28, p<.001$], but not Type 2 studies [$\chi^2 (1, K=18)=.13, p=.72$]. Finally, Type 2 and Type 3 studies also significantly differed [$\chi^2 (1, K=24)=36.23, p<.001$].

Discussion

This meta-analysis assessed (1) the effectiveness of psychological interventions in reducing PTSD symptoms in 2630 child and adolescent survivors of mass trauma; and (2) PTSD factors that may moderate treatment outcomes, including intervention package, treatment modality (individual vs. group), providers' level of training, intervention setting, parental involvement, participant age, length of therapy, intervention delivery timing, and the methodological rigor of the study.

In general, results suggest that psychological interventions are successful in ameliorating posttraumatic stress symptoms. The weighted mean effect size of PTSD reduction as a result of psychosocial treatment was large, yielding a 74 % improvement compared to baseline. Moreover, children and adolescents receiving any psychological intervention fared significantly better than those in control or waitlist groups with respect to PTSD symptoms, averaging 66 % more improvement than those not receiving treatment. Together these results suggest that natural recovery or regression to the mean cannot fully explain the positive outcome from these interventions [7•]. Thus there is clear evidence that child mental health interventions are effective. Future studies should provide evidence about the characteristics of those who do not improve in order to hone treatments to address the needs of all children and adolescents post disaster.

Although our meta-analysis includes several interventions with only two studies examining them (EMDR, Exposure, Eclectic with CBT), compelling evidence suggest that EMDR, Strict CBT, Exposure, and Eclectic with CBT intervention packages are all effective for children and adolescents post disaster. As EMDR, Exposure, and Eclectic with CBT demonstrated strong effectiveness based on only two studies each, more studies yielding positive effects are needed before such interventions can be recommended. Based on the studies included in this review, it may be prudent to use strict CBT for large-scale disaster responses until more is known. Strict CBT, which had a slightly lower effect size than EMDR and Exposure in this meta-analysis, was based on more studies (nine studies, across six research teams) involving more participants, suggesting that at this time, the evidence in its favor is more robust. Nevertheless, all interventions were effective and EMDR,

Exposure, and Strict CBT had the largest effect sizes and should be considered evidence-based practice. These findings and guidelines differ from those of the International Society of Traumatic Stress [72], which could not determine conclusive recommendations based on the available evidence at that time. However, the ISTSS guidelines were confined to interventions solely in the early phase, included data from studies involving accidents and had fewer studies on which to make a determination. Future research might examine what specific intervention components are shared across these treatments and dismantling efforts might determine which are most critical [7••]. Given the sizable public health demands of a disaster, distilling the necessary curative factors can be expected to enhance the cost-efficacy of interventions.

Several moderators were shown to affect treatment outcomes. Children receiving individual therapy had greater improvement than those in group interventions, although both formats yielded large effect sizes. Given that 43 % of the treatments were delivered more than a year after the event, it may be that more individualized treatment was necessary due to the chronicity of the PTSD symptoms. Similarly, while all providers were effective, those interventions carried out by mental health professionals yielded more improvement in PTSD than those carried out by other types of providers. These results need to be interpreted cautiously as only four studies examined non-mental health professionals. Outcomes varied according to the setting in which interventions were carried out such that the largest effect sizes were observed for interventions conducted in refugee camps, followed by interventions at health or mental health sites and finally, schools. Since only two studies in our sample were conducted in refugee camps, these results indicate that, at a minimum, interventions can be successfully delivered in temporary sheltering. Ultimately we need a larger set of studies to examine and untangle the interactions of format (group, individual), provider training, and setting.

Parental involvement in treatment enhanced outcome, but the effects of treatments not involving parents were still large. Thus, while parental engagement may be optimal it is not a necessity. In general, children ages 10 and above demonstrated greater treatment gains than children below 10, but all age groups benefited from treatment. It is noteworthy that effect sizes for all ages were of medium to large size (d range=0.91 to 1.30).

Optimal timing and length of post-disaster services continues to be a greatly debated topic [8••]. Strikingly, children who engaged in one half hour to nine hours of an intervention predicted better outcomes, but all the outcomes had medium to large effects (d range=0.95 to 1.25). It is unclear if this is an artifact due to the fact that those children and adolescents with greater needs required more time to address symptoms. Interventions delivered within four months of the event yielded the largest effect size, supporting the need for early interventions. Of the ten studies within four months, four initiated delivery within the first month, or the acute phase [73]. Thus we do not have enough information yet to make recommendations about optimal timing within the first four months and whether the acute interval is indeed an optimal time of receptiveness to secondary prevention strategies [73]. Strategies and funding to help researchers quickly assess acute pediatric post-disaster interventions are a high priority public health need.

Studies with more rigorous methodology were more likely to have stronger effect sizes. Strong effect sizes are noteworthy in light of concerns raised that the demands of conducting outcome research might detract from addressing and individualizing post-disaster needs [5]. These meta-analytic results suggest quite the contrary conclusion; despite the challenges of conducting clinical trials in post-disaster recovery environments, such efforts likely enhance outcomes. This conclusion is tempered by the fact that the inclusion criteria for this meta-analysis necessitated that all studies at minimum had some outcome data.

Our results bear consideration in light of several limitations. First and foremost, the number of studies meeting inclusion criteria was modest, especially within moderator subgroups. More robust analysis awaits accumulation of larger numbers of point estimates from independent samples. Moreover, many new recommended disaster-related interventions for children are not yet tested in scientific literature, nor explicitly linked to PTSD outcomes. For example the core principles of disaster interventions [74] are beyond the scope of this review. Second, as an extension of the first point, interactions among intervention packages and other moderators were precluded by the small k , warranting future examination as more studies become available for meta-analysis in this area. Finally, this review was confined to examination of PTSD symptoms only and did not include other important outcomes such as depression, anxiety, academic success, quality of life, and interpersonal functioning.

Conclusions

Meta-analytic results of the existing literature indicate that disaster interventions for children and adolescents clearly are efficacious. More outcomes research on emerging and existing interventions is needed to enhance public health interventions and address issues that cannot yet be determined based on the existing literature. Future research would benefit from including evaluations of cost-effectiveness and ease of dissemination.

References

Papers of particular interest, published recently, have been highlighted as:

• Of importance

•• Of major importance

1. Norris FH, et al. 60,000 disaster victims speak: Part I. An empirical review of the empirical literature, 1981–2001. *Psychiatry: Int Biol Process*. 2002; 65(3):207–39.
2. Gurwitch, RH.; Pfefferbaum, B.; Leftwich, MJT. Trauma Practice in the Wake of September 11, 2001. In: Gold, SN.; Faust, J., editors. *The Impact of Terrorism on Children: Considerations for a New Era*. New York: Haworth Press; 2002. p. 101-24.
- 3•. Rolfesnes ES, Idsoe T. School-based intervention programs for PTSD symptoms: a review and meta-analysis. *J Trauma Stress*. 2011; 24(2):155–65. This meta-analysis reviewed school-based interventions for PTSD resulting from a range of traumatic events. Findings were indicative of medium to large effect sizes in symptom reduction, suggesting that school-based interventions are effective in reducing PTSD symptoms in children and adolescents. [PubMed: 21425191]
4. Silverman WK, Pina AA, Viswesvaran C. Evidence-based psychosocial treatments for phobic and anxiety disorders in children and adolescents. *Clin Child Adolesc Psychol*. 2008; 37(1):105–30.

5. Pfefferbaum B, Newman E, Nelson SD. Mental Health Interventions for Children Exposed to Disasters and Terrorism. *J Child Adolesc Psychopharmacol*. 2014; 24(1):24–31. [PubMed: 24494722]
6. Pfefferbaum B, et al. Research methodology used in studies of child disaster mental health interventions for posttraumatic stress. *Comprehensive Psychiatry*. 2013
- 7•. Pfefferbaum B, et al. Child disaster mental health interventions, Part I: Techniques, outcomes, and methodological considerations. *Disaster Health*. 2014 This qualitative review reviews the existing literature with respect to therapy techniques, outcomes and methodology raising questions for future research. 10.4161/dish.27534
- 8•. Pfefferbaum B, et al. Child disaster mental health interventions. Part II: Timings Settings Deliv Approaches. 2014 This qualitative review describes the extant evidence base about optimal delivery and approaches to pediatric disaster-related interventions. 10.4161/dish.27535
9. Pfefferbaum B, et al. Universal Preventive Interventions for Children in the Context of Disasters and Terrorism. *Child Adolesc Psychiatr Clin N Am*. 2014; 23(2):363–82. [PubMed: 24656585]
10. Nathan, PE.; Gorman, JM. A guide to treatments that work. New York: Oxford University Press; 1998. A guide to treatments that work.
11. Brown EJ, Pearlman MY, Goodman RF. Facing fears and sadness: Cognitive-behavioral therapy for childhood traumatic grief. *Harv Rev Psychiatry*. 2004; 12(4):187–98. [PubMed: 15371061]
12. Cohen JA, et al. Treating traumatized children after Hurricane Katrina: Project Fleur de lis. *Clin Child Fam Psychol Rev*. 2009; 12(1):55–64. [PubMed: 19224365]
13. Goodman RF, et al. Letting the story unfold: A case study of client-centered therapy for childhood traumatic grief. *Harv Rev Psychiatry*. 2004; 12(4):199–212. [PubMed: 15371062]
14. Tufnell G. Eye movement desensitization and reprocessing in the treatment of pre-adolescent children with post-traumatic symptoms. *Clin Child Psychol Psychiatry*. 2005; 10(4):587–600.
15. Salloum A, Overstreet S. Grief and trauma intervention for children after disaster: Exploring coping skills versus trauma narration. *Behav Res Ther*. 2012; 50(3):169–79. [PubMed: 22317753]
16. Chapman L, et al. The effectiveness of art therapy interventions in reducing post traumatic stress disorder (PTSD) symptoms in pediatric trauma patients. *Art Therapy*. 2001; 18(2):100–4.
17. Cohen JA, Mannarino AP, Staron VR. A pilot study of modified cognitive-behavioral therapy for childhood traumatic grief (CBT-CTG). *J Am Acad Child Adolesc Psychiatry*. 2006; 45(12):1465–73. [PubMed: 17135992]
18. Kemp M, Drummond P, McDermott B. A wait-list controlled pilot study of eye movement desensitization and reprocessing (EMDR) for children with post-traumatic stress disorder (PTSD) symptoms from motor vehicle accidents. *Clin Child Psychol Psychiatry*. 2010; 15(1):5–25. [PubMed: 19923161]
19. Smith P, et al. Cognitive-behavioral therapy for PTSD in children and adolescents: a preliminary randomized controlled trial. *J Am Acad Child Adolesc Psychiatry*. 2007; 46(8):1051–61. [PubMed: 17667483]
20. Stallard P, Law FD. Screening and psychological debriefing of adolescent survivors of life-threatening events. *Br J Psychiatry*. 1993; 163:660–5. [PubMed: 8298836]
21. Stallard P, et al. A randomised controlled trial to determine the effectiveness of an early psychological intervention with children involved in road traffic accidents. *J Child Psychol Psychiatry*. 2006; 47(2):127–34. [PubMed: 16423143]
22. Van der Oord S, et al. Treatment of post-traumatic stress disorder in children using cognitive behavioural writing therapy. *Clin Psychol Psychother*. 2010; 17(3):240–9. [PubMed: 20013756]
23. Amaya-Jackson L, et al. Cognitive-behavioral treatment for pediatric posttraumatic stress disorder: Protocol and application in school and community settings. *Cogn Behav Pract*. 2003; 10(3):204–13.
24. Cohen JA, et al. Treatment of childhood traumatic grief: contributing to a newly emerging condition in the wake of community trauma. *Harv Rev Psychiatry*. 2004; 12(4):213–6. [PubMed: 15371063]
25. Fernandez I. EMDR as treatment of post-traumatic reactions: A field study on child victims of an earthquake. *Educ Child Psychol*. 2007; 24(1):65–72.

26. Field T, et al. Alleviating posttraumatic stress in children following Hurricane Andrew. *J Appl Dev Psychol.* 1996; 17(1):37–50.
27. Galante R, Foa D. An epidemiological study of psychic trauma and treatment effectiveness for children after a natural disaster. *J Am Acad Child Psychiatry.* 1986; 25(3):357–63.
28. Hardin SB, et al. Effects of a long-term psychosocial nursing intervention on adolescents exposed to catastrophic stress. *Issues Ment Health Nurs.* 2002; 23(6):537–51. [PubMed: 12217220]
29. Karairmak Ö, Aydin G. Reducing earthquake-related fears in victim and nonvictim children. *J Genet Psychol: Res Theory Hum Dev.* 2008; 169(2):177–85.
30. Karam EG, et al. Effectiveness and specificity of a classroom-based group intervention in children and adolescents exposed to war in Lebanon. *World Psychiatry: Off J World Psychiatr Assoc (WPA).* 2008; 7(2):103–9.
31. Mahmoudi-Gharaei J, et al. Group cognitive-behavior therapy and supportive art and sport interventions on Bam earthquake related post traumatic stress symptoms in children: a field trial. *Iran J Psychiatry.* 2009; (4):85–91.
32. Plummer CA, et al. Practice challenges in using psychological first aid in group format with children: a pilot study. *Brief Treat Crisis Interv.* 2009; 8:313–26.
33. Sahin NH, Yilmaz B, Batigun A. Psychoeducation for children and adults after the Marmara earthquake: An evaluation study. *Traumatology.* 2011; 17(1):41–9.
34. Salloum A, et al. Grief and trauma group therapy for children after Hurricane Katrina. *Soc Work Groups: J Community Clin Pract.* 2009; 32(1–2):64–79.
35. Shen Y-J. Short-term group play therapy with Chinese earthquake victims: Effects on anxiety, depression and adjustment. *Int J Play Ther.* 2002; 11(1):43–63.
36. Stubenbort K, Donnelly GR, Cohen JA. Cognitive-behavioral group therapy for bereaved adults and children following an air disaster. *Group Dyn: Theory Res Pract.* 2001; 5(4):261–76.
37. Tol WA, et al. Mediators and moderators of a psychosocial intervention for children affected by political violence. *J Consult Clin Psychol.* 2010; 78(6):818–28. [PubMed: 21114342]
38. Weems CF, et al. Effect of a school-based test anxiety intervention in ethnic minority youth exposed to Hurricane Katrina. *J Appl Dev Psychol.* 2009; 30(3):218–26.
39. Cohen, JA., et al. Cognitive-behavioral therapy for children and adolescents. In: Foa, EB., et al., editors. *Effective treatments for PTSD: Practice guidelines from the International Society for Traumatic Stress Studies.* 2. Guilford Press; New York, NY US: 2009. p. 223-244.
40. Goodman RF, Brown EJ. Service and science in times of crisis: Developing, planning, and implementing a clinical research program for children traumatically bereaved after 9/11. *Death Stud.* 2008; 32(2):154–80. [PubMed: 18693386]
41. Smith, P.; Dyregrov, A.; Yule, W. *Children and disaster: Teaching Recovery Techniques.* Bergen: Children and War Foundation; 1999.
42. Giannopoulou I, Dikaikou A, Yule W. Cognitive-behavioural Group Intervention for PTSD Symptoms in Children Following the Athens 1999 Earthquake: A Pilot Study. *Clin Child Psychol Psychiatry.* 2006; 11(4):543–53. [PubMed: 17163223]
43. Brown EJ, et al. Matching Interventions to Children's Mental Health Needs: Feasibility and Acceptability of a Pilot School-Based Trauma Intervention Program. *Educ Treat Child.* 2006; 29(2):257–86.
44. Catani C, et al. Treating children traumatized by war and tsunamis: A comparison between exposure therapy and meditation-relaxation in North-East Sri Lanka. *BMC Psychiatry.* 2009;9. [PubMed: 19292899]
45. Chemtob CM, Nakashima JP, Hamada RS. Psychosocial intervention for postdisaster trauma symptoms in elementary school children: a controlled community field study. *Arch Pediatr Adolesc Med.* 2002; 156(3):211–6. [PubMed: 11876663]
46. CATS Consortium . Implementation of CBT for youth affected by the World Trade Center disaster: Matching need to treatment intensity and reducing trauma symptoms. *J Trauma Stress.* 2010; 23(6): 699–707. [PubMed: 21171130]
47. de Roos C, et al. A randomised comparison of cognitive behavioural therapy (CBT) and eye movement desensitisation and reprocessing (EMDR) in disaster-exposed children. *European Journal Of Psychotraumatology.* 2011;2.

48. Jaycox LH, et al. Children's mental health care following Hurricane Katrina: A field trial of trauma-focused psychotherapies. *J Trauma Stress*. 2010; 23(2):223–31. [PubMed: 20419730]
49. Salloum A, Overstreet S. Evaluation of individual and group grief and trauma interventions for children post disaster. *J Clin Child Adolesc Psychol*. 2008; 37(3):459–507.
50. Berger R, Gelkopf M. School-based intervention for the treatment of tsunami-related distress in children: A quasi-randomized controlled trial. *Psychother Psychosom*. 2009; 78(6):364–71. [PubMed: 19738402]
51. Cain DS, et al. Weather the storm: Persistent effects and psychological first aid with children displaced by hurricane Katrina. *J Child Adolesc Trauma*. 2010; 3(4):330–43.
52. Chemtob CM, Nakashima J, Carlson JG. Brief treatment for elementary school children with disaster-related posttraumatic stress disorder: A field study. *J Clin Psychol*. 2002; 58(1):99–112. [PubMed: 11748599]
53. Gilboa-Schechtman E, et al. Prolonged exposure versus dynamic therapy for adolescent PTSD: a pilot randomized controlled trial. *J Am Acad Child Adolesc Psychiatry*. 2010; 49(10):1034–42. [PubMed: 20855048]
54. Goenjian AK, et al. Outcome of psychotherapy among early adolescents after trauma. *Am J Psychiatry*. 1997; 154(4):536–42. [PubMed: 9090342]
55. Goenjian AK, et al. A prospective study of posttraumatic stress and depressive reactions among treated and untreated adolescents 5 years after a catastrophic disaster. *Am J Psychiatry*. 2005; 162(12):2302–8. [PubMed: 16330594]
56. Lesmana CBJ, et al. A spiritual-hypnosis assisted treatment of children with PTSD after the 2002 Bali terrorist attack. *Am J Clin Hypn*. 2009; 52(1):23–34. [PubMed: 19678557]
57. Mahmoudi-Gharaei J, et al. The effects of a short-term cognitive behavioral group intervention on Bam earthquake related PTSD symptoms in adolescents. *Iran J Psychiatry*. 2009; 4:79–84.
58. March JS, et al. Cognitive-behavioral psychotherapy for children and adolescents with posttraumatic stress disorder after a single-incident stressor. *J Am Acad Child Adolesc Psychiatry*. 1998; 37(6): 585–93. [PubMed: 9628078]
59. Ronan KR, Johnston DM. Behaviourall-basedinterventions for children following volcanic eruptions: an evaluation of effectiveness. *Disaster Prevention Manag*. 1999; 8(3):169–76.
60. Scheeringa MS, et al. Trauma-focused cognitive-behavioral therapy for posttraumatic stress disorder in three-through six year-old children: a randomized clinical trial. *J Child Psychol Psychiatry Allied Discip*. 2011; 52(8):853–60. A representative study of the sample reviewed in the present meta-analytic study. This study examines the efficacy and feasibility of a trauma-focused cognitive-behavioral therapy in young children exposed to wide variety of traumas.
61. Shoohtary MH, Panaghi L, Moghadam JA. Outcome of cognitive behavioral therapy in adolescents after natural disaster. *J Adolesc Health: Off Publ Soc Adolesc Med*. 2008; 42(5):466–72.
62. Taylor LK, Weems CF. Cognitive-behavior therapy for disaster-exposed youth with posttraumatic stress: results from a multiple-baseline examination. *Behav Ther*. 2011; 42(3):349–63. [PubMed: 21658519]
63. Vijayakumar L, et al. Do all children need intervention after exposure to tsunami? *Int Rev Psychiatry (Abingdon, England)*. 2006; 18(6):515–22.
64. Vila G, Porche L-M, Mouren-Simeoni M-C. An 18-month longitudinal study of posttraumatic disorders in children who were taken hostage in their school. *Psychosom Med*. 1999; 61(6):746–54. [PubMed: 10593625]
65. Wolmer L, Laor N, Yazgan Y. School reactivation programs after disaster: could teachers serve as clinical mediators? *Child Adolesc Psychiatric Clin N Am*. 2003; 12(2):363–81.
66. Wolmer L, et al. Teacher-mediated intervention after disaster: A controlled three-year follow-up of children's functioning. *J Child Psychol Psychiatry*. 2005; 46(11):1161–8. [PubMed: 16238663]
67. Cohen, J. Statistical power analysis for the behavioral sciences. Hillsdale: Lawrence Erlbaum Associates, Inc; 1977. rev. ed
68. Hunter, JE.; Schmidt, FL. Methods of meta-analysis: Correcting error and bias in research findings. 2. Thousand Oaks: Sage Publications, Inc; 2004. Methods of meta-analysis: Correcting error and bias in research findings.

69. Rosenthal R, Rubin DB. A simple, general purpose display of magnitude of experimental effect. *J Educ Psychol.* 1982; 74(2):166–9.
70. Aguinis H, Sturman MC, Pierce CA. Comparison of three meta-analytic procedures for estimating moderating effects of categorical variables. *Organ Res Methods.* 2008; 11(1):9–34.
71. Card, NA. *Applied meta-analysis for social science research.* New York: Guilford Press; 2012. *Applied meta-analysis for social science research.*
72. Foa, EB.; Keane, TM.; Friedman, MJ.; Cohen, JA., editors. *Effective treatments for PTSD: Practice guidelines from the International Society for Traumatic Stress Studies.* Guilford Press; New York, NY US: 2009. International Society for Traumatic Stress Studies, Guideline 2: Acute interventions for children and adolescents; p. 542-545.
73. Litz, BT.; Maguen, S. Early intervention for trauma. In: Friedman, MJ.; Keane, TM.; Resick, PA., editors. *Handbook of PTSD: Science and practice.* New York: Guilford Press; 2007. p. 306-29.
74. Hobfoll SE, et al. Five essential elements of immediate and mid-term mass trauma intervention: Empirical evidence. *Psychiatry: Int Biol Process.* 2007; 70(4):283–315.

Table 1

Studies included in the meta-analysis

Study	N	Average age	% Male	Intervention package	Control	Outcome measure
Berger & Gelkopf (2009) [50]	166	11.5	52.4	Eclectic+CBT	Yes	UCLA-PTSD
Brown et al. (2006) [43]	63	10.3	54.0	Strict CBT; group	No	CPSS
Brown et al. (2006) [43]	15	10.3	24.2	Strict CBT; individual	No	CPSS
Cain et al. (2010) [51]	99	9.5	47.0	Psychological First Aid	No	CPTS-RI
Catani et al. (2009) [44]	15	11.6	62.5	Exposure	No	UPID
Catani et al. (2009) [44]	16	12.3	46.7	Relaxation	No	UPID
CATS Consortium (2010) [46]	239	11.6	43.0	Strict CBT; TFCTB	No	PTSD-RI
CATS Consortium (2010) [46]	67	11.6	55.0	CBT+Grief; Brief CBT	No	PTSD-RI
Chemtob et al. (2002) [52]	32	8.4	31.0	EMDR	No	CRI
Chemtob et al. (2002) [45]	176	8.2	38.6	Eclectic, no CBT; group	Yes	KRI
Chemtob et al. (2002) [45]	39	8.2	38.6	Eclectic, no CBT; individual	Yes	KRI
de Roos et al. (2010) [47]	26	10.2	50.0	EMDR	Yes	PTSD-RI
de Roos et al. (2010) [47]	26	10.0	61.5	Strict CBT	Yes	PTSD-RI
Gilboa-Schechtman et al. (2010) [53]	38	14.1	37.0	Exposure	Yes	CPSS
Goenjian et al. (1997) [54]	64	13.2	34.4	Eclectic+CBT	Yes	CPTSD-RI
Goenjian et al. (2005) [55]	63	16.0	40.0	Eclectic+CBT	Yes	CPTSD-RI
Jaycox et al. (2010) [48]	58	11.6	44.1	Strict CBT; CBITS	No	UPID
Jaycox et al. (2010) [48]	60	11.6	44.1	Strict CBT; TFCTB	No	UPID
Lesmana et al. (2009) [56]	226	9.8	64.6	SHAT	Yes	DSM-IV-TR
Mahmoudi-Gharaei et al. (2009) [57]	85	14.6	25.9	Eclectic+CBT	Yes	PSS
March et al. (1998) [58]	17	12.0	30.0	Strict CBT	No	CAPS-CA
Ronan & Johnston (1999) [59]	113	10.5	46.0	Eclectic+CBT	Yes	RI
Salloum & Overstreet (2008) [49]	28	10.0	67.8	Eclectic+CBT; individual	No	UCLA-PTSD
Salloum & Overstreet (2008) [49]	28	10.0	57.1	Eclectic+CBT; group	No	UCLA-PTSD
Scheeringa et al. (2011) [60•]	25	5.3	66.2	Strict CBT	Yes	PAPA PTSD
Shooshtray et al. (2008) [61]	168	15.5	47.4	Eclectic+CBT	Yes	IES-R
Taylor & Weems (2011) [62]	7	10.0	33.0	Strict CBT	No	PTSD-RI
Vijayakumar et al. (2006) [63]	135	12.2	47.0	Eclectic+CBT	Yes	CPTSD-RI

Study	N	Average age	% Male	Intervention package	Control	Outcome measure
Vila et al. (1999) [64]	47	7.5	53.0	Debriefing/Crisis Intervention	No	IES
Wolmer et al. (2003) [65]	202	8.2	44.0	Eclectic+CBT	No	CPTSD-RI
Wolmer et al. (2005) [66]	287	11.5	39.4	Eclectic+CBT	Yes	CPTSD-RI

Note. CAPS-CA=Clinician Administered PTSD Scale Child and Adolescent Version; CBITS=Cognitive-Behavioral Intervention for Trauma in Schools; CBT=Cognitive Behavioral Therapy; CPSS=Child PTSD Symptoms Scale; CPTSD-RI=Child PTSD Reaction Index; CPTS-RI=Child Post-Traumatic Stress Reaction Index; DSM-IV-TR=Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision; EMDR=Eye Movement Desensitization and Reprocessing; IES=Impact of Events Scale; IES-R=Impact of Events Scale-Revised; KRI=Kauai Recovery Index; PAPA PTSD=The Preschool Age Psychiatric Assessment PTSD module; PSS=PTSD Symptoms Scale; PTSD-RI=PTSD Reaction Index; RI=Reaction Index; SHAT=Spiritual-Hypnosis Assisted Treatment; TF-CBT=Trauma-Focused CBT; UCLA-PTSD=The University of California at Los Angeles PTSD Reaction Index; UPID=UCLA PTSD index for DSM-IV

Table 2

Meta-analytic results for treatment effects (N=2630)

Condition	K	Total N	Mean N	N-wt'd mean d	N-wt'd Var(d)	Var(e)	Resid. Var(d')	SD(d')	% Var(d) due to Var(e)	80 % Cred. Int.	
										lower	upper
Intervention group outcomes relative to controls	14	1617	115.5	.74	.39	.04	.35	.59	9.8	-.02	1.49
Pre-post intervention change only	30	2343	78.1	1.13	.54	.06	.48	.69	11.4	.25	2.02
Intervention package											
All	25	1889	75.6	1.11	.57	.06	.51	.71	11.1	.20	2.02
Eclectic with CBT	10	1052	105.2	1.07	.72	.04	.68	.82	6.1	.02	2.13
Strict CBT	9	510	56.7	1.25	.29	.09	.20	.44	30.7	.68	1.82
EMDR	2	58	29.0	2.15	.24	.23	.01	.09	96.6	2.03	2.26
Exposure	2	54	27.0	1.56	.07	.21	.00	.00	298.6	1.56	1.56
Eclectic with no CBT	2	215	107.5	.56	.01	.04	.00	.00	338.5	.56	.56
Individual vs. group/modality											
Individual treatment	14	633	45.2	1.38	.22	.11	.10	.32	52.3	.97	1.80
Group treatment	16	1710	106.8	1.04	.62	.04	.58	.76	6.9	.06	2.01
Provider											
Mental health professional	26	1825	70.2	1.19	.38	.07	.31	.56	18.1	.47	1.90
Teacher or other school personnel	4	518	129.5	.93	1.04	.03	1.00	1.00	3.4	-.35	2.21
Setting											
All	25	1871	74.8	1.18	.56	.06	.50	.71	11.5	.27	2.08
School	18	1445	80.3	1.11	.64	.06	.58	.76	9.2	.13	2.08
Health or mental health site	5	173	34.6	1.31	.24	.15	.09	.30	62.5	.93	1.69
Refugee camp	2	253	126.5	1.47	.21	.04	.17	.42	19.1	.94	2.00
Parental involvement											
Parents involved	11	855	77.7	1.30	.53	.06	.47	.69	11.9	.42	2.18
Parents not involved	19	1488	78.3	1.04	.51	.06	.45	.67	11.6	.17	1.90
Sample mean age											
5.3 – 9.83	8	846	105.8	.91	.50	.04	.46	.68	8.5	.04	1.77
10 – 11.6	14	912	65.1	1.30	.42	.08	.34	.58	18.3	.55	2.05
12 – 16	8	585	73.1	1.19	.65	.07	.58	.76	10.2	.21	2.17

Condition	K	Total N	Mean N	N-wt'd mean d	N-wt'd Var(d)	Var(e)	Resid. Var(d')	SD(d')	% Var(d) due to Var(e)	80 % Cred. Int. lower	upper
Hours of therapy											
0.5 – 4.5 hours	10	788	78.8	1.18	.30	.06	.24	.49	20.5	.56	1.80
6 – 9 hours	10	806	80.6	1.25	.31	.06	.25	.50	19.4	.61	1.90
10 – 18 hours	10	749	74.9	.95	.98	.06	.92	.96	6.2	-.27	2.18
Intervention delivery timing											
0 – 4 months	10	673	67.3	1.47	.39	.08	.31	.55	20.2	.76	2.18
4.5 – 12 months	7	780	111.4	.76	.29	.04	.25	.50	13.8	.13	1.40
35 – 69 months	13	890	68.5	1.20	.66	.07	.59	.77	10.7	.21	2.18
Nathan & Gorman Criteria											
Type 1	6	531	88.5	1.20	.29	.05	.24	.49	18.6	.57	1.83
Type 2	12	1147	95.6	1.22	.56	.05	.51	.71	9.0	.31	2.13
Type 3	12	665	55.4	.93	.63	.08	.55	.74	13.1	-.02	1.87

Note. K=number of studies included in given meta-analysis; Var(e)=variance expected due to sampling error; Resid. Var(d')=N-wt'd Var(d) – Var(e); SD(d')=square-root of Resid. Var(d'); 80 % Cred. Int. = 80 % credibility interval; 80 % of population ds are expected to fall within this interval. CBT= Cognitive Behavioral Therapy; EMDR=Eye Movement Desensitization and Reprocessing