Mini-Laparotomy for Superior Mesenteric Artery Aneurysm Due to Takayasu's Arteritis

Takatsugu Matsumoto, Mitsuru Ishizuka, Yukihiro Iso, Junji Kita, Keiichi Kubota

Department of Gastroenterological Surgery, Dokkyo Medical University, Tochigi, Japan

Superior mesenteric artery aneurysm (SMAA) is reported to be the third-most common type of visceral aneurysm (VA), accounting for 5% of all VAs. The etiology of SMAA is commonly thought to be infection, and it usually exists in the proximal part of the superior mesenteric artery, which is suitable for endovascular treatment. We herein report an extremely rare case of the distal part of SMAA caused by Takayasu's arteritis (TA), which was successfully resected using a mini-laparotomy method without impairing the intestinal blood supply. A 51-year-old woman was admitted to our hospital with sustained fever and lower back pain. Physical examination showed that she had a discrepancies in blood pressure between both arms. Contrast-enhanced whole-body computed tomography showed stenosis of the thoracic aorta and an aneurysm located in the distal part of the superior mesenteric artery. The diameter of the aneurysm was 4.5 cm. The aneurysm was resected via 4-cm mini-laparotomy, and the vascularity of the intestine was successfully preserved. The postoperative course was uneventful, and the patient was diagnosed as having TA based on both clinical and pathologic findings. Additional corticosteroid therapy was started to treat the arteritis, and at 3-month follow-up she was without critical incidents. Mini-laparotomy is a safe and less-invasive approach to treat SMAA, especially when the lesion is located in the distal part of the artery.

Key words: Superior mesenteric artery aneurysm – Takayasu's arteritis – Mini-laparotomy

Visceral aneurysms (VAs) are rare, with an incidence of 0.01% to 2% in routine autopsies.^{1,2} However, the recent widespread use of computed tomography (CT) has led to frequent incidental detection of them.³ Superior mesenteric

artery aneurysm (SMAA) is reported to be the third-most common type of VA,⁴ accounting for 5% of all VAs.⁵ Although SMAA shows few specific symptoms, several reports have revealed that it has a very high risk of rupture and mortality.^{4,5} SMAA is

Corresponding author: Takatsugu Matsumoto, MD, Department of Gastroenterological Surgery, Dokkyo Medical University, 880 Kitakobayashi, Mibu, Tochigi 321-0293, Japan.

Tel.: +81 282 87 2158; Fax: +81 282 86 6317; E-mail: takatsugu_m_475@yahoo.co.jp

Table 1 Laboratory data on admission^a

Variables	Measured value	Range
AST, U/L	16	8–38
ALT, U/L	12	4–44
ALP, U/L	128	104–338
LDH, U/L	<u>293</u>	106–211
T-Bil, mg/dL	0.6	0.1–1.0
TP, g/dL	6.6	6.5–8.1
Alb, g/dL	<u>2.5</u>	3.9–4.9
Na, mEq/L	138	135–151
K, mEq/L	4.0	3.3–4.8
Cl, mEq/L	100	98–108
UN, mg/dL	<u>6</u>	7–21
Cre, mg/dL	0.51	0.4–0.8
WBC, /μL	6800	3200–8500
Hb, g/dL	<u>7.7</u>	11.0–14.8
Plt, 10 ⁴ /μL	21.3	16.4–35.8
PT, %	75	>70
APTT, seconds	34.1	25–40
Fbg, mg/dL	359	150–400
CRP, mg/dL	<u>4.34</u>	<0.3
ESR, mm/h	<u>64</u>	3–15

Alb, albumin; ALP, alkaline phosphatase; ALT, alanine aminotransferase; APTT, activated partial thromboplastin time; AST, aspartate aminotransferase; Cre, creatinine; CRP, C-reactive protein; ESR, erythrocyte sedimentation rate; Fbg, fibrinogen; Hb, hemoglobin; LDH, lactate dehydrogenase; PLT, platelet; PT, prothrombin time; T-Bil, total bilirubin; TP, total protein; UN, urea nitrogen; WBC, white blood cells.

^aUnderlines show abnormal data.

commonly located in the proximal part of the SMA,^{4,6} which is suitable for endovascular treatment (ET).^{7,8} Here, we report a rare case of SMAA associated with Takayasu's arteritis (TA), located in the distal part of the SMA and treated by surgical resection using the mini-laparotomy method.

Case Report

A 51-year-old woman who had no significant medical history was admitted to our hospital with sustained fever and lower back pain. Physical examination revealed a bruit in her right subclavian artery and a difference in blood pressure between both arms. The back examination revealed that she had spontaneous pain on her lumbar spine, which was worsened by percussion. The abdominal examination was unremarkable. Laboratory data demonstrated that she had moderate anemia, a slightly elevated serum C-reactive protein level, and an elevated erythrocyte sedimentation rate (Table 1). She was also found to carry the following human leukocyte antigens (HLA): A24, B52, B60, CW12, and CW10. Autoantibody tests including antineu-

Fig. 1 Contrast-enhanced CT scan demonstrates an aneurysm in the superior mesenteric artery (arrow).

trophilic cytoplasmic antibody (ANCA) or rheumatoid factor were negative. Her serum complement levels were normal. The result of blood culture revealed that she had bacteremia resulting from *Streptococcus agalactiae* infection. However, transthoracic and transesophageal echocardiography demonstrated that there were no signs of vegetation in the heart or major vessels. Magnetic resonance imaging (MRI) showed edematous changes in the lumbar disks. Therefore, she was first diagnosed as having purulent diskitis and secondary bacteremia. Then, she was treated with ampicillin/sulbactam for 8 weeks to treat diskitis and the bacteremia. During the treatment, the back pain was gradually improved, but the fever still continued. To seek the cause of sustained fever, contrast-enhanced whole-body CT was performed, and it was revealed that her thoracic aorta was mildly stenosed and an aneurysm in the distal part of the SMA was incidentally detected (Fig. 1). Angiography also revealed an aneurysm in the distal part of the SMA (Fig. 2). No other arterial abnormality was revealed by both CT and angiography.

Because the aneurysm was very close to the marginal artery of the ileum, we selected surgical resection rather than endovascular treatment (ET) to preserve the vascularity of the ileum.

A mini-laparotomy was performed using a 4-cm incision with the Gelpert's double-ring wound retractor (Alexis, Applied Medical, Rancho Santa Margarita, California), and an aneurysmal mass with 4.5-cm diameter was found in the distal part of the SMA (Fig. 3). We could resect the aneurysm and preserve the ileal vascularity by performing careful dissection from the surrounding mesenteric tissue.

Fig. 2 Selective angiography of the SMA reveals that there is a saccular aneurysm in the distal part of the SMA (arrow).

The postoperative course was uneventful. She was diagnosed as having TA based on clinical findings in accordance with the American College of Rheumatology criteria for the classification of TA.⁹ The resected aneurysmal wall was markedly thickened macroscopically, and microscopic examinations showed that the inflammatory cells diffusely infiltrated the media, which were compatible with

Fig. 3 A 3.5 × 2.5 × 4.5-cm aneurysmal mass is located in the distal branch of the SMA (arrow).

Fig. 4 (A) Microscopic findings show marked thickness of the aneurysmal wall [H&E staining (×40)] and (B) infiltration of lymphocytes, plasma cells, and histiocytes [H&E staining (×100)].

TA (Fig. 4). After surgery, she was administered methylprednisolone, 5 mg/d, to treat TA. Consequently, the back pain and fever completely resolved, and the serum inflammatory markers also returned to normal. At follow-up 3 months after surgery, she had no postoperative complications. She is now treated as an outpatient by the rheumatologist and continues to receive methylprednisolone 5 mg/d.

Discussion

The etiology of SMAA is commonly thought to be infection.¹⁰ A variety of disorders are responsible for it, including atherosclerosis, medial dysplasia, or vasculitis.^{4,11} In the present case, the pathologic features of the resected specimen suggested that TA

was a major cause of the aneurysmal formation. TA is an uncommon chronic vasculitis first reported by Takayasu *et al* in 1908.¹² Although the pathogenesis of TA is not fully understood, cell-mediated autoimmune mechanisms are thought to be of primary importance. Patients with TA present initially with fever and anorexia, followed by multiple arterial occlusive symptoms depending on the location of disease involvement.¹³ The symptoms are sometimes vague when the activity of the TA is mild and chronic.

In this report, the patient was diagnosed as having TA because she fulfilled 3 of 6 diagnostic criteria for TA⁹: the presence of the right subclavian bruit, discrepancies in blood pressure between both arms, and arteriographic abnormalities. As TA mainly affects the aorta and its primary branches,¹⁴ and the lesions are usually stenotic,¹⁵ it is rare for TA to cause the peripheral arterial aneurysm. A search of the Medline database using the key words "Takayasu's arteritis," "visceral aneurysm," and "mesenteric aneurysm," revealed that this is the first case of peripheral SMAA associated with TA (www.pubmed.gov in the public domain).

In common, surgical treatment of VA includes aneurysmal exclusion or excision and, if necessary, reconstruction of the vessels or resection of the ischemic organ.^{3,4} Although a laparoscopic approach is reported to be feasible, especially for splenic artery aneurysm,^{2,16,17} the incision to extract the specimen is essential. In this case, the aneurysm was 4.5 cm in diameter, and intracorporeal mobilization was unnecessary because the lesion was located in the distal part of the SMA. Thus, we considered that aneurysmal excision could be performed via a single 4-cm incision with a wound retractor. In addition, by using mini-laparotomy, the surgeon could confirm the arterial pulsation at their finger and control unexpected bleeding more easily and safely.

ET is a choice of treatment for VA.^{18,19} ET contains transcatheter embolization and covered stent placement, which can be performed under local anesthesia without abdominal incision. Therefore, it is widely adopted for the treatment of VAs including SMAA, especially for patients with poor surgical risk. Morbidity of surgical treatment is reported to be 0% to 5%, with excellent long-term durability.¹⁶ However, success rates of ET in the literature have varied from 75% to 100%, with morbidity rates ranging from 14% to 25%.⁵ Incomplete aneurysm exclusion requires further intervention or surgical treatment. In addition, it is reported that ET is not feasible when the lesion has several collateral arteries, a large diameter, or a short distance to the visceral organ.^{8,20}

In this case, the aneurysm was located in the near side of both marginal arteries and intestinal wall, and the patient had no particular surgical risk.

Although the natural history of VA is still unclear, several series have documented that patients treated for VA have a good prognosis.^{4,6-8,21} For example, Marone *et al* have reported that the overall 10-year survival rate after the treatment of VA is 68%.⁸

Conclusion

We have described the first case of peripheral SMAA associated with Takayasu's arteritis that was successfully resected using a mini-laparotomy method without impairing the vascularity of the ileum.

Mini-laparotomy is a safe and less-invasive approach to treat SMAA, especially when the lesion is located in the distal part of the artery.

Acknowledgments

We have no conflicts of interest to declare. We received no funding or grant support for this study.

References

1. Carr SC, Mahvi DM, Hoch JR, Archer CW, Turnipseed WD. Visceral artery aneurysm rupture. *J Vasc Surg* 2001;**33**(4):806-811
2. Carmeci C, McClenathan J. Visceral artery aneurysms as seen in a community hospital. *Am J Surg* 2000;**179**(6):486-489
3. Panayiotopoulos YP, Assadourian R, Taylor PR. Aneurysms of the visceral and renal arteries. *Ann R Coll Surg Engl* 1996;**78**(5):412-419
4. Van Petersen A, Meerwaldt R, Geelkerken R, Zeebregts C. Surgical options for the management of visceral artery aneurysms. *J Cardiovasc Surg (Torino)* 2011;**52**(3):333-343
5. Gehlen JM, Heeren PA, Verhagen PF, Peppelenbosch AG. Visceral artery aneurysms. *Vasc Endovascular Surg* 2011;**45**(8):681-687
6. Stone WM, Abbas M, Cherry KJ, Fowl RJ, Gloviczki P. Superior mesenteric artery aneurysms: is presence an indication for intervention? *J Vasc Surg* 2002;**36**(2):234-237
7. Sachdev U, Baril DT, Ellozy SH, Lookstein RA, Silverberg D, Jacobs TS *et al*. Management of aneurysms involving branches of the celiac and superior mesenteric arteries: a comparison of surgical and endovascular therapy. *J Vasc Surg* 2006;**44**(4):718-724
8. Marone EM, Mascia D, Kahlberg A, Brioschi C, Tshomba Y, Chiesa R. Is open repair still the gold standard in visceral artery aneurysm management? *Ann Vasc Surg* 2011;**25**(7):936-946

9. Arend WP, Michel BA, Bloch DA, Hunder GG, Calabrese LH, Edworthy SM *et al.* The American College of Rheumatology 1990 criteria for the classification of Takayasu arteritis. *Arthritis Rheum* 1990;**33**(8):1129–1134
10. Friedman SG, Pogo GJ, Moccio CG. Mycotic aneurysm of the superior mesenteric artery. *J Vasc Surg* 1987;**6**(1):87–90
11. de Leeuw K, Goorhuis JF, Tielliu IF, Symoens S, Malfait F, de Paepe A *et al.* Superior mesenteric artery aneurysm in a 9-year-old boy with classical Ehlers-Danlos syndrome. *Am J Med Genet A* 2012;**158A**(3):626–629
12. Numano F. Introductory remarks for this special issue on Takayasu arteritis. *Heart Vessels* 1992;**7**(1):3–5
13. Kerr GS, Hallahan CW, Giordano J, Leavitt RY, Fauci AS, Rottem M *et al.* Takayasu arteritis. *Ann Int Med* 1994;**120**(11):919–929
14. Hata A, Noda M, Moriwaki R, Numano F. Angiographic findings of Takayasu arteritis: new classification. *Int J Cardiol* 1996;**54**(suppl):S155–S163
15. Numano F, Okawara M, Inomata H, Kobayashi Y. Takayasu's arteritis. *Lancet* 2000;**356**(9234):1023–1025
16. Pulli R, Dorigo W, Troisi N, Pratesi G, Innocenti AA, Pratesi C. Surgical treatment of visceral artery aneurysms: a 25-year experience. *J Vasc Surg* 2008;**48**(2):334–342
17. Reardon PR, Ota E, Craig ES, Matthews BD, Reardon MJ. Laparoscopic resection of splenic artery aneurysms. *Surg Endosc* 2005;**19**(4):488–493
18. Mendonca CT, Weingartner J, de Carvalho CA, Costa DS. Endovascular treatment of contained rupture of a superior mesenteric artery aneurysm resulting from neurofibromatosis type I. *J Vasc Surg* 2010;**51**(2):461–464
19. Ferrero E, Ferri M, Viazzo A, Robaldo A, Carbonatto P, Pecchio A, *et al.* Visceral artery aneurysms, an experience on 32 cases in a single center: treatment from surgery to multilayer stent. *Ann Vasc Surg* 2011;**25**(7):923–935
20. Ferrero E, Viazzo A, Ferri M, Robaldo A, Piazza S, Berardi G *et al.* Management and urgent repair of ruptured visceral artery aneurysms. *Ann Vasc Surg* 2011;**25**(7):981, 981.e7–981.e11
21. Huang Y-K, Hsieh H-C, Tsai F-C, Chang S-H, Lu M-S, Ko P-J. Visceral artery aneurysm: risk factor analysis and therapeutic opinion. *Eur J Vasc Endovasc Surg* 2007;**33**(3):293–301