Published in final edited form as:

Arch Toxicol. 2015 March ; 89(3): 459–483. doi:10.1007/s00204-014-1264-0.

Mechanisms of Acute Kidney Injury Induced by Experimental *Lonomia obliqua* Envenomation

Markus Berger¹, Lucélia Santi², Walter O. Beys-da-Silva², Fabrício Marcus Silva Oliveira³, Marcelo Vidigal Caliar³, John R. Yates 3rd², Maria Aparecida Ribeiro⁴, and Jorge Almeida Guimarães^{1,*}

¹Laboratório de Bioquímica Farmacológica, Centro de Biotecnologia, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, RS, Brazil

²Department of Chemical Physiology, The Scripps Research Institute, La Jolla, CA, United States

³Laboratório de Protozooses, Departamento de Patologia Geral, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, MG, Brazil

⁴Laboratório de Fisiologia Renal, Departamento de Fisiologia e Biofísica, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, MG, Brazil

Abstract

Background—*Lonomia obliqua* caterpillar envenomation causes acute kidney injury (AKI), which can be responsible for its deadly actions. This study evaluates the possible mechanisms involved in the pathogenesis of renal dysfunction.

Methods—To characterize *L. obliqua* venom effects we subcutaneously injected rats and examined renal functional, morphological and biochemical parameters at several time points. We also performed discovery based proteomic analysis to measure protein expression to identify molecular pathways of renal disease.

Results—*L. obliqua* envenomation causes acute tubular necrosis, which is associated with renal inflammation; formation of hematic casts, resulting from intravascular hemolysis; increase in vascular permeability and fibrosis. The dilation of Bowman's space and glomerular tuft is related to fluid leakage and intra-glomerular fibrin deposition, respectively, since tissue factor procoagulant activity increases in the kidney. Systemic hypotension also contributes to these alterations and to the sudden loss of basic renal functions, including filtration and excretion capacities, urinary concentration and maintenance of fluid homeostasis. In addition, envenomed kidneys increases expression of proteins involved in cell stress, inflammation, tissue injury, heme-induced oxidative stress, coagulation and complement system activation. Finally, the localization of the venom in renal tissue agrees with morphological and functional alterations, suggesting also a direct nephrotoxic activity.

*Corresponding author: Prof. Dr. Jorge Almeida Guimarães, Laboratório de Bioquímica Farmacológica, Centro de Biotecnologia (UFRGS). Av. Bento Gonçalves, 9500, Cep 91501-970, Porto Alegre, RS, Brasil. Tel.: +55 51 33086068. Fax +55 51 33087309. guimar@cbiot.ufrgs.br.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Conclusions—Mechanisms of *L. obliqua*-induced AKI are complex involving mainly glomerular and tubular functional impairment and vascular alterations. These results are important to understand the mechanisms of renal injury and may suggest more efficient ways to prevent or attenuate the pathology of *Lonomia*'s envenomation.

Keywords

Venom; *Lonomia*; Renal; Acute Kidney Injury; Nephrotoxicity; Acute Tubular Necrosis

1. Introduction

Accidents caused by venomous animals (mainly snakes, spiders, scorpions, bees, wasps and caterpillars) are a costly and critically important public health problem. Despite of this, public health authorities, nationally and internationally, have given little attention to this problem worldwide (Warrell, 2010; Williams *et al.*, 2010). Consequentially, the morbidity and mortality associated with envenomation cases produce a great impact on the population and healthcare systems. One of the most important and lethal effects of these animal venoms is nephrotoxicity and a broad clinical spectrum of renal function impairment has been reported in human and experimental models of envenomation (Sitprija, 2006; Berger *et al.*, 2012). As kidneys are highly vascularized organs and have the ability to concentrate substances into urine they are particularly susceptible to venom toxins. The most common clinical renal manifestation seen in human patients is acute tubular necrosis, but all renal structures may be involved. Thus, the occurrences of acute tubulointerstitial nephritis, renal cortical necrosis, mesangiolysis, vasculitis, glomerulonephritis, proteinuria, hematuria, hemoglobinuria and myoglobinuria have also been described (Sitprija, 2006).

Lonomia obliqua caterpillars are well known in southern Brazil where they cause severe hemorrhagic syndrome characterized by perturbed coagulation, ecchymosis, acute kidney injury (AKI) and generalized hemorrhage. Since the 1980's there has been a considerable increase in the number of hemorrhagic incidents in rural areas of the southernmost Brazilian states of Rio Grande do Sul, Santa Catarina and Paraná. The origin of this epidemic is not clear, but can be partially attributed to recent deforestation, as well as to a progressive reduction in the number of natural predators. Usually, accidents occur when the victim, unknowingly leans against a tree trunk containing hundreds of caterpillars and comes into contact with the caterpillar's venomous bristles, which are chitinous evaginations of cuticle. Often, the caterpillar is crushed, the bristles are broken and venomous secretions, including hemolymph, penetrate the human skin (Veiga *et al.*, 2001). The venom is composed of several active constituents with procoagulant, fibrinogenolytic, proteolytic and hemolytic activities (Pinto *et al.*, 2010). Although consumptive coagulopathy secondary to intravascular disseminated coagulation is commonly observed in human and experimental animals, AKI is the leading cause of death from *L. obliqua* envenomation (Zannin *et al.*, 2003; Gamborgi *et al.*, 2006; Berger *et al.*, 2010).

Early in the 1980's, the first registered cases of *L. obliqua*-induced hemorrhagic syndrome indicated that 18 % of envenomed patients had developed AKI. The mortality rate in these patients reached 50 % (Duarte *et al.*, 1990; Duarte *et al.*, 1994). However, a lower incidence

(5.2 %) was observed in the Brazilian state of Rio Grande do Sul from 1989 to 1995, when only 15 of 286 envenomed patients developed AKI (Duarte *et al.*, 1997). Another study analyzing a larger group of 2,067 envenomed patients in the Santa Catarina state in Brazil (from 1989 to 2003) reported that 39 victims (1.9 %) developed AKI (with serum creatinine levels 1.5 mg/dL). Eleven (32 %) of these patients were treated with dialysis and four (10.3 %) developed chronic renal injury (CRI). All victims with AKI presented concomitantly coagulation disturbances and hematuria and/or hemoglobinuria. Seven deaths (4 %) occurred during this period (Gamborgi *et al.*, 2006). An important conclusion of this work is that, even after the introduction of antivenom therapy (with anti-*Lonomia* serum) in 1995, there was no reduction in the incidence of AKI, despite the significant decrease in the number of deaths and patients who developed CRI (Gamborgi *et al.*, 2006). In fact, recently we observe that antivenom treatment was able to reduce creatinine and urea levels of rats only if administered 2 h post venom injection. Serotherapy after 6 h of envenomation fails to neutralize the rising in biochemical markers of renal injury (Berger *et al.*, 2013). Since the average time elapsed between the contact of a person with caterpillars and an appropriate medical care can vary from 19 to 37 h (Zannin *et al.*, 2003), it seems imperative to achieve a better understanding of the mechanisms involved in venom-induced AKI. It is clear that having such knowledge available it will then make possible to develop new efficient treatments in order to avoid or at least to reduce the progression of renal disease in *Lonomia*'s and other kind of animal envenoming.

The risk of conducting early renal biopsies in human patients, due to coagulation disturbances inherent to the envenomation, has made it difficult to analyze the acute kidney pathological alterations. There are only two case reports in the literature describing alterations of *Lonomia*-induced AKI. The main findings were oliguria, high levels of serum creatinine, thickening of the Bowman's capsule, focal tubular atrophy and acute tubular necrosis (Burdmann *et al.*, 1996; Fan *et al.*, 1998). Since no experimental studies were available until now, the contribution of several factors possibly associated with AKI, such as hemodynamic changes, vascular permeability alterations, hemolysis, tubular obstruction, glomerular fibrin deposition and even a direct venom nephrotoxicity, remain obscure in *Lonomia*-induced AKI.

In an attempt to better understand the progression of renal disease commonly observed after the contact with *L. obliqua* caterpillars, we have focused on the action of venom in the kidney. Therefore, an experimental rat model was used in order to characterize changes in renal function, tubular hydroelectrolytic transport, histopathology and hemodynamics.

2. Materials and Methods

2.1 Reagents

Evans blue dye, purified coagulation factors (VII, IX and X) and molecular weight standards used in SDS-PAGE and western-blot were purchased from Sigma-Aldrich (Saint Louis, MO, USA). Chromogenic substrate for factor Xa (S2222, Bz-Ile-Glu-Gly-Arg-pNa) was obtained from Chromogenix (Milano, Italy). Ketamine and xylazine were from Syntec, São Paulo, Brazil. *L. obliqua* antivenom (antilonomic serum - ALS), provided by the Butantan Institute (São Paulo, Brazil), was used as primary antibody for the detection of toxins in

urine and renal tissue. ALS is a horse-derived concentrate of purified polyclonal antibodies (IgG) that had been raised against *L. obliqua* bristle extract (Rocha-Campos *et al.*, 2001). Nonspecific background staining in immunohistochemical reactions was blocked using Ultra V Block reagent (Thermo Fisher Scientific, Waltham, MA, USA).

2.2 Venom

L. obliqua caterpillars were kindly provided by the Centro de Informações Toxicológicas (CIT), Porto Alegre, Rio Grande do Sul, Brazil. The specimens used in this study were collected in the cities of Bom Princípio and Progresso, both located in Rio Grande do Sul, Brazil. *L. obliqua* venom was obtained by homogenizing the bristles in cold phosphate-buffered saline (PBS), pH 7.4, as previously described (Berger *et al.*, 2010). The venom obtained following this procedure was designated as *Lonomia obliqua* Bristle Extract (LOBE). The protein content of the LOBE samples was determined using a BCA assay kit (Pierce, Rockford, USA) and the aliquots were stored at -80°C prior to use. The total number of caterpillars used for bristle extract preparation was 124 specimens and the protein concentration of the LOBE samples was 4.10 mg/mL. The total amount of venom extracted per caterpillar was 2.4 mg. All of the LOBE samples had similar *in vitro* pro-coagulant activities and the protein pattern for each sample as monitored by SDS-PAGE and gel filtration chromatography (Pinto *et al.*, 2006; Berger *et al.*, 2010) were also similar.

2.3 Ethical statements

All procedures involving animals were carried out in accordance with the Guiding Principles for the Use of Animals in Toxicology (International Society of Toxicology, <http://www.toxicology.org>) and the Brazilian College of Animal Experimentation (COBEA). The experimental protocol was approved by the ethical committee on research animal care of the Federal University of Rio Grande do Sul, Brazil (register number **2008177/2009**) and by the Institute's Animal Ethics Committee of the Federal University of Minas Gerais, Brazil (protocol **177/2008**).

2.4 Experimental protocol

2.4.1 Animals—Adult male Wistar rats, weighing 250–300 g, were supplied by the central animal facility of our institution. They were housed in standard conditions within a temperature controlled room ($22\text{--}23^{\circ}\text{C}$, on a 12 h light/dark cycle, with the lights on at 7:00 am) and had free access to water and food.

2.4.2 Selection of the venom dose—The severity of the natural envenoming is related mainly to the number of caterpillars involved as well as to the intensity of the exposure, since the venom is present not only in the caterpillar's bristles but also in their skin and hemolymph (Veiga *et al.*, 2001). Considering that accidents with medical importance involves contact with a colony containing at least 40 to 50 caterpillars (Gamborgi *et al.*, 2006) and that during venom extraction, after removal of all spicules, each caterpillar produces approximately 2.4 mg of venom, the total amount of venom injected in an individual weighing 70 kg can reach up to 1.4 – 1.7 mg/kg. In fact, these doses were calculated based on an artificial method of venom extraction, in which the caterpillar's bristle was macerated in a solution buffer. Thus, in a real envenomation situation the total

amount of venom transferred is probably lower than the amount calculated. In an attempt to reproduce the clinical conditions observed in a real envenomation we selected doses of 1.0 and 1.5 mg/kg injected subcutaneously into rats. These doses were also in accordance with the amount of venom used in other studies to induce coagulopathy and test the efficacy of antilonomic serum (Berger *et al.*, 2010; Dias da Silva *et al.*, 1996; Rocha-Campos *et al.*, 2001).

2.4.3 Venom administration—To follow the time course of kidney pathophysiological alterations, we used an experimental model of envenomation in rats. For this purpose, animals were divided into three groups ($n = 6/\text{group}$): The control animals (CTRL) were injected subcutaneously (s.c.) with 100 μL of sterile PBS solution and the experimental animals received a s.c injection containing 1.0 mg or 1.5 mg of LOBE per kg of body weight in a final volume of 100 μL . Immediately after treatments, the animals were distributed individually into metabolic cages, allowing quantitative urine collections and measurement of water intake. At several time points post-venom injection (2, 6, 12, 24, 48 and 96 h), blood, urine and kidneys were obtained for biochemical, histopathological and immunohistochemical analyses.

2.5 Sample preparation

Blood was collected in conscious rats through the caudal vein in 1:10 (v/v) 3.8 % trisodium citrate. Plasma was obtained by centrifugation at $1500 \times g$ for 10 min and stored at -80°C prior to use. Urine samples were also centrifuged at $2500 \times g$ for 5 min and the supernatants stored at the same conditions. After blood collection, animals from the different groups were anesthetized by intraperitoneal (i.p.) injection of a mixture of ketamine (75 mg/kg) and xylazine (10 mg/kg). Then, an intracardiac perfusion was performed through the left ventricle with PBS solution and a circulatory circuit was opened by an incision in the right atrium, to ensure the elimination of intravascular blood. Immediately after perfusion, kidneys were quickly removed. One of them was fixed for histological analysis and the other was frozen in liquid nitrogen and stored at -80°C for the measurement of tissue factor activity and proteomic analysis.

2.6 Biochemical measurements

Urinary and plasma levels of creatinine, urinary γ -glutamyl transferase (γ -GT) activity and proteinuria were determined by spectrophotometry (Turner SP-830 plus Barnstead, Dubuque, Iowa, USA) using commercially available kits (BioClin/Quibasa, Belo Horizonte, Brazil). Plasma and urinary concentrations of Na^+ and K^+ were measured by flame photometry (CELM 180; Belo Horizonte, Minas Gerais, Brazil). Osmolality was determined in plasma and urine samples by cryoscopic osmometry using a Micro-Osmometer 3320 (Advanced Instruments, Norwood, Massachusetts, USA). Urine proteins were also analyzed by gel electrophoresis which was performed according to Laemmli (1970). Urine samples from animals of different times post-venom injection were diluted ($10 \times$) and aliquots of 10 μL were submitted to SDS-PAGE on 8-20 % gradient gels under reducing conditions. Toxins excreted in urine were detected by Western-blot as previously described (Pinto *et al.*, 2006). Aliquots containing 50 μg of protein were separated by SDS-PAGE and transferred to PVDF membranes. Toxins were recognized using as a primary antibody an equine anti-

LOBE IgG (ALS) diluted 1:100 and as a secondary antibody a peroxidase-labeled anti-horse IgG diluted 1:1000.

2.7 Renal function parameters

At the time intervals mentioned above the following renal function parameters were determined: Glomerular filtration rate (GFR), osmolar clearance (C_{osm}), water free clearance ($C_{\text{H}_2\text{O}}$), fractional water excretion ($\text{FE}_{\text{H}_2\text{O}}$), fractional sodium excretion (FE_{Na^+}) and fractional potassium excretion (FE_{K^+}). GFR (expressed as mL/min/100 g of body weight) was estimated by the creatinine clearance (C_{cr}), using the standard formula: $C_{\text{cr}} = U_{\text{cr}} \cdot V / P_{\text{cr}}$, where U_{cr} is the urinary creatinine concentration, V is the urinary output and P_{cr} is the plasma creatinine concentration. C_{osm} (expressed as mL/min) was calculated as $C_{\text{osm}} = U_{\text{osm}} / P_{\text{osm}} \cdot V$, whereas U_{osm} and P_{osm} are the urinary and plasma osmolalities, respectively. Values of $C_{\text{H}_2\text{O}}$ (mL/min) and $\text{FE}_{\text{H}_2\text{O}}$ (%) were obtained respectively from the equations: $C_{\text{H}_2\text{O}} = V - C_{\text{osm}}$ and $\text{FE}_{\text{H}_2\text{O}} = V / \text{GFR} \cdot 100$. FE_{Na^+} and FE_{K^+} (expressed as %) were calculated according to the equation: $\text{FE} = \text{UE} / \text{PF} \cdot 100$. UE represents the urinary excretion of each ion and PF is the amount filtered in plasma (both expressed as nmol/min).

2.8 Hemodynamic parameters

Systemic arterial pressure was measured in conscious rats by an indirect tail-cuff method using an electrosphygmomanometer (LE 5001, Harvard Apparatus, Holliston, Massachusetts, USA) combined with a pneumatic pulse transducer/amplifier, which provides output signals proportional to cuff pressure and amplified Korotkoff sounds. Three consecutive readings of mean arterial pressure and heart rate were recorded before blood collection for each animal in each time post-venom injection.

2.9 Renal vascular permeability

The extravasation of Evans blue dye into the kidney was used as an index of increased vascular permeability (Pompermayer *et al.*, 2005). Rats received Evans blue dye (30 mg/kg) intravenously (1 mL/kg) via caudal vein 10 min prior to LOBE (1.5 mg/kg, s.c.) or PBS (100 μL , s.c.) injection. After 12 or 24 h, animals were anesthetized and perfused as described above to remove the intravascular Evans Blue. Then, the kidneys were quickly removed, weighed and allowed to dry for 24 h at 40 °C. The dry weight was determined and Evans blue dye extracted in 2.5 mL of 1 % formamide (48 h at 40 °C). The absorbance of extracted solution was measured in triplicate using a microplate reader spectrophotometer (SpectraMAX, Molecular Devices Co., Sunnyvale, USA) and the amounts of Evans blue dye were calculated by a standard curve made with known concentrations of Evans Blue. Results are presented as the amount of Evans blue dye extravasated (μg) per 100 mg of kidney tissue.

2.10 Renal tissue factor activity

Renal tissue factor (TF) was measured indirectly based on its ability to form a complex with factor VIIa (TF/FVIIa) to activate factor IX and X (Morrissey, 1995). Briefly, the kidneys were collected as described above, homogenized in cold PBS solution containing 1 % Triton X-100 and centrifuged at 9500 $\times$ g for 15 min. Samples of supernatants (with 10 μg of

protein) were incubated with a concentrate mixture of FVII+FIX+FX (total of 15 µg) in 20 mM Tris-HCl, pH 7.4 containing 10 mM of CaCl₂ for 10 min at 37 °C. Activated factor Xa (FXa) produced during the reaction was detected by the addition of a specific chromogenic substrate (0.2 mM S2222). The kinetics of *p*-nitroaniline release was monitored at 405 nm for 30 min in a final volume of 100 µL using a microplate reader spectrophotometer (SpectraMAX, Molecular Devices Co., Sunnyvale, USA). Each sample was measured in triplicate and results expressed as µmol of FXa generated per min per mg of kidney tissue.

2.11 Histology and immunohistochemistry

For renal histopathology, kidneys were collected as described above, sectioned sagittally and fixed in 10 % buffered formaldehyde, pH 7.2. After processing in alcohol and xylol, the organs were included in paraffin and 4 µm thick sections were obtained and stained with Haematoxylin and Eosin (H&E), Periodic Acid-Schiff (PAS) or Picro-Sirius reagent.

For immunohistochemical analysis of venom distribution in renal tissue, the kidney sections were deparaffinized and treated with 10 % hydrogen peroxide solution in methanol for 15 min to block endogenous peroxidase activity. Nonspecific binding sites were blocked with Ultra V Block reagent followed by incubation overnight at 4 °C with an equine anti-LOBE IgG (ALS) diluted 1:250 in PBS. After a washing step, the sections were incubated with the secondary antibody (a biotinylated anti-horse IgG produced in goat, diluted 1:100) and streptavidin diluted 1:100. The chromogenic reaction was developed by incubating the sections with 0.05 % diaminobenzidine solution and 0.2 % hydrogen peroxide. The progress of the reaction was monitored by light microscopy and stopped by washing the slides. Finally, the sections were counterstained with diluted Harris's haematoxylin. Control reactions were done by incubating kidney sections from non-envenomed rats with equine anti-LOBE IgG (ALS) under the same conditions as described above. No positive reactions were observed in these sections. Some sections were also used as negative controls. In these cases, the primary antibody was substituted by PBS.

2.12 Morphometric analysis

Glomerular morphological alterations were quantified by computer-assisted morphometric analysis using a method previously described (Caliari, 1997). Briefly, images with 40 × magnification of 30 glomeruli from each animal of different groups were randomly digitalized using a JVC TK1270/RGB microcamera (Tokyo, Japan). The KS300 software coupled to a Carl Zeiss image analyzer (Carl Zeiss, Oberkochen, Germany) was used to measure the total area (µm²) of each glomerulus, glomerular tuft and Bowman's space. The pixels of whole glomerulus and those corresponding to the glomerular tuft were selected and used for the generation of a binary image and subsequent calculation of the corresponding areas. The area of Bowman's space was obtained by the difference between the whole glomerular area and tuft area. The regions of positive immunohistochemical reaction (expressed as µm²) were also measured and used as a parameter to quantify venom distribution in the renal tissue.

2.13 Renal tissue proteomics

2.13.1 Sample preparation for mass spectrometry—The kidneys from controls and envenomed animals (at 24 h post-venom injection) were collected as described above, homogenized in cold PBS solution containing 1 % Triton X-100 with protease inhibitor cocktail (Halt protease inhibitor cocktail, Thermo, Rockford, USA) and centrifuged at $9500 \times g$ for 15 min. The resulting supernatants were completely lyophilized and stored at -80°C until use. Lyophilized samples were resuspended in water and precipitated using methanol/chloroform protocol. After precipitation samples were dried at room temperature and resuspended in 8 M urea. Each sample containing 100 μg of protein was reduced with 5 mM tris-2-carboxyethyl-phosphine (TCEP) at room temperature for 20 min and alkylated with 10 mM iodoacetamide at room temperature in the dark for 20 min. After reduction and alkylation, proteins were digested with 2 μg of trypsin (Promega, Madison, WI) by incubation at 37°C during 16 h. Samples were freeze-dried at -80°C and, after thaw, formic acid to a final concentration of 5% was added. Samples were centrifuged at 14,000 rpm for 20 min and the supernatant was collected and stored at -80°C .

2.13.2 MudPIT—The protein digest was pressure-loaded into a 250- μm i.d capillary packed with 2.5 cm of 5- μm Luna strong cation exchanger (SCX) (Whatman, Clifton, NJ) followed by 2 cm of 3- μm Aqua C18 reversed phase (RP) (Phenomenex, Ventura, CA) with a 1 μm frit. The column was washed with buffer containing 95 % water, 5 % acetonitrile, and 0.1 % formic acid. After washing, a 100- μm i.d. capillary with a 5- μm pulled tip packed with 11 cm of 3- μm Aqua C18 resin (Phenomenex, Ventura, CA) was attached via a union according to Klein et al. (2012). The entire split-column was placed in line with an Agilent 1100 quaternary HPLC (Palo Alto, CA) and analyzed using a modified 11-step separation as described previously (Washburn *et al.*, 2001). The buffer solutions used were 5 % acetonitrile, 0.1 % formic acid (Buffer A), 80 % acetonitrile, 0.1 % formic acid (Buffer B), and 500 mM ammonium acetate, 5% acetonitrile and 0.1% formic acid (Buffer C). Step 1 consisted of a 70-min gradient from 0–100 % (vol/vol) buffer B. Steps 2–10 had a similar profile with the following changes: 5-min in 100 % (vol/vol) buffer A, 3-min in X % (vol/vol) buffer C, a 6-min gradient from 0 to 15 % (vol/vol) buffer B, and a 85-min gradient from 15–100 % (vol/vol) buffer B. The 3-min buffer C percentages (X) were 10, 20, 30, 40, 50, 60, 70, 80, 90, and 100 % (vol/vol), respectively, for the 10-step analysis.

2.13.3 LTQ-Orbitrap—As peptides eluted from the microcapillary column, they were electrosprayed directly into a LTQ-Orbitrap (Thermo Fisher) with the application of a distal 2.4-kV spray voltage. Full MS spectra were acquired in profile mode, with a mass range of 400 – 1600 in the Orbitrap analyzer with resolution set at 60,000 followed by 10 data-dependent MS/MS spectra at 35 % normalized collision energy was repeated continuously throughout each step of the multidimensional separation. Minimal signal for fragmentation was set to 1,000. Dynamic exclusion was enabled with a repeat count of 1, duration of 30.00 s, list size of 500, exclusion duration of 180.00 s, and exclusion mass with high/low of 1.5 m/z. Application of mass spectrometer scan functions and HPLC solvent gradients were controlled by the Xcalibur data system.

2.13.4 Analysis of tandem mass spectra—MS/MS spectra were analyzed using the following software analysis protocol. Protein identification and quantification analysis were done with Integrated Proteomics Pipeline (IP2, Integrated Proteomics Applications, Inc. www.integratedproteomics.com/). Tandem mass spectra were extracted into ms2 files from raw files using RawExtract 1.9.9 (McDonald *et al.*, 2004) and were searched using ProLuCID algorithm (Xu *et al.*, 2006). MS/MS spectra remaining after filtering were searched with the ProLuCID algorithm against the EBI-IPI_rat_3.30_06-28-2007 concatenated to a decoy data base in which the sequence for each entry in the original data base was reversed (Peng *et al.*, 2003). Searches were performed with cysteine carbamidomethylation as a fixed modification. ProLuCID results were assembled and filtered using the DTASelect program (Tabb *et al.*, 2002) using two SEQUEST (Eng *et al.*, 1994) defined parameters: the cross-correlation score (XCorr) and normalized difference in cross-correlation scores (DeltaCN) to achieve a false discovery rate of 1%. The following parameters were used to filter the peptide candidates: -p 1 -y 1 --trypstat --fpf 0.01 --dm. Also, we used 50ppm as precursor tolerance, fragment mass tolerance of 600ppm, 3 as number of isotopic peaks and unlimited missed cleavages were allowed.

2.13.5 Assessing differential expression and exclusive proteins /

Bioinformatic tools—The software PatternLab (Carvalho *et al.* 2008, 2012) was used to identify exclusive proteins found in the control and treated conditions. PatternLab's Approximately Area Proportional Venn Diagram (AAPVD) module was used for pinpointing proteins uniquely identified in each condition. The following parameter was used: proteins that were not detected in at least 2 out of 3 runs per condition were not considered. G-test spectral counting quantitation was performed in a pair-wise comparison between the two groups, as previously reported (Ambatipudi *et al.* 2009). Proteins are considered differentially expressed with $P < 0.1$. Ingenuity Pathway Analysis tool (Ingenuity Systems; <http://www.ingenuity.com>) was used to generate functional annotations of identified proteins in known molecular pathways and/or biologic function in disease. The significance of the canonical pathways and biologic function defined by identified proteins was measured in two ways: (1) The number of proteins identified from the data set that map to a known pathway or function in disease and (2) a P value (Fisher exact test) determining the probability that the association between the proteins in the data set and the canonical pathway or function in disease is explained by chance alone.

2.14 Statistical Analyses

Results are expressed as mean $\pm$ SE. When appropriate, statistical comparisons were done by using one or two-way analysis of variance followed by the Bonferroni's test. A P value of less than 0.05 was chosen to establish significance. Statistical analysis was performed using GraphPad Prism (GraphPad Software Inc., San Diego, CA, USA).

3. Results

3.1 Renal function

To follow renal alterations in rats several physiological parameters were measured at different times post *Lonomia obliqua* venom injection. The results are presented in Fig. 1.

Between 2 and 6 h after LOBE administration (doses of 1 mg/kg or 1.5 mg/kg, s.c.), the animals showed signs of acute toxicity, including progressive weakness, lethargy and dyspnea. Compared to PBS-treated animals there was a reduction of body weight mainly at 6 h in rats that received the higher dose of LOBE (1.5 mg/kg, s.c.) (Fig. 1A). The weight loss was accompanied by a significant reduction in water intake between 6 and 48 h and by an increase in urinary output at 6 and 12 h for both tested doses (Fig. 1B). The polyuria was maximal at 6 h, when urinary output increased from 0.58 ± 0.08 mL/2h in PBS-treated animals to 1.80 ± 0.16 mL/2 h ($p < 0.05$) in rats injected with LOBE (1.5 mg/kg, s.c.). At the same time, animals that received a dose of 1.0 mg/kg had a urinary output significantly lower (0.9 ± 0.2 mL/2h, $p < 0.05$) compared to those treated with 1.5 mg/kg (Fig. 1C).

Important glomerular and tubular functions related to control of fluid filtration, water and electrolyte balance were severely impaired. Despite the polyuria, *L. obliqua*-induced AKI is associated with a marked reduction in the glomerular filtration rate (GFR). In the first 24 h of envenomation, a rapid decrease of GFR was observed in animals treated with both doses of LOBE. At 48 h envenomed rats (injected with 1.5 mg/kg of LOBE) presented values of GFR 16 times lower than controls at the same time. This remarkable effect of the venom on GFR was observed for as long as 96 hours (Fig. 1D). Similarly, the kidney's ability to concentrate urine (a primordial tubular function) was also impaired, since the density and osmolality of urine had a reduction in values between 2 and 48 h (Fig. 1E and F). The decrease in urinary osmolality was accompanied by a significant increase in the plasma osmolality observed mainly in rats injected with the higher dose of LOBE (Fig. 1G). Consistent with these results the osmolar clearance (C_{osm}) was lower in envenomed animals, while the free water clearance (C_{H_2O}) increased, which imply that the kidney is producing dilute urine through the excretion of solute-free water (Fig. 1H and I). As expected, with the loss of ability to retain water, the fractional excretion of water (FE_{H_2O}) increased from 0.33 ± 0.03 % in PBS-treated animals to 29.1 ± 4.9 % ($p < 0.05$) in rats injected with LOBE (1.5 mg/kg, s.c.) at 48 h (Fig. 1J). Likewise, tubular ability to conserve and maintain the electrolytic balance was lost throughout the envenoming period. The fractional excretion of sodium and potassium (FE_{Na^+} and FE_{K^+}) was high between 24 and 48 h, indicating a marked impairment in the tubular reabsorption of filtered Na^+ and K^+ (Fig. 1K and L). At 96 h most of the renal function parameters (excepting GFR) tends to return to normal levels (similar from that obtained to non-envenomed rats).

3.2 Proteinuria

Most of envenomed animals produced dark-brown-colored urine in the period of 6 to 12 h, indicating the occurrence of hematuria and/or hemoglobinuria. The presence of intact and fragmented erythrocytes, epithelial cells and leukocytes were observed in the urinary sediment by light microscopy (not shown). Consistent with these observations, *L. obliqua* venom induced massive acute proteinuria, which was maximal at 6 and 12 h post-venom injection (Fig. 2A). As expected, the group treated with 1.5 mg/kg of LOBE had more severe proteinuria compared to animals treated with 1.0 mg/kg, but for both groups, urinary protein excretion decreased progressively between 12 and 48 h, reaching levels similar to controls at 96 h. Analysis of urine by SDS-PAGE revealed the presence of several proteins with molecular weights ranging from 10 to 75 kDa (Fig. 2B). A greater variety of bands

were evident mainly at 6 and 12 h after envenomation, confirming this time interval as crucial to the development of glomerular injury. The two most prominent bands (observed around of 70 and 15 kDa) have molecular weights that match to serum albumin (68 kDa) and subunits of hemoglobin (16 kDa). The presence of venom excreted in urine was verified by western-blot. As shown in Fig. 2C, we detected at least 5 bands in urine from venom-treated rats at 6 and 12 h, which specifically react with antibodies raised against *L. obliqua* toxic proteins. These bands have molecular weights around of 70-60, 50, 37, 25 and 20 kDa (arrows in Fig. 2C) and were not recognized in urine from non-envenomed animals (controls). Several toxins in this range of molecular weights have already been identified through transcriptomic and proteomic analysis of LOBE. Some of them include lectins and c-type lectin-like proteins (70-60 kDa), serine-proteinases (50 kDa), cysteine-proteinases (37 kDa) and lipocalins (25-20 kDa) (Ricci-Silva *et al.*, 2008; Veiga *et al.*, 2005).

3.3 Renal histopathological alterations

Light microscopy of kidney biopsies from PBS-injected rats revealed a normal renal parenchyma (Fig. 3A). In contrast, envenomed animals showed progressive degenerative lesions compatible with acute tubular necrosis (ATN) (Fig. 3B–H). Increased acidophilia and dilation of renal tubules were observed between 6 and 48 h after venom administration (Fig. 3C–F). Loss of proximal brush border, cytoplasm vacuolation, and in some tubules, degeneration and desquamation of necrotic cells occurred between 12 and 48 h (Fig. 3D–F). The nuclei of the various proximal tubular cells at 24 and 48 h often showed pyknosis with clumping of chromatin material (Fig. 3E and F). In several tubules at 48 h, the renal epithelium was completely necrotic whereas the basement membrane was either intact or disrupted by tubular necrosis (Fig. 3F). Consistent with the histological signs of tubular injury, γ -GT urinary activity — which is considered an efficient biomarker for early diagnosis of ATN (Guder and Ross, 1984) — increased in a dose- and time-dependent manner up to 48 h. At 96 h, γ -GT activity decreased, but remained significantly high in animals treated with 1.5 mg/kg of LOBE. Histologically, the signs of lesions were much less marked and the proximal and post-proximal tubular epithelia assumed a normal appearance at 96 h (Fig. 3G and H).

Besides the evidence of degenerative lesions, proximal and distal tubules also had swollen lumens. Hyaline and hematic casts and cellular debris were found within tubules, obstructing their lumens (Figs. 3C–F and 4). Hyaline casts (predominantly formed by a protein-rich material) were more prevalent at 6 and 12 h (Figs. 3C–D and 4A), while hematic casts (predominantly formed by fragmented or intact erythrocytes) and cellular debris have appeared commonly at 12, 24 and 48 h (Figs. 3D–F and 4A–B).

An intense inflammatory response characterized by edema, cellular infiltration and fibrosis was observed mainly at 48 h and 96 h (Figs. 3F and 5). Inflammatory cell infiltrate and edema were detected at 48 h in regions of extensive necrosis (Fig. 3F) and within glomeruli (Fig. 5C). At 96 h, despite signs of tubular regeneration, the inflammatory infiltrate increased significantly, probably to help in tissue repair (Fig. 5B). Light microscopy of sections stained with picrosirius revealed extensive peritubular collagen deposition at 96 h after administration of *L. obliqua* venom (Fig. 5E) compared to PBS-treated rats (Fig. 5D),

indicating fibrosis. Foci of interglomerular collagen deposition were also observed in the renal cortex at the same time (Fig. 5F).

3.4 Glomerular morphometric alterations and renal tissue factor activity

Glomerular alterations in envenomed animals were associated with lobulation of the capillary tufts, dilation of glomeruli, deposition of a hyaline material inside the Bowman's space and distention of Bowman's space (Figs. 3D, 6A and 7B). Nodules that formed a dense, strongly-stained mesangial matrix, evidenced by the presence of a PAS positive stain in the capillary tufts was also observed between 2 and 12 h post-venom (Fig. 6A).

Consistent with histological observations, computer-assisted morphometric analysis has indicated a significant increase in glomerular area at all-time intervals examined. There was also an increase in the areas of glomerular tuft and Bowman's space, which confirms that the capillaries and Bowman's space are dilated (Fig. 6B). The enlargement of Bowman's space observed at 12 h possibly is related to the presence of a hyaline material (rich in plasma proteins) found in several glomeruli at this time (details in Figs 3D and 7B).

Since intra-glomerular fibrin deposition can impair renal function and *L. obliqua* venom is able to activate the coagulation system both *in vitro* and *in vivo* (Berger *et al.*, 2010), we decide to measure the levels of renal tissue factor (TF) activity during envenomation. TF is a transmembrane enzyme activator that triggers the coagulation cascade, generating activated factor X and fibrin. As showed in Fig. 6C, renal TF activity increased rapidly from the first 2 h, reaching levels 2 times higher than control values at 6 hours. At 12 and 24 h post-venom injection this level remained significantly high, but decreased progressively thereafter (Fig. 6C). Interestingly, the rise in renal TF activity was coincident with the presence of PAS positive deposits (which stain specifically glycoproteins) within the capillary tufts at 2-12 h (Fig. 6A).

3.5 Hemodynamics and renal vascular permeability

Mean arterial blood pressure was lower in animals that received LOBE compared to basal levels in controls (Table 1). Sustained hypotension was detected between 24 and 96 h in rats treated with the both doses of venom. The maximum decrease occurred at 24 h. In contrast, the heart rate increased significantly at 12 and 96 h in animals injected with 1 mg/kg of LOBE and at 48 h in animals injected with 1.5 mg/kg. Hypotension was accompanied by an increase in renal vascular permeability (Fig. 7). Kidney blood vessels were hyperemic and signs of plasma leakage, migration of inflammatory cells and interstitial edema were also evident after 12 h of venom administration (Fig. 7A). As mentioned above, several glomeruli at 12 h have their Bowman's space filled with a protein-rich material that had extravasated from glomerular capillaries (Fig. 7B). Confirming these observations, a marked increase of vascular permeability in the kidney, as measured by the extravasation of Evans blue dye, was detected at 12 and 24 h after LOBE injection (1.5 mg/kg, s.c) (Fig. 7C).

3.6 Immunohistochemical detection of venom in the renal tissue

Venom distribution and its binding to renal structures was investigated by submitting kidney biopsies from venom-treated and control rats to immunohistochemistry using anti-LOBE IgG that reacts specifically with venom toxins (Fig. 8). Positive immunohistochemical

reaction was found in cortical and medullar regions of kidneys from rats injected with 1.5 mg/kg of LOBE (Fig. 8D and C). Venom was detected in glomerular capillaries, Bowman's capsule, proximal and distal tubules and in intra-tubular casts. Staining for venom was intense in tubular brush border at 2 and 6 h and was also present in cells of tubules in degeneration at 48 h (Fig. 8B-D). Generally, tubules stained more than glomeruli. There was no immunoreactivity for venom in the renal structures of PBS-treated rats (controls) (Fig. 8A). The highest levels of venom (estimated by the area of positive immunohistochemical reaction) were detected in the renal tissue at 2 h (Fig. 8F). After that, venom immunoreactivity decreased progressively until 96 h when the staining was weak (Fig. 8E and F). *L. obliqua* toxins were also detected in other kidney structures such as in perivascular connective tissue of blood vessels, endothelium and smooth muscle of arteries and veins (Fig. 9A and B).

3.7 Kidney proteins differentially expressed during envenomation

To gain further mechanistic insight which drive venom-induced kidney disease, we applied a semi-quantitative discovery based shotgun proteomic approach to identify the proteins differentially expressed in the kidney of envenomed animals. For this purpose, the proteomic data was acquired by tandem mass spectrometry with subsequent quantification, analysis of differential protein expression, validation, and functional annotations in order to identify the involved molecular pathways. The proteins identified consist of two classes: (1) those that were differentially expressed and met our significance criteria and (2) proteins that were uniquely identified in kidneys of rats injected with PBS or LOBE (1.5 mg/kg) after 24h. Figure 10A represents differentially expressed and unique proteins *via* a Venn diagram. Overlaps between PBS- and LOBE-treated animals represent significant differentially expressed proteins and non-overlapping portions of the diagram represent unique protein identifications. A total of 779 proteins were identified. Twenty-four proteins were exclusively identified in envenomed animals; 169 were exclusively identified in control rats and 586 were common proteins in both treatments. Among these 586 proteins, 138 (23.5 %) were identified as being differentially expressed.

Ingenuity Pathway Analysis was used for functional annotations and revealed several key protein categories and pathways significantly enriched in the differentially expressed proteins. Thus, the sets of proteins were assigned to either biologic function in disease or canonical signaling pathways. Through these analyses it was evident that the set of proteins identified in control kidneys displayed healthy biological functions such amino acid metabolism and small-molecule biochemistry associated with normal renal metabolic pathways (Fig. 10B). On the other hand, protein expression shifts toward “cellular distress” functions in the kidney of envenomed rats (Fig. 10C). In this case, the expression profiles were considerably enriched for proteins that belong to acute phase inflammatory response signaling, LXR/RXR activation (involved in retinoic acid-mediated gene activation triggered by inflammatory stimulus), oxidative stress response pathways such as that mediated by the nuclear erythroid-related factor 2 (NRF2) and coagulation and complement systems (involved in thrombosis, fibrosis, inflammation and vascular alterations) (Fig. 10C).

Accordingly to our functional and histopathological results, several novel proteins related to renal disease were identified in the kidney of LOBE-treated animals. Proteins associated with tubular and glomerular injury, necrosis, inflammation and fibrosis were up-regulated or unique in envenomed rats (Fig. 10D). Some of these molecules, mainly those that were related to renal disease or identified in canonical pathways, are listed in Table 2. A complete list with differentially expressed and unique proteins of envenomed and control animals are shown in supplemental Table 1. It is worth mentioning, that proteins functionally linked to tissue injury (markers of necrosis and/or apoptosis), osmotic and oxidative stress, electrolytic imbalance, acute phase inflammatory response, fibrosis and thrombosis were expressed in the kidneys of LOBE-treated rats (Table 2). Proteins of the kallikrein-kinin and complement systems which are related to hypotension and control of vascular permeability were also found to be up-regulated in envenomed animals (Table 2). Our proteomic data also confirm that an important mechanism of *Lonomia*-induced AKI is mediated by hemoglobin and the release of its degradation products: free heme and iron. In fact, hemoglobin, hemopexin and ferritin were up-regulated and the expression of heme oxygenase-1 was uniquely induced in the kidneys of envenomed rats (Table 2). As a result of heme and iron release, several antioxidant enzymes were found to be up-regulated, indicating the generation of reactive oxygen species (Table 2).

4. Discussion

AKI is frequently described and is life threatening in several cases of snake and arthropod envenomation (Sitprija, 2006; Berger *et al.*, 2012). Particularly in *Lonomia obliqua* envenomation, AKI is the main cause of death and its mechanisms are completely unknown until now. In this work we use an *in vivo* experimental model to characterize the *L. obliqua*-induced AKI. According to our results, the pathophysiological mechanism seems to be complex and multifactorial involving four main issues: (1) vascular abnormalities; (2) tubular and glomerular alterations; (3) renal inflammation and (4) a direct venom cytotoxic activity.

4.1 Vascular abnormalities

Rats injected with *L. obliqua* venom presented important hemodynamic alterations characterized by systemic hypotension and increased heart rate. The time of maximal decrease in blood pressure was coincident with the reduction of GFR and the impairment of renal function. In addition, an increase of renal vascular permeability and edema was also observed at the same period of envenomation. These findings are important because systemic vasodilation is associated with the decrease in GFR and can lead to renal hypoperfusion and ischemia in different pathological conditions (Schrier *et al.*, 2004). Specifically in this type of envenomation, one mechanism that could contribute to vasodilation is the activation of kallikrein-kinin system (KKS). It is known that *L. obliqua* venom has toxins (kallikrein activators and kininogenases) able to activate plasma pre-kallikrein and directly release bradykinin (BK) from low molecular weight kininogen (LMWK) (Pinto *et al.*, 2010; Bohrer *et al.*, 2007). Interestingly, our results indicated that the expression of LMWK (the main substrate of tissue kallikrein and venom kininogenases) is up-regulated in the kidneys of envenomed animals, which may favor the generation of BK.

The immediate consequence of intravascular activation of KKS is a fall in systemic blood pressure and in peripheral tissues is edema and erythema formation (Bohrer *et al.*, 2007). The participation of BK was already confirmed, since the hypotensive and edematogenic responses elicited by LOBE were inhibited by HOE-140, a B2 receptor antagonist (Bohrer *et al.*, 2007). In agreement with these experimental observations, hypotension and reduced plasma levels of pre-kallikrein are common features observed in patients (Zannin *et al.*, 2003), which support the evidence that KKS is activated during envenomation and is clinically relevant. Moreover, the data presented here also suggest that KKS activation may be involved in venom-induced AKI.

4.2 Tubular and glomerular alterations

Despite the scarce clinical case-reports, the main pathological finding obtained from kidney biopsies is acute tubular necrosis (ATN). Experimental animals also showed histological alterations compatible with ATN such as loss of proximal brush border, cytoplasm vacuolation, pyknotic nuclei, degeneration and desquamation of necrotic cells. These necrotic cells exfoliating into the lumen due to either cell death or defective cell-to-cell or cell-to-basement membrane adhesion, can obstruct the flow of filtrated fluid and give rise to a back pressure limiting glomerular filtration (Trof *et al.*, 2006). Accordingly, in this work, markers of ATN were detected in urine (urinary γ -GT) and several proteins related to tubule injury were found to be up-regulated (Orm-1, YBX1, Alb, CK-B, FABP3 and NHE-RF1) or uniquely expressed (NGAL) in the kidneys of envenomed rats. Proteins linked to apoptosis (CytC, S100-A8 and MFP-1) were also up-regulated. The occurrence of glomerular dysfunction was evident, because envenomed rats presented massive proteinuria, serum Alb had a 2.1-fold increase in the kidneys and a band corresponding to the molecular weight of Alb was detected in urine. In addition, our results confirm that hematuria and hemoglobinuria are predominant characteristics of *L. obliqua*-induced AKI. Indeed, the venom has strong *in vitro* and *in vivo* hemolytic activity and a phospholipase A2 responsible for this effect was already isolated (Seibert *et al.*, 2006; Seibert *et al.*, 2010). Intact erythrocytes were found within the tubules forming intra-tubular casts and, as a result of intravascular hemolysis, different Hb subunits were detected in urine and kidneys. It is known that the formation of Hb deposits may be toxic to renal tubules due to heme cytotoxicity (Zager, 1996). Once reabsorbed by the proximal tubular cells the heme porphyrin ring is rapidly catabolized by Hmox-1 yielding equimolar amounts of free iron, biliverdin and carbon monoxide (Camara and Soares, 2005). Free iron then up-regulates intracellular ferritin expression, a key defense mechanism against iron-induced tissue damage. However, in the presence of large amounts of Hb the levels of released iron also increase, saturating the binding capacity of ferritin. Thus, the iron not removed by ferritin binding is able to readily accept and donate electrons and greatly facilitates free radical production (Zager, 1996; Khan, 2009; Zager *et al.*, 2012). As proteins related to canonical pathways of oxidative stress were identified exclusively in the kidneys of LOBE-treated rats, we believe that heme cytotoxicity plays a significant role in *L. obliqua*-induced AKI. Consistent with this, several anti-oxidant enzymes (Prdx-5, Prdx-2, GST, DJ-1 and SOD) and proteins associated with heme and iron metabolism (Hmox-1, ferritin, Hpx and Cygb) were up-regulated or uniquely expressed during envenomation, suggesting that heme is effectively metabolized, generating free radicals and inducing oxidative damage to proteins,

lipids and DNA. In fact, in previous experiments we demonstrated that LOBE induces kidney DNA damage leading to double-strand breaks and formation of oxidized purines and pyrimidines (Berger *et al.*, 2013). Probably this is associated with the increased expression of Rad50, a double-strand break repair protein, detected in the present work.

As a consequence of ATN, the excretion of Na^+ and K^+ increased significantly during envenomation. Injured proximal tubule cells have alterations in the actin and microtubule cytoskeletal networks that lead to a redistribution of Na^+/K^+ ATPase from the basolateral to the apical membrane, contributing to a decrease in Na^+ transport and reabsorption (Thadhani *et al.*, 1996). It was reported that FE_{Na^+} and FE_{K^+} also increase in envenomation caused by *Bothrops* snakes (Boer-Lima *et al.*, 1999; Linardi *et al.*, 2011). Despite the redistribution of Na^+/K^+ ATPase to the apical membrane, Linardi *et al.* (2011) reported an increase in expression and activity of Na^+/K^+ ATPase during envenomation by *Bothrops alternatus* and suggest that it is a protective mechanism triggered in response to natriuresis with the aim to preserve renal function during acute damage. In agreement with this observation, we found an increase in expression of type-3 Na^+/H^+ exchanger regulatory cofactor, which is important in Na^+ and HCO_3^- reabsorption in proximal tubule cells (Trof *et al.*, 2006). Likewise Na^+/K^+ , water excretion increased significantly and envenomed animals had polyuria. Several parameters measured, such as urinary density and osmolality, C_{osm} , $C_{\text{H}_2\text{O}}$ and $\text{FE}_{\text{H}_2\text{O}}$, indicate that kidneys from envenomed rats are producing dilute urine through the excretion of solute-free water. Possibly, the presence of intratubular Na^+ not reabsorbed by proximal tubule cells may contribute to the increased water excretion and both, Na^+ and water rejection, may also be associated with the fall in blood pressure.

Regarding the glomerular dysfunctions, a valuable hypothesis that should be considered is the deposition of fibrin clots in glomerular capillaries. Indeed, the most potent activity of LOBE *in vitro* is the procoagulant activity (Donato *et al.*, 1998; Veiga *et al.*, 2003). *In vivo*, LOBE also causes activation of coagulation and fibrinolysis leading to a consumptive coagulopathy characteristic of this type of envenomation. Two enzymes responsible for the venom procoagulant activity, activators of prothrombin and factor X, have already been isolated (Alvarez-Flores *et al.*, 2006; Reis *et al.*, 2006). Besides the direct effect on coagulation factors, LOBE is also able to induce a procoagulant profile in endothelial cells in culture through an up-regulation of TF expression (Pinto *et al.*, 2008). Confirming these results obtained in endothelial cells, envenomed kidneys showed an increase in TF procoagulant activity. In addition, the expression of known markers of disseminated intravascular coagulation and thrombosis (serpin 1 alpha-1-antiproteinase, PAI-1 and HC-2) increased in the kidneys of venom-treated rats. The gamma and beta chains of fibrinogen were detected solely in envenomed kidneys, which are probably related to the PAS positive stain observed in glomeruli and is suggestive of fibrin formation.

4.3 Renal Inflammation

Intense inflammatory response is a common feature in *L. obliqua* envenomation. Pain and edema are the most characteristic initial symptoms observed at the local site of contact (Zannin *et al.*, 2003). Usually signs of systemic inflammation with neutrophilic leukocytosis, cell infiltrate and edema have also been described in lungs, kidney and heart of

experimental animals (Berger *et al.*, 2013). The inflammatory response is accompanied by the production of several cytokines (TNF, IL-1 β , IL-8, IL-6, CCL2 and CXCL1), vasoactive mediators (BK, histamine, prostaglandins and nitric oxide), adhesion molecules (E-selectin, VCAM-1 and ICAM-3) and an increase in leukocyte rolling and adhesion to the endothelium (Alvarez-Flores *et al.*, 2006; Bohrer *et al.*, 2007; Pinto *et al.*, 2008; Berger *et al.*, 2010; Nascimento-Silva *et al.*, 2012). Specifically in the kidney, it was observed an up-regulation of several proteins related to acute phase inflammatory signaling, nephritis, inflammatory cell infiltration, increase in vascular permeability, glomerular dilation, distention of Bowman's space and interstitial edema. Kidney sections stained with picrosirius revealed extensive collagen deposition in cortical periglomerular and peritubular regions and proteins involved with fibrosis, such as Cygb, PAI-1, complement C3 and complement factor D were identified. Enhanced deposition of extracellular matrix (ECM) proteins in renal tissue has been observed in response to a variety of stimuli, including TGF- β , TNF- α , IL-1, several adhesion molecules and chemoattractants. These stimuli can also increase the levels of tissue inhibitors of matrix metalloproteinases, thereby attenuating ECM turnover, and thus, favoring the deposition of collagen and other matrix proteins (Eddy, 1996; Pawluczyk and Harris, 1998).

Besides fibrosis, activated complement components (C3 and factor D) may also contribute to the alterations in vascular permeability and acts as a chemoattractant for neutrophils (Turnberg *et al.*, 2006). Moreover, complement activation by albumin is a powerful underlying mechanism of tubular and interstitial injury via cytotoxic, proinflammatory, and fibrogenic effects, which often occur in renal diseases where proteinuria is present (Portella *et al.*, 2013). In an experiment conducted in proximal tubular cells incubated with serum proteins, *in vitro* complement activation was observed, which could be associated with changes in the cytoskeleton, production of superoxide anion, hydrogen peroxide, and proinflammatory cytokines, such as IL-6 and TNF- α (Abbate *et al.*, 2006).

4.4 Direct venom cytotoxic activity

As the kidneys are highly vascularized organs, they are particularly susceptible to direct venom toxicity (Sitprija, 2006). Renal epithelial cells in culture and isolated perfused kidneys have been used to characterize the direct cytotoxic effects of different venoms including bee, snake and spider venoms. The most common isolated toxins, which are nephrotoxic, belong to the classes of metalloproteinases, serine-proteinases, C-type lectins, phospholipases A2, sphingomyelinases D and L-amino acid oxidases (Berger *et al.*, 2012). In the case of *L. obliqua* some toxins belonging to these classes have already been isolated (Pinto *et al.*, 2010); however, their effects on renal cells are unknown. Our results provide some evidence that the whole venom probably has a direct nephrotoxic effect, since immunohistochemical staining confirmed the presence of venom in renal tissue, with stronger staining in the initial 6 h after venom administration followed by a progressive decrease thereafter. The venom was rapidly excreted in urine, because at least 5 bands which specifically react with antibodies raised against *L. obliqua* toxins were detected in urine. Interestingly, the positive immunohistochemical reaction for venom in different kidney structures agreed with the morphological and histological damage caused by the venom in these anatomical regions and indicated that there was a close correlation between the sites of

venom localization and subsequent tissue injury. Previous studies are consistent with our observations. Using immunochemical and radio-labeling methods to analyze venom biodistribution in rats, the highest quantities of LOBE were detected in kidneys, blood and urine (Rocha-Campos *et al.*, 2001; Da Silva *et al.*, 2004).

5. Conclusion

In this work a rat experimental model was used to study the progression of renal disease during *Lonomia obliqua* envenomation. According to our results, the pathophysiological mechanism involved in *L. obliqua*-induced AKI seems to be multifactorial where events such as systemic hypotension and fibrin deposition contribute to renal hypoperfusion, tubular necrosis and the sudden loss of basic renal functions, including filtration and excretion capacities, urinary concentration and maintenance of body fluid homeostasis. In addition, when compared to control rats, the kidneys from envenomed animals showed to be increasingly enriched for stress-related proteins, which are commonly associated to inflammation, tissue injury, heme-induced oxidative stress, coagulation and complement systems activation. Finally, the localization of the venom in renal tissue agreed with morphological and functional alterations, suggesting a close correlation between venom tissue levels and renal damage. Thus, the different mechanisms as well as the renal injury biomarkers identified here can be useful and guide further experiments to the discovery of alternative forms of treatment to *L. obliqua*-induced AKI.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by funding and fellowships from the Brazilian Agencies: Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Ministério da Educação, Brazil (CAPES-MEC) - Edital Toxinologia (Processo: 23038.006277/2011-85) and Conselho Nacional de Desenvolvimento Científico e Tecnológico, Ministério da Ciência e Tecnologia, Brazil (CNPq-MCT). This work was also supported by NIH grant P41 GM103533 (to JRY). We thank Dr Jeffrey N. Savas for his advice on the mass spectrometric analysis and for a critical reading of the manuscript.

References

- Abbate M, Zoja C, Remuzzi G. How does proteinuria cause progressive renal damage? J Am Soc Nephrol. 2006; 17:2974–2984. [PubMed: 17035611]
- Alvarez-Flores MP, Fritzen M, Reis CV, Chudzinski-Tavassi AM. Losac, a factor X activator from *Lonomia obliqua* bristle extract: its role in the pathophysiological mechanisms and cell survival. Biochem Biophys Res Commun. 2006; 343:1216–1223. [PubMed: 16597435]
- Ambatipudi KS, Lu B, Hagen FK, Melvin JE, Yates JR. Quantitative analysis of age specific variation in the abundance of human female parotid salivary proteins. J Proteome Res. 2009; 8:5093–5102. [PubMed: 19764810]
- Basnakian AG, Kaushal GP, Shah SV. Apoptotic pathways of oxidative damage to renal tubular epithelial cells. Antioxid Redox Signal. 2002; 4(6):915–924. [PubMed: 12573140]
- Berger M, Reck J Jr, Terra RM, Pinto AF, Termignoni C, Guimarães JA. *Lonomia obliqua* caterpillar envenomation causes platelet hypoaggregation and blood incoagulability in rats. Toxicon. 2010; 55(1):33–44. [PubMed: 19577588]

- Berger, M.; Vieira, MAR.; Guimaraes, JA. Acute Kidney Injury Induced by Snake and Arthropod Venoms. In: Polenakovic, M., editor. Renal Failure - The Facts. InTech-Open; Rijeka, Croatia: 2012. p. 157-186. ISBN: 978953-51-0630-2
- Berger M, Beys-da-Silva W, Santi L, de Oliveira IM, Jorge PM, Henriques JAP, Driemeier D, Vieira MAR, Guimarães JA. Acute *Lonomia obliqua* caterpillar envenomation-induced physiopathological alterations in rats: Evidence of new toxic venom activities and the efficacy of serum therapy to counteract systemic tissue damage. *Toxicon*. 2013; 74:179–192. [PubMed: 23994591]
- Bern M, Goldberg D, McDonald WH, Yates JR 3rd. Automatic quality assessment of peptide tandem mass spectra. *Bioinformatics*. 2004; 20:i49–54. [PubMed: 15262780]
- Boer-Lima PA, Gontijo JA, da Cruz-Höfling MA. Histologic and functional renal alterations caused by *Bothrops moojeni* snake venom in rats. *Am J Trop Med Hyg*. 1999; 61(5):698–706. [PubMed: 10586897]
- Bohrer CB, Reck J Jr, Fernandes D, Sordi R, Guimarães JA, Assreuy J, Termignoni C. Kallikrein-kinin system activation by *Lonomia obliqua* caterpillar bristles: involvement in edema and hypotension responses to envenomation. *Toxicon*. 2007; 49:663–639. [PubMed: 17188732]
- Bowler RP, Duda B, Chan ED, Enghild JJ, Ware LB, Matthay MA, Duncan MW. Proteomic analysis of pulmonary edema fluid and plasma in patients with acute lung injury. *Am J Physiol Lung Cell Mol Physiol*. 2004; 286(6):L1095–104. [PubMed: 14742308]
- Brocker C, Lassen N, Estey T, Pappa A, Cantore M, Orlova VV, Chavakis T, Kavanagh KL, Oppermann U, Vasiliou V. Aldehyde dehydrogenase 7A1 (ALDH7A1) is a novel enzyme involved in cellular defense against hyperosmotic stress. *J Biol Chem*. 2010; 285(24):18452–63. [PubMed: 20207735]
- Burdmann EA, Antunes I, Saldanha LB, Abdulkader RC. Severe acute renal failure induced by the venom of *Lonomia* caterpillars. *Clinical Nephrology*. 1996; 46(5):337–339.
- Caliari, MV. Princípios de Morfometria Digital: KS300 para iniciantes. UFMG. , editor. 1997. p. 149
- Camara NO, Soares MP. Heme oxygenase-1 (HO-1), a protective gene that prevents chronic graft dysfunction. *Free Radic Biol Med*. 2005; 38(4):426–435. [PubMed: 15649645]
- Caplin B, Wang Z, Slaviero A, Tomlinson J, Dowsett L, Delahaye M, Salama A, Wheeler DC, Leiper J. Alanine-glyoxylate aminotransferase-2 metabolizes endogenous methylarginines, regulates NO, and controls blood pressure. *Arterioscler Thromb Vasc Biol*. 2012; 32(12):2892–900. [PubMed: 23023372]
- Carvalho PC, Fischer JS, Chen EI, Yates JR 3rd, Barbosa VC. PatternLab for proteomics: a tool for differential shotgun proteomics. *BMC Bioinformatics*. 2008; 9:316. [PubMed: 18644148]
- Carvalho PC, Fischer JS, Xu T, Yates JR 3rd, Barbosa VC. PatternLab: from mass spectra to label-free differential shotgun proteomics. *Curr Protoc Bioinformatics*. 2012 Chapter 13:Unit13.19.
- Chandler RJ, Zerfas PM, Shanske S, Sloan J, Hoffmann V, DiMauro S, Venditti CP. Mitochondrial dysfunction in mut methylmalonic acidemia. *FASEB J*. 2009; 23(4):1252–1261. [PubMed: 19088183]
- Chang SH, Phelps PC, Berezesky IK, Ebersberger ML Jr, Trump BF. Studies on the mechanisms and kinetics of apoptosis induced by microinjection of cytochrome c in rat kidney tubule epithelial cells (NRK-52E). *Am J Pathol*. 2000; 156(2):637–649. [PubMed: 10666393]
- Chen G, Paka L, Kako Y, Singhal P, Duan W, Pillarisetti S. A protective role for kidney apolipoprotein E: Regulation of mesangial cell proliferation and matrix expansion. *J Biol Chem*. 2001; 276:49142–49147. [PubMed: 11579084]
- Chinnaiyan AM, Huber-Lang M, Kumar-Sinha C, Barrette TR, Shankar-Sinha S, Sarma VJ, Padgaonkar VA, Ward PA. Molecular signatures of sepsis: multiorgan gene expression profiles of systemic inflammation. *Am J Pathol*. 2001; 159(4):1199–1209. [PubMed: 11583946]
- Da Silva GH, Panunto PC, Hyslop S, Da Cruz-Höfling MA. Immunochemical detection of *Lonomia obliqua* caterpillar venom in rats. *Microsc Res Tech*. 2004; 65:276–281. [PubMed: 15662625]
- Devarajan P, Krawczeski CD, Nguyen MT, Kathman T, Wang Z, Parikh CR. Proteomic identification of early biomarkers of acute kidney injury after cardiac surgery in children. *Am J Kidney Dis*. 2010; 56(4):632–642. [PubMed: 20599305]

- Dias da Silva W, Campos CM, Gonçalves LR, Sousa-e-Silva MC, Higashi HG, Yamagushi IK, Kelen EM. Development of an antivenom against toxins of *Lonomia obliqua* caterpillars. *Toxicon*. 1996; 34:1045–1049. [PubMed: 8896196]
- Donato JL, Moreno RA, Hyslop S, Duarte A, Antunes E, Le Bonniec BF, Rendu F, de Nucci G. *Lonomia obliqua* Caterpillar Spicules Trigger Human Blood Coagulation via Activation of Factor X and Prothrombin. *Thromb Haemost*. 1998; 79:539–542. [PubMed: 9531036]
- Duarte AC, Caovilla JJ, Lorini JD, Mantovani G, Sumida J, Manfre PC, Silveira RC, de Moura SP. Insuficiência renal aguda por acidentes com lagartas. *J Bras Nefrol*. 1990; 12:184–187.
- Duarte AC, Crusius PS, Pires CAL. Insuficiência renal aguda nos acidentes com *lonomia obliqua*. *Nefrol Lat Americana*. 1994; 1(1):38–40.
- Duarte, AC. Síndrome Hemorrágica causada por larvas de mariposa do gênero *Lonomia*: estudo clínico-epidemiológico. Universidade Federal do Rio Grande do Sul; Porto Alegre: 1997. Dissertação
- Eng JK, McCormack AL, Yates JR. An approach to correlate tandem mass-spectral data of peptides with amino-acid-sequences in a protein database. *J Am Soc Mass Spectrom*. 1994; 5(11):976–989. [PubMed: 24226387]
- Eddy AA. Molecular insights into renal interstitial fibrosis. *J. Am. Soc. Nephrol*. 1996; 7:2495–2508. [PubMed: 8989727]
- Eddy AA. Proteinuria and interstitial injury. *Nephrol Dial Transplant*. 2004; 19(2):277–281. 2004. [PubMed: 14736944]
- Fan HW, Cardoso JL, Olmos RD, Almeida FJ, Viana RP, Martinez AP. Hemorrhagic syndrome and acute renal failure in a pregnant woman after contact with *Lonomia* caterpillars: a case report. *Rev Inst Med Trop São Paulo*. 1998; 40(2):119–120. [PubMed: 9755567]
- Feng MG, Navar LG. Afferent arteriolar vasodilator effect of adenosine predominantly involves adenosine A2B receptor activation. *Am J Physiol Renal Physiol*. 2010; 299(2):F310–5. [PubMed: 20462966]
- Gamborgi GP, Metcalf EB, Barros EJG. Acute renal failure provoked by toxin from caterpillars of the species *Lonomia obliqua*. *Toxicon*. 2006; 47:68–74. [PubMed: 16310819]
- Godoy JR, Oesteritz S, Hanschmann EM, Ockenga W, Ackermann W, Lillig CH. Segment-specific overexpression of redoxins after renal ischemia and reperfusion: protective roles of glutaredoxin 2, peroxiredoxin 3, and peroxiredoxin 6. *Free Radic Biol Med*. 2011; 51(2):552–561. [PubMed: 21586322]
- Guder WG, Ross BD. Enzyme distribution along the nephron. *Kidney Int*. 1984; 26:101–111. [PubMed: 6094907]
- Khan FY. Rhabdomyolysis: a review of the literature. *Netherland Journal of Medicine*. 2009; 67(9): 272–283.
- Klein MI, Xiao J, Lu B, Delahunty CM, Yates JR 3rd, Koo H. Streptococcus mutans protein synthesis during mixed-species biofilm development by high-throughput quantitative proteomics. *PLoS One*. 2012; 7:e45795. [PubMed: 23049864]
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 1970; 227:680–685. [PubMed: 5432063]
- Larsen R, Gozzelino R, Jeney V, Tokaji L, Bozza FA, Japiassú AM, Bonaparte D, Cavalcante MM, Chora A, Ferreira A, Marguti I, Cardoso S, Sepúlveda N, Smith A, Soares MP. A central role for free heme in the pathogenesis of severe sepsis. *Sci Transl Med*. 2010; 2(51):51ra71.
- Linardi A, Rocha e Silva TA, Miyabara EH, Franco-Penteado CF, Cardoso KC, Boer PA, Moriscot AS, Gontijo JA, Joazeiro PP, Collares-Buzato CB, Hyslop S. Histological and functional renal alterations caused by *Bothrops alternatus* snake venom: expression and activity of Na⁺/K⁺-ATPase. *Biochimica et Biophysica Acta*. 2011; 1810(9):895–906. [PubMed: 21704674]
- McDonald WH, Tabb DL, Sadygov RG, MacCoss MJ, Venable J, Graumann J, Johnson JR, Cociorva D, Yates JR. MS1, MS2, and SQT-three unified, compact, and easily parsed file formats for the storage of shotgun proteomic spectra and identifications. *Rapid Commun Mass Spectrom*. 2004; 18(18):2162–2168. [PubMed: 15317041]
- Morrissey JH. Tissue factor modulation of factor VIIa activity: use in measuring trace levels of factor VIIa in plasma. *Thromb Haemost*. 1995; 74(1):185–188. [PubMed: 8578455]

- Mimura I, Nangaku M, Nishi H, Inagi R, Tanaka T, Fujita T. Cytoglobin, a novel globin, plays an antifibrotic role in the kidney. *Am J Physiol Renal Physiol*. 2010; 299(5):F1120–1133. [PubMed: 20719976]
- Nascimento-Silva V, Rodrigues da Silva G, Moraes JA, Cyrino FZ, Seabra SH, Bouskela E, Almeida Guimarães J, Barja-Fidalgo C. A pro-inflammatory profile of endothelial cell in *Lonomia obliqua* envenomation. *Toxicon*. 2012; 60(1):50–60. [PubMed: 22779081]
- Nishi H, Inagi R, Kawada N, Yoshizato K, Mimura I, Fujita T, Nangaku M. Cytoglobin, a novel member of the globin family, protects kidney fibroblasts against oxidative stress under ischemic conditions. *Am J Pathol*. 2011; 178(1):128–139. [PubMed: 21224051]
- Pawluczyk IZA, Harris KPG. Cytokine interactions promote synergistic fibronectin accumulation by mesangial cells. *Kidney Int*. 1998; 54:62–70. [PubMed: 9648064]
- Peng J, Elias JE, Thoreen CC, Licklider LJ, Gygi SP. Evaluation of multidimensional chromatography coupled with tandem mass spectrometry (LC/LC-MS/MS) for large-scale protein analysis. The yeast proteome. *J. Proteome Res*. 2003; 2:43–50. [PubMed: 12643542]
- Pinho FM, Zanetta DM, Burdmann EA. Acute renal failure after *Crotalus durissus* snakebite: a prospective survey on 100 patients. *Kidney International*. 2005; 67(2):659–667. [PubMed: 15673314]
- Pinto AFM, Silva KRLM, Guimaraes JA. Proteases from *Lonomia obliqua* venomous secretions: comparison of procoagulant, fibrin(ogen)olytic and amidolytic activities. *Toxicon*. 2006; 47:113–121. [PubMed: 16360724]
- Pinto AFM, Dragulev B, Guimarães JA, Fox JW. Novel perspectives in the pathogenesis of *Lonomia obliqua* caterpillar envenomation based on assessment of host response by gene expression analysis. *Toxicon*. 2008; 51:1119–1128. [PubMed: 18367225]
- Pinto AFM, Berger M, Reck J Jr, Terra RM, Guimarães JA. *Lonomia obliqua* venom: *In vivo* effects and molecular aspects associated with the hemorrhagic syndrome. *Toxicon*. 2010; 56(7):1103–1112. [PubMed: 20114060]
- Portella VG, Silva-Filho JL, Landgraf SS, de Rico TB, Vieira MA, Takiya CM, Souza MC, Henriques MG, Canetti C, Pinheiro AA, Benjamim CF, Caruso-Neves C. Sepsis-surviving mice are more susceptible to a secondary kidney insult. *Crit Care Med*. 2013; 41(4):1056–1068. [PubMed: 23385098]
- Pompermayer K, Souza DG, Lara GG, Silveira KD, Cassali GD, Andrade AA, Bonjardim CA, Passaglio KT, Assreuy J, Cunha FQ, Vieira MA, Teixeira MM. The ATP-sensitive potassium channel blocker glibenclamide prevents renal ischemia/reperfusion injury in rats. *Kidney Int*. 2005; 67(5):1785–1796. [PubMed: 15840025]
- Reis CV, Andrade SA, Ramos OH, Ramos CR, Ho PL, Batista IF, Chudzinski-Tavassi AM. Lopap, a prothrombin activator from *Lonomia obliqua* belonging to the lipocalin family: recombinant production, biochemical characterization and structure-function insights. *Biochem. J*. 2006; 398(2):295–302. [PubMed: 16734589]
- Revelo MP, Federspiel C, Helderman H, Fogo AB. Chronic allograft nephropathy: Expression and localization of PAI-1 and PPAR-gamma. *Nephrol Dial Transplant*. 2005; 20:2812–2819. [PubMed: 16221712]
- Ricci-Silva ME, Valente RH, León IR, Tambourgi DV, Ramos OH, Perales J, Chudzinski-Tavassi AM. Immunochemical and proteomic technologies as tools for unravelling toxins involved in envenoming by accidental contact with *Lonomia obliqua* caterpillars. *Toxicon*. 2008; 51:1017–1028. [PubMed: 18342903]
- Rocha-Campos AC, Gonçalves LR, Higashii HG, Yamagushi IK, Fernandes I, Oliveira JE, Ribela MT, Sousa-e-Silva MC, Dias da Silva W. Specific heterologous F(ab')₂ antibodies revert blood incoagulability resulting from envenoming by *Lonomia obliqua* caterpillars. *Am. J. Trop. Med. Hyg*. 2001; 64:283–289. [PubMed: 11463118]
- Russo LM, Sandoval RM, McKee M, Osicka TM, Collins AB, Brown D, Molitoris BA, Comper WD. The normal kidney filters nephrotic levels of albumin retrieved by proximal tubule cells: retrieval is disrupted in nephrotic states. *Kidney Int*. 2007; 71(6):504–513. [PubMed: 17228368]

- Safadi FF, Thornton P, Magiera H, Hollis BW, Gentile M, Haddad JG, Liebhaber SA, Cooke NE. Osteopathy and resistance to vitamin D toxicity in mice null for vitamin D binding protein. *J Clin Invest.* 1999; 103(2):239–251. [PubMed: 9916136]
- Schrier RW, Wang W, Poole B, Mitra A. Acute renal failure: definitions, diagnosis, pathogenesis, and therapy. *J Clin Invest.* 2004; 114(1):5–14. [PubMed: 15232604]
- Seibert CS, Oliveira MRL, Gonçalves LRC, Santoro ML, Sano-Martins IS. Intravascular hemolysis induced by *Lonomia obliqua* caterpillar bristle extract: an experimental model of envenomation in rats. *Toxicon.* 2004; 44:793–799. [PubMed: 15500855]
- Seibert CS, Tanaka-Azevedo AM, Santoro ML, Mackessy SP, Torquato RJS, Lebrun I, Tanaka AS, Sano-Martins IS. Purification of a phospholipase A₂ from *Lonomia obliqua* caterpillar bristle extract. *Biochem. Biophys. Res. Commun.* 2006; 342:1027–1033. [PubMed: 16510123]
- Shashidharamurthy R, Mahadeswaraswamy YH, Ragupathi L, Vishwanath BS, Kemparaju K. Systemic pathological effects induced by cobra (*Naja naja*) venom from geographically distinct origins of Indian peninsula. *Exp Toxicol Pathol.* 2010; 62(6):587–592. [PubMed: 19733040]
- Sitprija V. Snakebite nephropathy. *Nephrology (Carlton).* 2006; 11(5):442–448. [PubMed: 17014559]
- Son D, Kojima I, Inagi R, Matsumoto M, Fujita T, Nangaku M. Chronic hypoxia aggravates renal injury via suppression of Cu/Zn-SOD: a proteomic analysis. *Am J Physiol Renal Physiol.* 2008; 294(1):F62–72. [PubMed: 17959751]
- Stefanska B, Huang J, Bhattacharyya B, Suderman M, Hallett M, Han ZG, Szyf M. Definition of the landscape of promoter DNA hypomethylation in liver cancer. *Cancer Res.* 2011; 71(17):5891–903. [PubMed: 21747116]
- Tabb DL, McDonald WH, Yates JR 3rd. DTASelect and Contrast: tools for assembling and comparing protein identifications from shotgun proteomics. *J Proteome Res.* 2002; 1:21–26. [PubMed: 12643522]
- Takatsuka H, Nakajima T, Nomura K, Wakae T, Toda A, Itoi H, Okada M, Misawa M, Hara H. Heparin cofactor II as a predictor of thrombotic microangiopathy after bone marrow transplantation. *Hematology.* 2006; 11(2):101–103. [PubMed: 16753849]
- Thadhani R, Pascual M, Bonventre JV. Acute renal failure. *N Engl J Med.* 1996; 334(22):1448–1460. [PubMed: 8618585]
- Thukral SK, Nordone PJ, Hu R, Sullivan L, Galambos E, Fitzpatrick VD, Healy L, Bass MB, Cosenza ME, Afshari CA. Prediction of nephrotoxicant action and identification of candidate toxicity-related biomarkers. *Toxicol Pathol.* 2005; 33(3):343–355. [PubMed: 15805072]
- Tofovic SP, Kusaka H, Li P, Jackson EK. Effects of adenosine deaminase inhibition on blood pressure in old spontaneously hypertensive rats. *Clin Exp Hypertens.* 1998; 20(3):329–44. [PubMed: 9605386]
- Togawa A, Miyoshi J, Ishizaki H, Tanaka M, Takakura A, Nishioka H, Yoshida H, Doi T, Mizoguchi A, Matsuura N, Niho Y, Nishimune Y, Nishikawa SI, Takai Y. Progressive impairment of kidneys and reproductive organs in mice lacking Rho GDIalpha. *Oncogene.* 18(39):5373–5380. [PubMed: 10498891]
- Trof RJ, Di Maggio F, Leemreis J, Groeneveld AB. Biomarkers of acute renal injury and renal failure. *Shock.* 2006; 26(3):245–53. [PubMed: 16912649]
- Turnberg D, Lewis M, Moss J, Xu Y, Botto M, Cook HT. Complement activation contributes to both glomerular and tubulointerstitial damage in adriamycin nephropathy in mice. *J Immunol.* 2006; 177:4094–4102. [PubMed: 16951374]
- Veiga ABG, Blochtein B, Guimarães JA. Structures involved production, secretion and injection of the venom produced by the caterpillar *Lonomia obliqua* (Lepidoptera, saturniidae). *Toxicon.* 2001; 39:1343–1351. [PubMed: 11384722]
- Veiga ABG, Pinto AFM, Guimarães JA. Fibrinogenolytic and procoagulant activities in the hemorrhagic syndrome caused by *Lonomia obliqua* caterpillars. *Thrombosis Research.* 2003; 111(1-2):95–101. [PubMed: 14644086]
- Veiga ABG, Ribeiro JM, Guimarães JA, Francischetti IM. A catalog for the transcripts from the venomous structures of the caterpillar *Lonomia obliqua*: identification of the proteins potentially involved in the coagulation disorder and hemorrhagic syndrome. *Gene.* 2005; 355:11–27. [PubMed: 16023793]

- Warrell DA. Snake bite. *Lancet*. 2010; 375(9708):77–88. [PubMed: 20109866]
- Washburn MP, Wolters D, Yates JR 3rd. Large-scale analysis of the yeast proteome by multidimensional protein identification technology. *Nature Biotechnology*. 2001; 19:242–247.
- Williams D, Gutiérrez JM, Harrison R, Warrell DA, White J, Winkel KD, Gopalakrishnakone P. Global Snake Bite Initiative Working Group; International Society on Toxinology. The Global Snake Bite Initiative: an antidote for snake bite. *Lancet*. 2010; 375(9708):89–91. [PubMed: 20109867]
- Xu T, Venable JD, Park SK, Cociorva D, Lu B. ProLuCID, a fast and sensitive tandem mass spectrometry-based protein identification program. *Mol Cell Proteomics*. 2006; 5:S174.
- Zager RA. Rhabdomyolysis and myohemoglobinuric acute renal failure. *Kidney International*. 1996; 49(2):314–326. [PubMed: 8821813]
- Zager RA, Johnson AC, Becker K. Renal cortical hemopexin accumulation in response to acute kidney injury. *Am J Physiol Renal Physiol*. 2012; 303(10):F1460–472. [PubMed: 22993068]
- Zannin M, Lourenço DM, Motta G, Costa LRD, Grando M, Gamborgi GP, Noguti MA, Chudzinski-Tavassi AM. Blood coagulation and fibrinolytic factors in 105 patients with hemorrhagic syndrome caused by accidental contact with *Lonomia obliqua* caterpillar in Santa Catarina, Southern Brazil. *Thromb. Haemost.* 2003; 89:355–364. [PubMed: 12574817]

Figure 1. Functional parameters during *L. obliqua*-induced AKI

Rats were injected subcutaneously with PBS (controls - CTRL) or LOBE (1.5 or 1.0 mg/kg). After different times post administration the following parameters were determined: **(A)** Body weight, **(B)** water intake, **(C)** urine outoput, **(D)** glomerular filtration rate (GFR), **(E)** urinary density, **(F)** urine osmolality (urine osm), **(G)** plasma osmolality (plasma osm), **(H)** osmolar clearance (C_{osm}), **(I)** water free clearance (C_{H₂O}), **(J)** fractional water excretion (FE_{H₂O}), **(K)** fractional sodium excretion (FE_{Na⁺}) and **(L)** fractional potassium excretion (FE_{K⁺}). Data are presented as means $\pm$ SE (n=6/group). Significant differences: * $p < 0.05$ vs. CTRL and § $p < 0.05$ vs. LOBE (1.0 mg/kg, s.c.).

Figure 2. Proteinuria

(A) Rats were injected subcutaneously with PBS (controls - CTRL) or LOBE (1.5 or 1.0 mg/kg). After different times post administration the protein levels in urine were measured. Data are presented as means $\pm$ SE (n=6/group). Statistical differences of * $p < 0.05$ vs. CTRL and § $p < 0.05$ vs. LOBE (1.0 mg/kg, s.c.) were considered significant. (B) Representative urine samples from CTRL and rats treated with LOBE (1.5 mg/kg, s.c.) were analyzed by SDS-PAGE (8-20 %) under reducing conditions. (C) Toxins excreted in urine were detected by western-blot. Samples of urine from CTRL and rats treated with LOBE (1.5 mg/kg, s.c.) at 6 and 12 h post-venom injection were separated by SDS-PAGE and different toxins were detected by immunoreaction with polyclonal antibodies against LOBE. Toxins present in crude bristle extract are also showed (LOBE). The arrows indicate bands detected in urine samples at 6 and 12 h of envenomation. Molecular weight (MW) standards were shown on the left of figure B and C.

Figure 3. *L. obliqua* envenomation induces acute tubular necrosis

Representative kidney sections from control (CTRL) or envenomed animals (injected with 1.5 mg/kg, s.c.) are presented (A-G). Note the normal morphology of kidney from CTRL animal (A) in comparison to the progressive degenerative lesions of venom-treated rats (B-G). Increased acidophilia, dilation of renal tubules, loss of proximal brush border, cytoplasm vacuolation, nuclear pyknosis and desquamation of necrotic cells can be observed (D-F). Hyaline (arrowheads in C, D and G) and hematic (arrows in D) casts are also present inside renal tubules. Arrowheads in E indicate necrotic cells. Asterisks in D and F indicate the presence of a hyaline material within the Bowman's space and an inflammatory cell infiltrate and edema, respectively. All sections were stained with H&E. Magnification: 10 $\times$. (H) Levels of urinary γ -glutamyl transferase (γ -GT) activity were measured in rats injected with PBS or LOBE (1.5 or 1.0 mg/kg) at different times post-administration. Data are presented as means $\pm$ SE (n=6/group). Statistical differences of * p < 0.05 vs. CTRL and § p < 0.05 vs. LOBE (1.0 mg/kg, s.c) were considered significant.

Figure 4. *L. obliqua* envenomation induces renal tubular obstruction

Representative kidney sections from animals injected with LOBE (1.5 mg/kg, s.c.) showing details of tubular obstruction by hyaline and hematic casts and cellular debris at 12 h (**A**) and 24 h (**B**) post-venom injection. Hyaline casts (black arrowheads) are formed by a protein-rich material (predominantly serum albumin and hemoglobin), while hematic casts (black arrows) are formed by fragmented or intact erythrocytes. Due to tubular necrosis, the basement membrane in some tubules is disrupted, resulting in detachment of necrotic cells into the lumen (white arrowheads). All sections were stained with H&E. Magnification: 4 × (panel A) and 20 × (panel B).

Figure 5. *L. obliqua* envenomation induces renal inflammation and fibrosis

Light micrographs showing a marked inflammatory cell infiltrate (arrowheads) in the tubulo-interstitial region at 96 h (B) and glomerulus at 48 h (C) after LOBE injection (1.5 mg/kg, s.c.). There were no signs of inflammation in control (CTRL) animals (A). It was also observed an extensive peritubular (E) and interglomerular (F) collagen deposition at 96 h (regions stained in red), indicating fibrosis. Arrowheads in these panels indicate inflammatory infiltrate. There were no signs of fibrosis in CTRL rats (D). Stain: H&E (panels A-C) and picrosirius (panels D-F). Magnification: 10 × (panel A, B, D and E), 20 × (panel F) and 40 × (panel C).

Figure 6. Glomerular alterations

A. Light micrographs showing a time dependent increase in the deposition of a PAS positive material in glomerular capillaries of LOBE-injected animals (1.5 mg/kg, s.c.) in comparison to controls. Also note the increase in glomerular size. All sections were stained with PAS. Magnification: 40 ×. **B.** Thirty glomeruli from each animal injected with the dose of 1.5 mg/kg were used to quantify the mean area of glomerulus, glomerular tuft and Bowman's space of animals treated with the dose of 1.5 mg/kg. Data are presented as means ± SE (n=6/group). Statistical differences of * $p < 0.05$ in comparison to the respective control (C) were considered significant. **C.** Renal tissue factor activity was measured in control (C) and envenomed (1.5 mg/kg, s.c.) animals by generation of factor Xa (FXa). Data are presented as means ± SE (n=6/group). Statistical differences of * $p < 0.05$ were considered significant in comparison to the respective control.

Figure 7. Renal vascular permeability

Representative micrographs of a kidney blood vessel (A) and glomerulus (B) from animals injected with LOBE (1.5 mg/kg, s.c.) after 12 h of envenomation. Note the vascular leakage and edema (asterisks) and migration of inflammatory cells to damaged tissue (arrowheads). Also, the presence of a hyaline material inside the Bowman's space (asterisk in B) was associated with the increase in glomerular area observed at this time. All sections were stained with H&E. Magnification: 10 × (panel A) and 40 × (panel B). C. Evaluation of changes in renal vascular permeability were assessed by Evans blue dye extravasation. Results are expressed as µg Evans blue dye per 100 mg of renal tissue from control (CTRL) and LOBE-treated (1.5 mg/kg, s.c.) rats at 12 and 24 h post-venom administration. Data are presented as means ± SE (n=6/group). Statistical comparisons are indicated.

Figure 8. Immunohistochemical detection of *L. obliqua* venom in renal tissue

Positive immunohistochemical reaction was found in cortical and medullary regions of kidneys from rats injected with 1.5 mg/kg of LOBE (B-E). Venom was detected in glomerular capillaries (arrows in B), Bowman's capsule (arrow in the inset B), tubular brush border (arrows in the inset C), in intra-tubular casts (arrowheads in C) and also was present in cells of tubules in degeneration (arrowheads in the insets C and D). After 96 h the immunoreactivity for venom was weak and mainly localized in tubules (arrows in E). There was no immunoreactivity in the renal structures of control (CTRL) rats (A). Magnification: 10 × (panels A-E) and 40 × (insets in B-D) F. The amounts of venom detected in renal tissue was estimated by the area of positive immunohistochemical reaction. Thirty sections per rat were analyzed as described in material and methods. Data are presented as means ± SE (n=6/group).

Figure 9. Immunohistochemical detection of *L. obliqua* venom in renal vascular tissue
Positive immunohistochemical reaction was detected in renal arteries (**A**) and veins (**B**) of rats injected with LOBE (1.5 mg/kg, s.c.) at 6 h of envenomation. Note the presence of venom in perivascular connective tissue (arrows) and endothelium and smooth muscle cells (arrowheads). Magnification: 10 × (panel B) and 40 × (panel A).

Figure 10. Kidney proteins differentially expressed during *L. obliqua* envenomation

Unique or differentially expressed kidney proteins from control and envenomed (1.5 mg/kg, s.c.) animals were identified at 24 h post-venom injection by proteomic analysis. Exclusive and common proteins in each condition, as well as the total number of proteins identified are showed *via* a Venn diagram (A). Top canonical pathways of differentially expressed and unique proteins identified in control (CTRL) (B) and LOBE-treated (C) kidneys are shown. Those proteins functionally related to renal disease were also categorized accordingly to their roles in different types of renal pathologies (D). Each functional annotation is assigned to a significance score represented as *P* value (Fisher exact test) determining the probability that the association between the proteins in the data set and the canonical pathway or function in disease is explained by chance alone. The number of identified proteins (NIP) that belong to a particular canonical pathway or play a role in renal pathology is shown.

Table 1

Hemodynamic parameters

Parameter	Group	Time after envenomation (h)						
		0	2	6	12	24	48	96
MAP (mmHg)	CTRL	94.6±2.2	107.5±1.5	100.2±3.4	96.3±1.2	115±12.2	111.4±1.1	118±12.6
	LOBE (1.0 mg/kg, s.c.)	103±3.8	92.8±2.6	104.3±2.7	84.5±3.6	88±3.1 [*]	91.6±1.5 [*]	91.7±3.5 [*]
	LOBE (1.5 mg/kg, s.c.)	104±2.3	124.8±8.6 [§]	112.7±4.8	95.1±2.9	74.9±1.5 ^{*§}	81.5±1.1 [*]	95.1±1.4 [*]
HR (Beats/min)	CTRL	366±5.4	391±1.8	391±5.8	403±2.1	401±4.1	362±11.3	376±8.7
	LOBE (1.0 mg/kg, s.c.)	383±3.7	402±16.1	409±13.1	449±6.9 [*]	398±11.4	381±8.5	443±8.5 [*]
	LOBE (1.5 mg/kg, s.c.)	355±7.2	390±14.3	391±5.4	393±10.2 [§]	427±6.6 [§]	424±5.7 ^{*§}	369±3.3 [§]

MAP: Mean arterial pressure, HR: heart rate. Data are presented as means ± SE (n=6/group).

^{*} Statistical differences of $p < 0.05$ vs. CTRL (1.0 mg/kg, s.c) were considered significant.

[§] Statistical differences of $p < 0.05$ vs. LOBE (1.0 mg/kg, s.c) were considered significant.

Table 2
Unique and differentially expressed proteins identified in the kidneys of rats envenomed by *L. obliqua*

Accession number ^a	Protein description	Fold change ^b	p-value ^c	Biological Function and/or Participation in Disease	Reference
<i>Up-regulated</i>					
P24090	Alpha-2-HS-glycoprotein	6.034	0.004	Cysteine-type endopeptidase inhibitor activity/Acute-phase response/Regulation of inflammatory response/Up-regulation of Ahsg2 is associated with inflammation and tissue edema.	Bowler <i>et al.</i> , 2004
P02764	Alpha-1-acid glycoprotein (Orml)	5.093	0.001	Acute-phase inflammatory response to antigenic stimulus/ Up-regulated in response to LPS/ Orml is considered an early and accurate biomarker of AKI.	Devarajan <i>et al.</i> , 2010
P11517	Hemoglobin subunit beta-2 (HbB)	4.395	0.001	Oxygen transport/Up-regulation of Hb is nephrotoxic and associated with kidney injury in several diseases including sepsis, type 2 diabetes and <i>Crotalus</i> envenomation.	Zager, 1996; Pinho <i>et al.</i> , 2005; Larsen <i>et al.</i> , 2010
Q76MJ6	YBX1	3.37	0.04	DNA-binding/Upregulation of Ybx1 mRNA in kidney is associated with proximal tubular injury in male rat.	Thukral <i>et al.</i> , 2005
Q6LE95	Low molecular weight kininogen (LMWK)	3.282	0.001	Acute-phase inflammatory response/ Precursor of the active peptide bradykinin (BK)/Related to vasodilation, hypotension and increase of vascular permeability/Participates of the contact-phase reactions of blood coagulation and kallikrein-kinin system/ <i>L. obliqua</i> venom is able to release BK from LMWK.	Bohrer <i>et al.</i> , 2007; Pinto <i>et al.</i> , 2010
Q921A4	Cytoglobin (Cygb)	3.003	0.015	Heme and iron binding/Up-regulated in conditions of oxidative stress/Plays a role in the development of renal fibrosis/ Overexpression of Cygb improved histological renal injury, preserved renal function, and ameliorated fibrosis.	Mimura <i>et al.</i> , 2010; Nishi <i>et al.</i> , 2011
Q9R063	Peroxi-redoxin-5, mitochondrial (Pdx-5)	2.796	0.001	Involved in redox regulation of the cell/ Eliminating peroxides generated during metabolism/ Play a major role in the cellular response to oxidative stress/ Up-regulated in conditions of renal ischemia-reperfusion injury and hypoxia.	Godoy <i>et al.</i> , 2011
P70388	DNA repair protein (Rad50)	2.713	0.061	DNA repair/Plays a central role in double-strand break repair/Up-regulation of Rad50 mRNA is associated with hepatocellular carcinoma.	Stefanska <i>et al.</i> , 2011;
P02793	Ferritin light chain 1	2.563	0.001	Ferric iron binding/Iron homeostasis and metabolism.	Zager, 1996; Zager <i>et al.</i> , 2012
P08932	T-kininogen 2 (isoform 2 of LMWK)	2.128	0.001	Acute-phase inflammatory response/ Precursor of the active peptide bradykinin (BK)/Related to vasodilation, hypotension and increase of vascular permeability/Participates of the contact-phase reactions of blood coagulation and kallikrein-kinin system/ <i>L. obliqua</i> venom is able to release BK from LMWK.	Bohrer <i>et al.</i> , 2007; Pinto <i>et al.</i> , 2010
P02770	Serum Albumin (Alb)	2.096	0.018	The main protein of plasma/ Involved in molecular transport/Regulation of the colloidal osmotic pressure of blood/Up-regulation of serum albumin mRNA and albuminuria is associated with proximal tubular and glomerular injury and interstitial nephritis in male rat.	Thukral <i>et al.</i> , 2005
P09006	Serine protease inhibitor A3N	1.891	0.001	Acute-phase inflammatory response/ Induced by IL-1, IL-6 and interferon- γ / Up-regulated during sepsis.	Chinnaiyan <i>et al.</i> , 2001
P31977	Ezrin	1.706	0.032	Involved in connections of major cytoskeletal structures to the plasma membrane of	Thukral <i>et al.</i> , 2005;

Accession number ^a	Protein description	Fold change ^b	p-value ^c	Biological Function and/or Participation in Disease	Reference
P35704	Peroxisiredoxin-2 (Ptdx-2)	1.691	0.015	Involved in redox regulation of the cell/ Eliminating peroxides generated during metabolism/ Play a major role in the cellular response to oxidative stress/ Up-regulated in conditions of renal ischemia-reperfusion injury and hypoxia.	Trof <i>et al.</i> , 2006 Godoy <i>et al.</i> , 2011
Q6P9V2	Ferritin, heavy polypeptide 1	1.543	0.012	Stores iron/ Important for iron homeostasis.	Zager, 1996; Zager <i>et al.</i> , 2012
P14942	Glutathione S-transferase alpha-4 (GST)	1.516	0.032	Antioxidant enzyme/ Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles/It is a marker of proximal tubular injury.	Thukral <i>et al.</i> , 2005; Trof <i>et al.</i> , 2006
P01946	Hemoglobin subunit alpha-1/2 (HbA)	1.495	0.001	Oxygen transport/Up-regulation of Hb is nephrotoxic and associated with kidney injury in several diseases including sepsis, type 2 diabetes and <i>Crotalus</i> envenomation.	Larsen <i>et al.</i> , 2010; Pinho <i>et al.</i> , 2005; Zager, 1996
B1H2I6	Hemoglobin alpha, adult chain 2 (HbB)	1.495	0.001	Oxygen transport/Up-regulation of Hb is nephrotoxic and associated with kidney injury in several diseases including sepsis, type 2 diabetes and <i>Crotalus</i> envenomation.	Larsen <i>et al.</i> , 2010; Pinho <i>et al.</i> , 2005; Zager, 1996
P07335	Creatine kinase B-type (CK-B)	1.452	0.012	Creatine kinase isoenzymes play a central role in energy transduction in tissues. In the kidney localized primarily in the outer medulla in the thick ascending limb and distal convoluted tubule/ Marker of acute tubular injury.	Shashidharamurthy <i>et al.</i> , 2010
P01026	Complement C3 (Neutrophil chemotactic factor-1)	1.319	0.044	Plays a central role in the activation of the complement system/ Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways/ It is a mediator of local inflammatory process/ It induces the contraction of smooth muscle/ Increases vascular permeability/ Causes histamine release from mast cells and basophilic leukocytes/ Acts as a chemoattractant for neutrophils/ Complement activation increases tubulo-interstitial fibrosis of the kidney.	Tumberg <i>et al.</i> , 2006
P07483	Fatty Acid Binding Protein 3 (FABP3)	1.286	0.001	FABP3 are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters/ Upregulation of FABP3 mRNA in kidney is associated with proximal tubular injury in male rat.	Thukral <i>et al.</i> , 2005
Q9JJ19	Na ⁺ /H ⁺ exchange regulatory cofactor NHE-RF1	1.196	0.053	Sodium transporter localized in apical membrane of proximal tubular cells/Important in renal sodium transport and phosphate absorption/ It is a marker of tubular injury.	Trof <i>et al.</i> , 2006
P17475	Serpin 1 Alpha-1-antiproteinase	1.175	0.071	Acute-phase inflammatory response/ Response to hypoxia/ Inhibitor of serine proteases/ Has affinity for plasmin and thrombin/Serpine1 is involved in damage of tubulointerstitium and is up-regulated in cases of disseminated intravascular coagulation (DIC).	Eddy, 2004
P32038	Complement factor D (C3 convertase activator)	1.144	0.017	Plays a central role in the activation of the complement system/ Factor D cleaves factor B when the latter is complexed with factor C3b, activating the C3bbb complex, which then becomes the C3 convertase of the alternate pathway. Its function is homologous to that of C1s in the classical pathway/ Complement activation increases tubulo-interstitial fibrosis in the kidney.	Tumberg <i>et al.</i> , 2006

Accession number ^a	Protein description	Fold change ^b	p-value ^c	Biological Function and/or Participation in Disease	Reference
P04276	Vitamin D-binding protein (DBP)	1.088	0.011	It carries the vitamin D sterols and prevents polymerization of actin by binding its monomers. DBP associates with membrane-bound immunoglobulin on the surface of B-lymphocytes and with IgG Fc receptor on the membranes of T-lymphocytes/DBP knockout mice displayed decreased damage in renal cortex.	Safadi <i>et al.</i> , 1999
O88767	Protein DI-1	1.07	0.001	Protects cells against oxidative stress and cell death/ Eliminates hydrogen peroxide/ May act as an atypical peroxidase-like peroxidase that scavenges hydrogen peroxide.	Godoy <i>et al.</i> , 2011
P04041	Glutathione peroxidase 1	1.059	0.096	Antioxidant enzyme/ Protects the hemoglobin in erythrocytes from oxidative breakdown	Zager, 1996; Basnakian <i>et al.</i> , 2002; Khan, 2009
P62898	Cytochrome c; somatic (CytC)	1.003	0.004	Electron carrier protein/ Plays a role in apoptosis/ Suppression of the anti-apoptotic members or activation of the pro-apoptotic members of the Bcl-2 family leads to altered mitochondrial membrane permeability resulting in release of cytochrome c into the cytosol/ Increases apoptosis in kidney tubule epithelial cells.	Chang <i>et al.</i> , 2000
P20059	Hemopexin (Hpx)	0.951	0.065	Participate in heme metabolism/ Binds and neutralizes pro-oxidant free heme/Potentially impacts heme-iron-mediated tubular injury.	Zager <i>et al.</i> , 2012
P07632	Superoxide dismutase [cuzn] (SOD)	0.606	0.001	Antioxidant enzyme/ Destroys radicals which are normally produced within the cells and which are toxic to biological systems	Zager, 1996; Basnakian <i>et al.</i> , 2002; Khan, 2009
<i>Down-regulated</i>					
Q63213	Alpha-2u globulin	5.33	0.001	Associated with proximal tubule necrosis and protein droplet accumulation in male rats/ It is down-regulated in chronic hypoxia in the kidney.	Son <i>et al.</i> , 2008
P97615	Thioredoxin, mitochondrial	4.301	0.008	Has an anti-apoptotic function/ Plays an important role in the regulation of mitochondrial membrane potential/ Response to hypoxia/ Response to oxidative stress.	Godoy <i>et al.</i> , 2011
Q03626	Murine globulin-1	3.268	0.045	Peptidase inhibitor activity/ Acute-phase inflammatory response/ It is down-regulated by up to 70% during acute inflammation or tumor development.	Thukral <i>et al.</i> , 2005
Q64565	Alanine--glyoxylate aminotransferase 2, mitochondrial	3.112	0.08	Enzyme responsible by the metabolism of dimethylarginine (ADMA)/ ADMA is a potent inhibitor of nitric-oxide (NO) synthase/ This activity provides mechanism through which the kidney regulates blood pressure/ Its inhibition decreases the production of NO.	Caplin <i>et al.</i> , 2012
D3ZKG1	Methylmalonyl-CoA mutase, mitochondrial (Mut)	2.55	0.05	Involved in the degradation of several amino acids, odd-chain fatty acids and cholesterol via propionyl-CoA to the tricarboxylic acid cycle/ Knockout mouse for Mut gene displayed increased tubulointerstitial nephritis.	Chandler <i>et al.</i> , 2009
Q62980	Heparin sulfate proteoglycan 2 (HSPG2)	2.391	0.001	Cell adhesion protein/ Increase glomerular and mesangial kidney proliferation/ Positive regulation of endothelial cell proliferation/ Response to hypoxia.	Chen <i>et al.</i> , 2001
P11232	Thioredoxin	2.191	0.009	Participates in various redox reactions through the reversible oxidation of its active center/ Cell redox homeostasis.	Godoy <i>et al.</i> , 2011

Accession number ^a	Protein description	Fold change ^b	p-value ^c	Biological Function and/or Participation in Disease	Reference
Q920P6	Adenosine deaminase	2.139	0.019	Catalyzes the hydrolytic deamination of adenosine/ Its inhibition leads to accumulation of adenosine, which has a potent afferent arteriolar vasodilator effect through A _{2B} receptors located in glomerular vessels/ Adenosine also has a platelet aggregation inhibitory activity.	Feng and Navar, 2010/ Tofovic <i>et al.</i> , 1998/
Q64057	Alpha-aminoacidic semialdehyde dehydrogenase (ALDH)	2.083	0.05	Multifunctional enzyme mediating important protective effects/ Protects against hyperosmotic stress/ Protects cells from oxidative stress by metabolizing a number of lipid peroxidation-derived aldehydes.	Brocker <i>et al.</i> , 2010
P98158	Low-density lipoprotein receptor-related protein 2 (Lrp2)	1.786	0.03	Acts together with cubilin to mediate HDL endocytosis/Downregulation of Lrp2 protein in apical pole from kidney proximal tubule is associated with nephrosis in male rat.	Russo <i>et al.</i> , 2007
Q5XI73	Rho GDP dissociation inhibitor alpha (ARHGDIa)	0.522	0.005	Regulates the GDP/GTP exchange reaction of the Rho proteins by inhibiting the dissociation of GDP from them, and the subsequent binding of GTP to them/ Knockout mouse for ARHGDIa displayed increased tubulointerstitial nephritis and degeneration of renal tubular epithelial cells.	Togawa <i>et al.</i> , 1999
<i>Unique proteins</i>					
P30152	Neutrophil gelatinase-associated lipocalin (NGAL)	-	-	Iron-trafficking protein involved in multiple processes such as apoptosis, innate immunity and renal development/ Involved in apoptosis due to interleukin-3 (IL3) deprivation/ It is a marker and predictor of renal injury/ Up-regulation of NGAL is associated with ischemic acute kidney injury.	Trof <i>et al.</i> , 2006
P06762	Heme oxygenase 1 (Hmox-1)	-	-	Heme oxygenase cleaves the heme ring to form biliverdin. Biliverdin is subsequently converted to bilirubin by biliverdin reductase.	Canara and Soares, 2005; Zager, 1996
P01048	Rat T-kininogen 1 (Isoform 1 of LMWK)	-	-	Acute-phase inflammatory response/ Precursor of the active peptide bradykinin (BK)/Related to vasodilation, hypotension and increase of vascular permeability/Participates of the contact-phase reactions of blood coagulation and kallikrein-kinin system/ <i>L. obliqua</i> venom is able to release BK from LMWK.	Bohrer <i>et al.</i> , 2007; Pinto <i>et al.</i> , 2010
P14480-1	Isoform 1 of Fibrinogen beta chain (FbB)	-	-	Precursor of fibrin clots/ Polymerize into fibrin and acting as a cofactor in platelet aggregation/ Conversion of fibrinogen to fibrin is triggered by thrombin/ Up-regulation of Fb mRNA in kidney is associated with proximal tubular injury and intra-glomerular fibrin deposition.	Thukral <i>et al.</i> , 2005
P20961	Plasminogen activator inhibitor 1 (PAI-1)	-	-	Serine protease inhibitor. This inhibitor acts as 'bait' for tissue plasminogen activator, urokinase, protein C and matrix metalloproteinase-3/ It's the major control point in the regulation of fibrinolysis/ Up-regulation of PAI-1 is associated with renal thrombosis and fibrosis in cases of chronic allograft nephropathy.	Revelo <i>et al.</i> , 2005
P50115	Protein S100-A8	-	-	Calcium- and zinc-binding protein/ Plays a prominent role in the regulation of inflammatory processes and immune response/ Its proinflammatory activity includes recruitment of leukocytes, promotion of cytokine and chemokine production, and regulation of leukocyte adhesion and migration. Can induce cell death via autophagy and apoptosis and this occurs through the cross-talk of mitochondria and lysosomes via reactive oxygen species (ROS).	Basnakian <i>et al.</i> , 2002

Accession number ^a	Protein description	Fold change ^b	p-value ^c	Biological Function and/or Participation in Disease	Reference
P02680-1	Gamma-B of Fibrinogen gamma chain (FbG)	-	-	Precursor of fibrin clots/ Polymerize into fibrin and acting as a cofactor in platelet aggregation/ Conversion of fibrinogen to fibrin is triggered by thrombin/ Up-regulation of Fb mRNA in kidney is associated with proximal tubular injury and intra-glomerular fibrin deposition.	Thukral <i>et al.</i> , 2005
Q64268	Heparin cofactor 2 (HC-2)	-	-	Thrombin inhibitor/ It is activated by the glycosaminoglycans, heparin or dermatan sulfate. In the presence of the latter, HC-II becomes the predominant thrombin inhibitor in place of antithrombin III (AT)/ HC-II -levels increase in thrombosis and intravascular disseminated coagulation (DIC).	Takatsuka <i>et al.</i> , 2006
P84817	Mitochondrial fission 1 protein (MFP-1)	-	-	Promotes the fragmentation of the mitochondrial network and its perinuclear clustering/ Can induce cytochrome c release from the mitochondrion to the cytosol, ultimately leading to apoptosis. Also mediates peroxisomal fission.	Basnakian <i>et al.</i> , 2002

Proteins functionally related to renal disease were selected and their specific role in renal pathology was reviewed based on literature data. Complete proteome analysis was included in supplemental table 1-3.

^a Accession number in UniProtKB/Swiss-Prot data base.

^b Log(2) ratio change.

^c Spec count G-test p-value (proteins were considered differentially expressed with $p < 0.1$).