Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/jcot

Original Article

Evaluation of pain in bilateral total knee replacement with and without tourniquet; a prospective randomized control trial

Nishikant Kumar M.S. (Ortho), DNB (Ortho), MNAMS^{a,*},
 Chandrashekhara Yadav M.S. (Ortho)^b, Swapnil Singh M.S.^b,
 Ashok Kumar M.S., MRCS^b, Aruljothi Vaithlingam M.S.^b,
 Sanjay Yadav M.S.^b

^a Senior Research Associate, AIIMS, India^b Professor Orthopaedics, All India Institute of Medical Sciences, AIIMS, New Delhi, India

ARTICLE INFO

Article history:

Received 3 September 2014

Accepted 22 January 2015

Available online 24 February 2015

Keywords:

Total knee arthroplasty

Bilateral

Thigh pain

Tourniquet

ABSTRACT

Aim: Thigh pain following tourniquet application is a common complaint in early post operative period following total knee arthroplasty.

Method: Post operative Thigh pain was evaluated in 30 consecutive simultaneous bilateral total knee arthroplasty patients between July 2013 and January 2014. Patient thigh pain was evaluated with the VAS score. The scale was applied on first, second, third day & second and six weeks after surgery.

Result: There were statistically significant difference in VAS score in non-tourniquet group on first, second, third post operative day. We did not find statistically significant difference at Second and Six weeks post operatively.

Conclusion: This Randomized trial demonstrates that non-tourniquet use in TKA has less early postoperative pain and leads to better recovery.

Copyright © 2015, Delhi Orthopaedic Association. All rights reserved.

1. Introduction

Tourniquet pain is one of the most intriguing pains for the anesthesiologist and also a cause of concern for the orthopaedic surgeons.

Tourniquets have been used in total knee replacements since the procedure was first introduced. Tourniquet use during TKA surgery decreases total blood loss, creates a bloodless field and decreases operative time.¹

Complications reported as a consequence of tourniquet application are skin injury [skin abrasions, blisters, breaks & pressure necrosis], nerve injury, post tourniquet syndrome, Deep venous thrombosis, Postoperative pain, wound healing disorders and early infections.^{2–12} Pain is one of the most important outcome measures that contribute to patient dissatisfaction after TKA. Thigh pain has been a very common early postoperative complaint in our patients after having total knee Arthroplasty (TKA) using a tourniquet. The purpose of this study was to evaluate the hypothesis that tourniquet is

* Corresponding author. c/o Bipin Kumar; Sector-4/D, Quarter No. 1038, Bokarosteel city, Jharkhand, India.

E-mail address: knishikant@gmail.com (N. Kumar).

<http://dx.doi.org/10.1016/j.jcot.2015.01.095>

0976-5662/Copyright © 2015, Delhi Orthopaedic Association. All rights reserved.

associated with more postoperative thigh pain as compared with no use of tourniquet in total knee replacement.

2. Material and methods

Thirty consecutive patients undergoing bilateral primary total knee replacement at this institution were included in the study. Patients with severe cardiac comorbidities or neurologic problems were excluded.

Every patient signed an informed consent before being included in the study. All surgeries were performed by the same surgical team with standard technique. Epidural anesthesia was used in all the patients.

Both knees were prepared at the same time and a single set of instruments were used. One knee was operated first and then other by senior author [CSY]. All patients received perioperative antibiotics [Amoxicillin-clavulanic acid 1.2 gm $\times$ 10].

The thigh that will receive the tourniquet pressure will be randomized according to a coin toss just prior to the start of surgery. The tourniquet cuff used was an 85 cm long and 8.5 cm wide. One soft roll pad was applied between the skin and the cuff.

In Thigh 1 [tourniquet used side] was inflated to a pressure of systolic blood pressure plus 100 mm Hg and was released after the first quadriceps stitch. Hemostasis was achieved before closure. The wound was closed after wound irrigation and then elastic bandages were applied. In Thigh 2, the tourniquet was wrapped around the thigh but was not inflated during the surgery. In both thighs, the midline skin incision and a Mid vastus approach was used. More electrocautery was used to facilitate the surgery in patients without a tourniquet. All drill holes in the distal femur were filled with an autogenous bone plug. Suction drains were inserted before closure. Postoperative collection in drain was noted. Drain was removed when Postoperative collection was less than 100 ml for 12 h. Average time of drain removal was 48 h in both groups. After closure the knee was placed in a compressive dressing after the application of sterile cotton rolls and crepe bandage to the limb from groin to just above ankle. The knee was immobilised in extension. Postoperatively, analgesia was

given with epidural morphine 50 μ g/kg along with 0.1% bupivacaine in 10 ml normal saline. Along with that IV Diclofenac sodium was used twice daily for five days postoperatively and then shifted to oral formulation accordingly. Active isometric quadriceps and continuous passive movement were started on the second post-operative day, and walking with full weight bearing permitted as tolerated under the supervision of a physiotherapist. Patient thigh pain was evaluated with the VAS score. The pain scale was applied on the first, second, and third day, as well as 2 and 6 weeks after surgery.

For VAS score a card board was placed on patient's bed side of A4 size with double-side print or photocopy the next two diagrams ensuring that the lines are exactly 10 cm in length and superimposed.

3. Result & analysis

Lower limb pain in the thirty consecutive eligible patients undergone simultaneous bilateral primary TKAs were noted. There were 9 men and 21 women with a mean age of 58 years [range,45–69].

The tourniquet was used on left side in 14 cases (46.7%) and on right side in 16 cases (53.3%). There were no statistical differences in operating times in both thighs [THIGH 1 = 68.35 $\pm$ 3.75 min; THIGH 2 = 69.1 $\pm$ 3.04 min].

Mean VAS Score noted on first, second, third days and second, sixth weeks after surgery were 5.75, 4.4, 3.35, 1.7 and 1.05 in THIGH 1 and 3.95, 2.7, 2.05, 1.25, 1 in THIGH 2 respectively (Table 1, Fig. 1).

There were statistically significant difference in VAS score between THIGH 1 and THIGH 2 on the first day [$p < 0.001$], second day [$p < 0.001$], third day [$p < 0.001$] postoperatively. We did not find statistically significant difference at second and six weeks after surgery (Table 1 and Fig. 1).

4. Discussion

Pneumatic tourniquets have been associated with many different complications. The tourniquet causes tissue ischemia underneath and distal to the cuff, resulting in a variety of metabolic, cellular, and microvascular changes

Table 1 – VAS score in both the thigh with tourniquet and without tourniquet (VAS 1, 2, 3, 4, and 5 are VAS on 1st, 2nd, 3rd day, second week and sixth week postop.)

Mean VAS score	Tourniquet	No tourniquet	P-value
	Mean $\pm$ SD	Mean $\pm$ SD	
VAS 1	5.75 $\pm$ 1.45	3.95 $\pm$ 1.09	0.001
VAS 2	4.4 $\pm$ 1.75	2.7 $\pm$ 0.97	0.001
VAS 3	3.35 $\pm$ 1.42	2.05 $\pm$ 0.75	0.010
VAS 4	1.7 $\pm$ 0.65	1.25 $\pm$ 0.44	0.141
VAS 5	1.1 $\pm$ 0.30	1.0 $\pm$ 0.0	>0.05

that become more severe with prolonged tourniquet inflation.

Experimental and clinical evidence suggest that the injury is greater in the compressed and ischemic tissues underneath the cuff than in the strictly ischemic distal tissues.⁶ The degree of muscle injury induced by tourniquet compression and ischemia is related to a complex interaction of both the magnitude and duration of tissue compression.¹³ The combined effect of muscle ischemia, edema and microvascular congestion leads to the “post-tourniquet syndrome”.⁷ This syndrome was first described by Bruner (1951) and is characterized by weakness, stiffness, edema, dysesthesia (a condition in which an unpleasant sensation is produced by ordinary stimuli), and pain in the extremity.¹³ The recovery of knee flexion in the early postoperative period has been shown to be significantly better when a tourniquet has not been used.^{9,14,15} Skeletal muscle may be more sensitive to an ischemic condition than peripheral nerves.¹⁶

The use of the tourniquet is considered to be a risk for nerve injury in the affected extremity. There may be different reasons for this, but the injury is not considered to be mainly due to ischemia distal to the tourniquet cuff, but depends primarily on the combination of ischemia, pressure forces, and shear forces underneath the inflated cuff.^{4–6} A compressive injury can lead to different degrees of disordered nerve function and symptoms, ranging from paresthesias to complete sensory loss and from slight muscle weakness to paralysis.¹⁷ Large myelinated fibers are more sensitive to compression than small non-myelinated fibers. However, smaller non-myelinated fibers may be more sensitive to ischemia.¹⁸

Electromyographic (EMG) abnormalities were reported in 71% of patients after lower-extremity tourniquet use. The

Fig. 1 – VAS in both the thigh with tourniquet and without tourniquet.

evidence of denervation lasted from two to six months. EMG abnormalities correlated with impaired postoperative function and delayed recovery. Patients undergoing lower limb surgery without tourniquet use shown less postoperative pain.^{9,19,20}

Lower tourniquet cuff pressures have also been suggested to decrease postoperative pain.^{21,22} When a standard cuff pressure (350 mm Hg) was compared with 100 mm Hg plus systolic blood pressure (mean cuff pressure 230 mm Hg) the results showed significantly less postoperative thigh pain during the first three days of the lower pressure.²¹

Pain relief in the immediate postoperative period following TKA is crucial in facilitating early recovery. Prompt patient satisfaction will avoid a delay in physical therapy and will reduce hospitalization time.

5. Conclusion

This Randomised trial demonstrates that non-tourniquet use in TKA has less early postoperative pain and leads to better recovery. This may be due to absence of local compression and ischemic damage to underlying quadriceps muscle.

Conflicts of interest

All authors have none to declare.

REFERENCES

1. Tai TW, Chang CW, Lai KA, Lin CJ, Yang CY. Effects of tourniquet use on blood loss and soft-tissue damage in total knee arthroplasty: a randomized controlled trial. *J Bone Joint Surg Am.* 2012;94:2209–2215.
2. Din R, Geddes T. Skin protection beneath the tourniquet. A prospective randomized trial. *ANZ J Surg.* 2004;74:721–722.
3. Choudhary S, Koshy C, Ahmed J, Evans J. Friction burns to thigh caused by tourniquet. *Br J Plast Surg.* 1998;51:142–143.
4. Nitz AJ, Dobner JJ. Upper extremity tourniquet effects in carpal tunnel release. *J Hand Surg Am.* 1989;14:499–504.
5. Nitz AJ, Matulionis DH. Ultrastructural changes in rat peripheral nerve following pneumatic tourniquet compression. *J Neurosurg.* 1982;57:660–666.
6. Ochoa J, Fowler TJ, Gilliatt RW. Anatomical changes in peripheral nerves compressed by a pneumatic tourniquet. *J Anat.* 1972;113(Pt 3):433–455.
7. Kam PC, Kavanagh R, Yoong FF. The arterial tourniquet: pathophysiological consequences and anaesthetic implications. *Anaesthesia.* 2001;56:534–545.
8. Hernandez AJ, Almeida AM, Favaro E, Sguizzato GT. The influence of tourniquet use and operative time on the incidence of deep vein thrombosis in total knee arthroplasty. *Clinics (Sao Paulo).* 2012;67:1053–1057.
9. Abdel-Salam A, Eyres KS. Effects of tourniquet during total knee arthroplasty. A prospective randomised study. *J Bone Joint Surg Br.* 1995;77:250–253.
10. Clarke MT, Longstaff L, Edwards D, Rushton N. Tourniquet-induced wound hypoxia after total knee replacement. *J Bone Joint Surg Br.* 2001;83:40–44.
11. Smith TO, Hing CB. Is a tourniquet beneficial in total knee replacement surgery? A meta-analysis and systematic review. *Knee.* 2010;17:141–147.

12. Maffulli N, Testa V, Capasso G. Use of a tourniquet in the internal fixation of fractures of the distal part of the fibula. A prospective, randomized trial. *J Bone Joint Surg Am.* 1993;75:700–703.
13. Pedowitz RA, Gershuni DH, Schmidt AH, Friden J, Rydevik BL, Hargens AR. Muscle injury induced beneath and distal to a pneumatic tourniquet: a quantitative animal study of effects of tourniquet pressure and duration. *J Hand Surg Am.* 1991;16:610–621.
14. Li B, Wen Y, Wu H, Qian Q, Lin X, Zhao H. The effect of tourniquet use on hidden blood loss in total knee arthroplasty. *Int Orthop.* 2009;33:1263–1268.
15. Wakankar HM, Nicholl JE, Koka R, D'arcy JC. The tourniquet in total knee arthroplasty. A prospective, randomised study. *J Bone Joint Surg Br.* 1999;81:30–33.
16. Sapega AA, Heppenstall RB, Chance B, Park YS, Sokolow D. Optimizing tourniquet application and release times in extremity surgery. A biochemical and ultrastructural study. *J Bone Joint Surg Am.* 1985;67:303–314.
17. Lundborg G, Dahlin LB. Anatomy, function, and pathophysiology of peripheral nerves and nerve compression. *Hand Clin.* 1996;12:185–193.
18. Dahlin LB, Shyu BC, Danielsen N, Andersson SA. Effects of nerve compression or ischaemia on conduction properties of myelinated and non-myelinated nerve fibres. An experimental study in the rabbit common peroneal nerve. *Acta Physiol Scand.* 1989;136:97–105.
19. Konrad G, Markmiller M, Lenich A, Mayr E, Ruter A. Tourniquets may increase postoperative swelling and pain after internal fixation of ankle fractures. *Clin Orthop Relat Res.* 2005;433:189–194.
20. Ledin H, Aspenberg Per, Good L. Tourniquet use in total knee replacement does not improve fixation, but appears to reduce final range of motion. *Acta Ortop.* 2012;83:499–503.
21. Worland RL, Arredondo J, Angles F, Lopez-Jimenez F, Jessup DE. Thigh pain following tourniquet application in simultaneous bilateral total knee replacement arthroplasty. *J Arthroplasty.* 1997;12:848–852.
22. Younger AS, McEwen JA, Inkpen K. Wide contoured thigh cuffs and automated limb occlusion measurement allow lower tourniquet pressures. *Clin Orthop Relat Res.* 2004;428:286–293.