Published in final edited form as:

J Hosp Med. 2015 May ; 10(5): 311–313. doi:10.1002/jhm.2342.

Insights into Inpatients with Poor Vision: A High Value Proposition

Valerie G. Press, MD, MPH¹, Madeleine I. Matthiesen, MD², Alisha Ranadive, BS³, Seenu M. Hariprasad, MD⁴, David O. Meltzer, MD, PhD¹, and Vineet M. Arora, MD, MAPP¹

¹Department of Medicine, University of Chicago, Chicago, IL

²Medicine-Pediatrics, Massachusetts General Hospital, Boston, MA

³Pritzker School of Medicine, University of Chicago, Chicago, IL

⁴Department of Ophthalmology, University of Chicago, Chicago, IL

Abstract

Background—Vision impairment is an under-recognized risk factor for adverse events among hospitalized patients, yet vision is neither routinely tested nor documented for inpatients. Low-cost (\$8 and up) non-prescription ‘readers’ may be a simple, high-value intervention to improve inpatients’ vision. We aimed to study initial feasibility and efficacy of screening and correcting inpatients’ vision.

Methods—From June 2012 through January 2014 we began testing whether participants’ vision corrected with non-prescription lenses **for eligible participants** failing a vision screen (Snellen chart) performed by research assistants (RAs). Descriptive statistics and tests of comparison, including t-tests and chi-squared tests, were used when appropriate. All analyses were performed using Stata version 12 (StataCorps, College Station, TX).

Results—Over 800 participants’ vision was screened (n=853). Older (≥ 65 years; 56%) participants were more likely to have insufficient vision than younger (<65 years; 28%; p<0.001). Non-prescription readers corrected the majority of eligible participants’ vision (82%, 95/116).

Discussion—Among an easily identified sub-group of inpatients with poor vision, low-cost ‘readers’ successfully corrected most participants’ vision. Hospitalists and other clinicians working in the inpatient setting can play an important role in identifying opportunities to provide high-value care related to patients’ vision.

Background

Vision impairment is an under-recognized risk factor for adverse events among hospitalized patients.¹⁻³ Inpatients with poor vision are at increased risk for falls and delirium^{1,3} and have more difficulty taking medications.⁴⁻⁵ They may also be at-risk for being unable to read critical health information, including consent forms and discharge instructions, or decreased quality-of-life, such as simply ordering food from menus. Yet, vision is neither routinely

tested, nor documented, for inpatients. Low-cost (\$8-and-up) non-prescription reading glasses, known as ‘readers,’ may be a simple, high-value intervention to improve inpatients’ vision. We aimed to study initial feasibility and efficacy of screening and correcting inpatients’ vision.

Methods

From June 2012 through January 2014, research assistants (RAs) identified eligible (adult [18 years], English speaking) participants daily from electronic medical records as part of an ongoing study of general medicine inpatients measuring quality-of-care at the University of Chicago Medicine.⁶ RAs tested visual acuity using Snellen pocket charts (participants wore corrective lenses if available). Readers were tested with sequential fitting (+2/+2.25/+2.75/+3.25) until vision corrected (sufficient vision: at-least 20/50 acuity in one eye),⁷ for **eligible participants**. Eligible participants included those with insufficient vision who were not already wearing corrective lenses, **and no** documented blindness or medically severe vision loss (for whom non-prescription readers would be unlikely to correct vision deficiencies; e.g., cataracts, glaucoma). The study was approved by the University of Chicago Institutional Review Board (IRB #9967).

Of note, while readers are typically used in populations over 40 years, readers were fitted for all participants to assess their utility for any hospitalized adult patient. Upon completing the vision screening and readers interventions, participants received instruction on how to access vision care and how to obtain readers (if they corrected vision) after hospital discharge.

Descriptive statistics and tests of comparison, including t-tests and chi-squared tests, were used when appropriate. All analyses were performed using Stata version 12 (StataCorps, College Station, TX).

Results

Over eight-hundred participants’ vision was screened (n=853); the majority were female (56%, 480/853), African-American (76%, 650/853), with a mean age of 53.4 years (SD 18.7), consistent with our study site’s demographics. Over one-third (36%, 304/853) of participants had insufficient vision. Older (65 years) participants (56%, 136/244) were more likely to have insufficient vision than younger participants (28%, 168/608; p<0.001).

Participants with insufficient vision were wearing their own corrective lenses during the testing (150/304, 49%), did not use corrective lenses (53/304, 17%), or were without available corrective lenses (99/304, 33%) or. (Figure 1a)

One-hundred sixteen of 304 participants approached for the readers intervention were eligible (n=112 reported medical eye disease; n=65 were wearing lenses; n=11 refused or were discharged before intervention implementation).

Non-prescription readers corrected the majority of eligible participants’ vision (82%, 95/116). Most participants’ (81/116, 70%) vision corrected using the two lowest calibration

readers (+2/+2.25); another 14 participants' (12%) vision corrected with higher strength lenses (+2.75/+3.25). (Figure 1b)

Discussion

We found that over one-third of the inpatients we examined have poor vision. Furthermore, among an easily identified sub-group of inpatients with poor vision, low-cost 'readers' successfully corrected most participants' vision. While preventive health is not commonly considered an inpatient issue, hospitalists and other clinicians working in the inpatient setting can play an important role in identifying opportunities to provide high-value care related to patients' vision.

Several important ethical, safety, and cost considerations related to these findings exist. Hospitalized patients commonly sign written informed consents, therefore due diligence to ensure patients' ability to read and understand the forms is imperative. Further, inpatient delirium is common, particularly among older patients.⁸ Existing or new onset delirium occurs in up to 24-35% of elderly inpatients.⁸ Vision is an important risk factor for multifactorial inpatient delirium, and early vision correction has been shown to improve delirium rates, as part of a multi-component intervention.⁹ Hospital-related patient costs per delirium episode have been estimated at \$16,303-\$64,421.¹⁰ The cost of a multi-component intervention was \$6,341 per case of delirium prevented.⁹ While only one potentially critical component, the cost of readers (\$8+) would pale in comparison.¹ Vision screening takes approximately 2.25 plus 2-6 minutes for the readers' assessment, with little training and high fidelity. Therefore, this easily implemented, potentially cost-saving, intervention targeting inpatients with poor vision may improve patient safety and quality-of-life in the hospital and even after discharge.

Limitations of the study include considerations of generalizability, as participants were from a single, urban academic medical center. Additionally, long-term benefits of the readers intervention were not assessed in this study. Finally, RAs provided the assessments, therefore further work is required to determine costs of efficient large-scale clinical implementation through nurse-led programs.

Despite these study limitations, the surprisingly high prevalence of poor vision among inpatients is a call to action for hospitalists. Future work should investigate the impact and cost of vision correction on hospital outcomes such as patient satisfaction, reduced rehospitalizations, and decreased delirium.¹¹

Acknowledgements

We would like to thank several individuals for their assistance with this project. Andrea Flores, MA, Senior Programmer, helped with programming and data support. Kristin Constantine, BA, Project Manager, for help developing and implementing the database for this project and Edward Kim, BA, Project Manager for help with management of the database and data collection. We would like to thank Ainoa Coltri and the Hospitalist Project research assistants for assistance with data collection. We would like to thank Frank Zdravetz, MPH for assistance with the creation of figures and Nicole Twu, MS for assistance with the project. We would like to thank other students who helped to collect data for this project, including Allison Louis, Victoria Moreira and Esther Schoenfeld.

Conflicts of Interest and Financial Disclosures

Dr. Press is supported by a career development award from the National Heart Lung and Blood Institute (NIH K23HL118151). A pilot award from The Center on the Demography and Economics of Aging (CoA, National Institute of Aging P30 AG012857) supported this project. Dr. Matthiesen and Ms. Ranadive received support from the Summer Research Program funded by the National Institutes on Aging Short-Term Aging-Related Research Program (T35AG029795). Dr. Matthiesen also received funding from the Calvin Fentress Fellowship Program. Dr. Hariprasad reports being a consultant or participating on a speaker's bureau for Alcon, Allergan, Regeneron, Genentech, Optos, OD-OS, Bayer, Clearside biomedical, and ocular Therapeutix. Dr. Meltzer received funding from the National Institutes on Aging Short-Term Aging-Related Research Program (T35AG029795), and from the Agency for Healthcare Quality and Research through the Hospital Medicine and Economics Center for Education and Research in Therapeutics (U18 HS016967-01), and from the National Institute of Aging through a Midcareer Career Development Award (K24 AG031326-01), from the National Cancer Institute (KM1 CA156717) and from the National Center for Advancing Translational Science (2UL1TR000430-06). Dr. Arora received funding from the National Institutes on Aging Short-Term Aging-Related Research Program (T35AG029795), and National Institutes on Aging (K23AG033763).

REFERENCES

1. Oliver D, Daly F, Martin FC, McMurdo ME. Risk factors and risk assessment tools for falls in hospital in-patients: a systematic review. *Age Ageing*. 2004; 33(2):122–30. [PubMed: 14960426]
2. Press VG, Shapiro MI, Mayo AM, Meltzer DO, Arora VM. More than meets the eye: relationship between low health literacy and poor vision in hospitalized patients. *J Health Commun*. 2013; 18(Suppl 1):197–204. [PubMed: 24093356]
3. Inouye SK, Zhang Y, Jones RN, Kiely DK, Yang F, Marcantonio ER. Risk factors for delirium at discharge: development and validation of a predictive model. *Arch Intern Med*. 2007; 167(13):1406–13. [PubMed: 17620535]
4. Press VG, Arora VM, Shah LM, et al. Misuse of Respiratory Inhalers in Hospitalized Patients with Asthma or COPD. *J Gen Intern Med*. 2011; 26(6):635–42. [PubMed: 21249463]
5. Beckman AG, Parker MG, Thorslund M. Can elderly people take their medicine? *Patient Educ Couns*. 2005; 59(2):186–91. [PubMed: 16257624]
6. Meltzer D, Morrison Manning, W.G. et al. Effects of physician experience on costs and outcomes on an academic general medicine service: Results of a trial of hospitalists. *Ann Intern Med*. 2002; 137(11):866–74. [PubMed: 12458986]
7. Kaiser PK. Prospective evaluation of visual acuity assessment: a comparison of Snellen versus ETDRS charts in clinical practice (An AOS Thesis). *Trans Am Ophthalmol Soc*. 2009; 107:311–24. [PubMed: 20126505]
8. Levkoff SE, Evans DA, Liptzin B, Cleary PD, Lipsitz LA, Wetle TT, Reilly CH, Pilgrim DM, Schor J, Delirium Rowe J. The occurrence and persistence of symptoms among elderly hospitalized patients. *Arch Intern Med*. 1992; 152(2):334–40. [PubMed: 1739363]
9. Inouye SK, Bogardus ST Jr, Charpentier PA, Leo-Summers L, Acampora D, Holford TR, Cooney LM Jr. A multicomponent intervention to prevent delirium in hospitalized older patients. *N Engl J Med*. 1999; 340(9):669–76. [PubMed: 10053175]
10. Leslie DL, Marcantonio ER, Zhang Y, Leo-Summers L, Inouye SK. One-year health care costs associated with delirium in the elderly population. *Arch Intern Med*. 2008; 168(1):27–32. [PubMed: 18195192]
11. Whitson HE, Whitaker D, Potter G, et al. A Low-Vision Rehabilitation Program for Patients With Mild Cognitive Deficits. *JAMA Ophthalmol*. 2013; 131(7):912–9. [PubMed: 23619914]

Figure 1.

The proportion of patients screened with insufficient vision (**A**) and the proportion of eligible patients with vision corrected by readers (**B**). Note: Percentages may not add to 100 due to rounding.