CORRESPONDENCE

Long-Term Opioid Use in Non-Cancer Pain

by PD Dr. med. Winfried Häuser, Dr. med. Fritjof Bock,
Dr. med. Peter Engeser, Prof. Dr. med. Dr. rer. nat. Thomas Tölle,
Dipl.-Psych. Anne Willweber-Strumpf, Prof. Dr. med. Frank Petzke
in issue 43/2014

An Alarming Development

The “Long-Term Opioid Use in Non-Cancer Pain” guideline (1) was revised to provide a rational framework for the controversially discussed use of opioid analgesic drugs in the treatment of non-cancer pain. One of the key messages of the updated guideline is that approximately 25 percent of patients with osteoarthritis, diabetic polyneuropathy, post-herpetic neuralgia, and chronic low back pain benefit from long-term opioid treatment.

The number of opioid drug prescriptions has, as is generally known, significantly increased. Given the high prevalence and vague definition of chronic back pain, this development is alarming. In the United States, the number of deaths caused by opioid analgesic drugs has meanwhile surpassed those for heroin and cocaine significantly (2).

One third of the patients receiving long-term opioid treatment for non-cancer pain will develop an opioid use disorder (3). Both these findings relate to the fact that there is a gap between real-world patient care and the world of scientific studies and that these drugs are used in a too broad and uncritical manner. Opioid analgesics do not only carry risks for the patients who have been prescribed these drugs; they also link the legal world of medicine with the illegal world of street drugs, as they move from older patients who receive the prescriptions to younger drug addicts (secret “diversion” in the family, sale on the black market, “recycling” of spent opioid patches). It is imperative to filter out those patients in whom opioid analgesics are contraindicated (e.g. because of previous addictions of any kind, including smoking) or who require precautionary measures.

During “participative decision-making” (1), the physician must seek the agreement of the patient to discontinue the medication if the treatment targets are not met—and the physician has to execute on this, if necessary (4). Here, taking a thorough medical history is crucial which covers previous treatment experiences, personality and mental health comorbidities (especially addictions). The key recommendations fall short of adequately addressing these aspects.

Should all of this be ignored, we may face a development similar to that in the United States—and in gerontopsychiatry we will be confronted with the cumulative final common path of this problem.

DOI: 10.3238/arztebl.2015.0251a

REFERENCES

1. Häuser W, Bock F, Engeser P, Tölle T, Willweber-Strumpf A, Petzke F: Clinical practice guideline: Long-term opioid use in non-cancer pain. *Dtsch Arztebl Int* 2014; 111: 732–40.
2. Manchikanti L, Atluri S, Hansen H, et al.: Opioids in chronic non-cancer pain: have we reached a boiling point yet? *Pain Physician* 2014; 17: E1–10.
3. Juurlink DN, Dhalla IA: Dependence and addiction during chronic opioid therapy. *J Med Toxicol* 2012; 8: 393–9.
4. Ballantyne JC, LaForge KS: Opioid dependence and addiction during opioid treatment of chronic pain. *Pain* 2007; 129: 235–55.

Dr. med. Dirk K. Wolter

Psykiatrien i Region Syddanmark
Gerontopsykiatrisk Afdeling Haderslev, Haderslev, Dänemark
dirk.wolter@rsyd.dk

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

Dr. Wolter criticizes that the key recommendations of the guidelines do not adequately cover measures to filter out patients in whom the use of opioid analgesics is contraindicated because of previous addictions or who require special precautionary measures for that reason.

The long version of the guidelines contains the recommendations to take a psychosocial history (including addiction history) and to rely on participative decision-making when deciding on the indication for opioid treatment (1). Discontinuation of opioid analgesics if treatment targets are not met as well as gradual discontinuation of opioid treatment when patients misuse the prescribed opioid analgesics despite supportive treatment by an addiction specialist are actually key recommendations of the guideline. Furthermore, the long version of the guidelines contains recommendations regarding the management of patients with current substance abuse and regarding opioid withdrawal (1). Therefore, the authors of the guidelines believe that the addiction medicine aspects are adequately covered in the guideline.

There is no evidence to support an “opioid epidemic” in Germany. In the United States, opioid prescriptions climbed sharply by 400 percent between 2000 and 2010, accompanied by a rise in the numbers of opioid-associated hospital admissions and deaths (2). In Germany, long-term prescriptions of opioid analgesics for non-cancer pain increased by 23 percent during the same period (1). An expert group established by the Federal Government of Germany analyzed data from various sources (surveys of pharmacists and addiction recovery centers, health insurance data, Federal Criminal Police Office data) and found no evidence of tilidine or tramadol abuse in any significant numbers (3).

Case series of German pain centers revealed no problem behavior with regard to the way patients on long-term opioid treatment took their medication (1). Thus, no evidence is available to support the claim that one third of the patients receiving opioid analgesics for the treatment of non-cancer pain suffer from an “opioid use disorder”.

In line with Dr. Wolter’s concerns, the recommendations of the guidelines are intended to prevent an “opioid epidemic” from happening in Germany.

DOI: 10.3238/arztebl.2015.0251b

REFERENCES

1. Häuser W, Bock F, Hege-Scheuing G, et al.: Empfehlungen der aktualisierten Leitlinie LÖNTS – Langzeitanwendung von Opioiden bei nicht-tumorbedingten Schmerzen. Schmerz 2015; DOI 10.1007/s00482-014-1463-x.

2. Sullivan MD, Howe CQ: Opioid therapy for chronic pain in the United States: Promises and perils. Pain 2013; 154: 94–100.
3. Radbruch L, Glaeske G, Grond S, et al.: Topical review on the abuse and misuse potential of tramadol and tilidine in Germany. Subst Abuse 2013; 34: 313–20.
4. Häuser W, Bock F, Engeser P, Tölle T, Willweber-Strumpf A, Petzke F: Clinical practice guideline: Long-term opioid use in non-cancer pain. Dtsch Arztebl Int 2014; 111: 732–40.

PD Dr. med. Winfried Häuser

Innere Medizin I
Klinikum Saarbrücken gGmbH, Saarbrücken
whaeuser@klinikum-saarbruecken.de

Conflict of interest statement

PD Dr. Häuser owns mutual stock funds that may contain pharmaceutical company stock. He has been paid for serving on an advisory board for Daiichi Sankyo. He has received lecture honoraria from the Abbott, Janssen-Cilag, MSD, Sharp & Dohme, and Pfizer companies.