Published in final edited form as:

Tetrahedron Lett. 2015 March 11; 56(11): 1441–1444. doi:10.1016/j.tetlet.2015.01.188.

Synthesis of 1,2-dihydro-2-oxo-4-quinolinyl phosphates from 2-acyl-benzoic acids

Xinhua He^{a,b}, Tharcilla Aglio^a, Jeffrey R. Deschamps^c, Rachita Rai^a, and Fengtian Xue^{a,*}

^aDepartment of Pharmaceutical Sciences, University of Maryland School of Pharmacy, 20 Penn Street, Baltimore, Maryland 21201

^cNaval Research Laboratory, Code 6930, 4555 Overlook Ave., Washington, DC 20375

Abstract

We report a facile synthesis of 1,2-dihydro-2-oxo-4-quinolinyl phosphates (**1a-l**) starting from 2-acyl-benzoic acids (**2a-l**) in the presence of phosphoryl azides via a one-pot cascade reaction involving a Curtius rearrangement, an intramolecular nucleophilic addition of the enol carbon to the isocyanate intermediate, and an addition-elimination of the enol oxygen to the phosphoryl azide. During the reaction three new bonds are formed under mild conditions to yield 1,2-dihydro-2-oxo-4-quinolinyl phosphates in modest yields.

Keywords

Phosphoric ester; Quinolinone; 2-Acyl-benzoic acid; Curtius rearrangement; Phosphoryl azide

The 1,2-dihydro-2-oxo-4-quinolinyl phosphate (DOQP) system has found broad application in different areas. Phosphoric ester are present in medicinally active synthetic therapeutic agents including antibacterial and antifungal agents,¹ and as ATP kinase inhibitors for the treatment of cancer.² The 3-carboxanilide derivatives of DOQP have been used as prodrugs of aromatic hydrocarbon receptor (AhR) activators for the treatment of autoimmune disorders such as multiple sclerosis (MS).³ In addition, DOQP have been developed as phosphoryl transfer agents that can assist the transformation of AMP into ATP.⁴ The quinolinone group of DOQP is a well-documented scaffold that is present in numerous biologically active molecules such as farnesyltransferase inhibitors,⁵ β 2 agonists,⁶ 5TH antagonists,⁷ and immunomodulators.^{1b} Moreover, in the field of organic synthesis, enol phosphates are one of the most widely used starting material for Suzuki-Miyaura cross-coupling.⁸

© 2015 Published by Elsevier Ltd.

*Correspondence to Department of Pharmaceutical Sciences, University of Maryland School of Pharmacy, 20 Penn Street, Baltimore, Maryland 21201 Phone: 410-706-8521 fxue@rx.umaryland.edu.

^bCurrent Address: Beijing Institute of Pharmacology and Toxicology, 27 Taiping Road, Haidian District, Beijing, 100850

Supplementary data Supplementary data associated with this article can be found in the online version.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A common synthetic route to DOQP is via the Perkow reaction, in which the starting material α -halo ketones⁹ or α -OTf ketones in a modified Perkow reaction,¹⁰ is treated with a trialkyl phosphite ester to generate the enol phosphate product (Scheme 1). Although this method is widely used for the preparation of DOQP, it suffers from major drawbacks. First, the scope of the Perkow reaction is limited to trialkyl phosphite, therefore, compounds such as diphenyl phosphate cannot be synthesized using this method. Moreover, the preparation of starting material α -halo ketone **1** takes multiple steps. In the course of our research on novel PTP1B inhibitors for Type 2 diabetes (T2D), we performed a series of reactions intended to synthesize a collection of compounds based on a DOQP core.

Curtius rearrangement is widely used for the formation of isocyanate starting with acid azide.¹¹ The resulting isocyanate can react *in situ* with various nucleophiles such as amino groups or hydroxy groups to yield ureas and carbamates, respectively. Motivated by this cascade process, we hypothesize that when treated with an azide source (e.g., phosphoryl azide), 2-acylbenzoic acid **2** can undergo the Curtius rearrangement to generate isocyanate **3**, which cyclizes spontaneously *in situ* via the enol carbon atom to give intermediate **4**. Tautomerization of ketone **4** yields 4-hydroxyquinolinone **5**. The hydroxyl group of **5** attacks a second equiv of phosphoryl azide to generate aryl enol phosphate **1** (Scheme 2).

To test our hypothesis, in initial studies 2-acetylbenzoic acid (**2a**) was chosen as the starting material to optimize the reaction conditions for the formation of 2-hydroxyquinolin-4-yl diphenyl phosphate (**1a**) in the presence of 2 equiv of diphenyl phosphoryl azide (DPPA, Table 1). Using inorganic bases K₂CO₃ (entry 1) and Cs₂CO₃ (entry 2), no obvious formation of compound **1a** can be detected. However when tertiary amines triethylamine (Et₃N, entry 3) or diisopropylethylamine (DIPEA, entry 4) were used, we were able to isolate the desired product **1a** with 17% and 16% yields, respectively, along with small amounts of 4-methylene-1,4-dihydro-2H-benzo[d][1,3]oxazin-2-one **3a**. It is noted that compound **3a** decomposed as a solution in deuterated chloroform (CDCl₃). No obvious reaction was observed when pyridine was used as the based (entry 5). An additional equivalence of DPPA was not beneficial for the reaction (entry 6). After screening solvents for the reaction (entries 7-11), we found that the 1,2-dichloroethane (DCE) gave the highest yields (entry 11). Next, we kept the solvent of the reaction as DCE and re-visited the base used in the reaction (entries 12-16). We found that DIPEA gave the highest yield for reactions in DCE (entry 12). The reaction temperature was also studied (entries 17-19). It turned out that the original choice of 70 °C was optimal for the formation of compound **1a**. We noticed that at 110 °C a much more complex reaction was achieved with less compound **1a** formed (entry 20). When one equivalence of DPPA was used (entry 21), 4-hydroxyquinolin-2(1H)-one (**4** in Scheme 2) was isolated as the major product.¹²

Next we studied the scope of the reaction using the optimized conditions (Table 2). When 2-propionylbenzoic acid (**2b**) was treated with two equiv of DPPA at 70 °C for 12 h, the desired quinolinyl phosphate **1b** was isolated in modest yields in both toluene (entry 1) and DCE (entry 2). The chemical structure of compound **1b** has been confirmed by single crystal X-ray analysis (Figure 1). Decreased yields were observed when 2-pentanoylbenzoic acid (**2c**) was used as the starting material (entries 3-4), which is likely due to the unfavorable steric effect of the propyl group at the R₂ position of **2c**. The synthesis of starting material

2c has been reported previously.¹³ Effects of substitution on the phenyl ring have also been studied (entries 5-12). When 2-acetyl-3-chlorobenzoic acid (**2d**)¹³ and 2-acetyl-3-methylbenzoic acid (**2e**)¹³ were used as the starting material, low yields were obtained for reactions in both toluene and DCE (entries 5-8). The low yields are likely due to the steric hindrance of the 3-chloro or 3-methyl groups during the nucleophilic addition of the enol intermediate to the second equivalence of DPPA. Increased yields were achieved when the substituent on the phenyl ring moved further from the enol group (entries 9-12), with the formation of compounds **1f** and **1g** in modest yields. Other phosphoryl azides have also been used in the study (entries 13-22). 2-Propionylbenzoic acid reacted in the presence of diethyl phosphoryl azide¹⁴ (DEPA) to give compound **1h** in modest yields (entries 13-14). When compounds **2a** and **2b** were treated with 2-azido-5,5-dimethyl-1,3,2-dioxaphosphinane 2-oxide,¹⁴ the desired product **1i** and **1j** were isolated in modest yields, respectively (entries 15-18). Similarly modest yields were obtained when phenyl substituted starting material **2f** and **2g** were treated 2-azido-5,5-dimethyl-1,3,2-dioxaphosphinane 2-oxide (entries 19-22). The overall isolated yields of the listed reactions was not high, however, considering the complexity of the three-step cascade reaction and the mild reaction condition involved to perform the reaction, the current method represents a practical alternative method for the preparation of DOQP.

In summary, we have disclosed a practical method to synthesize DOQP starting from readily available starting material 2-acyl-benzoic acids. The reaction proceeds smoothly in various azide sources at 70 °C to give a collection of DOQP derivatives in modest yields.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We gratefully acknowledge funding from the Department of Pharmaceutical Sciences, University of Maryland School of Pharmacy. The X-ray crystallographic work was supported by NIDA through Interagency Agreement #Y1-DA1101 with the Naval Research Laboratory (NRL).

References

- (1) (a). Jhajharia P, Samota MK, Seth G. Phosphorus, Sulfur Silicon Relat. Elem. 2013; 188:1361.(b) Kieseier BC, Wiendl H. Cns Drugs. 2007; 21:483. [PubMed: 17521228]
- (2). Charrier JY, S. C. Davis CT, Studley J. PCT Int. Appl. 2013049719. 2013
- (3). Pettersson L. WO 2012050500 A1. 2012
- (4). Moriguchi T, Okada K, Seio K, Sekine M. Lett. Org. Chem. 2004; 1:140.
- (5). Caraglia M, Marra M, Viscomi C, D'Alessandro AM, Budillon A, Meo G, Arra C, Barbieri A, Rapp UR, Baldi A, Tassone P, Venuta S, Abbruzzese A, Tagliaferri P. Int. J. Cancer. 2007; 121:2317. [PubMed: 17657738]
- (6). Matera MG, Cazzola M. Drugs. 2007; 67:503. [PubMed: 17352511]
- (7). McCort G, Hoornaert C, Aletru M, Denys C, Duclos O, Cadilhac C, Guilpain E, Dellac G, Janiak P, Galzin AM, Delahaye M, Guilbert F, O'Connor S. Bioorg. Med. Chem. 2001; 9:2129. [PubMed: 11504649]
- (8) (a). Oppedisano A, Prandi C, Venturello P, Deagostino A, Goti G, Scarpi D, Occhiato EG. J. Org. Chem. 2013; 78:11007. [PubMed: 24083469] (b) Leidy MR, Hoffman JM, Pongdee R.

- Tetrahedron Lett. 2013; 54:6889. [PubMed: 24999287] (c) Nicolaou KC, Yu RC, Shi L, Cai Q, Lu M, Heretsch P. Org. Lett. 2013; 15:1994. [PubMed: 23550898] (d) Scarpi D, Bartali L, Casini A, Occhiato EG. Eur. J. Org. Chem. 2013; 1306
- (9) (a). Borowitz IJ, Yee KC, Crouch RK. The Journal of Organic Chemistry. 1973; 38:1713.(b) Lichtenthaler FW. Chem. Rev. 1961; 61:607.(c) Moinizadeh N, Klemme R, Kansy M, Zimmer R, Reissig HU. Synthesis-Stuttgart. 2013; 45:2752.(d) Paleta O, Pomeisl K, Kafka S, Klasek A, Kubelka V. Beilstein J. Org. Chem. 2005; 1
- (10). Pomeisl K, Kvicala J, Paleta O, Klasek A, Kafka S, Kubelka V, Havlicek J, Cejka J. Tetrahedron. 2007; 63:10549.
- (11). Brase S, Gil C, Knepper K, Zimmermann V. Angew. Chem. Int. Ed. 2005; 44:5188.
- (12). Ishida T, Kikuchi S, Yamada T. Org. Lett. 2013; 15:3710. [PubMed: 23819443]
- (13). Barcia JC, Cruces J, Estevez JC, Estevez RJ, Castedo L. Tetrahedron Lett. 2002; 43:5141.
- (14). Kim SH, Jung DY, Chang S. J. Org. Chem. 2007; 72:9769. [PubMed: 17979288]

Figure 1.
Single crystal structure of compound **1b**.

A. Previous methods **B. Current work** **Scheme 1.**

Synthetic methods to DOQP. (A) Perkow reaction and (B) current work.

Scheme 2. Synthetic plan to aryl enol phosphates **1** from 2-acyl-benzoic acids **2**.

Table 1Optimization of Conditions for 2-Hydroxyquinolin-4-yl Diphenyl Phosphate **1a**^a

entry	DPPA, equiv	solvent	base ^b	t, °C	yield 1a , % ^c
1	2.0	toluene	K ₂ CO ₃	70	< 5
2	2.0	toluene	Cs ₂ CO ₃	70	< 5
3	2.0	toluene	Et ₃ N	70	17
4	2.0	toluene	DIPEA	70	16
5	2.0	toluene	pyridine	70	< 5
6	3.0	toluene	Et ₃ N	70	18
7	2.0	DCM	Et ₃ N	70	15
8	2.0	CDCl ₃	Et ₃ N	70	19
9	2.0	<i>t</i> -BuOH	Et ₃ N	70	< 5
10	2.0	THF	Et ₃ N	70	6
11	2.0	DCE	Et ₃ N	70	34
12	2.0	DCE	DIPEA	70	47
13	2.0	DCE	DMAP	70	< 5
14	2.0	DCE	pyridine	70	< 5
15	2.0	DCE	K ₂ CO ₃	70	< 5
16	2.0	DCE	Cs ₂ CO ₃	70	< 5
17	2.0	DCE	DIPEA	40	14
19	2.0	DCE	DIPEA	90	46
20	2.0	DCE	DIPEA	110	26 ^d
21	1.0	DCE	DIPEA	70	< 5

^aThe reaction was carried out in a pressure tube with 1.0 mmol of **2a**, 2.0 mmol of DPPA and 2.0 mmol of base in 15 mL of solvent for overnight.^bK₂CO₃ = potassium carbonate, Cs₂CO₃ = cesium carbonate, Et₃N = triethylamine, DIPEA = diisopropyl ethylamine.^cDetermined by ¹H NMR using pyridin-3-ylmethanol as an internal standard.^dTLC indicated a complicated reaction significant amount of side products was observed.

Table 2Conditions and Yields for 4-(Phosphonoxy)-2(1*H*)-quinolinones **1b-l**^a

entry	substrate 2	azide	product 1	solvent	yield ^c , %
1		DPPA		toluene	40
2		DPPA		DCE	33
3		DPPA		toluene	15
4		DPPA		DCE	19
5		DPPA		toluene	8
6		DPPA		DCE	15
7		DPPA		toluene	28
8		DPPA		DCE	10
9		DPPA		toluene	38
10		DPPA		DCE	46
11		DPPA		toluene	27
12		DPPA		DCE	50
13		DPPA		toluene	25
14		DPPA		DCE	15

entry	substrate 2	azide	product 1	solvent	yield ^c , %
15				toluene	40
16				DCE	32
17				toluene	25
18				DCE	22
19				toluene	24
20				DCE	12
21				toluene	27
22				DCE	37

^aThe reaction was carried out with 1.0 mmol of **2b-l**, 2.0 mmol of (R₃O)₂PON₃, 2.0 mmol of DIPEA in 15.0 mL of DMF at 70 °C for 16 h.

^bThe NMR spectra of known compounds are identical to the reported ones in the corresponding references.

^cIsolated yield.