Published in final edited form as:

Policy Insights Behav Brain Sci. 2014 October ; 1(1): 239–247. doi:10.1177/2372732214549328.

Psychological Aspects of Contraception, Unintended Pregnancy, and Abortion

Julia R. Steinberg¹ and Lisa R. Rubin²

¹Department of Psychiatry, University of California, San Francisco, CA, USA

²Department of Psychology, New School for Social Research, New York, NY, USA

Abstract

The knowledge of important biopsychosocial factors linking women's reproductive health and mental health is increasing. This review focuses on psychological aspects of contraception, unintended pregnancy, and abortion because these are common reproductive health experiences in U.S. women's lives. This review addresses the mental-health antecedents and consequences of these experiences, mostly focusing on depression and depressive symptoms before and after unintended pregnancy and contraception. As mental-health antecedents, depressive symptoms predict contraceptive behaviors that lead to unintended pregnancy, and mental-health disorders have been associated with having subsequent abortions. In examining the mental-health consequences, most sound research does not find abortion or contraceptive use to cause mental-health problems. Consequently, evidence does not support policies based on the notion that abortion harms women's mental health. Nevertheless, the abortion-care setting may be a place to integrate mental-health services. In contrast, women who have births resulting from unintended pregnancies may be at higher risk of postpartum depression. Social policies (e.g., paid maternity leave, subsidized child care) may protect women from mental-health problems and stress of unplanned children interrupting employment, education, and pre-existing family care responsibilities.

Keywords

unintended pregnancy; abortion; contraception; depressive symptoms

Introduction

Women's reproductive health across the life span encompasses adolescent sexual health, pubertal timing, sexuality, sex education, menstrual cycle, unintended pregnancy and abortion, contraception, body image, pregnancy, childbirth, miscarriage, assisted

© The Author(s) 2014

Corresponding Author: Julia R. Steinberg, Department of Psychiatry, University of California, San Francisco, 3333 California St., Ste. 465, Box 0848, San Francisco, CA 94143-0848, USA. Julia.Steinberg@ucsf.edu.

Tweet: Contraception, unintended pregnancy, abortion, and psychological health: Does science inform reproductive policies?

Declaration of Conflicting Interests: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

reproductive technologies, menopause, and reproductive aging. Highlighting the importance of this topic, the World Health Organization (2009) reviewed research at the intersection of mental health and reproductive health. Here we provide a more critical, in-depth, and up-to-date focus on psychological aspects of contraception, unintended pregnancy, and abortion. As their psychology cannot be understood outside of their sociopolitical context, we focus on the U.S. context, where policies and clinical practice have much potential to be informed by scientific research.

Unintended pregnancy, abortion, and contraception are common experiences in U.S. women's lives. Approximately half (51%) of pregnancies in the United States are unintended (Finer & Zolna, 2014), one of the highest of developed nations (Singh, Sedgh, & Hussain, 2010); 30% of U.S. women will have an abortion by the age of 45 (Jones & Kavanaugh, 2011); and 99% of U.S. women who have sex will use contraception at some point in their reproductive life spans (Mosher & Jones, 2010). Few consider psychological aspects of each of these common experiences in women's lives, particularly mental-health-related aspects. In this review, the first two sections on contraception and unintended pregnancy focus on depression or depressive symptoms, because this is one of the most common and debilitating mental-health conditions among reproductive-aged women (National Comorbidity Survey Replication, 2007; U.S. Burden of Disease Collaborators, 2013). Moreover, in reproductive experiences, depression's role is better understood than other psychological conditions. Because abortion and mental-health research has been conducted for approximately 50 years, broader mental-health outcomes appear here.

Contraception, unintended pregnancy, and abortion, though distinct, are interrelated. Forty-three percent of unintended pregnancies are due to inconsistent or incorrect contraceptive use, and another 52% are due to non-use (Sonfield, Hasstedt, & Gold, 2014). Many unintended pregnancies end in abortion. Specifically, 40% of unintended pregnancies, excluding miscarriages, end in abortion, and 95% of abortions result from unintended pregnancies (Finer & Zolna, 2011).

Psychological Aspects of Contraception

Some psychological research examines whether hormonal contraception causes mood disorders, namely depression. In the 60s and 70s, when oral contraceptive pills used higher dose hormones, using oral contraception briefly seemed to increase depressive symptoms (Parry & Rush, 1979). In contrast, recent studies with current hormonal contraceptives, which use lower doses, find no association or that hormonal contraceptive use lowers depressive symptoms (Keyes et al., 2013; O'Connell, Davis, & Kerns, 2007). However, because 95% of unintended pregnancies are due to flaws in contraceptives use, the role of psychological health in contraceptive behaviors is also crucial. Research has begun to examine whether depressive symptoms influence women's contraceptive behaviors.

Depressive Symptoms' Influence on Contraceptive Behaviors

Psychology research previously focused on understanding risky sexual behavior, such as not using condoms or birth control without a significant focus on mental-health variables (Lopez, Tolley, Grimes, Chen, & Stockton, 2013; Miller, 1986; Reid & Aiken, 2011). This

research is now considering how depressive symptoms influence condom use or contraceptive behaviors. Experiencing more depressive symptoms can lead to contraceptive behaviors that are proximate determinants of unintended pregnancy (Alvy et al., 2011; Carvajal et al., 2014; Hall, Moreau, Trussell, & Barber, 2013a, 2013b; Hall, White, Rickert, Reame, & Westhoff, 2012).

Among adolescent girls or young adult women, more depressive symptoms are associated with not using contraception, inconsistent or incorrect use, and earlier discontinuation (Garbers, Correa, Tobier, Blust, & Chiasson, 2010; Hall et al., 2013a, 2013b; Hall et al., 2012). For instance, women with more depressive symptoms were less likely to leave a reproductive health visit with any method at all (Garbers et al., 2010). However, other work has found no association between depression and contraceptive behaviors among women (Callegari et al., 2014; Faisal-Cury, Menezes, & Huang, 2013; Shrier, Harris, Sternberg, & Beardslee, 2001), or that worse mental health (including depressive symptoms and diagnosis) was associated with using or choosing more effective contraceptive methods (Callegari et al., 2014; Steinberg et al., 2013). These different findings may be due to how the outcome was assessed or the chronic versus contextual nature of the depressive symptoms that were experienced. Depression and depressive symptoms may be associated with one behavior set in the context of a routine reproductive health visit, and another behavior set in the context of an acute stressor, such as when facing an unintended pregnancy. Further research is needed to fully understand individual and situational influences on contraceptive decisions and behaviors. Considering the significant body of research linking depression with cognition and behavior (Alvy et al., 2011; Bandura, 1997; Carvajal et al., 2014; Cléry-Melin et al., 2011; Hall et al., 2012; Jeong & Cranney, 2009; Joorman, Teachman, & Gotlib, 2009; Matt, Vázquez, & Campbell, 1992; Philipson, Wakefield, & Kasparian, 2011), psychological studies of depression and contraceptive use might investigate the influence of depressive symptoms on memory and information-processing of contraceptive side effects, motivation to use contraceptive consistently and correctly, and contraceptive self-efficacy, among other variables, to elucidate mechanisms linking depression with contraceptive use.

Policy and Clinical Implications

Depressive symptoms do not necessarily lead to selecting less effective contraception (Steinberg et al., 2013), so clinicians should not assume that depressed women will not be interested in effective methods. Nevertheless, because more depressive symptoms do sometimes lead to non-use, incorrect or inconsistent use, and discontinued use, encouraging long-acting reversible methods (e.g., intrauterine device or implant) may best help women with depressive symptoms protect themselves. Furthermore, policies that provide free Food and Drug Administration (FDA)-approved contraception may remove obstacles among women with more depressive symptoms. Consequently, the passage of the Patient Protection and Affordable Care Act of (2010) may enable women with more depressive symptoms to use more effective contraception.

Psychological Aspects of Unintended Pregnancy

A 2010 Institute of Medicine report on women's health identified unintended pregnancy as one research area showing little progress, in part because the proportion of pregnancies that are unintended—approximately half—has remained high and steady for 20 years (Guttmacher Institute, 2013). Here we follow the standard definition of unintended pregnancy: pregnancies earlier than women desired or not wanted ever in the future (Mosher, Jones, & Abma, 2012).

Psychological Effects of Unintended Pregnancy

Preventing unintended pregnancy not only supports women in fulfilling their reproductive goals but also avoids negative outcomes for mothers, children, and families, and saves costs to society (Gipson, Koenig, & Hindin, 2008; Singh & Darroch, 2012; Sonfield, Hasstedt, Kavanaugh, & Anderson, 2013). The research reflects the outcomes associated with two choices for women facing an unintended pregnancy, namely, mental-health effects of births and of abortions resulting from unintended pregnancies.

Births from unintended pregnancy connect with antepartum and postpartum depression (C. T. Beck, 2001; Lancaster et al., 2010). Factors common among those having unintended pregnancies and postpartum depression (e.g., poverty, being single, or being in a violent relationship) could drive this association (Abbasi, Chuang, Dagher, Zhu, & Kjerulff, 2013). Alternatively, women who decide to carry unintended pregnancies to term may experience more stress, disappointment, or ambivalence from the unanticipated interruptions to educational, career, or other life aspirations; this in turn may lead to more depressive symptoms during and after pregnancy. Because women who have unintended pregnancies are disproportionately poor and unmarried (Finer & Zolna, 2014), a large group may also experience financial strain from a birth due to an unintended pregnancy. Not all U.S. women receive leave after giving birth, so if a woman takes maternity leave, she may lose her job (Family and Medical Leave Act of 1993, 2006). In addition, because maternity leave is unlikely to be paid (Abt Associates, 2012), financial strain may lead to shorter maternity leave, associated with maternal depression (Feldman, Sussman, & Zigler, 2004).

Policy Implications of Unintended Pregnancy's Consequences

One policy to promote mental health after an unintended pregnancy is requiring paid maternity leave and subsidizing child care, particularly for those most in need (Finer & Zolna, 2014; Guttmacher Institute, 2013). All developed nations, except the United States, have paid maternity leave, and longer leave times (Livingston, 2013). Many also have government-subsidized child care, so women may return to work after childbirth. The 1993 Family and Medical Leave Act (2006) requires that *eligible* employers allow 12 weeks of *unpaid* maternity leave, though some employers that are not required to provide maternity leave do, and some provide *paid* maternity leave. Indeed, 21% of U.S. employers provide paid maternity leave (Abt Associates, 2012, Table 2.5.3). Approximately 35% of U.S. employees' workplaces offer paid maternity leave to most or all female employees (Abt Associates, 2012, Table 2.5.2). Of women who reported being employed during their last pregnancy, approximately 70% reported taking maternity leave (Child Health USA, 2013).

The “White House Summit on Working Families” (June 2014) and the related Council of Economic Advisors (CEA; 2014) report support improved federal policies. Although the CEA (2014) report highlights the impact of leave and flexibility on health, it does not mention mental health. However, policies requiring paid maternity leave and supporting child care could psychologically benefit women, enabling them to stay in the workforce, relieving stress of lost wages, avoiding separation from their newborn just after birth, and ensuring child care upon returning to the workplace. Stress, disappointment, and depression from having to halt a career or education may be averted if affordable and reliable child care was available.

Another way to promote mental health may be to assess pregnancy intention in the prenatal setting. Women who report that their pregnancy was not wanted could be screened for mental-health symptoms at various prenatal visits, enabling early intervention if needed.

Psychological Antecedents of Unintended Pregnancy

Many factors are associated with unintended pregnancy. Much U.S. research has focused on sociodemographic factors (Finer & Zolna, 2011, 2014). For instance, over the past 20 years, unintended pregnancy has increasingly concentrated among poor and low-income women (Guttmacher Institute, 2013). Thus, providing no-cost contraception, particularly long-acting reversible methods (e.g., intrauterine devices, implants) reduces the number of unintended pregnancies (Foster et al., 2011; Peipert, Madden, Allsworth, & Secura, 2012). But it does not eliminate them because methods are used inconsistently, used incorrectly, discontinued, or not chosen (Finer & Sonfield, 2013; Montouchet & Trussell, 2013). Understanding other factors, including psychological ones, that lead to unintended pregnancy is important for prevention.

Psychological health is one such factor contributing to unintended pregnancy. Some research finds that more depressive symptoms link to unintended pregnancies (Hall, Kusunoki, Gatny, & Barber, 2014; Nelson & Lepore, 2013), but not always (Tenkku et al., 2009). There are various reasons why depressive symptoms may influence unintended pregnancy. Women with more depressive symptoms may engage in contraceptive behaviors that put them at risk of unintended pregnancy (see previous section; for example, Hall et al., 2012). Women with more depressive symptoms may be more likely to have violent or controlling partners (Devries et al., 2013; Lehrer, Buka, Gortmaker, & Shrier, 2006), thereby having less reproductive autonomy and more likelihood of being subjected to reproductive coercion, leading to unintended pregnancy (Clark, Allen, Goyal, Raker, & Gottlieb, 2014; Miller et al., 2014).

Since the findings on the relationship between depressive symptoms and unintended pregnancy are disparate, this relationship may be more complicated than a linear one.¹ Consequently, the field is too nascent for drawing policy implications.

Does Psychological Health Influence Decision-Making Around an Unintended Pregnancy?

Although some research has investigated women's reasons for having abortions (Finer, Frohworth, Dauphinee, Singh, & Moore, 2005; Wokoma, Jampala, Bexhell, Guthrie, & Lindow, 2014), little research has examined what influences women's decision-making at

the time an unintended pregnancy occurs or how women make decisions about whether to terminate versus carry an unintended pregnancy to term. Some women may have decided before they are pregnant what they would do if faced with an unplanned pregnancy. Other women may decide what to do only when faced with it (Lydon, Dunkel-Schetter, Cohan, & Pierce, 1996). Just as depressive symptoms are associated with other health decisions and behaviors (DiMatteo, Lepper, & Croghan, 2000), including contraceptive decisions and behaviors (Hall et al., 2012; Steinberg et al., 2013), depressive symptoms may influence women's decision to carry versus terminate an unintended pregnancy.²

Psychological Aspects of Abortion

Research on psychological aspects of abortion revolves around two main questions: (a) Does abortion harm women's mental health? and (b) What personal and contextual factors influence post-abortion adjustment? Published literature answering the former question often uses trauma theory (Lee, 2003; Reardon, 1987; Robinson, Stotland, Russo, Lang, & Occhiogrosso, 2009; Speckhard, 1985). An alternative, the common-risk-factors approach (Steinberg & Finer, 2011), is largely a methodological and statistical argument: Women who have abortions have an increased prevalence of subsequent mental-health problems because of "third variables" (common risk factors). Studies answering the latter question, what influences post-abortion adjustment, usually use a stress and coping perspective (Adler, 1975; Major et al., 1990; Major, Richards, Cooper, Cozzarelli, & Zubek, 1998).

Public and scientific interest in abortion and mental health grew after President Reagan commissioned Surgeon General C. Everett Koop to write a report on the health effects of having an abortion. Instead of submitting a report to President Reagan, Koop wrote a letter stating "... the data do not support the premise that abortion does or does not cause or contribute to psychological problems" (Koop, 1989, p. 174) and testified at a congress subcommittee that possible psychological problems resulting from abortion are "miniscule from a public health perspective" (H.R., 1989).

Does Abortion Harm Women's Mental Health?

Since President Reagan's request and Surgeon General Koop's letter, three reviews by academic or professional organizations have examined evidence regarding whether abortion harms women's mental health (Adler et al., 1992; Major et al., 2009; National Collaborating Centre for Mental Health, 2011). All conclude that abortion (relative to birth of an unwanted or unintended pregnancy) does not increase women's risk of subsequent mental-health problems. Of others reviews (Charles, Polis, Sridhara, & Blum, 2008; Coleman, 2011; Robinson et al., 2009; Steinberg & Russo, 2009), only one concluded that the evidence demonstrate abortion harms women's mental health (Coleman, 2011). Because this article had several fatal methodological flaws and violated guidelines for meta-analysis, its conclusions have been rendered invalid (Kendall, Bird, Cantwell, & Taylor, 2012; Steinberg, Trussell, Hall, & Guthrie, 2012).

As mentioned above, two frameworks appear in the published literature. The first draws from trauma theory, and postulates that abortion is akin to war or rape, is necessarily traumatic, and is likely to lead to mental-health problems resembling post-traumatic stress

disorder (Lee, 2003; Reardon, 1987; Robinson et al., 2009; Speckhard, 1985). This framework originally found support from interviews with small numbers of women specifically recruited because they deemed their abortion a highly stressful experience (Speckhard, 1985). Since then, quantitative studies of poor quality have been published documenting the trauma of having an abortion (Major et al., 2009; National Collaborating Centre for Mental health, 2011; Steinberg & Russo, 2009). One example of a poor quality study (Coleman, Coyle, Shuping, & Rue, 2009) that has received much scientific scrutiny, been used to inform policy, and attracted the attention of the media concluded that abortion led to 12 different mental-health problems in the National Comorbidity Survey (Coleman et al., 2009, 2011; Planned Parenthood v. Rounds, 2012). Although the study (Coleman et al., 2009, 2011) reported that mental health was assessed after the abortion, this was not the case (Steinberg & Finer, 2012). Instead, *lifetime* mental-health diagnoses and *lifetime* abortion experience were correlated with one another, meaning that this study did not assess whether the abortion came before or after the mental-health outcome.

Sounder research examines the association between abortion versus birth and *subsequent* mental health, controlling as needed for other confounding factors. The simple association between abortion and subsequent mental-health disorders—including mood, anxiety, substance-use disorders, and suicidal ideation—appears open to alternative explanations (Steinberg, Becker, & Henderson, 2011; Steinberg & Finer, 2011; Steinberg, McCulloch, & Adler, 2014; Steinberg & Russo, 2008; Taft & Watson, 2008). In models that control for confounding factors—such as prior mental health, prior or current adverse experiences, and sociodemographic factors—the association between abortion and subsequent mental health evaporates for mood and anxiety disorders, and suicidal ideation (Fergusson, Horwood, & Boden, 2013; Steinberg et al., 2011; Steinberg & Finer, 2011; Steinberg et al., 2014; Steinberg & Russo, 2008; Taft & Watson, 2008). Furthermore, strong predictors of post-pregnancy (including post-birth and post-abortion) mental health were prior mental health, and adverse experiences. These results demonstrate that abortion is not a cause of these mental-health problems, and the simple association between abortion and these outcomes is spurious. For substance-use disorders, the association becomes non-significant when controlling for confounding factors (Fergusson et al., 2013), or reduces but remains significant (Steinberg & Finer, 2011; Steinberg et al., 2014). This association may have remained significant because important confounding factors (e.g., pregnancy intention, risk-taking personality) were not able to be controlled.

What Personal and Contextual Factors Influence Post-Abortion Adjustment?

From a stress and coping perspective, depending on personal, contextual, and social factors, a woman may perceive an abortion as more or less stressful (e.g., Major et al., 2000; Major et al., 1990; Major et al., 1998; Mueller & Major, 1989). How she perceives the experience then influences her coping strategies, and these coping strategies then influence subsequent adjustment. Some factors that predict better psychological adjustment include viewing the situation as less stressful, having social support, having more self-efficacy in coping, having a resilient personality, giving less meaning to the pregnancy, having had less intention to get pregnant, having no history of mental-health problems, and using coping strategies such as

acceptance of the situation and positive reframing (Major et al., 2000; Major et al., 1990; Major, Mueller, & Hildebrandt, 1985; Major et al., 1998).

Mental Health as an Antecedent to Abortion

Women who have abortions are more likely to have had prior mental-health problems (Steinberg & Finer, 2011; Steinberg et al., 2014; van Ditzhuijzen, ten Have, de Graaf, van Nijnatten, & Vollebergh, 2013). As noted, mental-health symptoms influence contraceptive behaviors and risk of unintended pregnancy. However, whether or not mental health influences pregnancy decision-making, once faced with an unintended pregnancy, remains unclear.

Policy and Clinical Implications Regarding the Psychology of Abortion

Generally, U.S. abortion policies are not informed by the psychological science. Some U.S. policies and court decisions assume that abortion harms women's mental health. For instance, eight states require that women seeking abortions be warned of a negative psychological effect from having an abortion. For example, in South Dakota, women are told that the medical risks of having an abortion include "depression and related psychological distress" and "suicide ideation and suicide" (South Dakota Codified Law, 2012). Although this law had been challenged in the courts, in 2012 the U.S. Court of Appeals for the Eighth Circuit upheld this law (*Planned Parenthood v. Rounds*). This ruling largely relied on findings from the aforementioned flawed Coleman et al.'s (2009) article. Furthermore, in the majority opinion of a 2007 Supreme Court decision to uphold a later abortion ban (*Gonzales v. Carhart*, 2007), Justice Kennedy stated " ... it seems unexceptionable to conclude ... Severe depression and loss of esteem can follow [an abortion]" (p. 29). These policies are not supported by the evidence-base in the abortion and mental-health field.

Besides policies, the psychological science should inform clinical practice. Although abortion does not cause mental-health problems, women having abortions are more likely to have had mental-health problems. Because women who have abortions are more likely to have histories of mental-health problems, and these mental-health problems may be untreated, the abortion-care setting may be a place to provide mental-health screening or interventions. Given that women seeking abortion are disproportionately low-income, this may be the only setting in which these women receive mental-health services.

Conclusion

Depression or depressive symptoms do not result from using hormonal contraception, nor are depression or other mental-health problems effects of having an abortion. Therefore, policies based on the notion that abortion harms women are unwarranted. Still, abortion clinics may be an appropriate setting to integrate mental-health services. In contrast, the science does connect unintended pregnancies resulting in births and subsequent depression. Consequently, policies that provide paid maternity leave and subsidized child care may relieve stress for many women having unintended pregnancies.

Research on depressive symptoms as an antecedent of contraception, unintended pregnancy, and abortion, is still emerging. However, most research finds depressive symptoms relate to contraceptive behaviors that lead to unintended pregnancy and abortion. Future research, beneficial to both psychology and reproductive health fields, could uncover mechanisms by which depressive symptoms lead to not using contraception, using it inconsistently or incorrectly, and discontinuing it. As well, such research could help understanding whether and how depressive symptoms or depression influences women's decision to carry to term or terminate an unintended pregnancy. Providing free contraception, as the Affordable Care Act does, may remove barriers perceived as more difficult by women with depressive symptoms.

Notes

1. It may be curvilinear, such that those with the least or no depressive symptoms and those with the most depressive symptoms are at lowest risk of unintended pregnancy, albeit for different reasons. Women with little or no depressive symptoms may be better able to protect themselves from unintended pregnancy by using contraception consistently and correctly, and not having relationship with controlling partners. Women with the most depressive symptoms may have little sexual desire and withdraw from social activities (American Psychiatric Association, 2013; Hamilton & Meston, 2013), thereby not engaging in frequent sexual activity with others and decreasing their chances of having an unintended pregnancy. Alternatively, some women with high levels of depressive symptoms may engage in sexual behaviors that lead to unintended pregnancies whereas others do not, in which case no relationship may be found between depressive symptoms and unplanned pregnancy.
2. Various social-psychological theories propose that affect, which is influenced by psychological health, influences peoples' decisions, judgments, and behaviors (Forgas, 1995; Kahneman, 2003; Schwarz & Clore, 2003; Singer & Salovey, 1988). For instance, women who are experiencing more depressive symptoms at the time of making a decision about an unintended pregnancy may be more likely to use this as information in their decision-making process (Forgas, 1995; Schwarz & Clore, 2003). Feelings of depression may be attributed to the unexpected pregnancy and possibility of having a child that is not wanted, leading women to choose to terminate their pregnancy. Alternatively, those with more mental-health symptoms may have more negative views of the world (Beck, 1987) and not want to bring a child into it. Future research that seeks to understand how women make decisions when faced with an unintended pregnancy is imperative to understand how to help women make the best decisions for themselves and to inform decision-making theories.

Acknowledgments

Funding: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by a National Institute of Child Health and Human Development (NICHD)/National Institutes of Health (NIH) K01 Career Development Award K01HD075834 awarded to Julia R. Steinberg.

References

- Abbasi S, Chuang CH, Dagher R, Zhu J, Kjerulff K. Unintended pregnancy and postpartum depression among first-time mothers. *Journal of Women's Health*. 2013; 22(5):412–416.
- Abt Associates. Cambridge, MA: Author; 2012. Family and Medical Leave in 2012: Technical Report. Retrieved from <http://www.dol.gov/asp/evaluation/fmla/FMLA-2012-Technical-Report.pdf>
- Adler NE. Emotional responses of women following therapeutic abortion. *American Journal of Orthopsychiatry*. 1975; 45:446–454. [PubMed: 1146976]
- Adler NE, David HP, Major BN, Roth SH, Russo NF, Wyatt GE. Psychological factors in abortion: A review. *American Psychologist*. 1992; 47:1194–1204. [PubMed: 1443858]
- Alvy LM, McKiran DJ, Mansergh G, Koblin B, Colfax GN, Flores SA, et al. Project Mix Study Group. Depression is associated with sexual risk among women who have sex with men, but is mediated by cognitive escape and self-efficacy. *AIDS and Behavior*. 2011; 15:1171–1179. [PubMed: 20217471]
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th. Washington, DC: Author; 2013.
- Bandura, A. Self-efficacy: The exercise of control. New York, NY: W.H. Freeman & Company; 1997.
- Beck AT. Cognitive models of depression. *Journal of Cognitive Psychotherapy*. 1987; 1(1):5–37.
- Beck CT. Predictors of postpartum depression: An update. *Nursing Research*. 2001; 50:275–285. [PubMed: 11570712]
- Callegari L, Zhao X, Nelson KM, Lehavot K, Bradley KA, Borrero S. Associations of mental illness and substance use disorders with prescription contraception use among women veterans. *Contraception*. 2014; 90:97–103. [PubMed: 24731860]
- Carvajal DN, Ghazarian SR, Crowne SS, Brown PB, Pokras OC, Duggan AK, Barnett B. Is depression associated with contraceptive motivations, intentions, and use among a sample of low-income Latinas? *Women's Health Issues*. 2014; 24:e105–e113. [PubMed: 24439935]
- Charles VE, Polis CB, Sridhara SK, Blum RW. Abortion and long-term mental health outcomes: A systematic review of the evidence. *Contraception*. 2008; 78:436–450. [PubMed: 19014789]
- Child Health USA. Washington, DC: Health Resources and Service Administration's Maternal and Child Health Bureau; 2013. Maternity leave. Retrieved from <http://mchb.hrsa.gov/chusa13/programs-policies/programs-policies.html>
- Clark LE, Allen RH, Goyal V, Raker C, Gottlieb AS. Reproductive coercion and co-occurring intimate partner violence in obstetrics and gynecology patients. *American Journal of Obstetrics & Gynecology*. 2014; 210(42):e1–e8.
- Cléry-Melin ML, Schmidt L, Lafargue G, Baup N, Fossati P, Pessiglione M. Why don't you try harder? An investigation of effort production in major depression. *PLoS One*. 2011; 6(8):e23178. [PubMed: 21853083]
- Coleman PK. Abortion and mental health: Quantitative synthesis and analysis of research published 1995–2009. *The British Journal of Psychiatry*. 2011; 199:180–186. [PubMed: 21881096]
- Coleman PK, Coyle CT, Shuping M, Rue VM. Induced abortion and anxiety, mood, and substance abuse disorders: Isolating the effects of abortion in the National Comorbidity Survey. *Journal of Psychiatric Research*. 2009; 43:770–776. [PubMed: 19046750]
- Coleman PK, Coyle CT, Shuping M, Rue VM. Corrigendum to “Induced abortion and anxiety, mood, and substance abuse disorders: Isolating the effects of abortion in the National Comorbidity Survey.”. *Journal of Psychiatric Research*. 2011; 45:1133–1134. *Journal of Psychiatric Research* 2009; 43:770–776.
- The Council of Economic Advisers. The economics of paid and unpaid leave. 2014. Retrieved from http://www.whitehouse.gov/sites/default/files/docs/leave_report_final.pdf
- Devries KM, Mak JY, Bacchus LJ, Child JC, Falder G, Petzold M, et al. Watts CH. Intimate partner violence and incident depressive symptoms and suicide attempts: A systematic review of longitudinal studies. *PLOS Medicine*. 2013; 10(5):e1001439. [PubMed: 23671407]

- DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for noncompliance with medical treatment: Meta-analysis of the effects of anxiety and depression on patient adherence. *Archives of Internal Medicine*. 2000; 160:2101–2107. [PubMed: 10904452]
- Faisal-Cury A, Menezes PR, Huang H. The relationship between perinatal psychiatric disorders and contraception use among postpartum women. *Contraception*. 2013; 88:498–502. [PubMed: 23507171]
- Family and Medical Leave Act of 1993, 29 U.S.C. §§ 2601–2654. 2006. Retrieved from <http://www.dol.gov/whd/regs/statutes/fmla.htm>
- Feldman R, Sussman AL, Zigler E. Parental leave and work adaptation at the transition to parenthood: Individual, marital, and social correlates. *Journal of Applied Developmental Psychology*. 2004; 25:459–479.
- Fergusson DM, Horwood LJ, Boden JM. Does abortion reduce the mental health risks of unwanted or unintended pregnancy? A reappraisal of the evidence. *Australia & New Zealand Journal of Psychiatry*. 2013; 47:819–827.
- Finer LB, Frohworth LF, Dauphinee LA, Singh S, Moore AM. Reasons U.S. women have abortions: Quantitative and qualitative perspectives. *Perspectives on Sexual and Reproductive Health*. 2005; 37:110–118. [PubMed: 16150658]
- Finer LB, Sonfield A. The evidence mounts on the benefits of preventing unintended pregnancy. *Contraception*. 2013; 87:126–127. [PubMed: 23305754]
- Finer LB, Zolna MR. Unintended pregnancy in the United States: Incidence and disparities, 2006. *Contraception*. 2011; 84:478–485. [PubMed: 22018121]
- Finer LB, Zolna MR. Shifts in intended and unintended pregnancies in the United States, 2001–2008. *American Journal of Public Health*. 2014; 104(Suppl. 1):S43–S48. [PubMed: 24354819]
- Forgas JP. Mood and judgment: The Affect Infusion Model (AIM). *Psychological Bulletin*. 1995; 117(1):39–66. [PubMed: 7870863]
- Foster DG, Biggs MA, Rostovtseva D, de Bocanegra HK, Darney PD, Brindis CD. Estimating the fertility effect of expansions of publicly funded family planning services in California. *Women's Health Issues*. 2011; 21:418–424. [PubMed: 21802962]
- Garbers S, Correa N, Tobier N, Blust S, Chiasson MA. Associations between symptoms of depression and contraceptive method choices among low-income women at urban reproductive health centers. *Maternal and Child Health Journal*. 2010; 14:102–109. [PubMed: 19067135]
- Gipson JD, Koenig MA, Hindin MJ. The effects of unintended pregnancy on infant, child, and parental health: A review of the literature. *Studies in Family Planning*. 2008; 39:18–38. [PubMed: 18540521]
- Gonzales v. Carhart Co., 550 U.S. 124 (2007)
- Guttmacher Institute. New York, NY: Author; 2013. Unintended pregnancy in the United States (Fact Sheet). Retrieved from <http://www.guttmacher.org/pubs/FB-Unintended-Pregnancy-US.html#6>
- Hall KS, Kusunoki Y, Gatny H, Barber J. The risk of unintended pregnancy among young women with mental health symptoms. *Social Science & Medicine*. 2014; 100:62–71. [PubMed: 24444840]
- Hall KS, Moreau C, Trussell J, Barber J. Role of young women's depression and stress symptoms in their weekly use and nonuse of contraceptive methods. *Journal of Adolescent Health*. 2013a; 53:241–248. [PubMed: 23582524]
- Hall KS, Moreau C, Trussell J, Barber J. Young women's consistency of contraceptive use—Does depression or stress matter? *Contraception*. 2013b; 88:641–649. [PubMed: 23850075]
- Hall KS, White KO, Rickert VI, Reame N, Westhoff C. Influence of depressed mood and psychological stress symptoms on perceived oral contraceptive side effects and discontinuation in young minority women. *Contraception*. 2012; 86:518–525. [PubMed: 22673038]
- Hamilton LD, Meston CM. Chronic stress and sexual function in women. *The Journal of Sexual Medicine*. 2013; 10:2443–2454. [PubMed: 23841462]
- H.R. (1989). 135 Cong. Rec. E 906.
- Institute of Medicine. Women's health research: Progress, pitfalls, and promise. Washington, DC: The National Academies Press; 2010.

- Jeong JM, Cranney J. Motivation, depression, and naturalistic time-based prospective remembering. *Memory*. 2009; 17(7):732–741. [PubMed: 19637094]
- Jones RK, Kavanaugh ML. Changes in abortion rates between 2000 and 2008 and lifetime incidence of abortion. *Obstetrics & Gynecology*. 2011; 117:1358–1366. [PubMed: 21606746]
- Joorman J, Teachman BA, Gotlib IH. Sadder and less accurate? False memory for negative material in depression. *Journal of Abnormal Psychology*. 2009; 118:412–417. [PubMed: 19413415]
- Kahneman D. A perspective on judgment and choice: Mapping bounded rationality. *American Psychologist*. 2003; 58:697–720. [PubMed: 14584987]
- Kendall T, Bird V, Cantwell R, Taylor C. To meta-analyse or not to meta-analyse: Abortion, birth and mental health. *The British Journal of Psychiatry*. 2012; 200:12–14. [PubMed: 22215864]
- Keyes KM, Cheslack-Postava K, Westhoff C, Heim CM, Haloossim M, Walsh K, Koenen K. Association of hormonal contraceptive use with reduced levels of depressive symptoms: A national study of sexually active women in the United States. *American Journal of Epidemiology*. 2013; 178:1378–1388. [PubMed: 24043440]
- Koop CE. The US Surgeon General on the health effects of abortion. *Population and Development Review*. 1989; 15:172–175. [PubMed: 11656046]
- Lancaster CA, Gold KJ, Flynn HA, Yoo H, Marcus SM, Davis MM. Risk factors for depressive symptoms during pregnancy: A systematic review. *American Journal of Obstetrics & Gynecology*. 2010; 202:5–14. [PubMed: 20096252]
- Lee, E. Abortion, motherhood, and mental health: Medicalizing reproduction in the United States and Great Britain. New York, NY: Aldine de Gruyter; 2003.
- Lehrer JA, Buka S, Gortmaker S, Shrier LA. Depressive symptomatology as a predictor of exposure to intimate partner violence among US female adolescents and young adults. *Archives of Pediatrics & Adolescent Medicine*. 2006; 160:270–276. [PubMed: 16520446]
- Livingston, G. Washington, DC: Pew Research Center; 2013. Among 38 nations, U.S. is the outlier when it comes to paid parental leave. Retrieved from <http://www.pewresearch.org/fact-tank/2013/12/12/among-38-nations-u-s-is-the-outlier-when-it-comes-to-offering-paid-parental-leave/>
- Lopez, LM.; Tolley, EE.; Grimes, DA.; Chen, M.; Stockton, LL. Theory-based interventions for contraception; Cochrane Database of Systematic Review. 2013. p. CD007249 Retrieved from <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD007249.pub3/abstract?sessionid=29C5B85483D479E853D02A54721539E8.f03t03>
- Lydon J, Dunkel-Schetter C, Cohan CL, Pierce T. Pregnancy decision making as a significant life event: A commitment approach. *Journal of Personality and Social Psychology*. 1996; 71(1):141–151. [PubMed: 8708996]
- Major B, Appelbaum M, Beckman L, Dutton MA, Russo NF, West C. Abortion and mental health: Evaluating the evidence. *American Psychologist*. 2009; 64:863–890. [PubMed: 19968372]
- Major B, Cozzarelli C, Cooper L, Zubek J, Richards C, Wilhite M, Gramzow RH. Psychological responses of women after first-trimester abortion. *Archives of General Psychiatry*. 2000; 57:777–784. [PubMed: 10920466]
- Major B, Cozzarelli C, Sciacchitano AM, Cooper ML, Test M, Mueller PM. Perceived social support, self-efficacy, and adjustment to abortion. *Journal of Personality and Social Psychology*. 1990; 59:452–463. [PubMed: 2231279]
- Major B, Mueller P, Hildebrandt K. Attributions, expectations, and coping with abortion. *Journal of Personality and Social Psychology*. 1985; 48:585–599. [PubMed: 3989663]
- Major B, Richards C, Cooper ML, Cozzarelli C, Zubek J. Personal resilience, cognitive appraisals, and coping: An integrative model of adjustment to abortion. *Journal of Personality and Social Psychology*. 1998; 74:735–752. [PubMed: 9523416]
- Matt GE, Vázquez C, Campbell WK. Mood-congruent recall of affectively toned stimuli: A meta-analytic review. *Clinical Psychology Review*. 1992; 12:227–255.
- Miller E, McCauley HL, Tancredi DJ, Decker MR, Anderson H, Silverman JG. Recent reproductive coercion and unintended pregnancy among female family planning clients. *Contraception*. 2014; 89:122–128. [PubMed: 24331859]
- Miller WB. Why some women fail to use their contraceptive method: A psychological investigation. *Family Planning Perspectives*. 1986; 18(1):27–32. [PubMed: 3803546]

- Montouchet C, Trussell J. Unintended pregnancies in England in 2010: Costs to the National Health Service (NHS). *Contraception*. 2013; 87:149–153. [PubMed: 22878145]
- Mosher, WD.; Jones, J. Vital Health Statistics, Series 23. Washington, DC: National Center for Health Statistics; 2010 Aug. Use of contraception in the United States: 1982–2008.
- Mosher, WD.; Jones, J.; Abma, JC. National Health Statistics Reports, No 55. Atlanta, GA: Centers for Disease Control and Prevention; 2012. Intended and unintended births in the United States: 1982–2010.
- Mueller P, Major B. Self-blame, self-efficacy, and adjustment to abortion. *Journal of Personality and Social Psychology*. 1989; 57(6):1059–1068. [PubMed: 2614659]
- National Collaborating Centre for Mental Health. Induced abortion and mental health: A systematic review of the mental health outcomes of induced abortion, including their prevalence and associated factors. London, England: Academy of Medical Royal Colleges; 2011.
- National Comorbidity Survey Replication. Table 2. 12-month prevalence of DSM- IV/WMH-CIDI disorders by sex and cohort. 2007. Retrieved from <http://www.hcp.med.harvard.edu/ncs/>
- Nelson DB, Lepore SJ. The role of stress, depression, and violence on unintended pregnancy among young urban women. *Journal of Women's Health*. 2013; 22:673–680.
- O'Connell K, Davis AR, Kerns J. Oral contraceptives: Side effects and depression in adolescent girls. *Contraception*. 2007; 75:299–304. [PubMed: 17362710]
- Parry BL, Rush AJ. Oral contraceptives and depressive symptomatology: Biologic mechanisms. *Comprehensive Psychiatry*. 1979; 20:347–358. [PubMed: 37042]
- Patient Protection and Affordable Care Act, 42 U.S.C. § 18001 (2010).
- Peipert JF, Madden T, Allsworth JE, Secura GM. Preventing unintended pregnancies by providing no-cost contraception. *Obstetrics & Gynecology*. 2012; 120:1291–1297. [PubMed: 23168752]
- Philipson S, Wakefield CE, Kasparian NA. Women's knowledge, beliefs, and information needs in relation to risks and benefits associated with use of the oral contraceptive pill. *Journal of Women's Health*. 2011; 20:635–642.
- Planned Parenthood v. Rounds Co., 668 F. 3d 889 (8th Cir. 2012).
- Reardon, DC. Aborted women: Silent no more. Chicago, IL: Loyola University Press; 1987.
- Reid AE, Aiken LS. Integration of five health behavior models: Common strengths and unique contributions to understanding condom use. *Psychology & Health*. 2011; 26:1499–1520. [PubMed: 21678166]
- Robinson GE, Stotland NL, Russo NF, Lang JA, Occhiogrosso M. Is there an “abortion trauma syndrome”? Critiquing the evidence. *Harvard Review of Psychiatry*. 2009; 17:268–290. [PubMed: 19637075]
- Schwarz N, Clore GL. Mood as information: 20 years later. *Psychological Inquiry: An International Journal for the Advancement of Psychological Theory*. 2003; 14:296–303.
- Shrier LA, Harris SK, Sternberg M, Beardslee WR. Associations of depression, self-esteem, and substance use with sexual risk among adolescents. *Preventive Medicine*. 2001; 33:179–189. [PubMed: 11522159]
- Singer JA, Salovey P. Mood and memory: Evaluating the network theory of affect. *Clinical Psychology Review*. 1988; 8:211–251.
- Singh, S.; Darroch, JE. New York, NY: Guttmacher Institute and United Nations Population Fund; 2012. Adding it up: Costs and benefits of contraceptive services—Estimates for 2012. Retrieved from <http://www.guttmacher.org/pubs/AIU-2012-estimates.pdf>
- Singh S, Sedgh G, Hussain R. Unintended pregnancy: Worldwide levels, trends, and outcomes. *Studies in Family Planning*. 2010; 41:241–250. [PubMed: 21465725]
- Sonfield, A.; Hasstedt, K.; Gold, RB. New York, NY: Guttmacher Institute; 2014. Moving forward: Family planning in the era of health reform. Retrieved from <http://www.guttmacher.org/pubs/family-planning-and-health-reform.pdf>
- Sonfield, A.; Hasstedt, K.; Kavanaugh, ML.; Anderson, R. New York, NY: Guttmacher Institute; 2013. The social and economic benefits of women's ability to determine whether and when to have children. Retrieved from www.guttmacher.org/pubs/social-economic-benefits.pdf
- South Dakota Codified Law, § 34–23A–10.1(1) e (i & ii), 2005, enacted 2012.

- Speckhard, AC. Doctoral dissertation. 1985. The psycho-social aspects of stress following abortion. Retrieved from University Microfilms International. No. 8519300)
- Steinberg JR, Becker D, Henderson JT. Does the outcome of a first pregnancy predict depression, suicidal ideation, or lower self-esteem? Data from the National Comorbidity Survey. *American Journal of Orthopsychiatry*. 2011; 81:193–201. [PubMed: 21486261]
- Steinberg JR, Finer LB. Examining the association of abortion history and current mental health: A reanalysis of the National Comorbidity Survey using a common-risk-factors model. *Social Science & Medicine*. 2011; 72:72–82. [PubMed: 21122964]
- Steinberg JR, Finer LB. Coleman, Coyle, Shuping, and Rue make false statements and draw erroneous conclusions in analyses of abortion and mental health using the National Comorbidity Survey. *Journal of Psychiatric Research*. 2012; 46:407–409. [PubMed: 22348853]
- Steinberg JR, McCulloch CE, Adler NE. Abortion and mental health: Findings from the National Comorbidity Survey-Replication. *Obstetrics & Gynecology*. 2014; 123(2, Pt. 1):263–270. [PubMed: 24402590]
- Steinberg JR, Russo NF. Abortion and anxiety: What's the relationship? *Social Science & Medicine*. 2008; 67:238–252. [PubMed: 18468755]
- Steinberg JR, Russo NF. Evaluating research on abortion and mental health. *Contraception*. 2009; 80:500–503. [PubMed: 19913142]
- Steinberg JR, Trussell J, Hall KS, Guthrie K. Fatal flaws in a recent meta-analysis on abortion and mental health. *Contraception*. 2012; 86:430–437. [PubMed: 22579105]
- Steinberg JR, Tschann JM, Henderson JT, Drey EA, Steinauer JE, Harper CC. Psychological distress and post-abortion contraceptive method effectiveness level chosen at an urban clinic. *Contraception*. 2013; 88:717–724. [PubMed: 24094755]
- Taft AJ, Watson LF. Depression and termination of pregnancy (induced abortion) in a national cohort of young Australian women: The confounding effect of women's experience of violence. *BMC Public Health*. 2008; 8:75–82. [PubMed: 18302736]
- Tenkku LE, Flick LH, Homan S, Loveland Cook CAL, Campbell C, McSweeney M. Psychiatric disorders among low-income women and unintended pregnancies. *Women's Health Issues*. 2009; 19:313–324. [PubMed: 19733801]
- U.S. Burden of Disease Collaborators. The state of US health, 1990-2010: Burden of diseases, injuries, and risk factors. *Journal of the American Medical Association*. 2013; 310:591–608. [PubMed: 23842577]
- van Ditzhuijzen J, ten Have M, de Graaf R, van Nijmegen CHCJ, Vollebergh WAM. Psychiatric history of women who have had an abortion. *Journal of Psychiatric Research*. 2013; 47:1737–1743. [PubMed: 23941742]
- Wokoma TT, Jampala M, Bexhell H, Guthrie KA, Lindow SW. Reasons provided for requesting a termination of pregnancy in the UK. *Journal of Family Planning & Reproductive Health Care*. 2014 Advance online publication. 10.1136/jfprhc-2013-100745
- World Health Organization. Mental health aspects of women's reproductive health: A global review of the literature. Geneva, Switzerland: Author; 2009.

Key Points

- Science provides evidence relevant to reproductive policies regarding contraception, unintended pregnancy, and abortion.
- More depressive symptoms predict contraceptive behaviors that lead to unintended pregnancy, and mental-health disorders predict subsequent abortions.
- Contraception and abortion themselves do not predict mental-health issues.
- Depression does follow unintended pregnancies ending in birth.