Published in final edited form as:

Am J Obstet Gynecol. 2015 May ; 212(5): 633.e1–633.e6. doi:10.1016/j.ajog.2015.01.005.

Pregnancy Induces Persistent Changes in Vascular Compliance in Primiparous Women

Erin A MORRIS, MD¹, Sarah A HALE, PhD¹, Gary J BADGER, MS², Ronald R MAGNESS, PhD³, and Ira M BERNSTEIN, MD¹

¹Obstetrics, Gynecology and Reproductive Sciences, University of Vermont, Burlington, VT

²Medical Biostatistics, University of Vermont, Madison WI

³Department of Obstetrics and Gynecology Perinatal Research Laboratories, University of Wisconsin, Madison WI

Abstract

Objective—Pregnancy induces rapid, progressive and substantial changes to the cardiovascular system. The low recurrence risk of preeclampsia, despite familial predisposition, suggests an adaptation associated with pregnancy that attenuates risk for subsequent preeclampsia. We aimed to evaluate the persistent effect of pregnancy on maternal cardiovascular physiology.

Study Design—Forty-five healthy nulliparous women underwent baseline cardiovascular assessment preconception and repeated an average of 30 months later. After baseline evaluation, 17 women (Preg) conceived singleton pregnancies and all delivered at term. The remaining 28 women comprised the non-pregnant control group (NP). We measured mean arterial blood pressure (MAP), cardiac output (CO), plasma volume (PV), pulse wave velocity (PWV), uterine blood flow (UBF), and flow-mediated vasodilation (FMD) at each visit.

Results—There was a significant decrease in mean arterial pressure from the prepregnancy visit to postpartum in women with an interval pregnancy (pregnancy: 85.3 ± 1.8 , postpartum: 80.5 ± 1.8 mm Hg), with no change in NP subjects (visit 1: 80.3 ± 1.4 , visit 2: 82.8 ± 1.4 mm Hg), ($p = .002$). Pulse wave velocity was significantly decreased in women with an interval pregnancy (pregnancy: 2.73 ± 0.05 , postpartum: 2.49 ± 0.05 m/s), as compared to those without an interval pregnancy (visit 1: 2.56 ± 0.04 , visit 2: 2.50 ± 0.04 m/s), ($p = .005$). We did not observe a residual effect of pregnancy on cardiac output, plasma volume, uterine blood flow or flow-mediated vasodilation.

© 2015 Published by Elsevier Inc.

Corresponding Author: Ira M. Bernstein, Department of Obstetrics, Gynecology and Reproductive Sciences, Smith 410, Fletcher Allen Health Care, 111 Colchester Avenue, Burlington, VT 05401, Phone: (802) 847-5112, Fax: (802) 847-2772, ira.bernstein@uvm.edu.

The authors report no conflict of interest.

Findings presented at the Society for Maternal-Fetal Medicine's (SMFM) 34th Annual Meeting – The Pregnancy Meeting™, New Orleans, LA, February 3–8, 2014.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conclusion—Our observations of decreased mean arterial pressure and reduced arterial stiffness following pregnancy suggest a significant favorable effect of pregnancy on maternal cardiovascular remodeling. These findings may represent a mechanism by which preeclampsia risk is reduced in subsequent pregnancies.

Keywords

cardiovascular remodeling; hypertension; preeclampsia; pregnancy; vascular stiffness

Introduction

Pregnancy is known to induce rapid, progressive and substantial changes to the cardiovascular system, ultimately facilitating successful pregnancy outcome. Women who develop hypertensive disorders during pregnancy are considered to have “failed” the cardiovascular stress test of pregnancy and likely represent a sub-population where cardiovascular accommodation was inadequate (1–4). Risk for preeclampsia, a failure of the pregnancy stress test, is highest for first pregnancies, and decreases with subsequent pregnancies, with the length of the inter-pregnancy interval being an important determinant of subsequent pregnancy-associated hypertension (5–8). This phenomenon suggests that there may be an adaptation associated with prior pregnancy that attenuates the risk for preeclampsia in subsequent pregnancies, and whose influence is time-limited.

Evaluation of the persistence of pregnancy-induced cardiovascular remodeling is limited. Few studies describe changes due to pregnancy with reference to prepregnancy values. Clapp and Capeless reported a persistent increase in cardiac output as well as decreased peripheral vascular resistance at one year postpartum, with greater changes from pre-pregnancy values in parous women (9). Our laboratory has evaluated the pattern of changes in blood pressure that accompany a first pregnancy compared with a subsequent pregnancy and found reduced mean arterial pressure prior to and throughout subsequent pregnancies compared with first pregnancies (10). In those observations, the interval of time between pregnancies was inversely related to the difference in mean arterial pressure between pregnancies, such that as time between pregnancies increased, the decrease in mean arterial pressure became smaller. This phenomenon was confirmed in a larger cohort where mean arterial pressure was found to be reduced in second pregnancies in a time-dependent fashion (11). These observations are consistent with reports that longer inter-pregnancy intervals are associated with increased risk for the recurrence of preeclampsia (12–15).

The current report examines broad cardiovascular indices, including measures of uterine blood flow (UBF) and flow-mediated vasodilation (FMD) in nulliparous women prior to pregnancy and approximately one year postpartum compared to nulliparous women without an interval pregnancy, with matched elapsed time points. We aimed to evaluate the effect of pregnancy on postpartum cardiovascular physiology. We hypothesized that pregnancy would be associated with a reduction in arterial stiffness that would persist postpartum.

Materials and Methods

Thirty four nulligravid women interested in conception were enrolled in this research study through an open advertisement. Women were provided with ovulation detection kits (Quidel Corporation, San Diego, CA) to assist with achieving a successful conception. Twenty-eight women, also nulligravid, not interested in conception, were recruited as the control/non-pregnant group. All subjects were young (18–40) healthy, nonsmokers with regular menstrual cycles at the time of enrollment. None of the women had a history of hypertension, autoimmune disease, diabetes or other disorders known to affect blood pressure. Subjects provided a list of current medications and supplements at each study visit. Subjects taking anti-hypertensives or other medications known to affect blood pressure were not eligible for study participation. A questionnaire completed at the time of the initial visit evaluated for a family history of hypertension, stroke, myocardial infarction, clotting disorder and type 2 diabetes.

Of the pregnancy group, thirty women subsequently conceived. Eight subjects conceived before baseline pre-pregnancy studies were performed; one subject had a first trimester miscarriage; one subject was lost to follow-up and three subjects did not return for their postpartum visit. The remaining 17 subjects, all of whom conceived singleton pregnancies, had complete prepregnancy assessments, term pregnancy outcomes and a postpartum visit, comprise the current report. One woman developed complicated hypertension (CH) during the third trimester with new-onset hypertension ($>140/90$ mmHg), elevated liver enzymes, elevated uric acid concentration (>5 mg/dl) and fetal growth restriction, and had iatrogenic delivery at 37 weeks. Women were enrolled consecutively over a 33 month period, from May 2004 through February 2007. Prior to each study visit subjects were provided with a 3500-mg sodium-balanced diet for 72 hours. Each subject was asked to abstain from alcohol and caffeine beginning at least 24 hours before the study and to avoid the use of decongestants and non-steroidal medications beginning at least 48 hours before the study. All study visits in non-pregnant women were performed during the follicular phase. The research protocols were approved by the University of Vermont Human Investigational Committees. All women studied provided written informed consent.

Each periodic assessment was conducted between 8 AM and 10 AM. Subjects were admitted to the University of Vermont Clinical Research Center on the day of the study after an overnight fast. For subjects' prepregnancy visit, first-void urine was obtained to confirm non-pregnant state. Following height and weight determination, subjects rested in the supine position for the remainder of the study and for a minimum of 30 minutes before physiologic assessment.

Blood pressure

Pulse pressure (PP) and mean arterial pressure (MAP) were measured by continuous non-invasive tonometric radial artery blood pressure monitoring, employing the Colin Pilot 9200 device (San Antonio, TX), with auto-standardization to brachial artery measurements.

Cardiac Output

Cardiac output was determined by Doppler echocardiographic examination. Doppler-derived forward stroke volume (FSV) across the aortic valve was calculated as the product of the left ventricular outflow tract area and the outflow tract velocity time integral as assessed by pulsed Doppler using previously described methods (9). Five complete spectral envelopes with the largest Doppler shift were recorded and averaged for each patient. The Doppler stroke volume was calculated as the product of the outflow tract area and the velocity time integral. CO is expressed as mL/minute after integration of stroke volume with pulse rate.

Plasma Volume

Plasma volume (PV) was calculated using the Evans Blue Dye method, as previously described (16). An 18-gauge intravenous saline lock was placed in the antecubital vein for baseline blood draw, administration of Evans blue dye and subsequent blood draws. The PV was reported in total mL, and corrected for body mass index (BMI).

Pulse wave velocity

Brachial pulse wave forms were obtained by Doppler ultrasound using a 10 MHz transducer. Time from EKG R wave to peak systolic flow in the brachial artery was used to determine pulse wave velocity (PWV), relative to the distance from the heart to brachial artery. (The distance from the heart to brachial artery was calculated post hoc as height*0.33.)

Uterine Blood Flow

Uterine blood flow was assessed using color Doppler ultrasound with an 8.0 MHz transvaginal transducer employing a Vivid 7 General Electric ultrasound unit, (Milwaukee WI). Uterine artery measurements were obtained lateral to the cervix at the level of the internal os. Vessel diameter was measured between the inner surfaces of the vessel walls during real time power color Doppler imaging. Five estimates of vessel diameter were made for each uterine artery. For determination of average mean velocity, angle correction was employed and all angles of sono-incidence were less than or equal to 60 degrees. Five velocity measurements were taken of each vessel and average mean velocity was calculated over an observation window of 1–3 minutes. Mean vessel diameter was used to calculate volumetric blood flow after integration with average mean velocity to express flow in mL/min: (time averaged mean velocity) X cross sectional area X 60. Right and left uterine artery blood flow calculations were added to obtain total uterine blood flow.

Flow-mediated vasodilation

These studies are functional in-vivo assessments of endothelial health as reflected by the ability of the endothelium to generate a shear stress-mediated vasodilatory signal in response to an acute increase in volumetric flow. Studies were conducted by establishing baseline brachial artery vessel diameter under direct visualization with a 10 MHz linear transducer employing a Vivid 7 General Electric ultrasound unit (Milwaukee, WI). A mean of 3 measurements was accepted as the best estimate of diameter. Visualization of vessels was made 2 finger breadths above the antecubital fossa of the arm. A blood pressure cuff was then placed across the distal extremity and inflated to 50 mmHg above systolic pressure for

a period of 5 minutes. Following deflation, vessel diameter (3 estimates at each time point) was measured at 50, 60, and 70 seconds and the mean of these measurements was accepted as the post-restriction maximal diameter. The vasodilatory response was calculated as the difference in diameters, pre and post restriction, divided by the baseline diameter, resulting in flow-mediated dilation percent (FMD%) and expressed as a function of directly measured shear stress as described below.

Shear stress measurements

Blood samples were collected in 8 ml EDTA tubes and sent to the laboratory of Dr. Ronald R. Magness at University of Wisconsin-Madison for analysis. Blood viscosities were determined using a cylindrical spindle digital Torque viscometer (Brookfield Engineering Labs, Inc., Middleboro, MA). Shear stress (SS) was calculated by the following formula: $4 * UBF * \text{viscosity} / \pi * r^3$. FMD% was divided by SS to determine FMD/SS.

Statistical methods

Baseline characteristics were compared between subjects with and without an interval pregnancy using two sample Student's t-tests. Repeated measures analyses of variance were used to compare physiologic measures across groups and over time. SAS statistical software (SAS Institute, Cary, NC) was used for all data analysis. Data are presented as means $\pm$ standard errors of the means.

Results

All interval pregnancies were singletons, and the majority of the subjects in both groups were Caucasian, 82% (37/45). Demographic characteristics are presented in Table 1. There were no significant differences between groups in age, BMI, cycle day studied or interval of time between visits. Women in the pregnancy group were examined an average of 14.4 ± 1.2 months postpartum. Family histories were similar between the two groups; 64.7% of Preg (11/17) and 50% of NP subjects (14/28, $p = .41$) reported a family history of hypertension in a first degree relative. None of the women reported taking anti-hypertensive medications or other medications known to affect blood pressure at the time of their evaluations.

There was a significant decrease in MAP from the prepregnancy visit to postpartum in women with an interval pregnancy ($p = .013$, Figure 1). There was no significant change in MAP over time in the nonpregnant group (group*time interaction, $p = .002$, Table 2). PWV decreased in both groups over time (time effect: $p < .0001$, Table 2). PWV was, however, significantly decreased in those women with an interval pregnancy as compared to those without an interval pregnancy (group*time interaction, $p = .005$, Figure 2). Pulse pressure decreased from prepregnancy to postpartum in women with an interval pregnancy and this trend was not seen in the nonpregnant group, but the group*time interaction did not reach statistical significance ($p = .17$, Table 2).

UBF significantly increased over time in both groups ($p < .0001$), and there was no significant difference in UBF between groups. Both PV/BMI and FMD/SS significantly decreased in both groups over time ($p = .04$ and $p = .02$, respectively) and there was no evidence of a group or group*time interaction (Table 2).

Comment

In this study of women examined prior to first pregnancy in addition to postpartum, we evaluated the effect of pregnancy on cardiovascular physiology and compared women with an interval pregnancy to a time-matched group of women who did not have an interval pregnancy. We found that women with an interval pregnancy had significantly decreased blood pressure and increased arterial compliance compared to their prepregnancy evaluation, when compared to those without an interval pregnancy. Our observations suggest a significant beneficial effect of pregnancy on cardiovascular remodeling that may contribute to the attenuation of preeclampsia risk in a subsequent pregnancy.

Few studies have evaluated postpartum cardiovascular measures relative to prepregnancy values. Clapp and Capeless found at one year postpartum, as compared to prepregnancy values, women had significantly higher cardiac output, due to both increased stroke volume and increased end-diastolic volume (9). Women in their study also demonstrated lower systemic vascular resistance at one year postpartum when combining primiparous with multiparous women, but showed no significant change in mean arterial pressure. We did not observe increased cardiac output relative to prepregnancy values. However, our group of women were nulliparous at study entry, and subsequent pregnancies may amplify the increase in cardiac output, as Clapp and Capeless found a significantly greater increase in cardiac output in multiparous as compared to primiparous women.

Preeclampsia risk is generally highest for first pregnancies, and decreases with subsequent pregnancies in the absence of a prior preeclamptic pregnancy (5–8). Furthermore, a longer time interval between subsequent pregnancies is associated with an increasing risk of preeclampsia, suggesting a protective effect of pregnancy that diminishes over time (11–15). Our data showing a decrease in MAP after pregnancy strongly suggests positive cardiovascular adaptation that may favor future pregnancy outcome and suggests a mechanism by which preeclampsia risk is reduced in subsequent pregnancies. This is further supported by earlier studies from our laboratory showing a strong negative correlation between length of interpregnancy time interval and reduction in mean arterial pressure between first and second pregnancies (10). These findings were reproduced in a larger cohort where MAP was approximately 2 mmHg lower in the second pregnancy with a very short interpregnancy interval and the decrease in MAP was nonexistent after two years (11).

Pulse wave velocity (PWV) is considered the gold-standard for measuring arterial stiffness and is correlated with future risk for cardiovascular disease and mortality (17,18). Several studies show increased PWV after clinical onset of PE (19–21). More recently, women destined to develop preeclampsia have been shown to have increased PWV as compared to normal pregnancies as early as 11 to 13 weeks gestation (22, 23). Our laboratory has previously reported decreased arterial compliance prior to pregnancy, manifested by increased pulse pressure and pulse wave velocity, in patients who later develop preeclampsia (24). To our knowledge, the current study is the first to describe attenuation of PWV by pregnancy as compared to a nonpregnant control cohort, and suggests that risk for preeclampsia as assessed by arterial compliance may be modified by pregnancy.

We did not observe a significant effect of pregnancy on uterine blood flow (UBF), although UBF increased significantly over time in both groups. UBF is known to fluctuate during the menstrual cycle, with the uterine arteries demonstrating a progressive increase in blood flow beginning after menstrual day 10 and continuing through the luteal phase (25). Although many studies have used Doppler ultrasound to evaluate uterine artery velocity waveforms in early and mid-pregnancy with the goal of predicting adverse pregnancy outcome, few have evaluated pre-pregnancy UBF. Hale et al (26) evaluated UBF and uterine artery resistance index (RI) pre-pregnancy and correlated this with early pregnancy values. They observed that pre-pregnancy UBF was positively correlated with UBF in early pregnancy, suggesting that uterine physiology and hemodynamics prior to pregnancy may be an important determinant of uterine physiology during pregnancy. To our knowledge, this is the first longitudinal report to evaluate UBF pre-pregnancy and in the postpartum period, in comparison to a time-matched nulliparous control group. We suspect the increase in UBF seen between visit 1 and visit 2 in both groups was at least in part due to timing during the menstrual cycle, as subjects on average were assessed later during their menstrual cycle at visit 2 (visit 1 cycle day: 8.9 ± 0.61 , visit 2 cycle day: 12.2 ± 0.96).

Impaired endothelial function as assessed by ultrasound measurement of flow mediated vasodilation (FMD) of the brachial artery in response to shear stress (SS) has been shown to be an independent predictor of future cardiovascular events (27, 28). Existing reports of FMD in the third trimester of normotensive pregnancy show mixed results, however impaired FMD in preeclamptic pregnancies is well-described (29, 30). We did not see evidence of pregnancy-induced change in FMD, though FMD/SS decreased over time in both groups. Consistent with this observation, in prior studies from our laboratory, FMD/SS was not found to be associated with subsequent preeclampsia when assessed prior to first pregnancy (31). Similarly, we did not see an effect of pregnancy on plasma volume when corrected for body mass index (PV/BMI), although PV/BMI decreased in both groups over time.

In this study we evaluated the effect of pregnancy on cardiovascular physiology. Our observations of decreased MAP and increased arterial compliance following pregnancy suggest a significant favorable effect of pregnancy on cardiovascular remodeling. These findings may represent a mechanism by which preeclampsia risk is reduced in subsequent pregnancies. Additional longitudinal prospective studies would be useful to explore this phenomenon in women who were destined to develop preeclampsia to evaluate their specific postpartum cardiovascular phenotypic adaptation and their time course.

Acknowledgments

Grant support: NIH R01 HL 071944 (IMB), M01 RR000109 (UVM GCRC), NIH P01 HD38843 (RRM) and NIH R01 HL 117341 (RRM)

References

1. Craici I, Wagner S, Garovic VD. Preeclampsia and future cardiovascular risk: formal risk factor or failed stress test? Therapeutic advances in cardiovascular disease. 2008 Aug; 2(4):249–59. [PubMed: 19124425]

2. Bernstein IM, Meyer MC, Osol G, Ward K. Intolerance to volume expansion: a theorized mechanism for the development of preeclampsia. *Obstetrics and gynecology*. 1998 Aug; 92(2):306–8. [PubMed: 9699772]
3. Myatt L, Webster RP. Vascular biology of preeclampsia. *Journal of thrombosis and haemostasis : JTH*. 2009 Mar; 7(3):375–84. [PubMed: 19087223]
4. Bilhartz TD, Bilhartz PA, Bilhartz TN, Bilhartz RD. Making use of a natural stress test: pregnancy and cardiovascular risk. *Journal of women's health (2002)*. 2011 May; 20(5):695–701.
5. McDonald SD, Best C, Lam K. The recurrence risk of severe de novo pre-eclampsia in singleton pregnancies: a population-based cohort. *BJOG : an international journal of obstetrics and gynaecology*. 2009 Nov; 116(12):1578–84. [PubMed: 19681845]
6. Van Rijn BB, Hoeks LB, Bots ML, Franx A, Bruinse HW. Outcomes of subsequent pregnancy after first pregnancy with early-onset preeclampsia. *American journal of obstetrics and gynecology*. 2006 Sep; 195(3):723–8. [PubMed: 16949403]
7. Basso O, Christensen K, Olsen J. Higher risk of pre-eclampsia after change of partner. An effect of longer interpregnancy intervals? *Epidemiology*. 2001; 12(6):624–629. [PubMed: 11679788]
8. Barton JR, Sibai BM. Prediction and prevention of recurrent preeclampsia. *Obstetrics and gynecology*. 2008 Aug; 112(2 Pt 1):359–72. [PubMed: 18669736]
9. Clapp JF 3rd, Capeless E. Cardiovascular function before, during, and after the first and subsequent pregnancies. *The American journal of cardiology*. 1997 Dec 1; 80(11):1469–73. [PubMed: 9399724]
10. Bernstein IM, Thibault A, Mongeon JA, Badger GJ. The influence of pregnancy on arterial compliance. *Obstetrics and gynecology*. 2005 Mar; 105(3):621–5. [PubMed: 15738034]
11. Mikolajczyk RT, Zhang J, Ford J, Grewal J. Effects of interpregnancy interval on blood pressure in consecutive pregnancies. *American journal of epidemiology*. 2008 Aug 15; 168(4):422–6. [PubMed: 18495629]
12. Basso O, Christensen K, Olsen J. Higher risk of pre-eclampsia after change of partner. An effect of longer interpregnancy intervals? *Epidemiology (Cambridge, Mass)*. 2001 Nov; 12(6):624–9.
13. Conde-Agudelo A, Belizan JM. Maternal morbidity and mortality associated with interpregnancy interval: cross sectional study. *BMJ (Clinical research ed)*. 2000 Nov 18; 321(7271):1255–9.
14. Skjaerven R, Wilcox AJ, Lie RT. The interval between pregnancies and the risk of preeclampsia. *The New England journal of medicine*. 2002 Jan 3; 346(1):33–8. [PubMed: 11778000]
15. Shachar BZ, Lyell DJ. Interpregnancy interval and obstetrical complications. *Obstetrical & gynecological survey*. 2012 Sep; 67(9):584–96. [PubMed: 22990461]
16. Bernstein IM, Damron D, Schonberg AL, Shapiro R. The relationship of plasma volume, sympathetic tone, and proinflammatory cytokines in young healthy nonpregnant women. *Reprod Sci*. 2009 Oct; 16(10):980–5. [PubMed: 19531800]
17. Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. *Journal of the American College of Cardiology*. 2010 Mar 30; 55(13):1318–27. [PubMed: 20338492]
18. Sonoda H, Takase H, Dohi Y, Kimura G. Factors associated with brachial-ankle pulse wave velocity in the general population. *J Hum Hypertens*. 2011
19. Kaihura C, Savvidou MD, Anderson JM, McEniery CM, Nicolaides KH. Maternal arterial stiffness in pregnancies affected by preeclampsia. *Am J Physiol Heart Circ Physiol*. 2009 Aug; 297(2):H759–H64. [PubMed: 19465544]
20. Robb AO, Mills NL, Din JN, Smith IB, Paterson F, Newby DE, et al. Influence of the menstrual cycle, pregnancy, and preeclampsia on arterial stiffness. *Hypertension*. 2009 Jun; 53(6):952–8. [PubMed: 19398652]
21. Savvidou MD, Kaihura C, Anderson JM, Nicolaides KH. Maternal arterial stiffness in women who subsequently develop pre-eclampsia. *PloS one*. 2011; 6(5):e18703. [PubMed: 21559278]
22. Khalil AA, Cooper DJ, Harrington KF. Pulse wave analysis: a preliminary study of a novel technique for the prediction of pre-eclampsia. *BJOG*. 2009 Jan; 116(2):268–76. discussion 76–7. [PubMed: 19076958]

23. Khalil A, Akolekar R, Syngelaki A, Elkhoul M, Nicolaides KH. Maternal hemodynamics at 11–13 weeks' gestation and risk of pre-eclampsia. *Ultrasound in obstetrics & gynecology*. 2012 Jul; 40(1):28–34. [PubMed: 22565361]
24. Hale S, Choate M, Schonberg A, Shapiro R, Badger G, Bernstein IM. Pulse pressure and arterial compliance prior to pregnancy and the development of complicated hypertension during pregnancy. *Reprod Sci*. 2010 Sep; 17(9):871–7. [PubMed: 20639475]
25. Bernstein IM, Ziegler WF, Leavitt T, Badger GJ. Uterine artery hemodynamic adaptations through the menstrual cycle into early pregnancy. *Obstetrics and gynecology*. 2002 Apr; 99(4):620–4. [PubMed: 12039123]
26. Hale SA, Schonberg A, Badger GJ, Bernstein IM. Relationship between prepregnancy and early pregnancy uterine blood flow and resistance index. *Reprod Sci*. 2009 Nov; 16(11):1091–6. [PubMed: 19657141]
27. Lerman A, Zeiher AM. Endothelial function: cardiac events. *Circulation*. 2005 Jan 25; 111(3):363–8. [PubMed: 15668353]
28. Maruhashi T, Soga J, Fujimura N, Idei N, Mikami S, Iwamoto Y, et al. Relationship between flow-mediated vasodilation and cardiovascular risk factors in a large community-based study. *Heart (British Cardiac Society)*. 2013 Dec; 99(24):1837–42. [PubMed: 24153417]
29. Brandao AH, Cabral MA, Leite HV, Cabral AC. Endothelial function, uterine perfusion and central flow in pregnancies complicated by Preeclampsia. *Arquivos brasileiros de cardiologia*. 2012 Oct; 99(4):931–5. [PubMed: 22983153]
30. Mori T, Watanabe K, Iwasaki A, Kimura C, Matsushita H, Shinohara K, et al. Differences in vascular reactivity between pregnant women with chronic hypertension and preeclampsia. *Hypertension research*. 2013 Sep 26.
31. Hale SA, Badger GJ, McBride C, Magness R, Bernstein IM. Prepregnancy Vascular Dysfunction in Women who Subsequently Develop Hypertension During Pregnancy. *Pregnancy hypertension*. 2013 Apr 1; 3(2):140–5. [PubMed: 23772359]

Figure 1. Changes in Mean Arterial Pressure

Bar graph showing the changes in mean arterial pressure (MAP) from visit 1 to visit 2 in women with and without an interval pregnancy. Data presented are means $\pm$ standard errors of the means. *Significant change in MAP from visit 1 to visit 2 at $P < .05$.

Figure 2. Changes in Pulse Wave Velocity

Bar graph showing the changes in pulse wave velocity (PWV) from visit 1 to visit 2 in women with and without an interval pregnancy. Data presented are means $\pm$ standard errors of the means. *Significant change in PWV from visit 1 to visit 2 at $P < .05$.

Table 1Demographic Characteristics of Study Volunteers^a

Characteristic	NP (n = 28) Mean ± SE	Preg (n = 17) Mean ± SE	P value
Age (years)	29.3 ± 1.1	29.9 ± 0.6	.63
BMI (kg/m ²)	22.4 ± 0.6	23.2 ± 0.8	.46
Visit 1/Prepregnancy cycle day	9.4 ± 0.7	8.1 ± 1.1	.36
Visit 2/Postpartum cycle day	12.5 ± 1.1	11.8 ± 1.7 ^b	.73
Interval between visits (months)	30.4 ± 1.3	29.3 ± 1.6	.59
Interval since delivery (months)		14.4 ± 1.2	

^aSignificance determined by Student's t-test.

^bOne woman had not resumed cycling at her postpartum visit, thus mean cycle day is based on n = 16.

Table 2
Comparison of physiologic variables between women with and without an interval pregnancy^a

Variable	Pregnant (Preg) N=17				Non-pregnant Controls (NP) N=28				<i>p-value</i> #	
	Prepregnancy	Postpartum	Visit 1	Visit 2	Group Effect	Time Effect	Group*Time Interaction			
<i>Physiologic Measures</i>										
MAP (mmHg)	85.3 ± 1.8	80.5 ± 1.8	80.3 ± 1.4	82.8 ± 1.4	.53	.29			.002	
PWV (m/s)	2.73 ± 0.05	2.49 ± 0.05	2.56 ± 0.04	2.50 ± 0.04	.17	<.001			.005	
PP (mmHg)	44.1 ± 2.2	39.6 ± 2.2	42.4 ± 1.7	42.9 ± 1.7	.71	.28			.17	
CO (L/min)	4.7 ± 0.2	4.7 ± 0.2	4.6 ± 0.2	4.4 ± 0.2	.35	.32			.46	
UBF (mL/min)	20.7 ± 4.0	38.5 ± 4.1	22.7 ± 3.1	34.7 ± 3.2	.84	<.001			.27	
PV/BMI	124.9 ± 5.1	117.3 ± 5.1	132.5 ± 3.9	120.6 ± 3.9	.27	.02			.61	
FMD/SS	0.33 ± .05	0.17 ± 0.06	0.22 ± .05	0.17 ± 0.05	.31	.04			.24	

^aData are presented as means ± SE.

Abbreviations: CO, cardiac output; FMD/SS, flow-mediated vasodilation/shear stress; MAP, mean arterial pressure; PP, pulse pressure; PV/BMI, plasma volume/body mass index; PWV, pulse wave velocity; UBF, uterine blood flow