Published in final edited form as:

Curr Vasc Pharmacol. 2014 ; 12(6): 779.

Commentary on the Special Issue on the Impact of Myogenic tone in Health and Disease

Richard J. Roman and Richard P. E. Van Dokkum

Department of Pharmacology, University of Mississippi Medical Center, Jackson, MS 39216 And
Dept. Clinical Pharmacology, University Medical Center Groningen, Antonius Deusinglaan 1,
9713 AV, Groningen

Abstract

Autoregulation is a vital homeostatic mechanism that helps maintain constant delivery of oxygen to organs despite fluctuations in arteriolar pressure. Autoregulation of blood flow to elevations in pressure is largely mediated by the myogenic response of small arteries and arterioles which constrict in response to elevations in distending pressure. There is now general agreement that the myogenic response is an intrinsic property of vascular smooth muscle cells in the vessel wall that involves depolarization and calcium influx through L-type voltage-gated calcium channels (VGCC), calcium/calmodulin-dependent phosphorylation of myosin light chain kinase and actin myosin-based contraction. Despite intensive investigation, however, the mechanotransduction events that initiate the myogenic response and the signaling pathways involved remain uncertain. This special issue on the Impact of Myogenic Tone in Health and Disease includes 9 papers that address current thought regarding the molecular mechanisms underlying myogenic control of vascular tone in the renal, cerebral and coronary circulations and the evidence that impairments in the myogenic response contribute to the development of vascular and end organ damage associated with hypertension, diabetes and aging.

Keywords

Kidney; afferent arteriole; glomerulus; myogenic response; tubuloglomerular feedback

Autoregulation of blood flow is a critical homeostatic mechanism that helps maintain delivery of oxygen to vital organs despite fluctuations in arteriolar pressure. In the renal circulation, it maintains the filtration of waste products during hypotensive episodes and prevents transmission of elevated pressures to the glomerular capillaries that promote the development of proteinuria and chronic kidney disease. Similarly, autoregulation of cerebral blood flow protects the brain from increases in capillary hydrostatic pressure, vascular damage and cerebral edema following acute elevations in arterial pressure and from ischemic injury in response to embolization or systemic hypotension.

Send Correspondence to: Richard J. Roman, Department of Pharmacology and Toxicology, University of Mississippi Medical Center, 2500 North State Street, Jackson, MS 39216.

The authors contributed equally to the content of this review.

CONFLICT OF INTEREST

The authors confirm that this article content has no conflicts of interest.

Autoregulation of cerebral blood flow in response to elevations in arterial pressure is largely mediated by myogenic constriction of small cerebral arteries and arterioles, whereas the response to reductions in pressure also involves the release of vasodilator metabolic mediators from the surrounding brain tissue. In the renal circulation, the myogenic response of the afferent arteriole accounts for most of the rapid compensation to elevations in arterial pressure whereas, tubuloglomerular feedback, a modified form of metabolic blood flow regulation, contributes to the response to reductions in pressure.

The myogenic response is an intrinsic property of the vascular smooth muscle cells in small arteries that constrict in response to elevations in transmural pressure. It can be readily demonstrated in arterioles *in vitro* devoid of endothelial or parenchymal influences. The phenomenon was first described by Sir William Bayliss over 100 years ago [1]. His original work was confirmed and extended through a series of papers by Bjorn Folkow [2, 3]. At physiological distending pressure, small arteries and arterioles constrict and develop myogenic tone [1]. This response sets the baseline levels of intracellular calcium and the membrane potential in the vessel wall and the basal level of vascular tone and vascular reactivity to neural and hormonal factors [4]. More recent work, has indicated that the myogenic response is triggered by complex interactions of extracellular matrix and integrins with cytoskeletal elements and activation of mechanosensitive ion channels either directly or secondary to the release of intracellular mediators [5]. Despite intensive investigation, however, the exact sequence of the mechanotransduction events that initiate the myogenic response and the signaling pathways involved remains uncertain. Five of the papers in this special issue on the Impact of Myogenic Tone in Health and Disease review up to the date information regarding the molecular mechanisms underlying myogenic control of vascular tone in the renal, cerebral and coronary circulations.

Other papers in this special issue summarizes the evidence that the myogenic response is impaired and contributes to the development of vascular and end organ damage associated with hypertension, diabetes and aging. In this regard, the myogenic responsiveness of the cerebral circulation is impaired following ischemic and hemorrhagic stroke and traumatic brain injury. It is shifted to higher pressures by chronic hypertension. Loss of myogenic responsiveness allow for greater transmission of the elevated systemic pressure to the small arterioles in the brain and promote vascular remodeling that increases the susceptibility to ischemic insults. Indeed, chronic hypertension is a major risk factor for small vessel disease in the cerebral circulation, which contributes to white matter damage, vascular dementia and declines in cognitive function.

In the kidney, the myogenic response and autoregulation of renal blood flow is typically intact in patients with essential hypertension and most do not develop renal injury. Similar renoprotection is seen in the spontaneous hypertensive rat model of hypertension and in angiotensin II-infused mouse models of hypertension. However, 4 of the papers in this issue reveal that autoregulation of renal blood flow is impaired in diabetes and in several genetic and experimental models of hypertension and that this contributes to the development of proteinuria and severe renal disease. Several of the mechanisms that are explored in these papers include alterations in vascular K channel function, increases in vascular reactive oxygen species, deficiency in the formation of 20-HETE and changes ATP-P2 receptor

function. One of the papers also presents evidence that chronic treatment of type II diabetic animals with a dipeptidyl peptidase inhibitor preserves the myogenic response in renal arterioles and opposes the development of renal injury.

A common theme that is presented by most of the papers in this Special Issue regarding the signaling mechanisms involved in the myogenic control of vascular tone is that it involves activation of phospholipase C, increased intracellular levels of IP₃, DAG and 20-HETE, calcium release from intracellular stores, opening of transient receptor potential channels, blockade of large conductance calcium activated K channels, membrane depolarization, calcium entry through L-type calcium channels, phosphorylation of myosin light chain kinase and sensitization of the contractile mechanism via activation of the rho kinase pathway. This detailed review of the literature is important in that it highlights several potential intervention points that might be used to restore the efficiency of the myogenic response in hypertension and diabetes. These include transient receptor channel agonists, K channel blockers, 20-HETE agonists, L-type calcium channel agonists and rho kinase agonists. The potential role of these compounds as novel therapeutic targets to oppose the development of stroke and renal disease in hypertension and diabetes are explored in several of the papers in this special issue.

Acknowledgments

This work was supported in part by grants HL-36279 and 29547.

References

1. Bayliss WM. On the local reactions of the arterial wall to changes of internal pressure. *J Physiol.* 1902; 28:220–231. [PubMed: 16992618]
2. Folkow B. Intravascular pressure as a factor regulating the tone of the small vessels. *Acta Physiol Scand.* 1949; 17:289–310. [PubMed: 18151727]
3. Folkow B. Description of the myogenic hypothesis. *Circ Res.* 1964; 15(suppl):279–87. [PubMed: 14206315]
4. Johnson, PC. The myogenic response. In: Bohr, DF.; Somlyo, AP.; Sparks, HV., Jr, editors. *The Handbook of Physiology; The Cardiovascular System; Vascular Smooth Muscle.* American Physiological Society; Bethesda, MD: 1980. p. 409-442.
5. Davis MJ, Hill MA. Signaling mechanisms underlying the vascular myogenic response. *Physiol Rev.* 1999; 79:387–423. [PubMed: 10221985]