Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/jor

Case Report

Shortening femoral osteotomy with stemmed resurfacing total knee arthroplasty for severe flexion contracture in Juvenile Rheumatoid Arthritis

Brock Kitchen^b, Hugo B. Sanchez^a, Russell A. Wagner^{a,*}^a University of North Texas Health Science Center, Department of Orthopaedic Surgery, John Peter Smith Hospital, Fort Worth, TX 76104, United States^b University of North Texas Health Science Center, Texas College of Osteopathic Medicine, Fort Worth, TX 76107, United States

ARTICLE INFO

Article history:

Received 5 December 2013

Accepted 30 May 2014

Available online 14 July 2014

Keywords:

Flexion contracture

Femoral osteotomy

Juvenile rheumatoid arthritis

Total knee arthroplasty

ABSTRACT

Juvenile Rheumatoid Arthritis (JRA) is a progressive disease characterized by pain, swelling, and loss of motion in the joints of adolescents. Total knee arthroplasty (TKA) can be indicated, during the adolescent years, in patients with advanced JRA to alleviate pain and improve function. Because of the relative infrequency of TKA in patients with JRA, evaluation of the type of TKA performed and the results merit review. This case report presents two distinct operations performed to obtain full extension. 1. Distal femoral resection with conversion to hinged arthroplasty. 2. Femoral shortening osteotomy with resurfacing TKA.

Copyright © 2014, Professor P K Surendran Memorial Education Foundation. Publishing Services by Reed Elsevier India Pvt. Ltd. All rights reserved.

1. Introduction

Juvenile Rheumatoid Arthritis (JRA) is a progressive disease characterized by pain, swelling, and loss of motion in the joints of adolescents. Total knee arthroplasty (TKA) can be indicated, during the adolescent years, in patients with advanced JRA to alleviate pain and improve function. Because of the relative infrequency of TKA in patients with JRA, evaluation of the type of TKA performed and the results merit

review. This case report presents a nineteen year old male with advanced JRA and severe knee flexion contractures who received bilateral knee replacement in an effort to achieve ambulation. Two distinct operations were performed to obtain full extension: 1. Distal femoral resection with conversion to hinged arthroplasty. 2. Femoral shortening osteotomy with total knee arthroplasty with revision components. Both methods resulted in intraoperative full extension of the knee, yet the second method preserved more bone stock and

* Corresponding author. Department of Orthopaedic Surgery, John Peter Smith Hospital, 1500 South Main Street, Fort Worth, TX 76104, United States. Tel.: +1 817 927 1370; fax: +1 817 927 3955.

E-mail address: rwagner@jpshealth.org (R.A. Wagner).
<http://dx.doi.org/10.1016/j.jor.2014.05.002>

0972-978X/Copyright © 2014, Professor P K Surendran Memorial Education Foundation. Publishing Services by Reed Elsevier India Pvt. Ltd. All rights reserved.

avoided the use of a hinge. The patient gave his consent for this information to be published.

2. Case presentation

A 19-year-old Caucasian male presented with severe bilateral knee flexion contractures due to long-standing Juvenile Rheumatoid Arthritis. He was diagnosed with JRA at age 6 and had been non-ambulatory since age 7. JRA had affected nearly all of the joints in his body, excluding his hips which remained relatively functional. His knees were not excessively painful, yet cumbersome and hindered his ability to function. Initially, he presented requesting bilateral amputations with the goal of using prosthetic legs to stand and ambulate.

Physical exam revealed bilateral knee flexion contractures of 120°, with further flexion upto 140°. Knees were not inflamed or tender. His hips had good painless range of motion and distal motor, sensation, and pulses were intact. Lateral radiographs of both knees revealed squaring of the distal femur which appeared that it would prevent extension (Fig. 1). Because of the severe flexion contractures, AP radiographs were distorted.

Initial considerations included gradual skeletal traction or external fixation as a measure to gain extension, but because gradual distraction via skeletal traction or serial casts is often limited by a boney block¹ and his radiographs revealed a significant boney block (Fig. 1) in the squaring of the distal femur, traction was thought to be more appropriately considered as a possible postoperative treatment. Conversely, knee replacement would provide reshaping of the distal femur, access to the joint, the opportunity to adjust the ligament balance, and treatment for the absence of cartilage. After several office visits discussing options and the significant risk of complications,^{2,3} the patient decided to proceed with bilateral knee replacements. The initial surgical plan was to perform a posterior stabilized type TKA with posterior soft tissue releases in a staged type manner as described by Clayton et al⁴ Bilateral arthroplasty was scheduled to avoid the difficulty of rehabilitating the first knee in the presence of a severe flexion contracture in the second knee. Because of the severe flexion

contractures, significant bone resection was anticipated. If extreme bone resection was required, a hinged knee arthroplasty would be used because of its effectiveness in correcting severe knee contractures.⁵

A midline incision was made over the right patella and a medial parapatellar approach was performed. Extensive erosive and fibrous changes were noted, including absolutely no cartilage throughout the joint and fibrous tissue between all of the compartments.^{2,3} The fibrous tissue was cleared and the patella subluxed. 14 mm was resected from the distal femur and then another 14 mm from the tibia so that the flexion gap would accommodate the trial components and provide additional extension. The box cut was performed and the posterior cruciate ligament, posterior capsule, and gastrocnemius were completely released. Next, trial femoral and tibial components were placed, resulting in good medial and lateral stability in flexion; however, extension was limited to about 90°. Posteromedial and posterolateral incisions were placed, and the hamstring tendons and popliteus were cut for Z-lengthening. At this point, the posterior aspect of the knee was completely free of soft tissue other than the skin and neurovascular structures, which limited extension at 45°. Gradual distraction with external fixation or a turnbuckle splint was considered. The decision was made to proceed with the rotating hinge arthroplasty. To accommodate the distal femoral replacement prosthesis and obtain extension, an additional 13½ cm of the distal femur was resected. The smallest stem available for the hinged prosthesis was 10 mm and this corresponded to the diameter of the femoral canal; however, the trial stem did not have a stable press-fit. Although cementing the stem required some reaming, thereby polishing the intramedullary canal and diminishing the strength of cemented fixation, the prosthesis was cemented in place. The construct resulted in full extension of the knee with minimal tension of the neurovascular bundle. Pulses were intact. With the preoperative patella baja and now loose quadriceps mechanism, a V-Y quadricepsplasty was performed.⁶

Although full extension of the right knee was achieved, the senior author was not completely satisfied with how the operation had evolved and decided on a different course of action for the left knee. The proximal tibia and distal femur were prepared for resurfacing type knee prosthesis in the same manner as on the right knee, with PCL and posterior capsular release. However, prior to releasing the hamstrings and rather than resecting 15 cm from the distal femur and using a hinged type knee prosthesis, a 15 cm intercalary segment was removed from the supracondylar area of the distal femur. Two step cuts were performed to remove the femoral segments and equalize the leg lengths. The osteotomy was then stabilized by the stemmed femoral component. At this point, only minor mediolateral balancing was necessary to obtain full extension. The femoral stem was cemented and bone from the resections was placed as a bone graft around the junction of the step cuts. A V-Y quadricepsplasty was performed as had been done to the right side. In order to close the skin, a Z-plasty was performed, bringing the lateral side flap proximally and the medial side distally and excising excess skin. Surgical time and blood loss were not individually recorded, but were roughly equivalent (Figs. 2 and 3).

Fig. 1 – Preoperative lateral radiographs of the right and left knees showing “squaring” of the distal femoral condyles, preventing extension.

3. Results/complications

Immediately postoperatively, he was unable to dorsiflex his ankles and toes and had altered sensation. These changes resolved in the early postoperative period. He was placed into knee immobilizers to limit flexion for 8 weeks in order to protect the quadriceps advancement. The patient gradually gained the ability to walk with the aid of a walker, progressed to a cane, and had improved range of motion of both knees. At 2 years, his range of motion was 20–75° on the right and 18–85 on the left. Leg lengths were roughly the same due to the same length of bone resected, but because of the residual flexion contracture this was hard to determine. A scanogram was not obtained to accurately determine leg lengths. The right knee with the hinge had intermittent significant anterior thigh pain located around the site of the femoral stem, while the left knee with the shortening had pain-free function.

4. Discussion

As many as 10% of patients affected by Juvenile Rheumatoid Arthritis suffer from progressive and disabling arthritis,⁷ often with both functional and pain limitations, despite optimal medical management. A relatively recent study conducted by Parvizi et al² concluded that total knee arthroplasty, in this subset of patients, provided excellent improvement in function & substantial pain relief. This study described the outcome in both functional and pain scores in 25 total knee arthroplasty in thirteen patients with JRA. The mean Knee Society function score improved from 14.8 to 39.2 at the time of latest follow-up, and the mean Knee Society pain score markedly improved from 27.6 to 88.3 points at time of latest follow-up. This study supports other studies,^{8,9} all of which, found similar patterns of improvements in function, pain, and sometimes range of motion. In conclusion they all

Fig. 2 – Postoperative lateral radiographs of the right and left knees showing the resection of the distal femur and rotating hinge type of total knee implants on the right knee and supracondylar shortening osteotomy with stemmed revision components on the left knee.

Fig. 3 – Postoperative AP radiographs of the right and left knees showing the resection of the distal femur and rotating hinge type of total knee implants on the right knee and supracondylar shortening osteotomy with stemmed revision components on the left knee.

substantiated the role of TKA in this particular subset of patients who despite medical management continue to suffer from progressive and disabling JRA.

Although function and pain is markedly improved with arthroplasty, full correction of flexion contractures has proven to be much more difficult. Massin et al¹⁰ recently reported the outcome of 107 TKA's performed to correct both flexion contracture of greater than 20° and knee stiffness. As a result of the arthroplasty, extension increased by an average of 20° with a residual flexion contracture of 7° in the flexion contracture only group. Firestone et al¹¹ has also showed that the degree of preoperative flexion contracture influenced the final extension recovery. In our review of the literature, we were unable to locate a surgical correction of a 120° flexion contracture. Scott et al¹² in a recent publication explained why full correction of such a severe flexion contractures have been virtually unattainable. He explains that correction of flexion contractures >45° are limited because it required as much as 13 mm of distal femoral resection and beyond this soon approaches the origins of the collateral ligaments. Because of this anatomic limitation the amount of literature regarding correction of flexion contracture >45° is relatively absent. We were also unable to locate a case in which a shortening femoral osteotomy was used to obtain extension of a flexion contracture.

This case outlines atleast some of the challenges and limitations in correction of such contractures and provides a novel and straight forward surgical solution in overcoming these challenges. Furthermore, performing two very different operations in the same patient to correct severe flexion contracture warrants comparison. The initial presentation of severe flexure contracture combined with obvious squaring of the distal femur led the senior author to initially consider a resurfacing type TKA. The completion of the first knee taught that the main limiting factor in correcting the long-standing knee contracture was the neurovascular bundle rather than joint soft tissue. As this problem is similar to the one faced when performing hip arthroplasty with severe dysplasia, it

seemed reasonable to apply the same shortening osteotomy technique in the knee. This would provide advantages over distal femur resection and use of a hinged TKA, retaining the natural integrity of the knee, and avoiding the mechanical complications of a hinge. The result of using the shortening femoral osteotomy proved to be superior to the distal femoral resection and hinged knee in this patient. The results in our patient showed substantial improvements in function, extension and range of motion, but a persistent lack of full extension. Furthermore, the durability of the knees in a 19 year old is in question. In the future, we will plan to use the shortening osteotomy in this situation as it preserves bone stock.

Conflicts of interest

All authors have none to declare.

REFERENCES

1. Sarokhan AJ, Scott RD, Thomas WH, Sledge CB, Ewald FC, Cloos DW. Total knee arthroplasty in juvenile rheumatoid arthritis. *J Bone Joint Surg Am.* 1983 Oct;65:1071–1080.
2. Parvizi J, Lajam CM, Trousdale RT, Shaughnessy WJ, Cabanela ME. Total knee arthroplasty in young patients with juvenile rheumatoid arthritis. *J Bone Joint Surg Am.* 2003 Jun;85:1090–1094.
3. Lu H, Mow CS, Lin J. Total knee arthroplasty in the presence of severe flexion contracture: a report of 37 cases. *J Arthroplasty.* 1999;14:775–780.
4. Clayton ML, Thompson TR, Mack RP. Correction of alignment deformities during total knee arthroplasties: staged soft-tissue releases. *Clin Orthop Relat Res.* 1986 Jan;202:117–124.
5. Tew M, Forster IW. Effects of knee replacement on flexion deformity. *J Bone Joint Surg Br.* 1987;69:395.
6. Aglietti P, Windsor RE, Buzzi R, Insall JN. Arthroplasty for the stiff or ankylosed knee. *J Arthroplasty.* 1989;4:1–5.
7. Cassidy JT. Juvenile rheumatoid arthritis. In: Kelly WN, Harris ED, Ruddy S, Sledge CB, eds. *Textbook of Rheumatology*. 3rd ed. Philadelphia: W.B. Saunders; 1989:1289–1311.
8. Palmer DH, Mulhall KJ, Thompson CA, Severson EP, Santos ER, Saleh KJ. Total knee arthroplasty in juvenile rheumatoid arthritis. *J Bone Joint Surg Am.* 2005 Jul;87:1510–1514.
9. Bellemans J, Victor J, Westhovens R, Dequeker J, Fabry G. Total knee arthroplasty in the young rheumatoid patient. *Acta Orthop Belg.* 1997 Sep;63:189–193.
10. Massin P, Petit A, Odri G, et al. Total knee arthroplasty in patients with greater than 20 degrees flexion contracture. *Orthop Traumatol Surg Res.* 2009 Jun;95:S7–S12.
11. Firestone TP, Krackow KA, Davis 4th JD, Teeny SM, Hungerford DS. The management of fixed flexion contractures during total knee arthroplasty. *Clin Orthop Relat Res.* 1992 Nov;284:221–227.
12. Scott RD. Correcting flexion contractures: getting it straight. *Orthopedics.* 2008 Sep;31:925–926.