Published in final edited form as:

Am J Surg. 2015 May ; 209(5): 913–919. doi:10.1016/j.amjsurg.2014.12.027.

The Impact of Delaying Elective Resection of Diverticulitis on Laparoscopic Conversion Rate

The Colorectal Writing Group for the SCOAP-CERTAIN Collaborative[§], Vlad V. Simianu, MD^{1,*}, Mika N. Sinanan, MD, PhD¹, Amir L. Bastawrous, MD, MBA², Richard P. Billingham, MD², Alessandro Fichera, MD¹, Michael G. Florence, MD², Daniel O. Herzig, MD³, Eric K. Johnson, MD⁴, Scott R. Steele, MD⁴, Richard C. Thirlby, MD⁵, and David R. Flum, MD, MPH^{1,*}

¹Department of Surgery, University of Washington, Seattle, WA

²Department of Surgery, Swedish Medical Center, Seattle, WA

³Department of Surgery, Oregon Health & Science University, Portland, OR

⁴Department of Surgery, Madigan Army Medical Center, Tacoma, WA

⁵Department of Surgery, Virginia Mason Medical Center, Seattle, WA

§

Summary

Delaying elective colectomy for diverticulitis may increase the difficulty of laparoscopic colectomy due to chronic inflammation. An increasing number of prior episodes of diverticulitis was not associated with higher conversion rates.

Introduction—Guideline-concordant delay in elective laparoscopic colectomy for diverticulitis may result in repeated bouts of inflammation. We aimed to determine whether conversion rates from elective laparoscopic colectomy are higher after multiple episodes of diverticulitis.

Methods—Prospective cohort study evaluating laparoscopic colectomy conversion rates for diverticulitis from 42 hospitals.

Results—Between 2010 and 2013, 1,790 laparoscopic colectomies for diverticulitis (mean age 57.8 ± 13 ; 47% male) resulted in 295 (16.5%) conversions. Conversion occurred more frequently in non-elective operations ($p < 0.001$) and with fistula indications ($p = 0.012$). Conversion rates decreased with surgeon case-volume ($p = 0.028$). Elective colectomy exclusively for episode-based

© 2015 Published by Elsevier Inc.

*Corresponding Author for Manuscript: Vlad V Simianu, MD, Surgical Outcomes Research Center (SORCE), UW Medical Center, Box 354808, 1107 NE 45th St., Suite 502, Seattle, WA 98105, Phone: (317) 445-7792, Fax: (206) 616-9032, vsimianu@uw.edu.

*Corresponding Author for Reprints: David R. Flum, MD MPH, Surgical Outcomes Research Center (SORCE), UW Medical Center, Box 354808, 1107 NE 45th St., Suite 502, Seattle, WA 98105, Phone: (206) 543-1664, Fax: (206) 616-9032, daveflum@uw.edu.

§The Surgical Care and Outcomes Assessment Program and The Comparative Effectiveness Research Translation Network (SCOAP-CERTAIN) Collaborative

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

indications (n=784) had a conversion rate of 12.9%. Increasing episodes of diverticulitis were not associated with higher conversion rates, even among surgeons with similar experience levels.

Discussions—Conversion from laparoscopic colectomy for diverticulitis did not increase after multiple episodes of diverticulitis. Delaying elective resection appears to not prevent patients from the benefits of laparoscopy.

Keywords

diverticulitis; laparoscopy; elective; colectomy; conversion; complication

Introduction

Acute diverticulitis is one of the most common indications for hospitalization related to the gastrointestinal (GI) tract in the United States (US), where it is estimated it will result in an estimated 300,000 admissions, 1.5 million days/year of inpatient care, and upwards of \$1.8 billion in healthcare costs in 2014.^{1,2} While diverticulitis is one of the leading indications for emergency colectomy and colostomy,^{3,4} most colectomies for diverticulitis are performed electively to prevent recurrence or progression of disease.

Laparoscopic colectomy for the management of diverticular disease was more widely adopted after publication of initial studies in colorectal cancer in the early 2000s,^{5,6} and contemporary evidence supports lower morbidity, shorter hospitalization, and higher patient satisfaction with the laparoscopic approach to diverticulitis.⁴ Accordingly, most modern professional guidelines, including the 2014 update from the American Society of Colon and Rectal Surgeons (ASCRS),¹ recommend a laparoscopic approach to colectomy for diverticulitis.

However, given an increasingly recognized disconnect between episodes of diverticulitis and disease progression and recurrence,⁴ the timing of elective colectomy has become less clear. The classic surgical dogma of operating after 2 episodes, maintained as recently as the 2000 ASCRS guidelines,⁷ or delay operating until 3 or more episodes of diverticulitis as per the 2006 guidelines,⁸ have given way to recommendations to avoid episode-based surgery altogether and consider each patient's need for elective colectomy on a case-by-case basis.¹

Whether this delay in operating until after multiple episodes of diverticulitis, potentially increasing inflammation and scarring, has impacted the ability to complete operations laparoscopically remains to be determined. More so than laparoscopy for malignancy, laparoscopy for diverticulitis entails technical challenges of inflammation and adhesions, and failed laparoscopy rates are as high as 20%.^{9,10} Conversion to an open operation negates the benefits of a laparoscopic approach, and there is growing interest in factors associated with failed laparoscopy for diverticulitis. In Washington State, where nearly half of all colon operations are performed laparoscopically,¹¹ we sought to describe the factors associated with conversion and the impact of delayed elective colectomy on conversion from laparoscopy.

Methods

This study was exempted from human subjects review by the University of Washington Human Subject Review Committee. The Comparative Effectiveness Research Translation Network (CERTAIN) provided research and analytic support to the Surgical Care and Outcomes Assessment Program (SCOAP).¹²

Data Sources and Definitions

The primary cohort was defined by consecutive patients who underwent laparoscopic colon resection for diverticulitis between January 1, 2010 to December 31, 2013 in 42 Washington State hospitals that participate in SCOAP. Sociodemographic, clinical, and operative details were extracted from inpatient medical records by trained chart abstractors at each clinical site. SCOAP metrics and data dictionary are available via a secure page at www.SCOAP.org. A modified Charlson comorbidity index for each patient was calculated.¹³

The SCOAP data collection platform for diverticulitis has been previously described,¹¹ and includes indications for the operation such as number of prior episodes of diverticulitis, chronic complications including gastrointestinal bleeding (GIB), stricture, and fistula, and an 'other' category to capture additional indications. Surgical approach was derived from the operative report and operating room logs looking for specific identification of open, laparoscopic, laparoscopic/hand-assisted, and laparoscopic/robotic-assisted surgical approaches. As in our prior definitions, the latter three categories were considered laparoscopic procedures on an intention to treat basis.¹⁴ Conversion was defined through operative reports indicating that opening the abdomen was necessary to complete the procedure. Operations included were right/transverse hemicolectomy, left hemicolectomy, low anterior resection (including sigmoidectomy), and total abdominal colectomy. Because of a recognized association between laparoscopic procedural volume and conversion rates,¹⁵ we describe the relationship between conversion rates with surgical volume. Overall rates of procedures and conversions at the surgeon level were acquired using a de-identified code unique to each surgeon in the database (n=198 surgeons).

The main outcome of interest for this study was rate of conversion from laparoscopy. In addition, we describe rates of in-hospital complications and composite of adverse events (CAE). In-hospital complications include SCOAP's standard measures of cardiac, pulmonary, renal, infectious, or other complications requiring non-operative intervention. CAE included any of these with the addition of re-operative interventions and in-hospital deaths.¹⁴

The quality of indications data improved dramatically at SCOAP hospitals contemporaneous to a statewide benchmarking and educational initiative.¹¹ In order to minimize bias from chronic complication indications and missing data, we defined a subgroup *a priori* to include only those patients who underwent elective laparoscopic colectomy for an episode-based indication and had non-missing data.

Statistical Analysis

Patient characteristics, operative indications, and outcomes were summarized using frequency distributions for categorical variables, and mean (SD) for continuous variables. We stratified our description by conversion from laparoscopy. Categorical variables were compared using Pearson χ^2 statistic. Continuous variables were compared using the Student's t-test. Linear and logistic regression models were used to evaluate the association of case volume (clustered at surgeon level) and prior episode number, respectively, on conversion from laparoscopy, adjusting for patient, clinical, and operative characteristics identified as statistically significant ($p < 0.05$) on univariate evaluation or identified as clinically important in previous studies. A p value of less than 0.05 was considered statistically significant. All analysis was performed using STATA version 13 (STATA Corp, College Station, Tex).

Results

Between 2010 and 2013, 49.5% ($n=1,790$) of all colectomies performed for diverticulitis in at Washington State's SCOAP hospitals were initiated laparoscopically and 16.5% ($n=295$) were converted. Patients undergoing colectomy had a mean age of 57.8 ± 13 years and 47% were male. The demographics, indications and short-term outcomes for this cohort are summarized in Table 1. Notably, conversions occurred more frequently in patients who had non-elective procedures, chronic complication indications for their operation, or had right/transverse hemicolectomy performed. Patients who had conversions had a longer length of stay, operative time, increased discharges to skilled nursing facilities, and more in-hospital complications and CAEs (all statistically significant at $p < 0.05$).

The surgeon-specific proportion of cases converted declined sharply with increasing case volume of laparoscopic colectomy ($p = 0.03$) (Figure 1). Operations performed for criteria concordant to the 2006 ASCRS guidelines (3+ episodes or chronic complication) did not correlate with surgical volume ($p=0.31$).

After adjusting for surgical volume, chronic complication indication, elective procedure, prior operation, BMI 30+, anatomic segment removed and year of operation, increasing episode number was not associated with higher likelihood of conversion ($p = 0.189$) (Table 2).

A subgroup of 784 patients underwent elective resection solely for episode-based indications (Table 3). These patients were younger (mean age of 55.3 ± 11 years) and had fewer comorbidities than the remainder of the cohort. The conversion rate in this group was 12.9%. After adjustment for surgical volume, prior operation, BMI 30+, anatomic segment removed and year of operation, increasing episode number was not associated with higher likelihood of conversion in this subgroup ($p=0.75$).

Comments

To our knowledge, this study represents the largest cohort of laparoscopic colectomy for diverticulitis in which conversion rates were evaluated. Between 2010 and 2013 at SCOAP

hospitals in Washington State, nearly 1,800 colectomies for diverticulitis were started laparoscopically, and 16.5% were converted. In the subgroup of patients undergoing elective colectomy for an episode-based indication, the rate of conversion was 12.9%. Conversion was higher in non-elective cases and those with complicated disease indications, and was inversely related to surgical volume. After adjustment for several factors known to be predictive of conversion including surgeon experience,^{16,17} increasing episodes of diverticulitis preceding surgery were not associated with increased conversion rate.

Laparoscopic colectomy was widely and rapidly adopted following publication of the safety and benefits of the procedure in colorectal cancer,^{5,6} and its application to diverticulitis has shown benefits in morbidity, hospitalization, cosmesis and overall patient satisfaction, as substantiated by several randomized control trials and a systematic review.^{4,18,19} However, laparoscopic colectomy for diverticulitis has unique technical difficulties that must be overcome to safely complete the operation, such as inflammation distorting landmark anatomy or fistulas to adjacent organs. While conversion rates for all indications for colorectal surgery are approximately 13% according to a large meta-analysis,²⁰ conversion rates specific for diverticulitis (which range between 2 and 20%)^{9,10} have been more difficult to define, in part due to challenges in the management of elective versus non-elective cases, as well as differences between uncomplicated and complicated diverticulitis. In addition, existing literature suggests that surgeon experience, BMI, prior surgery, and presence of complicated disease contribute to conversion from laparoscopy.^{16,17}

Our findings support the higher frequency of conversion in patients requiring urgent operation and those having complicated disease, specifically fistula.^{16,17} Additionally, our study demonstrates that higher case volume is associated with lower conversion rate, as conversion rates dropped as surgeons approached 30 cases. This number should be interpreted cautiously, as the case volume is limited to years in study and only to diverticulitis indications. Published literature estimates the learning curve for all laparoscopic colectomies to be between 30 and 50 cases¹⁵ and may be as high as 60 for diverticulitis.²¹ Much of what is known about learning curves for laparoscopy comes from single centers of excellence,^{15,21} while our cohort, including 198 surgeons across 42 hospitals, may be describing practice that is more generalizable to the wider surgical community. Conversion forfeits the benefits of laparoscopy,^{16,22} and our data confirm that converted cases had a longer operative time, hospitalization, and increased complications.

Overall, complicated diverticular disease is associated with a higher conversion rate than uncomplicated disease. This is not surprising as the most frequently described reasons for conversion are inflammation and adhesions, which occur more frequently when treating complicated diverticulitis. However, a higher rate of conversion with increasing prior episodes of diverticulitis, as would be suspected if multiple episodes were causing chronic inflammation, is not supported by our data. This finding is in line with other population-level studies suggesting that prior episodes are poor predictors of recurrence and progression of disease,^{1,4}

Our study has certain limitations. First, the SCOAP registry is procedural rather than disease-based, and subject to confounding bias. For example, it may be that surgeons with

higher surgical volume (and lower conversion rates) may be more likely to delay surgery until multiple episodes of diverticulitis. We did not, however, find an association between number of operations performed and proportion of cases concordant with 2006 ASCRS indications. In addition, the SCOAP data collection platform is based on operative reports, and there is heterogeneity in the technical aspects of how laparoscopic colectomy is performed and/or how individual surgeons describe these conversions (like hand-assistance, for example). Third, a substantial portion of patients was missing indications data. We have previously described the marked improvement in data quality in this cohort,¹¹ and note that we did not identify systematic differences in patients with and without indication data. Cognizant of this limitation in the data, we defined a subgroup with complete data for analysis, whose results parallel the patterns we saw in the larger cohort. Finally, we acknowledge that even this large cohort may be underpowered to make definitive claims about rates of conversion. As an estimate, to detect a difference between 20% and 10% conversion rates ($\alpha = 0.05$, power = 0.90) with various numbers of preceding episodes would require over 550 patients in each episode group.

Despite these limitations, the results from this statewide cohort of laparoscopic colectomy for diverticulitis add contemporary evidence to inform the management of a disease whose surgical indications are in evolution. Our study also demonstrates that conversion rates from laparoscopy did not increase after multiple episodes of diverticulitis, while confirming previously published associations of conversion from laparoscopy to complicated disease and surgeon experience. These data suggest that delaying elective resection does not prevent patients from benefiting from the advantages of laparoscopic surgery.

Acknowledgments

The authors would like to acknowledge the SCOAP Colorectal Writing Group for their ongoing and thoughtful input on conception and review of this manuscript.

Funding Disclosures: Research reported in this publication was supported by the National Institute Of Diabetes And Digestive And Kidney Diseases of the National Institutes of Health under Award Number T32DK070555. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The Surgical Care and Outcomes Assessment Program (SCOAP) is a Coordinated Quality Improvement Program of the Foundation for Health Care Quality. CERTAIN is a program of the University of Washington, the academic research and development partner of SCOAP.

References

1. Feingold D, Steele SR, Lee S, et al. Practice parameters for the treatment of sigmoid diverticulitis. *Dis Colon Rectum*. 2014; 57(3):284–294.10.1097/DCR.0000000000000075;10.1097/DCR.0000000000000075 [PubMed: 24509449]
2. Yen L, Davis KL, Hodkins P, Loftus EV, Erder MH. Direct costs of diverticulitis in a US managed care population. *american journal of pharmacy*. 2012; 4(5):e118–e129.
3. Anaya DA, Flum DR. Risk of emergency colectomy and colostomy in patients with diverticular disease. *Arch Surg*. 2005; 140(7):681–685.10.1001/archsurg.140.7.681 [PubMed: 16027334]
4. Morris AM, Regenbogen SE, Hardiman KM, Hendren S. Sigmoid diverticulitis: A systematic review. *JAMA*. 2014; 311(3):287–297. [PubMed: 24430321]
5. Clinical Outcomes of Surgical Therapy Study Group. A comparison of laparoscopically assisted and open colectomy for colon cancer. *N Engl J Med*. 2004; 350(20):2050–2059.10.1056/NEJMoa032651 [PubMed: 15141043]

6. Lacy AM, Garcia-Valdecasas JC, Delgado S, et al. Laparoscopy-assisted colectomy versus open colectomy for treatment of non-metastatic colon cancer: A randomised trial. *Lancet*. 2002; 359(9325):2224–2229.10.1016/S0140-6736(02)09290-5 [PubMed: 12103285]
7. Wong WD, Wexner SD, Lowry A, et al. Practice parameters for the treatment of sigmoid diverticulitis--supporting documentation. the standards task force. the american society of colon and rectal surgeons. *Dis Colon Rectum*. 2000; 43(3):290–297. [PubMed: 10733108]
8. Rafferty J, Shellito P, Hyman NH, Buie WD. Standards Committee of American Society of Colon and Rectal Surgeons. Practice parameters for sigmoid diverticulitis. *Dis Colon Rectum*. 2006; 49(7): 939–944.10.1007/s10350-006-0578-2 [PubMed: 16741596]
9. Reissfelder C, Buhr HJ, Ritz JP. Can laparoscopically assisted sigmoid resection provide uncomplicated management even in cases of complicated diverticulitis? *Surg Endosc*. 2006; 20(7): 1055–1059.10.1007/s00464-005-0529-3[doi] [PubMed: 16736310]
10. Jones OM, Stevenson AR, Clark D, Stitz RW, Lumley JW. Laparoscopic resection for diverticular disease: Follow-up of 500 consecutive patients. *Ann Surg*. 2008; 248(6):1092–1097.10.1097/SLA.0b013e3181884923[doi] [PubMed: 19092355]
11. Simianu VV, Bastawrous AL, Billingham RP, et al. Addressing the appropriateness of elective colon resection for diverticulitis: A report from the SCOAP CERTAIN collaborative. *Ann Surg*. 2014; 260(3):533–8. discussion 538-9. 10.1097/SLA.0000000000000894[doi] [PubMed: 25115429]
12. Devine EB, Alfonso-Cristancho R, Devlin A, et al. A model for incorporating patient and stakeholder voices in a learning health care network: Washington state's comparative effectiveness research translation network. *J Clin Epidemiol*. 2013; 66(8 Suppl):S122–9.10.1016/j.jclinepi.2013.04.007;10.1016/j.jclinepi.2013.04.007 [PubMed: 23849146]
13. Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. *J Clin Epidemiol*. 1992; 45(6):613–619. [PubMed: 1607900]
14. Kwon S, Billingham R, et al. Surgical Care and Outcomes Assessment Program (SCOAP) Collaborative. Adoption of laparoscopy for elective colorectal resection: A report from the surgical care and outcomes assessment program. *J Am Coll Surg*. 2012; 214(6):909–18.e1.10.1016/j.jamcollsurg.2012.03.010[doi] [PubMed: 22533998]
15. Dincler S, Koller MT, Steurer J, Bachmann LM, Christen D, Buchmann P. Multidimensional analysis of learning curves in laparoscopic sigmoid resection: Eight-year results. *Dis Colon Rectum*. 2003; 46(10):1371–8. discussion 1378-9. 10.1097/01.DCR.0000089054.22223.41[doi] [PubMed: 14530677]
16. Le Moine MC, Fabre JM, Vacher C, Navarro F, Picot MC, Domergue J. Factors and consequences of conversion in laparoscopic sigmoidectomy for diverticular disease. *Br J Surg*. 2003; 90(2):232–236.10.1002/bjs.4035[doi] [PubMed: 12555302]
17. Vaccaro CA, Rossi GL, Quintana GO, Soriano ER, Vaccarezza H, Rubinstein F. Laparoscopic colorectal resections: A simple predictor model and a stratification risk for conversion to open surgery. *Dis Colon Rectum*. 2014; 57(7):869–874.10.1097/DCR.0000000000000137[doi] [PubMed: 24901688]
18. Gervaz P, Inan I, Perneger T, Schiffer E, Morel P. A prospective, randomized, single-blind comparison of laparoscopic versus open sigmoid colectomy for diverticulitis. *Ann Surg*. 2010; 252(1):3–8.10.1097/SLA.0b013e3181dbb5a5[doi] [PubMed: 20505508]
19. Klarenbeek BR, Bergamaschi R, Veenhof AA, et al. Laparoscopic versus open sigmoid resection for diverticular disease: Follow-up assessment of the randomized control sigma trial. *Surg Endosc*. 2011; 25(4):1121–1126.10.1007/s00464-010-1327-0[doi] [PubMed: 20872022]
20. Miskovic D, Ni M, Wyles SM, Tekkis P, Hanna GB. Learning curve and case selection in laparoscopic colorectal surgery: systematic review and international multicenter analysis of 4852 cases. *Dis Colon Rectum*. 2012 Dec; 55(12):1300–10.10.1097/DCR.0b013e31826ab4dd[doi] [PubMed: 23135590]
21. Tekkis PP, Senagore AJ, Delaney CP, Fazio VW. Evaluation of the learning curve in laparoscopic colorectal surgery: Comparison of right-sided and left-sided resections. *Ann Surg*. 2005; 242(1): 83–91. doi:00000658-200507000-00012[pii]. [PubMed: 15973105]

22. Hassan I, Cima RR, Larson DW, et al. The impact of uncomplicated and complicated diverticulitis on laparoscopic surgery conversion rates and patient outcomes. *Surg Endosc.* 2007; 21(10):1690–1694.10.1007/s00464-007-9413-7[doi] [PubMed: 17593455]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Demographics, indications and outcomes, stratified by conversion.

		Not converted		Converted		Total cohort		
		N	%	N	%	N	%	
		1495	83.5	295	16.5	1790	100	p value
Mean age (SD)		57.5 (12.6)		59.1 (12.9)		57.8 (12.7)		0.054
<hr/>								
Sex								
M		707	47.3	135	45.8	842	47.0	0.63
F		788	52.7	160	54.2	948	53.0	
<hr/>								
White		1,307	87.4	254	86.1	1,561	87.2	0.52
<hr/>								
Comorbidities								0.29
0		1,108	74.1	214	72.5	1,322	73.9	
1		311	20.8	59	20.0	370	20.7	
2		59	4.0	15	5.1	74	4.1	
3		17	1.1	7	2.4	24	1.3	
<hr/>								
BMI 30+		620	41.5	115	39.0	735	41.1	0.43
<hr/>								
Indications								
<hr/>								
Non-elective		114	7.6	56	19.0	170	9.5	p<0.001
<hr/>								
Number of prior episodes*								p<0.001
None		115	12.1	43	23.4	158	13.9	
1		142	14.9	31	16.9	173	15.2	
2		137	14.4	27	14.7	164	14.5	
3 to 10		520	54.7	73	39.7	593	52.5	
> 10		37	4.0	10	5.4	47	4.0	
<hr/>								
Chronic complication**		238	16.1	65	22.1	303	17.1	0.012
<hr/>								
Colovesicular fistula		120	8.1	34	11.6	154	8.7	0.054
<hr/>								

	Not converted		Converted		Total cohort		p value
	N	%	N	%	N	%	
	1495	83.5	295	16.5	1790	100	
Current GI bleed	33	2.2	7	2.4	40	2.3	0.87
Stricture	61	4.1	17	5.8	78	4.4	0.20
Other fistula	36	3.1	15	7.1	51	3.8	0.01
Missing indication	453	30.3	81	27.5	534	29.8	0.33
Operation type							
Right Hemicolectomy	72	4.8	26	8.8	98	5.5	0.01
Left Hemicolectomy	450	30.1	83	28.1	533	29.8	0.50
Low Anterior Resection ***	973	65.1	203	68.8	1,176	65.7	0.22
Total Abdominal Colectomy	15	1	1	0.3	16	0.9	0.27
Prior surgery	509	34.1	117	40.0	626	35.0	0.17
Outcomes							
Mean OR time, min (SD)	174 (74)		197 (92)		178 (78)		p<0.001
Mean Length of Stay (SD)	5.2 (7.0)		7.2 (5.4)		5.6 (6.8)		p<0.001
Discharge home	1,430	95.7	266	90.2	1,696	94.8	0.002
In-hospital complication	90	6.0	49	16.6	139	7.8	p<0.001
CAE ****	140	9.4	70	23.7	210	11.7	p<0.001

* Number of episode % calculated from population with known episode information (total n = 1135, non-converted n =951, converted n=184)

** calculated as Pearson χ^2 for any chronic complication indication between converted and non-converted cases

*** Low Anterior Resection category includes sigmoidectomy

***** Composite Adverse Events (CAE), includes complications as well as re-operative interventions and in-hospital deaths

Table 2

Association between number of prior episodes and likelihood of conversion.

Conversion	Unadjusted Association				Adjusted Association *			
	OR	CI	P		OR	CI	P	
Prior episodes **								
None								
	REF	REF			REF	REF		
1	0.58	0.35	0.99	0.04	0.71	0.41	1.22	0.22
2	0.53	0.31	0.91	0.02	0.59	0.33	1.05	0.07
3 to 10	0.38	0.24	0.58	p<0.001	0.53	0.33	0.85	0.01
10+	0.72	0.33	1.58	0.42	0.99	0.44	2.25	0.99
Surgical volume								
	0.99	0.98	0.99	p<0.001	0.99	0.98	0.99	p<0.001
Chronic complication								
	1.48	1.09	2.02	0.01	1.37	0.96	1.95	0.08
Elective								
	0.35	0.25	0.50	p<0.001	0.43	0.29	0.65	p<0.001
BMI 30+								
	0.90	0.70	1.16	0.43	0.92	0.69	1.21	0.54
Prior operation								
	0.79	0.61	1.02	0.07	0.75	0.57	1.00	0.052
Right Hemicolectomy								
	1.91	1.20	3.05	0.01	1.63	0.93	2.83	0.086
Year								
	0.86	0.77	0.96	0.01	0.87	0.77	0.98	0.03

* Adjusted for all other variables listed

** Association listed for each category of prior episodes using no prior episodes as reference; overall associations for increasing category of episode are OR 0.91 (0.83, 0.99; p=0.026) and OR 0.94 (0.87, 1.03; p=0.189) for unadjusted and adjusted models, respectively.

Table 3
Elective surgery solely for episode-based indication: subgroup demographics and indications stratified by conversion.

		Not converted		Converted		p value
		N	%	N	%	
Mean age (SD)		683	87.12	101	12.88	
		55.0 (11.6)		56.9 (12.7)		0.13
<hr/>						
Sex						
		M	311	45.53	55	54.46
		F	372	54.47	46	45.54
<hr/>						
White			605	88.58	88	87.13
			88.58		87.13	
<hr/>						
Comorbidities						0.53
		0	513	75.11	70	69.31
		1	146	21.38	25	24.75
		2	20	2.93	5	4.95
		3	4	0.59	1	0.99
<hr/>						
BMI 30+			289	42.31	47	46.53
			42.31		46.53	
<hr/>						
Indications						
<hr/>						
Number of prior episodes						0.19
		1	91	13.32	17	16.83
		2	102	14.93	14	13.86
		3 to 10	459	67.2	61	60.4
		> 10	31	4.54	9	8.91
<hr/>						
Operation type						
<hr/>						
Right Hemicolectomy			21	3.07	5	4.95
			226	33.09	26	25.74
Left Hemicolectomy						0.14
			447	65.45	74	73.27
Low Anterior Resection *						0.12
			5	0.73	0	0
Total Abdominal Colectomy						0.39
<hr/>						

		Not converted		Converted		p value
		N	%	N	%	
Prior abdominal surgery		683	87.12	101	12.88	0.13
		231	33.82	42	41.58	

* Low Anterior Resection category includes sigmoidectomy