reporting medical mistakes. The survey, which was designed to adequately reflect gender and differing geographical areas and levels of seniority, included family doctors and hospital specialists.

The survey did not specifically ask about "near misses," but responses showed that only 15% of serious incidents resulting in death or serious disability were reported to existing schemes.

Almost all (97%) of the 2575 respondents felt that a reporting system would improve patient care, but eight out of 10 said they did not trust their employing NHS trust or the Department of Health to run it. Instead, almost all respondents favoured a system that was completely independent.

The National Patient Safety Agency began its roll-out of a new, confidential, and anonymous reporting system (the National Reporting and Learning System) across NHS trusts in England and Wales last November, with full implementation by the end of this year.

Caroline White *London*

Royal College concerned over transfer of out of hours GP services

Professor David Haslam, chairman of the council of the Royal College of General Practitioners, has expressed concerns to the parliamentary health select committee over the "potential crisis" in GP out of hours services when responsibility for night and weekend cover is transferred to primary care trusts on 31 December.

Commenting on the readiness of primary care trusts for their new role, Professor Haslam said, "I think there is a general underestimation of the complexity of what goes on in an out of hours consultation."

Dr Mark Reynolds, chairman of the National Association of GP Co-operatives, voiced similar concerns: "People in primary care trusts are taking this job on with no experience of it and think things will be fine."

But the NHS Alliance, the main organisation representing primary care trusts, submitted written evidence stating: "Most

trusts have the replacement of GP out-of-hours traditional arrangements in hand and are confident in alternative service provision."

Bruno Rushforth *BMJ*

Young black women have more aggressive breast cancers than white women

Breast tumours grow faster and are more aggressive in young black women than young white women, says new US research.

For social, economic, and cultural reasons breast cancer in African-American women is diagnosed later than in white women, giving poorer survival rates.

Tumours from 124 African-American women and 397 white women aged 20 to 54 years in Atlanta, Georgia, were examined for the proteins that control how quickly a cancer cell divides (*Cancer* 2004;100:2533-42).

Nearly one in five of the black women had cancer that was in stage III or higher, compared with only one in eight white women. Black women were also five times more likely to have high grade, aggressive tumours than white women.

"We looked at markers as they relate to race and to the kind of tumour these women presented with," said Dr Peggy Porter, one of the authors and a pathologist at the Fred Hutchinson Cancer Research Center, Seattle, who reviewed all the tumour specimens without knowing the race of the participants.

The odds of a breast carcinoma diagnosis at a younger age and at a later stage were higher for African-American women than for white women, it reports.

Black women were four times more likely than white women to have a high concentration of cyclin E, a factor that is associated with poorer survival. High concentrations of cyclin D are associated with better survival, but African-American women were only half as likely to have high cyclin D concentrations as white women, the research showed.

Janice Hopkins Tanne *New York*

United States brings in new rules to prevent surgical errors

Scott Gottlieb *New York*

New rules in the United States that came into effect this week require doctors and hospitals to take certain specified steps to prevent operating on the wrong body part or wrong patient.

Among the requirements, surgeons and nurses must take what is being referred to as "time out" before operating on patients, to confirm that the right patient is on the table and that doctors are operating on the right body part and the right side of the body.

Hospital regulators hope the rules, which came into effect on 1 July, will put an end to growing reports of operations on the wrong site or the wrong patient or that involved the wrong procedure.

"These should never happen," said Dr Dennis O'Leary, who heads the Joint Commission on Accreditation of Healthcare Organizations, the agency that accredits medical facilities in the United States, allowing them to continue to operate. The agency can revoke the accreditation of hospitals or other surgery sites that don't comply with the new safety steps.

No precise figures on exactly how many such mistakes occur are available, because the commission receives only voluntary reports. It is assumed that the reports are only a small percentage of the 70 million operations done in the United States each year.

More than 40 million in-patient surgical procedures and 31 million outpatient procedures are performed each year in the United States. The commission calculated that from 1995 to the end of January 2004 there were 300 cases of surgical errors involving the wrong part or side of the body, the wrong procedure, or the wrong patient.

Of the 300 cases some 59% involved surgeons operating on the wrong side of a patient. Ten per cent of the cases involved the wrong surgical procedure and 12% the wrong patient. Another 19% involved the wrong site on the body.

One of the new rules is that surgeons must literally sign the incision site—while the patient is awake and cooperating if possible—with a marker that will not wash off in the operating room.

The whole operating team must also stop all other work just before surgery begins and go through a checklist to ensure that the correct patient is on the table and that everyone—surgeons, nurses, anaesthetists, and technicians—agrees what procedure is being done, on what body part.

More than 40 medical organisations, including the American Academy of Orthopedic Surgeons, have now signed up to help the accreditation commission teach the new rules. □

A marked thoracotomy incision, signed by the surgeon, to show the correct side for the operation.