Published in final edited form as:

Curr Top Dev Biol. 2015 ; 111: 401–419. doi:10.1016/bs.ctdb.2014.11.012.

Developing and regenerating a sense of taste

Linda A. Barlow^{1,*} and Ophir D. Klein²

¹Department of Cell and Developmental Biology, Graduate Program in Cell Biology, Stem Cells and Development, and the Rocky Mountain Taste and Smell Center, University of Colorado, School Medicine, Anschutz Medical Campus, Aurora, CO 80045. USA

²Departments of Orofacial Sciences and Pediatrics, Program in Craniofacial and Mesenchymal Biology, and Institute for Human Genetics, University of California San Francisco, San Francisco, California 94143, USA

Abstract

Taste is one of the fundamental senses, and it is essential for our ability to ingest nutritious substances and to detect and avoid potentially toxic ones. Taste buds, which are clusters of neuroepithelial receptor cells, are housed in highly organized structures called taste papillae in the oral cavity. Whereas the overall structure of the taste periphery is conserved in almost all vertebrates examined to date, the anatomical, histological, and cell biological, as well as potentially the molecular details of taste buds in the oral cavity are diverse across species and even among individuals. In mammals, several types of gustatory papillae reside on the tongue in highly ordered arrangements, and the patterning and distribution of the mature papillae depends on coordinated molecular events in embryogenesis. In this review, we highlight new findings in the field of taste development, including how taste buds are patterned and how taste cell fate is regulated. We discuss whether a specialized taste bud stem cell population exists and how extrinsic signals can define which cell lineages are generated. We also address the question of whether molecular regulation of taste cell renewal is analogous to that of taste bud development. Finally, we conclude with suggestions for future directions, including the potential influence of the maternal diet and maternal health on the sense of taste in utero.

Taste is important for life. It serves as the gateway to substances that enter the body, allowing us to distinguish nutritious food items from potentially toxic ones. Classically, taste buds in the oral cavity, primarily on the tongue, were shown to detect 5 basic tastes: sour, salty, bitter, sweet and umami – savory or “deliciousness” in Japanese. More recently, fatty acids and calcium have emerged as potential tastants that can be sensed by taste bud cells (Iwata et al., 2014; Liman et al., 2014; Passilly-Degrace et al., 2014; Tordoff et al., 2008b; Tucker et al., 2014).

Among humans, taste preferences vary widely, and these preferences in turn influence dietary choices, which impact body weight and therefore health (Mennella, 2014). A key question is what underlies this variability. Not surprisingly, it appears that environmental,

* author for correspondence, Linda A. Barlow, Linda.barlow@ucdenver.edu, Voice: +1 303-724-3438, FAX: +1 303-724-3420, Mail: Dept CDB, MS 8101, 12801 E 17th Ave, Aurora CO 80045, USA.

genetic, and epigenetic mechanisms are at play. In mammals, including humans, the maternal diet during gestation and postnatal lactation is learned by her offspring. In humans, innervated and differentiated taste buds that are presumably functional are evident by 10–13 weeks of development (Bradley and Stern, 1967; Witt and Reutter, 1996, 1998). Throughout gestation, taste stimuli reach the amniotic fluid, which is continually swallowed by the fetus, and following birth, tastes of the maternal diet are evident in breast milk. This exposure heavily influences the dietary choices of offspring as they discover these new tastes (Beauchamp and Mennella, 2009; Mennella, 2014). However, maternal health also impacts the gestational experience, as it results in fetal metabolic programming via presumed epigenetic mechanisms (Dyer and Rosenfeld, 2011), which in the case of diabetic or obese mothers, can predispose offspring to diabetes and cardiovascular disease. Although conclusive studies regarding alterations in taste sensitivity in this context have not been performed, it is well known that diabetes and obesity affect taste preferences in adults. For example, in diabetic patients, taste responses, especially to sweet, are blunted (Wasalathanthri et al., 2014), and obese individuals also have decreased taste sensitivity (Stewart et al., 2010; Stewart et al., 2011).

The pattern of taste buds is established during embryogenesis, such that the first functional taste bud cells are specified during gestation and differentiate around birth. Whereas most sensory epithelia, such as hair cells of the inner ear and photoreceptors of the retina, have limited renewal potential, taste cells are remarkable in their ability to turn over rapidly and continuously throughout adult life (Beidler and Smallman, 1965; Farbman, 1980; Feng et al., 2014; Hamamichi et al., 2006; Perea-Martinez et al., 2013). Despite regular sensory cell replacement, the sense of taste is remarkably stable throughout life in healthy individuals. However, taste can be distorted or lost in cancer patients when these individuals are treated with chemotherapeutic drugs, and in head and neck cancer patients following targeted radiotherapy (Berteretche et al., 2004; Hong et al., 2009; Ruo Redda and Allis, 2006; Vissink et al., 2003). These treatments are thought to disrupt taste function by diminishing taste bud cell renewal (Nguyen et al., 2012, and references therein). Thus, we hypothesize that both regulation of taste bud development, including patterning and formation of the proper ratio of taste receptor cell types, and taste bud renewal, i.e., generation of functional taste cell types in the proper ratios with the proper timing, underlie variability in taste function and dysfunction.

In this review, we highlight new data in the context of the important open questions in the field rather than providing an exhaustive survey of the literature; for more comprehensive reviews on taste development, regeneration and function, please see (Kapsimali and Barlow, 2013), (Feng et al., 2014) and (Liman et al., 2014), respectively.

How are taste buds patterned?

Taste bud distribution is highly variable across vertebrate species, including in mammals (Jackowiak, 2006 and references therein), fish, amphibians and birds (Erdogan and Iwasaki, 2014; Finger, 1997; Northcutt, 2004). In addition, taste bud pattern varies even within a single species, including in humans (Fischer et al., 2013; Miller and Reedy, 1990) and

rodents (mouse inbred lines: Reiner et al., 2008; rat: Miller and Preslar, 1975; Tordoff et al., 2008a).

The initial pattern of taste buds on the tongue is evident at mid-gestation (at embryonic day (E) 12.0 in mice), when bilateral rows of columnar epithelial placodes (taste placodes) form in the otherwise cuboidal epithelium of the developing tongue (Farbman, 1965; Mistretta, 1972). Subsequently, placodes undergo morphogenesis into mushroom-shaped (fungiform) taste papillae, which house taste buds that differentiate at birth. Taste placodes express Sonic hedgehog (*Shh*) from the earliest stages of their development, and expression of *Shh* persists in the apices of papillae through the remainder of embryogenesis (Bitgood and McMahon, 1995; Hall et al., 1999; Jung et al., 1999). Lineage tracing of the *Shh*⁺ taste placodes, commencing at E12.5 or 13.5, reveals that these cells differentiate directly into the first taste bud cells at birth but do not contribute to the surrounding taste papillae (Thirumangalathu et al., 2009). Rather, we have proposed that *Shh*-descendent taste bud precursor cells may function as signaling centers to induce adjacent epithelial and mesenchymal cells to form taste papillae (Thirumangalathu et al., 2009).

A number of pathways regulate the initial patterning of taste placodes in rodents, including Wnt/ β -catenin (Iwatsuki et al., 2007; Liu et al., 2007), Bone Morphogenetic Proteins (Bmps) (Beites et al., 2009; Zhou et al., 2006), *Shh* (Hall et al., 2003; Liu et al., 2004; Mistretta et al., 2003), Epidermal Growth Factor (*Egf*) (Liu et al., 2008), and Fibroblast Growth Factors (*Fgfs*) (Kapsimali et al., 2011; Petersen et al., 2011; reviewed in Kapsimali and Barlow, 2013)(Table 1). In particular, normal activation of the Wnt/ β -catenin pathway within the developing lingual epithelium is required for formation of taste placodes, whereas ectopic activation of the pathway in the entire epithelium drives differentiation into enlarged *Shh*⁺ taste bud precursors embedded in oversized fungiform papillae (Liu et al., 2007).

Interestingly, Wnt ligands are expressed in both epithelial and mesenchymal compartments in the developing tongue (Iwatsuki et al., 2007; Liu et al., 2007; Liu et al., 2012), leaving open the question of which source(s) of Wnt protein are responsible for taste patterning. Recently, conditional deletion of *Wls* (also known as *Gpr177* or *Evi1*) from the early oral endoderm under the *Shh*^{GFP-Cre} allele (Harfe et al., 2004) revealed that epithelial Wnt production is required for taste placode initiation (Zhu et al., 2014). The *Wls* gene encodes an intracellular protein that enables Wnt ligand secretion (Banziger et al., 2006), and deletion of epithelial *Wls* which causes loss of epithelial Wnt ligand secretion, leads to absence of taste placode formation. One caveat to these studies however, is that *Shh* is expressed within the oral endoderm commencing at E9.5 (Echelard et al., 1993) and thus loss of *Wls* function using the *Shh*^{GFP-Cre} is also induced early with respect to tongue and taste placode formation, which begin at E11.5, (Kaufman, 1999) and E12.0 (Hall et al., 2003), respectively. Therefore, it is formally possible that early epithelial WLS function is required for development of epithelial competence to respond to later Wnt signals from the mesenchyme. Nonetheless these findings are consistent with experimental data from axolotl (salamander) embryos, where early taste bud specification and patterning are governed by mechanisms intrinsic to the epithelium and independent of oral mesenchyme (Barlow, 2001; Barlow and Northcutt, 1997; Parker et al., 2004). Thus, numerous pathways can affect taste

bud patterning, and subtle differences in timing, competence to receive the signal, and the strength of the signal may be equally influential.

In addition, there are significant distinctions in patterning of different regions of the tongue as well as patterning of taste papillae in different organisms. One area in which this has been explored is the difference between the small anterior fungiform papillae and the large posterior circumvallate papilla (CVP). Most studies of development and patterning have focused on the fungiform papillae; although there is a large literature concerning physiology, anatomy and cell biology of the adult CVP, only a limited number of developmental studies have been performed on this organ (see Kapsimali and Barlow, 2013 for review). A number of pathways that are known to regulate fungiform papillae pattern have no reported phenotype in the CVP, including the Shh (Mistretta et al., 2003), Bmp (Beites et al., 2009), and Wnt (Iwatsuki et al., 2007) pathways. In contrast, the Fgf pathway has been shown to be a critical regulator of CVP development in mice. A balance between Sprouty (Spry) genes and *Fgf10*, which respectively antagonize and activate receptor tyrosine kinase (RTK) signaling, regulate the number of CVPs (Petersen et al., 2011), such that in wild type mice, only a single CVP forms. Deletion of a single Sprouty family member, *Spry2*, resulted in duplication of the CVP, as a result of an increase in the size of the CV placode progenitor field. Combined deletion of two Sprouty genes in *Spry1*^{-/-};*Spry2*^{-/-} embryos led to the formation of multiple CVPs, demonstrating the redundancy of Sprouty genes in regulating the size of the progenitor field. By contrast, deletion of *Fgf10* led to absence of the CVP, thus identifying FGF10 as an inductive, mesenchyme-derived factor for taste papillae. Recently, the transcription factor Wilms' tumor 1 protein (WT1) was found to have a critical role in CVP development, as deletion of *Wt1* led to failure of CVP development (Gao et al., 2014). Several WT1 target genes that are members of canonical signaling pathways were identified, including *Lef1* from the Wnt pathway, *Ptch1* from the Shh pathway, and *Bmp4*. Interestingly, there are also some hints emerging that anterior vs posterior taste papillae may be regulated in opposite ways by specific signaling pathways, such as the Fgf pathway (Petersen et al., 2011). This may be due to origins in different germ layers, as the anterior tongue epithelium is derived from ectoderm whereas epithelium covering the posterior tongue has an endodermal origin (Adams, 1931; Barlow, 2000; Rothova et al., 2012). Of interest, while taste buds are restricted to the oral cavity of most vertebrates, several fishes have evolved external taste buds, including catfishes which have thousands of taste buds distributed in the head and trunk epithelium (Atema, 1971; Landacre, 1907; Northcutt, 2005). Their external location suggests that these taste buds must arise from surface ectoderm (Landacre, 1907), while oral taste buds of fish and amphibians originate primarily from endoderm (Barlow and Northcutt, 1995; Johnston, 1910). Thus the catfish would be an ideal model to test if and how embryonic origin affects the mode of taste bud development.

Regulation of taste cell fate

Taste buds comprise 60–100 elongate cells, which are classified into 3 morphological types and as many as 5 or more functional categories (Figure 1)(Feng et al., 2014; Liman et al., 2014). In animal models used in the majority of developmental studies including fish and rodents, taste cells differentiate postnatally (see references in Kapsimali and Barlow, 2013), such that in adults, the different taste cell types occur in proper ratios within each bud

(Chaudhari and Roper, 2010; Kim et al., 2003; Ma et al., 2007; Ohtubo and Yoshii, 2010). Generally, the Type I or glial-like cells are most common, followed by Type II sweet/bitter/umami detectors, and least frequent are the Type III sour receptor cells, although these ratios vary with respect to location in the tongue. For example, in rats and mice, Type III cells are more frequent in taste buds in the posterior CVP than in taste buds of the anterior fungiform papillae (Ma et al., 2007; Ohtubo and Yoshii, 2010). In mice, taste buds in the anterior tongue have more sweet cells per bud, whereas in posterior tongue, taste buds have more bitter cells (Kim et al., 2003; Tizzano et al., 2008). How and when these cell fates are assigned in embryos remains an important open question.

Not surprisingly, in light of its important role in fate decisions in many tissues, Notch function has been implicated in taste cell fate decisions during embryonic development (reviewed in Kapsimali and Barlow, 2013). In mice, Notch pathway genes are expressed in and around the developing CV papilla in the posterior tongue, and this expression begins after the papilla is specified (Seta et al., 2003). In late gestation embryos, *Mash1/Ascl1*, which is transcriptionally repressed by Notch signaling (Kageyama and Ohtsuka, 1999), is expressed in small numbers of epithelial cells in locations consistent with that of the first differentiated taste buds at early postnatal stages; this expression pattern suggested a role for *Ascl1* in the differentiation of one or more taste cell types. Indeed, genetic deletion of *Ascl1* results in loss of expression of numerous markers of Type III taste cells, supporting the hypothesis that *Ascl1* is required for Type III cell fate (Kito-Shingaki et al., 2014; Seta et al., 2011). Interestingly, Type III cells are the taste cells most similar to neurons, and *Mash1* is a proneural gene that drives expression of Notch ligands cell autonomously to activate Notch signaling in adjacent cells, but *Mash1* also directs fate cell autonomously while keeping neighbors in a stem cell state (Kageyama et al., 2005). In addition, the transcription factor *Hes1*, considered a primary Notch target gene (Ohtsuka et al., 1999), has been shown to repress differentiation of Type II taste cells, as excess Type II cells differentiate in *Hes1*^{-/-} taste papillae (Ota et al., 2009). Notch function in taste cell selection is conserved in zebrafish, where it again plays a role in specifying Type II-like versus Type III-like cell fates (Kapsimali et al., 2011).

In addition to components of the Notch pathway, another transcription factor, *Skn1a/Pou2f3* has been shown to be required for differentiation of Type II sweet/bitter/umami cells in adults, as taste buds in adult *Skn1a*^{-/-} mice lack Type II cells and have excess Type III cells (Matsumoto et al., 2011). If and how this transcription factor forms a gene regulatory network with *Ascl1* and/or *Hes1* remains to be explored. Finally, and intriguingly, genetic control of specification of the most common taste cell fate, the Type I glial-like cell, remains a mystery.

How can we link embryonic development and adult taste cell renewal?

As mentioned above, embryonic *Shh*⁺ placodes are taste bud precursors, which differentiate into the first taste bud cells. As taste bud cells renew, these first taste cells are ultimately replaced over time. However, embryonic *Shh*⁺ cells do not contribute to the stem cell pool that enables adult taste cell renewal, as all *Shh*-descendent taste bud cells are lost by 4 months postnatally (Thirumangalathu et al., 2009). Rather, in adults, taste receptor cells are

renewed from cytokeratin (K) 14+/K5+ basal keratinocytes adjacent to taste buds (Figure 2) (Okubo et al., 2009). K14+ lingual keratinocytes also give rise to the general epithelium of the tongue, which comprises non-taste filiform papillae (Hume and Potten, 1976; Mistretta, 1972). How K14+/K5+ stem cells are regulated to produce both taste buds and general epithelium is poorly understood. However, this population generates only non-taste epithelium in vitro (Luo et al., 2009), suggesting that either taste bud stem cells represent a rare or distinct population, or that key extrinsic signals are required that were absent under the culture conditions employed.

Intriguingly, other lingual epithelial stem populations that contribute to non-taste filiform papillae, but not to taste buds, have been identified. One gene that has been recently studied is *Bmi1* which together with K14 and K5 labels cells at the base of the interpapillary pit (Tanaka et al., 2013). These cells were reported to be unipotent stem cells for keratinized epithelial cells but not for taste bud cells. In vitro organoids could also be generated from single *Bmi1*-positive cells (Hisha et al., 2013). Similarly, lineage tracing using a *Tcf3^{CreER}* knock-in mouse model showed that *Tcf3* marks stem cells as well as transient progenitors and cells undergoing active differentiation in the tongue (Howard et al., 2014).

Is there a specialized taste bud stem cell population, or are extrinsic signals responsible for defining which cell lineages are generated, and when?

Recently, a new population of lingual stem cells has been identified that gives rise to both epithelium and taste buds in the large CVP situated at the midline of the posterior tongue. These cells express the leucine-rich repeat-containing G-protein coupled receptor 5 (LGR5), which is also expressed by the stem cells of the intestine and other organs (Barker et al., 2007; Ng et al., 2014; Plaks et al., 2013). In the CVP of mice carrying an *Lgr5* reporter allele (*Lgr5^{GFP-CreER}*; Barker et al., 2007), taste buds themselves are not *Lgr5^{GFP+}*; instead, GFP-bright epithelial cells are situated at the deepest portion of the papilla where taste buds are not present, while low expressing *Lgr5^{GFP+}* basal epithelial cells are found higher up in the papilla and adjacent to taste buds. Lineage tracing revealed that *Lgr5+* cells give rise all 3 taste cell types (Takeda et al., 2013; Yee et al., 2013), and that the deep *Lgr5^{GFP+}* cells are lineage labeled, and persist for months (Yee et al., 2013). Interestingly, both groups reported that *Lgr5* expression was not detectable in the anterior tongue by GFP expression, or by PCR for *Lgr5* transcripts, indicating that this stem cell population is specific to the posterior CVP. This result highlights again the distinction described above between ectodermally-derived anterior taste buds and endodermally-derived CVP taste buds, and it is intriguing in that *Lgr5* also marks a key stem cell population in the endodermally derived gut (see Barker et al., 2013 for review).

To date, an analogous stem cell population remains to be identified for fungiform taste buds. However, our recent data point to a different model, where specialized stem cells may not be required in the anterior tongue, but rather extrinsic signals can drive bipotent K14+ cells toward the taste fate (Castillo et al., 2014). Using an inducible gain-of-function approach, we over-expressed SHH (SHHcKI) in K14+ basal keratinocytes and found that ectopic taste

buds form throughout the non-taste epithelium and are interspersed among the endogenous fungiform papillae. Thus, SHH signals can induce cell type-replete taste bud differentiation from K14+ progenitors regardless of location. These results indicate that K14+ keratinocytes are broadly competent to generate taste receptor cells in response to SHH, revealing an unappreciated competency of the anterior tongue epithelium to build taste buds outside of fungiform papillae in the adult tongue. Expansion of taste fate in the lingual epithelium is well documented in the embryonic tongue, as all of the following promote the taste fate in lingual regions where taste buds do not form in controls: increased β -catenin in vitro or in vivo (Iwatsuki et al., 2007; Liu et al., 2007; Okubo et al., 2006), SHH inhibition in vitro (Hall et al., 2003; Liu et al., 2004; Mistretta et al., 2003), or genetic deletion of *Follistatin* in vivo (Beites et al., 2009). Interestingly, manipulation of each of these pathways was reported to have no impact on the development of the posterior CVP (Beites et al., 2009; Iwatsuki et al., 2007; Liu et al., 2004; Mistretta et al., 2003). Likewise, in adult mice, despite SHH overexpression, taste buds in the posterior CV papilla were only mildly affected (Castillo, Klein and Barlow, unpublished), again suggesting that SHH may function differently in ectodermally derived anterior versus endodermally derived posterior lingual epithelium even in adulthood.

Is molecular regulation of taste cell renewal analogous to that of taste bud development?

In contrast to development, during which several pathways have been shown to function in taste bud development, only Shh function has been examined thus far in adults. Specifically, ectopic overexpression of SHH promotes differentiation of cell type-replete taste buds (Castillo et al., 2014). This pro-taste function contrasts directly with the taste-repressive function of Shh signaling in development. This difference is likely due to differences in cells receiving SHH signals, as *Shh*+ cells in adult taste buds and in taste placodes appear quite similar (Miura and Barlow, 2010; Miura et al., 2006; Nakayama et al., 2008): both are specified, immediate postmitotic precursors of differentiated taste bud cells and both are SHH non-responsive, as they do not express *Ptch1* or *Gli1* (Hall et al., 1999; Liu et al., 2013; Miura et al., 2001; Miura et al., 2014; Thirumangalathu et al., 2009). In embryos, established *Shh*+ cells inhibit neighboring *Ptch1* and *Gli1* expressing epithelial cells (Hall et al., 1999) from acquiring a taste fate (Hall et al., 2003; Iwatsuki et al., 2007; Liu et al., 2004; Mistretta et al., 2003). However, in adult epithelium, while postmitotic precursors within taste buds also signal via SHH to surrounding *Gli1*+ and *Ptch1*+ taste papilla epithelial cells (see Figure 2)(Liu et al., 2013; Miura et al., 2004; Miura et al., 2001), SHH now promotes rather than represses taste fate (Castillo et al., 2014). When this shift in SHH function occurs may shed light on the timing of the transition from initial development to continual taste cell renewal.

In addition to *Ptch1* and *Gli1*, *Gli2* is expressed by basal keratinocytes adjacent to taste buds (a pattern similar to that of *Ptch1* and *Gli1*), as well as more broadly throughout the non-taste epithelium (Liu et al., 2013), suggesting that *Gli2* may play a role in taste cell renewal and underlie in part the ability of SHH to induce ectopic taste buds. Indeed, activation of a hyperactive *Gli2* allele, which is oncogenic in skin (Grachtchouk et al., 2011), abolishes

fungiform taste buds in adult mice; however, it is unclear if this is due to oncogenic growth or to a more direct effect on lingual epithelial cell fate (Liu et al., 2013).

Finally, in addition to a role in lingual epithelium, SHH signals are received in the mesenchymal compartment of taste papillae of both embryos and adults (Hall et al., 1999; Liu et al., 2013; Miura et al., 2001). In adults, SHH-receiving cells in the mesenchyme have been proposed to comprise a niche for taste bud cell maintenance (Liu et al., 2013), an idea that was suggested initially based on expression of *Bmp4* in the taste papilla mesenchyme (Nguyen and Barlow, 2010). Likewise, in the embryonic tongue, in addition to signaling to adjacent epithelium, *Shh*⁺ taste placodes signal to the subjacent mesenchymal compartment. This epithelial-to-mesenchymal Shh signaling may function in the extensive morphogenesis of taste papillae, comparable to the role of *Shh* in development of other epithelial appendages, such as teeth, feather and hair follicles (Chuong et al., 2000; Pispas and Thesleff, 2003), although this remains to be tested. As we mentioned above, taste buds in salamander embryos develop independently of oral mesenchyme. Cultured epithelial explants devoid of mesodermal and neural crest-derived cells develop cell type-replete taste buds (Barlow and Northcutt, 1997), but it is important to note that in axolotls taste buds do not reside in papillae, but rather are embedded in the oral epithelium (Fährmann, 1967; Northcutt et al., 2000; Toyoshima et al., 1987). Lingual taste papillae thus appear to be primarily a mammalian innovation, which may have evolved to prevent taste bud desiccation or protect taste buds from abrasive foodstuffs. Thus we propose that the primary event in taste development is specification of taste bud precursors, and that papillary development is secondary. Specifically, in both amphibians and mammals, we hypothesize that taste bud precursors are specified by epithelium-intrinsic processes, while in mammals these taste bud precursors in turn organize adjacent epithelium and mesenchyme to build taste papillae around them (Thirumangalathu et al., 2009). It remains to be determined if the canonical Shh signaling pathway, in addition to its role in taste placode patterning, also guides papilla morphogenesis directly and/or indirectly.

In closing, the overall structure of the taste periphery, which is composed of multicellular taste buds that detect primary tastants in the oral cavity, is conserved in all vertebrates examined to date, except the hagfishes (Braun, 1996; Braun, 1998; Finger, 1997), whereas the anatomical, histological, cell biological and molecular details of taste buds in the oral cavity are quite varied (Barlow, 1999; Jiang et al., 2012; Liman, 2012; Northcutt, 2004). These differences in structure and function have evolved as adaptations to the different diets consumed, but also indicate that this sensory system is quite flexible over evolutionary time scales. In addition, the pattern and cellular makeup of taste buds have been shown to vary within mammalian species, such as humans and mice, and in mouse and zebrafish embryos taste bud pattern and cell complement are easily manipulated experimentally via drug or genetic perturbations. This raises the question of how impressionable the sense of taste is in utero, in terms of exposure to maternal diet and to maternal overall health: can the taste bud array be permanently altered anatomically and at the level of taste cell fate decisions by gestational experience? And how might these changes in the taste periphery impact taste function and dietary selections and ultimately the health of offspring?

Acknowledgements

Thanks to Fernando Giraldez for his good humored comments during the writing of this manuscript, and to members of the Giraldez, Pujades and Alsina labs at the PRBB for their gracious hosting of LB while on sabbatical at the PRBB, Barcelona, Spain.

Supported by DC012675 and DC012383 to LAB and DE021420 to ODK.

References

- Adams AE. Some effects of removal of endoderm from the mouth region of early *Amblystoma punctatum* embryos. *J. exp. Zool.* 1931; 58:147–163.
- Atema J. Structures and functions of the sense of taste in the catfish (*Ictalurus natalis*). *Brain Behav. Evol.* 1971; 4:273–294. [PubMed: 5118142]
- Banziger C, Soldini D, Schutt C, Zipperlen P, Hausmann G, Basler K. Wntless, a conserved membrane protein dedicated to the secretion of Wnt proteins from signaling cells. *Cell.* 2006; 125:509–522. [PubMed: 16678095]
- Barker N, Tan S, Clevers H. Lgr proteins in epithelial stem cell biology. *Development.* 2013; 140:2484–2494. [PubMed: 23715542]
- Barker N, van Es JH, Kuipers J, Kujala P, van den Born M, Cozijnsen M, Haegebarth A, Korving J, Begthel H, Peters PJ, Clevers H. Identification of stem cells in small intestine and colon by marker gene Lgr5. *Nature.* 2007; 449:1003–1007. [PubMed: 17934449]
- Barlow, LA. The Biology of Amphibian Taste. In: Heatwole, H.; Dawley, E., editors. *Amphibian Biology: Sensory Perception*. 1999. p. 743–782.
- Barlow, LA. Taste buds in ectoderm are induced by endoderm: Implications for mechanisms governing taste bud development. In: Olsson, L.; Jacobson, C-O., editors. *Regulatory Processes in Development: The Legacy of Sven Hörstadius; Proceedings of the Wenner-Gren International Symposium*; Portland Press. 2000. p. 185–190.
- Barlow LA. Specification of pharyngeal endoderm is dependent on early signals from axial mesoderm. *Development.* 2001; 128:4573–4583. [PubMed: 11714682]
- Barlow LA, Northcutt RG. Embryonic origin of amphibian taste buds. *Dev. Biol.* 1995; 169:273–285. [PubMed: 7750643]
- Barlow LA, Northcutt RG. Taste buds develop autonomously from endoderm without induction by cephalic neural crest or paraxial mesoderm. *Development.* 1997; 124:949–957. [PubMed: 9056771]
- Beauchamp GK, Mennella JA. Early flavor learning and its impact on later feeding behavior. *J Pediatr Gastroenterol Nutr.* 2009; 48(Suppl 1):S25–S30. [PubMed: 19214055]
- Beidler LM, Smallman RL. Renewal of cells within taste buds. *J. Cell Biol.* 1965; 27:263–272. [PubMed: 5884625]
- Beites CL, Hollenbeck PL, Kim J, Lovell-Badge R, Lander AD, Calof AL. Follistatin modulates a BMP autoregulatory loop to control the size and patterning of sensory domains in the developing tongue. *Development.* 2009; 136:2187–2197. [PubMed: 19474151]
- Berteretche MV, Dalix AM, d'Ornano AM, Bellisle F, Khayat D, Faurion A. Decreased taste sensitivity in cancer patients under chemotherapy. *Support Care Cancer.* 2004; 12:571–576. [PubMed: 15179562]
- Bitgood MJ, McMahon AP. Hedgehog and Bmp genes are coexpressed at many diverse sites of cell-cell interaction in the mouse embryo. *Dev. Biol.* 1995; 172:126–138. [PubMed: 7589793]
- Bradley RM, Stern IB. The development of the human taste bud during the foetal period. *J. Anat.* 1967; 101:743–752. [PubMed: 6059822]
- Braun CB. The sensory biology of the living jawless fishes: A phylogenetic assessment. *Brain Behav. Evol.* 1996; 48:262–276. [PubMed: 8932867]
- Braun CB. Schreiner organs: a new craniate chemosensory modality in hagfishes. *J Comp Neurol.* 1998; 392:135–163. [PubMed: 9512266]

- Castillo D, Seidel K, Salcedo E, Ahn C, de Sauvage FJ, Klein OD, Barlow LA. Induction of ectopic taste buds by SHH reveals the competency and plasticity of adult lingual epithelium. *Development*. 2014; 141:2993–3002. [PubMed: 24993944]
- Chaudhari N, Roper SD. The cell biology of taste. *J Cell Biol*. 2010; 190:285–296. [PubMed: 20696704]
- Chuong CM, Patel N, Lin J, Jung HS, Widelitz RB. Sonic hedgehog signaling pathway in vertebrate epithelial appendage morphogenesis: perspectives in development and evolution. *Cell Mol Life Sci*. 2000; 57:1672–1681. [PubMed: 11130174]
- Dyer JS, Rosenfeld CR. Metabolic imprinting by prenatal, perinatal, and postnatal overnutrition: a review. *Seminars in reproductive medicine*. 2011; 29:266–276. [PubMed: 21769766]
- Echelard Y, Epstein DJ, St-Jacques B, Shen L, Mohler J, McMahon JA, McMahon AP. Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in the regulation of CNS polarity. *Cell*. 1993; 75:1417–1430. [PubMed: 7916661]
- Erdogan S, Iwasaki S. Function-related morphological characteristics and specialized structures of the avian tongue. *Annals of anatomy = Anatomischer Anzeiger : official organ of the Anatomische Gesellschaft*. 2014; 196:75–87. [PubMed: 24219998]
- Fährmann W. Licht- und elektronenmikroskopische Untersuchungen an der Geschmacksknospe des neotenen Axolotls (*Siredon mexicanum* Shaw). *Zeit. Microsk. Anat. Forsch*. 1967; 77:117–152.
- Farbman AI. Electron microscope study of the developing taste bud in rat fungiform papilla. *Dev. Biol*. 1965; 11:110–135. [PubMed: 14300090]
- Farbman AI. Renewal of taste bud cells in rat circumvallate papillae. *Cell Tissue Kinet*. 1980; 13:349–357. [PubMed: 7428010]
- Feng P, Huang L, Wang H. Taste bud homeostasis in health, disease, and aging. *Chem Senses*. 2014; 39:3–16. [PubMed: 24287552]
- Finger TE. Evolution of taste and solitary chemoreceptor cell systems. *Brain Behav Evol*. 1997; 50:234–243. [PubMed: 9310198]
- Fischer ME, Cruickshanks KJ, Schubert CR, Pinto A, Klein R, Pankratz N, Pankow JS, Huang GH. Factors related to fungiform papillae density: the beaver dam offspring study. *Chem Senses*. 2013; 38:669–677. [PubMed: 23821729]
- Gao Y, Toska E, Denmon D, Roberts SG, Medler KF. WT1 regulates the development of the posterior taste field. *Development*. 2014; 141:2271–2278. [PubMed: 24803588]
- Grachtchouk M, Pero J, Yang SH, Ermilov AN, Michael LE, Wang A, Wilbert D, Patel RM, Ferris J, Diener J, Allen M, Lim S, Syu LJ, Verhaegen M, Dlugosz AA. Basal cell carcinomas in mice arise from hair follicle stem cells and multiple epithelial progenitor populations. *J Clin Invest*. 2011; 121:1768–1781. [PubMed: 21519145]
- Hall JM, Bell ML, Finger TE. Disruption of sonic hedgehog signaling alters growth and patterning of lingual taste papillae. *Dev Biol*. 2003; 255:263–277. [PubMed: 12648489]
- Hall JM, Hooper JE, Finger TE. Expression of *Sonic hedgehog Patched* and *Gli1* in developing taste papillae of the mouse. *J. Comp. Neurol*. 1999; 406:143–155. [PubMed: 10096602]
- Hamamichi R, Asano-Miyoshi M, Emori Y. Taste bud contains both short-lived and long-lived cell populations. *Neuroscience*. 2006; 141:2129–2138. [PubMed: 16843606]
- Harfe BD, Scherz PJ, Nissim S, Tian H, McMahon AP, Tabin CJ. Evidence for an expansion-based temporal Shh gradient in specifying vertebrate digit identities. *Cell*. 2004; 118:517–528. [PubMed: 15315763]
- Hisha H, Tanaka T, Kanno S, Tokuyama Y, Komai Y, Ohe S, Yanai H, Omachi T, Ueno H. Establishment of a novel lingual organoid culture system: generation of organoids having mature keratinized epithelium from adult epithelial stem cells. *Scientific reports*. 2013; 3:3224. [PubMed: 24232854]
- Hong JH, Omur-Ozbek P, Stanek BT, Dietrich AM, Duncan SE, Lee YW, Lesser G. Taste and odor abnormalities in cancer patients. *J Support Oncol*. 2009; 7:58–65. [PubMed: 19408458]
- Howard JM, Nuguid JM, Ngole D, Nguyen H. Tcf3 expression marks both stem and progenitor cells in multiple epithelia. *Development*. 2014; 141:3143–3152. [PubMed: 25038042]
- Hume WJ, Potten CS. The ordered columnar structure of mouse filiform papillae. *J Cell Sci*. 1976; 22:149–160. [PubMed: 977666]

- Iwata S, Yoshida R, Ninomiya Y. Taste transductions in taste receptor cells: basic tastes and moreover. *Current pharmaceutical design*. 2014; 20:2684–2692. [PubMed: 23886388]
- Iwatsuki K, Liu HX, Gronder A, Singer MA, Lane TF, Grosschedl R, Mistretta CM, Margolskee RF. Wnt signaling interacts with Shh to regulate taste papilla development. *Proc Natl Acad Sci U S A*. 2007; 104:2253–2258. [PubMed: 17284610]
- Jackowiak H. Scanning electron microscopy study of the lingual papillae in the European mole (*Talpa europea*, L., Talpidae). *Anatomia, histologia, embryologia*. 2006; 35:190–195.
- Jiang P, Josue J, Li X, Glaser D, Li W, Brand JG, Margolskee RF, Reed DR, Beauchamp GK. Major taste loss in carnivorous mammals. *Proc Natl Acad Sci U S A*. 2012; 109:4956–4961. [PubMed: 22411809]
- Johnston JB. The limit between the ectoderm and entoderm in the mouth, and the origin of taste buds in Amphibians. *Am. J. Anat.* 1910; 10:41–67.
- Jung HS, Oropeza V, Thesleff I. Shh, Bmp-2, Bmp-4 and Fgf-8 are associated with initiation and patterning of mouse tongue papillae. *Mech Dev*. 1999; 81:179–182. [PubMed: 10330496]
- Kageyama R, Ohtsuka T. The Notch-Hes pathway in mammalian neural development. *Cell Res*. 1999; 9:179–188. [PubMed: 10520600]
- Kageyama R, Ohtsuka T, Hatakeyama J, Ohsawa R. Roles of bHLH genes in neural stem cell differentiation. *Exp Cell Res*. 2005; 306:343–348. [PubMed: 15925590]
- Kapsimali M, Barlow LA. Developing a sense of taste. *Semin Cell Dev Biol*. 2013; 24:200–209. [PubMed: 23182899]
- Kapsimali M, Kaushik AL, Gibon G, Dirian L, Ernest S, Rosa FM. Fgf signaling controls pharyngeal taste bud formation through miR-200 and Delta-Notch activity. *Development*. 2011; 138:3473–3484. [PubMed: 21791527]
- Kaufman, MH. *The Atlas of Mouse Development*. London: Academic Press; 1999.
- Kim MR, Kusakabe Y, Miura H, Shindo Y, Ninomiya Y, Hino A. Regional expression patterns of taste receptors and gustducin in the mouse tongue. *Biochem Biophys Res Commun*. 2003; 312:500–506. [PubMed: 14637165]
- Kito-Shingaki A, Seta Y, Toyono T, Kataoka S, Kakinoki Y, Yanagawa Y, Toyoshima K. Expression of GAD67 and Dlx5 in the taste buds of mice genetically lacking Mash1. *Chem Senses*. 2014; 39:403–414. [PubMed: 24682237]
- Landacre FL. On the place of origin and method of distribution of taste buds in *Ameiurus melas*. *J. Comp. Neurol*. 1907; 17:1–66.
- Liman ER. Changing senses: chemosensory signaling and primate evolution. *Advances in experimental medicine and biology*. 2012; 739:206–217. [PubMed: 22399404]
- Liman ER, Zhang YV, Montell C. Peripheral coding of taste. *Neuron*. 2014; 81:984–1000. [PubMed: 24607224]
- Liu F, Thirumangalathu S, Gallant NM, Yang SH, Stoick-Cooper CL, Reddy ST, Andl T, Taketo MM, Dlugosz AA, Moon RT, Barlow LA, Millar SE. Wnt-beta-catenin signaling initiates taste papilla development. *Nat Genet*. 2007; 39:106–112. [PubMed: 17128274]
- Liu HX, Ermilov A, Grachtchouk M, Li L, Gumucio DL, Dlugosz AA, Mistretta CM. Multiple Shh signaling centers, participate in fungiform papilla and taste bud, formation and maintenance. *Dev Biol*. 2013; 382:82–97. [PubMed: 23916850]
- Liu HX, Grosse AS, Iwatsuki K, Mishina Y, Gumucio DL, Mistretta CM. Separate and distinctive roles for Wnt5a in tongue, lingual tissue and taste papilla development. *Dev Biol*. 2012; 361:39–56. [PubMed: 22024319]
- Liu HX, Henson BS, Zhou Y, D'Silva NJ, Mistretta CM. Fungiform papilla pattern: EGF regulates inter-papilla lingual epithelium and decreases papilla number by means of PI3K/Akt, MEK/ERK, and p38 MAPK signaling. *Dev Dyn*. 2008; 237:2378–2393. [PubMed: 18729215]
- Liu HX, Maccallum DK, Edwards C, Gaffield W, Mistretta CM. Sonic hedgehog exerts distinct, stage-specific effects on tongue and taste papilla development. *Dev Biol*. 2004; 276:280–300. [PubMed: 15581865]
- Luo X, Okubo T, Randell S, Hogan BL. Culture of endodermal stem/progenitor cells of the mouse tongue. *In Vitro Cell Dev Biol Anim*. 2009; 45:44–54. [PubMed: 18830772]

- Ma H, Yang R, Thomas SM, Kinnamon JC. Qualitative and quantitative differences between taste buds of the rat and mouse. *BMC Neurosci.* 2007; 8:5. [PubMed: 17207280]
- Matsumoto I, Ohmoto M, Narukawa M, Yoshihara Y, Abe K. Skn-1a (Pou2f3) specifies taste receptor cell lineage. *Nat Neurosci.* 2011; 14:685–687. [PubMed: 21572433]
- Mennella JA. Ontogeny of taste preferences: basic biology and implications for health. *Am J Clin Nutr.* 2014; 99:704S–711S. [PubMed: 24452237]
- Miller IJ Jr, Reedy FE Jr. Variations in human taste bud density and taste intensity perception. *Physiol Behav.* 1990; 47:1213–1219. [PubMed: 2395927]
- Miller IJ, Preslar AJ. Spatial distribution of rat fungiform papillae. *Anat. Rec.* 1975; 181:670–684.
- Mistretta, CM. Topographical and histological study of the developing rat tongue, palate and taste buds. In: Bosma, JF., editor. *Third Symposium on Oral Sensation and Perception. The Mouth of the Infant.* Charles C. Thomas; Springfield, IL: 1972. p. 163-187.
- Mistretta CM, Liu HX, Gaffield W, MacCallum DK. Cyclopamine and jervine in embryonic rat tongue cultures demonstrate a role for Shh signaling in taste papilla development and patterning: fungiform papillae double in number and form in novel locations in dorsal lingual epithelium. *Dev Biol.* 2003; 254:1–18. [PubMed: 12606278]
- Miura H, Barlow LA. Taste bud regeneration and the search for taste progenitor cells. *Arch Ital Biol.* 2010; 148:107–118. [PubMed: 20830973]
- Miura H, Kato H, Kusakabe Y, Tagami M, Miura-Ohnuma J, Ninomiya Y, Hino A. A strong nerve dependence of sonic hedgehog expression in basal cells in mouse taste bud and an autonomous transcriptional control of genes in differentiated taste cells. *Chem Senses.* 2004; 29:823–831. [PubMed: 15574818]
- Miura H, Kusakabe Y, Harada S. Cell lineage and differentiation in taste buds. *Arch Histol Cytol.* 2006; 69:209–225. [PubMed: 17287576]
- Miura H, Kusakabe Y, Sugiyama C, Kawamatsu M, Ninomiya Y, Motoyama J, Hino A. Shh and Ptc are associated with taste bud maintenance in the adult mouse. *Mech Dev.* 2001; 106:143–145. [PubMed: 11472844]
- Miura H, Scott JK, Harada S, Barlow LA. Sonic hedgehog-expressing basal cells are general post-mitotic precursors of functional taste receptor cells. *Dev Dyn.* 2014; 243:1286–1297. [PubMed: 24590958]
- Nakayama A, Miura H, Shindo Y, Kusakabe Y, Tomonari H, Harada S. Expression of the basal cell markers of taste buds in the anterior tongue and soft palate of the mouse embryo. *J Comp Neurol.* 2008; 509:211–224. [PubMed: 18465790]
- Ng A, Tan S, Singh G, Rizk P, Swathi Y, Tan TZ, Huang RY, Leushacke M, Barker N. Lgr5 marks stem/progenitor cells in ovary and tubal epithelia. *Nat Cell Biol.* 2014; 16:745–757. [PubMed: 24997521]
- Nguyen HM, Barlow LA. Differential expression of a BMP4 reporter allele in anterior fungiform versus posterior circumvallate taste buds of mice. *BMC Neurosci.* 2010; 11:129. [PubMed: 20942907]
- Nguyen HM, Reyland ME, Barlow LA. Mechanisms of taste bud cell loss after head and neck irradiation. *J Neurosci.* 2012; 32:3474–3484. [PubMed: 22399770]
- Northcutt RG. Taste buds: development and evolution. *Brain Behav Evol.* 2004; 64:198–206. [PubMed: 15353910]
- Northcutt RG. Taste bud development in the channel catfish. *J Comp Neurol.* 2005; 482:1–16. [PubMed: 15612020]
- Northcutt RG, Barlow LA, Braun CB, Catania KC. Distribution and innervation of taste buds in the axolotl. *Brain Behav. Evol.* 2000; 56:123–145. [PubMed: 11124515]
- Ohtsuka T, Ishibashi M, Gradwohl G, Nakanishi S, Guillemot F, Kageyama R. Hes1 and Hes5 as notch effectors in mammalian neuronal differentiation. *Embo J.* 1999; 18:2196–2207. [PubMed: 10205173]
- Ohtubo Y, Yoshii K. Quantitative analysis of taste bud cell numbers in fungiform and soft palate taste buds of mice. *Brain Res.* 2010; 1367:13–21. [PubMed: 20971092]
- Okubo T, Clark C, Hogan BL. Cell lineage mapping of taste bud cells and keratinocytes in the mouse tongue and soft palate. *Stem Cells.* 2009; 27:442–450. [PubMed: 19038788]

- Okubo T, Pevny LH, Hogan BL. Sox2 is required for development of taste bud sensory cells. *Genes Dev.* 2006; 20:2654–2659. [PubMed: 17015430]
- Ota MS, Kaneko Y, Kondo K, Ogishima S, Tanaka H, Eto K, Kondo T. Combined in silico and in vivo analyses reveal role of Hes1 in taste cell differentiation. *PLoS Genet.* 2009; 5:e1000443. [PubMed: 19343206]
- Parker MA, Bell M, Barlow LA. Cell contact-dependent mechanisms specify taste bud number and size during a critical period early in embryonic development. *Dev Dyn.* 2004; 230:630–642. [PubMed: 15254897]
- Passilly-Degrace P, Chevrot M, Bernard A, Ancel D, Martin C, Besnard P. Is the taste of fat regulated? *Biochimie.* 2014; 96:3–7. [PubMed: 23933093]
- Perea-Martinez I, Nagai T, Chaudhari N. Functional cell types in taste buds have distinct longevities. *PLoS One.* 2013; 8:e53399. [PubMed: 23320081]
- Petersen CI, Jheon AH, Mostowfi P, Charles C, Ching S, Thirumangalathu S, Barlow LA, Klein OD. FGF signaling regulates the number of posterior taste papillae by controlling progenitor field size. *PLoS Genet.* 2011; 7:e1002098. [PubMed: 21655085]
- Pispa J, Thesleff I. Mechanisms of ectodermal organogenesis. *Dev. Biol.* 2003; 262:195–362. [PubMed: 14550785]
- Plaks V, Brenot A, Lawson DA, Linnemann JR, Van Kappel EC, Wong KC, de Sauvage F, Klein OD, Werb Z. Lgr5-expressing cells are sufficient and necessary for postnatal mammary gland organogenesis. *Cell reports.* 2013; 3:70–78. [PubMed: 23352663]
- Reiner DJ, Jan TA, Boughter JD Jr, Li CX, Lu L, Williams RW, Waters RS. Genetic analysis of tongue size and taste papillae number and size in recombinant inbred strains of mice. *Chem Senses.* 2008; 33:693–707. [PubMed: 18653645]
- Rothova M, Thompson H, Lickert H, Tucker AS. Lineage tracing of the endoderm during oral development. *Dev Dyn.* 2012; 241:1183–1191. [PubMed: 22581563]
- Ruo Redda MG, Allis S. Radiotherapy-induced taste impairment. *Cancer Treat Rev.* 2006; 32:541–547. [PubMed: 16887272]
- Seta Y, Oda M, Kataoka S, Toyono T, Toyoshima K. Mash1 is required for the differentiation of AADC-positive type III cells in mouse taste buds. *Dev Dyn.* 2011; 240:775–784. [PubMed: 21322090]
- Seta Y, Seta C, Barlow LA. Notch-associated gene expression in embryonic and adult taste papillae and taste buds suggests a role in taste cell lineage decisions. *J Comp Neurol.* 2003; 464:49–61. [PubMed: 12866128]
- Stewart JE, Feinle-Bisset C, Golding M, Delahunty C, Clifton PM, Keast RS. Oral sensitivity to fatty acids, food consumption and BMI in human subjects. *Br J Nutr.* 2010; 104:145–152. [PubMed: 20196892]
- Stewart JE, Feinle-Bisset C, Keast RS. Fatty acid detection during food consumption and digestion: Associations with ingestive behavior and obesity. *Prog Lipid Res.* 2011; 50:225–233. [PubMed: 21356242]
- Takeda N, Jain R, Li D, Li L, Lu MM, Epstein JA. Identifies Progenitor Cells Capable of Taste Bud Regeneration after Injury. *PLoS One.* 2013; 8:e66314. [PubMed: 23824276]
- Tanaka T, Komai Y, Tokuyama Y, Yanai H, Ohe S, Okazaki K, Ueno H. Identification of stem cells that maintain and regenerate lingual keratinized epithelial cells. *Nat Cell Biol.* 2013; 15:511–518. [PubMed: 23563490]
- Thirumangalathu S, Harlow DE, Driskell AL, Krimm RF, Barlow LA. Fate mapping of mammalian embryonic taste bud progenitors. *Development.* 2009; 136:1519–1528. [PubMed: 19363153]
- Tizzano M, Dvoryanchikov G, Barrows JK, Kim S, Chaudhari N, Finger TE. Expression of Galpha14 in sweet-transducing taste cells of the posterior tongue. *BMC Neurosci.* 2008; 9:110. [PubMed: 19014514]
- Tordoff MG, Alarcon LK, Lawler MP. Preferences of 14 rat strains for 17 taste compounds. *Physiol Behav.* 2008a; 95:308–332. [PubMed: 18639567]
- Tordoff MG, Reed DR, Shao H. Calcium taste preferences: genetic analysis and genome screen of C57BL/6J×PWK/PhJ hybrid mice. *Genes, brain, and behavior.* 2008b; 7:618–628.

- Toyoshima K, Miyamoto K, Shimamura A. Fine structure of taste buds in the tongue, palatal mucosa and gill arch of the axolotl, *Ambystoma mexicanum*. *Okajimas Folia Anat. Jpn.* 1987; 64:99–110. [PubMed: 3431793]
- Tucker RM, Mattes RD, Running CA. Mechanisms and effects of "fat taste" in humans. *Biofactors.* 2014; 40:313–326. [PubMed: 24591077]
- Vissink A, Jansma J, Spijkervet FK, Burlage FR, Coppes RP. Oral sequelae of head and neck radiotherapy. *Crit Rev Oral Biol Med.* 2003; 14:199–212. [PubMed: 12799323]
- Wasalathanthri S, Hettiarachchi P, Prathapan S. Sweet taste sensitivity in pre-diabetics, diabetics and normoglycemic controls: a comparative cross sectional study. *BMC endocrine disorders.* 2014; 14:67. [PubMed: 25123551]
- Witt M, Reutter K. Embryonic and early fetal development of human taste buds: a transmission electron microscopical study. *Anat. Rec.* 1996; 246:507–523. [PubMed: 8955790]
- Witt M, Reutter K. Innervation of developing human taste buds. An immunohistochemical study. *Histochem Cell Biol.* 1998; 109:281–291. [PubMed: 9541477]
- Yee KK, Li Y, Redding KM, Iwatsuki K, Margolskee RF, Jiang P. Lgr5-EGFP Marks Taste Bud Stem/Progenitor Cells in Posterior Tongue. *Stem Cells.* 2013; 31:992–1000. [PubMed: 23377989]
- Zhou Y, Liu HX, Mistretta CM. Bone morphogenetic proteins and noggin: Inhibiting and inducing fungiform taste papilla development. *Dev Biol.* 2006; 297:198–213. [PubMed: 16828469]
- Zhu X, Liu Y, Zhao P, Dai Z, Yang X, Li Y, Qiu M, Zhang Z. Gpr177-mediated Wnt Signaling is Required for Fungiform Placode Initiation. *J Dent Res.* 2014; 93:582–588. [PubMed: 24736288]

Figure 1.

Taste buds comprise 3 morphological cell types, I, II and III, which are recognized by their expression of specific marker proteins. A. Type I cells express NTPdase2, which localizes to cell membranes that tightly wrap other cells within the bud, such that individual NTPdase2+ cells are not discernable (see Miura et al., 2014 for detailed explanation). B. Type II cells express PLC β 2 in the cytosol, and the protein is excluded from nuclei of PLC β 2+ taste cells (*). C. NCAM marks the surface of Type III taste cells (*), as well as some nerve fibers

extending into buds (arrows). All panels are images of taste buds from the circumvallate papilla of mice. Apical taste pores are indicated with arrowheads. Scale bar is 20 μ m.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2.

Schematic diagram of taste bud cell types and lineage relationships. Type I (yellow), II (blue) and III (red) taste cells are elongate, postmitotic (open nuclei) cells, which arise from proliferating, K5+/K14+ keratinocytes (green, mitotic nuclei) outside of taste buds. These progenitors also give rise to non-taste epithelial cells (orange). Following their final division, cells destined to become taste cells enter taste buds and become taste bud precursors, which are *Shh*+ ovoid cells (lavender) located in the basal compartment of the taste bud. *Shh*+ cells are postmitotic, and differentiate into each of the 3 taste cell types. Black arrows indicate observed lineage relationships among cell types (see text for details), but are not absolute, nor exhaustive.

Table 1

Summary of the function of major signaling pathways in embryonic taste bud development.

Pathway	Demonstrated functions	References
Wnt/β-catenin	- Promotes taste fate in vivo and in vitro - Required for taste fate in vivo	(Iwatsuki et al., 2007) (Liu et al., 2007)
BMP	- Prior to taste placode specification, BMPs promote taste fate - Following placode specification, BMPs repress taste fate in vitro - Loss of mesenchymal follistatin promotes taste fate in vivo	(Zhou et al., 2006) (Beites et al., 2009)
SHH	- Shh represses taste fate in vitro - Inhibition of Shh in vitro expands taste fate	(Iwatsuki et al., 2007) (Hall et al., 2003) (Mistretta et al., 2003)
FGF	- Loss of Spry1/2 promotes taste fate - Loss of mesenchymal FGF10 represses taste fate	(Petersen et al., 2011)
EGF	- EGF represses taste fate in vitro - Inhibition of EGF promotes taste fate in vitro	(Liu et al., 2008)