Poster presentation

Open Access

Predicting spike activity in neuronal cultures

Tayfun Gürel^{*1,2}, Ulrich Eger^{1,3}, Steffen Kandler^{1,3}, Luc De Raedt^{1,5} and Stefan Rotter^{1,4}

Address: ¹Bernstein Center for Computational Neuroscience (BCCN), Albert-Ludwigs-University Freiburg, Germany, ²Machine Learning Lab, Institute for Computer Science, Albert-Ludwigs-University Freiburg, Germany, ³Institute for Biology 3, Albert-Ludwigs-University Freiburg, Germany, ⁴Institute for Frontier Areas of Psychology and Mental Health, Freiburg, Germany and ⁵Department of Computer Science, Katholieke Universiteit Leuven, Belgium

Email: Tayfun Gürel^{*} - guerel@informatik.uni-freiburg.de

^{*} Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, **8**(Suppl 2):P185 doi:10.1186/1471-2202-8-S2-P185

© 2007 Gürel et al; licensee BioMed Central Ltd.

Discovering the functional connectivity and modeling the dynamics of neuronal networks is essential to understand neural information processing. Here we focus on neuronal cultures of neocortical tissue, which are closed system in vitro neural networks. Recordings of spontaneous activity from neuronal cultures using multi-electrode array (MEA) technology have revealed that the activity is composed of irregular network-wide bursts of spikes, even in the absence of any external stimulation [1]. Although it is reasonable to think of 'spontaneous' fluctuations which start a burst in these cultures, the spatio-temporal spread of activity is nevertheless generated and shaped by the underlying network. It is then an interesting problem to characterize the underlying synaptic connectivity based on activity measurements. This knowledge will help us understanding how a given anatomical structure generates different activity patterns, and hence would be a significant step towards understanding structure-function relations in the neural networks of the brain.

Activity dynamics in neuronal cultures display both non-linear and non-stationary characteristics. Noise is another innate property of those networks causing high variability of the activity. The combination of these properties suggests the use of automated adaptive methods, (i.e. machine learning algorithms) to infer appropriate models of the activity dynamics. Specifically, we propose an algorithm to learn a predictive computational model of spon-

aneous activity in neuronal cultures. The learned model may also be regarded as an abstraction of the underlying effective network connectivity, i.e. its functional connectivity. Although similar functional connectivity models have been described previously [2,3], we take a steepest descent approach to learn functional connectivity, which naturally allows for online learning and, hence, is able to capture network plasticity, i.e. changes in the structure. We use the log-likelihood of point processes as a criterion for optimization. This approach has previously been suggested to analyse neural receptive field plasticity [4]. Here we apply it to multi-channel recordings from neuronal cultures, and demonstrate its use for learning functional connectivity and predicting upcoming spike activity. A ROC curve analysis of our experiments shows that this online approach predicts upcoming spike activity very well.

Acknowledgements

This work was funded by the German Federal Ministry of Education and Research (BMBF grant 01GQ0420 to BCCN Freiburg).

References

1. Marom S, Shahaf G: **Development, learning and memory in large random networks of cortical neurons: lessons beyond anatomy.** *Q Rev Biophys* 2002, **35**(1):63-87.
2. Chornoboy ES, Schramm LP, Karr A: **Maximum likelihood identification of neural point process systems.** *Biol Cybern* 1988, **59**:265-275.

3. Okatan M, Wilson MA, Brown EN: **Analyzing functional connectivity using a network likelihood model of ensemble neural spiking activity.** *Neural Comput* 2005, **17**:1927-1961.
4. Brown EN, Ngyuen DP, Wilson LM, Solo V: **An analysis of neural receptive field plasticity by point process adaptive filtering.** *Proc Natl Acad Sci USA* 2001, **98**:12261-12266.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp