Published in final edited form as:

J Adolesc Health. 2014 December ; 55(6 0): S6–S12. doi:10.1016/j.jadohealth.2014.07.021.

Respondent-driven sampling for an adolescent health study in vulnerable urban settings: a multi-country study

Michele R. Decker, ScD¹, Beth Marshall, DrPH¹, Mark Emerson, BS¹, Amanda Kalamar¹, Laura Covarrubias, MSPH¹, Nan Astone, PhD², Ziliang Wang, MS³, Ersheng Gao, MD, MPH³, Lawrence Mashimbye, MSc⁴, Sinead Delany-Moretlwe, MD, PhD⁴, Rajib Acharya, PhD⁵, Adesola Olumide, MPH⁶, Oladosu Ojengbede⁶, and Robert Blum, MD, PhD, MPH¹

¹Johns Hopkins Bloomberg School of Public Health, Department of Population, Family and Reproductive Health, Baltimore, MD, USA

²Urban Institute, Washington, DC, USA

³Department of Epidemiology and Social Science Research on Reproductive Health, Shanghai Institute of Planned Parenthood Research

⁴Wits Reproductive Health & HIV Institute, School of Clinical Medicine, University of the Witwatersrand, Johannesburg

⁵Population Council, New Delhi, India

⁶Institute of Child Health, College of Medicine, University of Ibadan/University College Hospital Ibadan

Abstract

The global adolescent population is larger than ever before and is rapidly urbanizing. Global surveillance systems to monitor youth health typically use household- and school-based recruitment methods. These systems risk not reaching the most marginalized youth made vulnerable by conditions of migration, civil conflict and other forms of individual and structural vulnerability. We describe the methodology of the Well Being of Adolescents in Vulnerable Environments (WAVE) survey, which used respondent-driven sampling (RDS) to recruit male and female youth aged 15 to 19 years and living in economically distressed urban settings in Baltimore, USA, Johannesburg, South Africa, Ibadan, Nigeria, Delhi, India and Shanghai, China (migrant youth only) for a cross-sectional study. We describe a shared recruitment and survey administration protocol across the five sites, present recruitment parameters, and illustrate challenges and necessary adaptations for use of RDS with youth in disadvantaged urban settings. We describe the reach of RDS into populations of youth who may be missed by traditional household-based and school-based sampling. Across all sites, an estimated 9.6% were unstably

© 2014 Published by Society of Adolescent Health and Medicine.

Corresponding Author: Michele R. Decker, ScD, Assistant Professor, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe St.; E4142, Baltimore, MD, 21205, USA, +1 410-502-2747, mdecker@jhsph.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

housed; among those enrolled in school, absenteeism was pervasive with 29% having missed over 6 days of school in the past month. Overall findings confirm the feasibility, efficiency and utility of RDS in quickly reaching diverse samples of youth, including those both in and out of school and those unstably housed, and provide direction for optimizing RDS methods with this population. In our rapidly urbanizing global landscape with an unprecedented youth population, RDS may serve as a valuable tool in complementing existing household- and school-based methods for health-related surveillance that can guide policy.

Keywords

adolescent health; respondent-driven sampling; gender differences

Introduction

Globally, The current population of youth ages 10–24 is larger than ever before in history, with an estimated one in four individuals worldwide within this age group; the vast majority live in low and middle income nations.¹ While adolescents are considered a healthy population, youth suffer morbidity and mortality stemming from intentional and unintentional injury, and sexual and reproductive health issues spanning sexually transmitted infections (STI) including HIV, and unintended pregnancy.^{1,2} Health risk behavior in the forms of tobacco, alcohol and other substance use, as well as lack of physical activity, can initiate and intensify during this life stage and impart risk for subsequent chronic health conditions.^{1,2}

The 2012 global synthesis of adolescent health² illustrates that much of our knowledge of population-level adolescent health is derived from household-based and school-based surveillance systems, including the household-based Demographic and Health Survey (DHS) and Multiple Indicator Cluster Survey (MICS). Both systems include adolescents ages 15 and over. the international Health Behaviour in School-Aged Children (HBSC) and the Global School Health Survey (GSHS) include younger adolescents up to age 15. Despite the insights and international comparisons enabled by these systems,^{3–9} significant gaps remain in our understanding of global adolescent health. The school-based sampling strategies of the HBSC and GSHS, while efficient, risk biased estimates in settings with chronic school absenteeism and high drop-out rates. These biases are particularly salient for women and girls, who are less likely to be in school in some settings.^{2,10} The household-based sampling of DHS and MICS risks overlooking The most vulnerable youth who risk housing instability for reasons of poverty, institutionalization, migration, civil conflict and other forms of individual and structural instability. The health of youth who risk being missed by these surveillance systems is likely compromised. Experiences of housing instability, food insecurity, poverty and gender inequity constitute powerful social determinants of health.¹¹ Adequate housing protects urban children from disease, injuries and accidents; living environments are closely linked with psychological stress and overall health.¹²

The confluence of heightened risk and limited access to health services for youth who are out of school and unstably housed, coupled with the lack of globally comparable data on health outcomes and determinants using samples that adequately include these populations, has prompted calls for the exploration of respondent-driven sampling (RDS)^{2,13} to reach a broader spectrum of youth, much in the way that it has been applied to other marginalized populations often made invisible through more traditional sampling methods. RDS¹⁴ is a chain-based recruitment method that has been widely adopted for HIV-related research¹⁵ with marginalized, hidden populations including injection drug users (IDU), men who have sex with men (MSM), and female sex workers (FSWs).¹⁶ More recently, the method has been adopted for other hidden populations, such as migrant workers,¹⁷ street children and homeless youth.^{20,21}

We undertook a comparative investigation of adolescent health spanning domains of substance use, sexual and reproductive health, mental health, and gender-based violence, in distressed urban setting in five countries using RDS. Here, We describe recruitment in these five geo-cultural settings, including challenges and solutions, and illustrate the extent to which RDS reaches into populations of youth who risk exclusion from household-based and school-based surveillance systems.

Methods

In 2013, the cross-sectional Well Being of Adolescents in Vulnerable Environments (WAVE) study surveyed youth aged 15 to 19 years living in economically distressed urban settings in Baltimore, USA; Johannesburg, South Africa; Ibadan, Nigeria; Delhi, India; and Shanghai, China (migrant youth only, as they constitute a particularly vulnerable population in this setting). RDS was selected for this study population given feasibility concerns for household based sampling or school based sampling, e.g., inaccessible households (e.g., locked apartment buildings in Johannesburg) and the many migrant youth not enrolled in school (Shanghai). A shared protocol enabled cross-site comparability. At each site, an extensive formative phase was conducted, including key-informant interviews, community mapping, focus groups and in-depth interviews.²⁰ In accordance with recommendations for formative research prior to RDS implementation,^{21,22} this phase informed RDS acceptability and logistics; we assessed evaluated youths network properties including subgroupings and the level of networking within and across subgroupings, identified necessary seed characteristics and potential seeds, and refined survey domains, consistent with formative RDS recommendations.

Eligible seeds and participants were adolescents aged 15–19 and residing in the geographically-designated target communities at each site. Adolescent seed participants were purposefully selected to serve as the initial contacts for recruiting from the target population through the formative phase, and through existing relationships with youth, community advisory boards, and other service providers at each site. Seeds represented diverse school enrollment, gender, employment, and residential statuses. Consistent with RDS methods,¹⁵ seeds and subsequent recruits were provided with up to three recruitment coupons each to recruit additional adolescents into the study until recruitment goals were reached. Each coupon had an expiration date, after which it could not be redeemed. Coupon

expiration dates were used to control recruitment pace and to end recruitment when the sample size was achieved. Controlling chain referral recruitment through restricting the number of referrals per person encourages long recruitment chains and reach into the target population. Coupons were identifiable by sequential numbers which linked recruits to their recruiters, enabling creation of recruitment chains. Several steps ensured the tracing of these links, including duplicative documentation of coupon links in both hard copy recruitment logs as well as electronically collected survey data. Procedures were piloted extensively in the Baltimore, MD site, with iterative discussion of pilot findings and challenges with partners at all sites. All other sites were able to transition directly into data collection from the pilot phase once logistics were finalized. Many made minor procedural adaptations through the pilot to maximize participant comfort, minimize coupon tampering, and ensure capacity to link participants with recruiters via coupon number documentation.

Consistent with RDS methods, participants self-reported the size of their social network to account for potential bias due to differences in selection probability for participants with larger versus smaller networks. To improve accuracy,²¹ network size questions were asked sequentially and structured to ensure reciprocity in social ties. The sequence was: how many youth [age and location eligibility]...do you know, ...do you know that know you, ... do you know that know you and that you have seen in the past two weeks, ... do you know that know you and that you have seen and talked to in the past two weeks, with the final question serving as the participant's network size.

When a seed or recruit presented for data collection, staff assessed eligibility, obtained informed consent including parental consent for youth under age 18, and oriented participants to survey procedures. To maximize confidentiality and minimize bias, the anonymous survey was self-administered directly in audio computer-assisted self interview instrument (ACASI), which has been demonstrated to enhance accuracy in reporting on sensitive topics among many populations.²³, and can overcome literacy issues through the audio component. On occasion, participants requested research staff assistance. Youth aged 17 years and younger were required to obtain parent/guardian consent to participate. All study procedures were approved by Institutional Review Boards at Johns Hopkins Bloomberg School of Public Health and at each participating site.

The survey

The current manuscript describes recruitment parameters following RDS reporting recommendations,²⁴ included domains of physical, mental, and sexual and reproductive health, violence, family, community as well as challenges and school environments, and demographics. It was developed in English, professionally translated and back-translated into local languages at each site, and piloted with native speakers to ensure comprehension. Discrepancies were resolved through an iterative process. Here we present data on housing instability and school enrollment. housing instability was defined as reporting not having a regular place to stay, OR having a regular place to stay but staying there on average, four or less nights per week in the past 30 days, and having stayed in four or more places in the past week. School enrollment was assessed via a single item, "Are you currently in school?" Among participants in school, past month absenteeism was assessed via a single item,

“About how many days of school did you miss in the last 30 days, even if it was just part of a day?”; responses were categorized for analysis.

The current manuscript describes recruitment parameters following RDS reporting recommendations. We describe challenges and adaptations across settings based on our monthly research team meetings, as well as a brief survey on challenges and adaptations completed by research staff in each site following data collection. We also present the prevalence of housing instability and school absenteeism within our sample, to approximate our reach into populations of youth who may be missed by household or school-based surveys. Prevalence estimates for each outcome were generated by site in STATA both overall and by gender; school enrollment estimates were stratified by age group. Analysis procedures accounted for the non-independence of observations using complex SVY procedures and were weighted using the RDSII estimator²⁵. After identifying differences in age distribution across sites, we developed a post-stratification age weight which is used in conjunction with the RDSII weight.

Results

Recruitment parameters

Recruitment parameters are presented in Table 1. A total of 44 (range 5–14) seeds were enrolled across the sites, and 5175 (range 657–1347) recruitment coupons distributed. Among returned coupons (2391 total), some (39) were ineligible or opted not to participate (3). We enrolled a total of 2393 participants across sites including seeds (range 455–500). The sample was reduced to 2339 (range 438–500 per site) after identifying a small number of duplicate, non-complete and otherwise inaccurate surveys. Overall data quality was high, with the majority of participants in each site providing complete data.

Implementation challenges and solutions

Several challenges arose in implementing RDS with youth; Table 2 summarizes challenges and adaptations.

Pace, ineligible individuals, and coupon misuse—One of the most consistent challenges was the slow start of recruitment, followed by the rapid pace of RDS once implemented. Figures 1-5 illustrate recruitment pace by gender in each site. Baltimore and Johannesburg in particular experienced a slow start to recruitment, followed by rapid redemption of coupons and participant enrollment. Sites adapted to the pace by shortening the expiration time frame for coupons in the beginning of data collection to quicken the pace, and then by lengthening the expiration time to slow the pace of data collection. Coupon expiration time frames ranged from 3 – 14 days with most sites settling on 7 days for the bulk of their data collection.

The other consistent challenge was young people trying to gain access to the study outside of the eligibility requirements. Most sites experienced young people suspected of being over or under the age eligibility trying to enroll. In turn, prospective participants in Delhi and Shanghai were required to produce identification and/or proof of age. In the other sites where documented proof of age was less common, young people were asked to provide their

birth date, name of/year in school or even contact information for a parent or guardian who could confirm age. Site staff also stressed that the secondary incentives were contingent on recruiting eligible participants. Research staff in Shanghai, Baltimore, and Johannesburg noticed instances of participants selling their own coupons, or collecting coupons from other participants to recruit more respondents. In Johannesburg, the coupon was revised to state “NOT FOR SALE” and in Baltimore and Shanghai, staff stressed that coupons were to be distributed only by the participant. In cases of suspected misuse, coupons were cancelled or not distributed.

Baltimore, Shanghai and Johannesburg each had at least one recruitment chain that grew at a much more rapid pace than the others, and at least one seed that was not productive. These sites addressed the nonproductive seeds by recruiting additional seeds. In Baltimore, outreach to non-productive seeds was also attempted to encourage coupon distribution. Shanghai and Baltimore handled their large, fast-growing recruitment chains differently. In Shanghai, the research team established a waiting period before the coupons were active. In Baltimore, the team limited the number of coupons distributed to respondents in the fast-growing chain while exploring potential misuse, and ultimately identified stopped distributing coupons to respondents in one branch after confirming coupon misuse.

In Baltimore, Johannesburg, and Shanghai, research staff noted a handful of participants who attempted to complete the survey twice. In Baltimore, a core research staff was familiar with participant faces and able to stop repeat enrollees, and in one case identifying a repeat enrollee who made it through screening so that their second round of data could be dropped from the data set. In Shanghai, similarly, staffing consistency enabled participant familiarity that deflected any attempted repeat efforts. In Johannesburg, the team implemented a computer-based fingerprint biometric system to record all participants, which allowed them to identify repeat attempts before they were consented.

staff in Baltimore, Ibadan, and Shanghai noticed a handful of participants not answering the survey questions seriously and attributed this to participants who were rushing through the survey to join their friends who often were waiting onsite or just outside the interview venue. All sites intervened with respondents who were observed to be mindlessly clicking through the survey, who were informed that they would not receive recruitment coupons if they appeared to complete the survey haphazardly. In Baltimore and Shanghai, the researchers did not allow youth to recruit on the same day that they received coupons. The Baltimore team also discouraged participants who had already taken the survey from waiting for their friends. In Shanghai, an appointment system was established to minimize waiting times. Analytically, a protocol for identifying incomplete and nonsensical data was developed to identify and eliminate poor responders.

Differential recruitment pace by gender—In Each site, recruitment fluctuated on a daily basis. Recruitment was faster for male participants than for female participants. Even with female seeds, the recruitment of male participants outpaced the recruitment of female participants. Near the end of data collection, some sites restricted coupons by gender in an effort to achieve a more gender-balanced sample. For example in Baltimore, towards the end of data collection as coupons were distributed, participants were informed that they could

only recruit young women and in Johannesburg, respondents were allowed to recruit only one young man with their three coupons. In Shanghai and Ibadan, recruitment moved at a faster pace for those enrolled in school compared to those not enrolled. To slow the pace of student enrollment, coupons distributed to students had a seven day waiting period before they were active.

Reach into unstably housed and schooled populations of youth

Housing instability in the current sample ranged widely from just under 2% to nearly 17% (Table 3). In total across all sites an estimated 9.61% were unstably housed. School enrollment naturally varied by age in all sites. In Baltimore, nearly all 15 year olds were enrolled in school while more than 40% of 18 year olds were not in school. In Shanghai, where the sample of adolescents was migrant adolescents, nearly 30% of those age 15 were not in school and 97% were not in school by age 19. Among those enrolled, school absenteeism was pervasive; in total across sites 29.4% had missed six or more days of school in the past month, ranging from less than 1% in Shanghai to upwards of 64% in Ibadan.

Discussion

Our study represents the first attempt at a multi-country RDS survey on adolescent health in distressed urban environments. It was successful in achieving recruitment goals and ensuring consistent procedures across sites. Lessons learned provide direction for optimizing RDS procedures with similar populations.

RDS recruitment enabled rapid recruitment of a diverse sample of youth in each of the five study sites. Our implementation challenges, including ensuring eligibility criteria, coupon tampering and misuse and pacing, were strikingly similar to those identified across a variety of other settings and populations. For example, entrepreneurial individuals seeking to evade eligibility criteria or attempting duplicate enrollment to receive more incentives has been previously noted, as have research “hustlers” who sell, duplicate, or otherwise attempt to misuse coupons.^{21,26,27} It is possible that our target population of adolescents in distressed environments may be uniquely susceptible to some of these issues due to the disadvantage communities in which they reside.

gender differences in recruitment patterns were notable, with women recruiting more slowly than their male counterparts, and young men recruiting men to a greater extent than women, suggesting that their networks are more accessible. This created challenges in gender balance across the sample and concern for small cell sizes for the women’s sample. By their nature, past RDS studies with MSM and FSWs are gender-stratified and thus do not allow observations on potential gender differences in recruitment pace. The consistency of our findings on gender recruitment pace across sites is striking, and strongly suggests gender as a critical dimension in RDS recruitment. Gender-stratification of coupons could mitigate this issue in future mixed-gender RDS research.

evidence of inclusion of unstably housed (nearly 10%) and out of school youth (12–38%) suggests that RDS enabled the study to reach into adolescent populations that may have

been missed by traditional household- or school-based survey methods. For example, in household surveys, a respondent to the survey is selected after his/her household is selected, usually from a complete listing of households in the survey target area. Using this sampling strategy, the household is the primary sampling unit and thus any respondent who is not stably housed might be missed, biasing the results of the survey. In school-based surveys, respondents are usually selected to participate from a listing of all students in a school or a particular classroom. With this sampling strategy, any potential respondent must be enrolled in school and be present on the day of interview (not absent and regularly attending school). Any adolescent who has dropped out of school, has graduated, or is frequently truant risks being missed by school-based sampling. RDS can offer a means of overcoming these limitations; in turn, it can complement the knowledge gleaned from school- and household-based surveillance systems for youth, particularly where concerns for housing instability and out-of-school youth are pervasive.

Housing instability and missed school estimates should be considered with some caution, despite measurement piloting. In Delhi sample, the 17% housing instability estimate was felt to be unrealistically high. Cultural differences were felt to have prompted misinterpretation in the assessment, particularly as it relates to what constitutes “home”. For example, child of migrants may not have considered Delhi as a permanent home and in turn responded having lived away from home. Youth who co-reside with their parents are strictly supervised such that the very concept of staying away from their parents was alien to them. In addition, due to delays, some data collection overlapped with holiday breaks, particularly in Ibadan. This introduces potential inaccuracies in housing instability and missed school days, where participants may have included school days missed due to being on vacation, and/or housing instability due to traveling during this time.

While findings indicate that RDS enabled inclusion of respondents who risk being missed in school based survey or a household survey, a more rigorous evaluation of reach into these populations would require a comparative design. The broader generalizability of our sample is unknown. It is possible that unknown barriers restricted participation and potentially created bias. The advantages of data collection via ACASI may vary across contexts,^{23,28–30} which is a challenge for its use in a protocol shared across geo-cultural settings. Several limitations of RDS also warrant consideration in interpreting the current results and in considering the value of RDS for youth in distressed urban settings. Some evidence indicates that RDS sampling inference methods can fail to mitigate bias - with youth in particular under-represented. This unmitigated bias can emerge where specific sub-populations are over- or under-recruited by all groups, rather than specific groups, and where under-represented groups have network sizes comparable to those of their non-under-represented peers. These and other evaluations of RDS highlight the need to optimize the inferences that can be made using this method, and underscore the value of formative research before implementing RDS. Many of the ethical issues associated with RDS are unique to marginalized populations more so than the method itself. One exception is that of monetary incentives; where RDS is used in HIV research, obtaining test results is a dominant participation motivation. By contrast, participation motivations remain unclear for studies that lack additional benefits, such as ours, highlighting the need to ensure that monetary incentives are sufficient, but not so high as to coerce participation. Finally

geospatial elements can influence RDS recruitment; for example residential location and proximity to study site relates to recruitment probability in some settings, and geospatial bottlenecks can also occur, rendering some geographic areas effectively excluded. While we did not collect the identifiers, e.g., geocoding data, necessary to evaluate potential geospatial biases, our study sites represented smaller, predesignated geographic areas within large cities.

Overall, our findings illustrate the feasibility, efficiency, and utility of RDS for adolescent surveillance in distressed urban environments across multiple geocultural settings. Our study is timely given the rapid urbanization and growth of the youth population globally, coupled with recognition of the social-structural hazards of vulnerable urban environments. RDS may well serve as a valuable recruitment tool to monitor health and well-being among young people in these urban environments, particularly where they risk being overlooked by household and school-based surveillance systems. It is critical to ensure the voices and experiences of homeless, marginally and unstably housed, and out-of-school youth in the surveillance that guides health-related policy.

Acknowledgements

This research was supported by Young Health Programme, a partnership between AstraZeneca, Johns Hopkins Bloomberg School of Public Health and Plan International, a leading global children's charity.

REFERENCES

1. Sawyer SM, Afifi RA, Bearinger LH, et al. Adolescence: a foundation for future health. *Lancet*. 2012 Apr 28; 379(9826):1630–1640. [PubMed: 22538178]
2. Patton GC, Coffey C, Cappa C, et al. Health of the world's adolescents: a synthesis of internationally comparable data. *Lancet*. 2012 Apr 28; 379(9826):1665–1675. [PubMed: 22538181]
3. Gray N, Azzopardi P, Kennedy E, Willersdorf E, Creati M. Improving adolescent reproductive health in Asia and the Pacific: do we have the data? A review of DHS and MICS surveys in nine countries. *Asia-Pacific journal of public health / Asia-Pacific Academic Consortium for Public Health*. 2013 Mar; 25(2):134–144. [PubMed: 21807622]
4. Reynolds HW, Wong EL, Tucker H. Adolescents' use of maternal and child health services in developing countries. *International family planning perspectives*. 2006 Mar; 32(1):6–16. [PubMed: 16723297]
5. Akmatov MK. Child abuse in 28 developing and transitional countries--results from the Multiple Indicator Cluster Surveys. *International journal of epidemiology*. 2011 Feb; 40(1):219–227. [PubMed: 20943933]
6. Kennedy E, Gray N, Azzopardi P, Creati M. Adolescent fertility and family planning in East Asia and the Pacific: a review of DHS reports. *Reproductive health*. 2011; 8:11. [PubMed: 21545708]
7. Magadi MA, Agwanda AO, Obare FO. A comparative analysis of the use of maternal health services between teenagers and older mothers in sub-Saharan Africa: evidence from Demographic and Health Surveys (DHS). *Social science & medicine* (1982). 2007 Mar; 64(6):1311–1325. [PubMed: 17174017]
8. Peltzer K. Early sexual debut and associated factors among in-school adolescents in eight African countries. *Acta paediatrica* (Oslo, Norway : 1992). 2010 Aug; 99(8):1242–1247.
9. Roberts C, Currie C, Samdal O, Currie D, Smith R, Maes L. Measuring the health and health behaviours of adolescents through cross-national survey research: recent developments in the Health Behaviour in School-aged Children (HBSC) study. *Journal of Public Health*. 2007; 15(3): 179–186.
10. Unicef. *The State of the World's Children 2011: Adolescence-an Age of Opportunity*. 2011

11. Marmot M, Friel S, Bell R, Houweling TA, Taylor S. Closing the gap in a generation: health equity through action on the social determinants of health. *Lancet*. 2008 Nov 8; 372(9650):1661–1669. [PubMed: 18994664]
12. Unicef. The State of the World's Children 2012: Children in an Urban World. New York, NY: 2012 Feb. 2012
13. Woan J, Lin J, Auerswald C. The health status of street children and youth in low- and middleincome countries: a systematic review of the literature. *The Journal of adolescent health : official publication of the Society for Adolescent Medicine*. 2013 Sep; 53(3):314–321. e312. [PubMed: 23706729]
14. Heckathorn DD. Respondent-driven sampling: a new approach to the study of hidden populations. *Social problems*. 1997:174–199.
15. Magnani R, Sabin K, Saidel T, Heckathorn D. Review of sampling hard-to-reach and hidden populations for HIV surveillance. *Aids*. 2005 May; 19(Suppl 2):S67–S72. [PubMed: 15930843]
16. Decker MR, Wirtz AL, Moguilnyi V, et al. Female Sex Workers in Three Cities in Russia: HIV Prevalence, Risk Factors and Experience with Targeted HIV Prevention. *AIDS and behavior*. 2013 Aug 9.
17. Qiu P, Yang Y, Ma X, et al. Respondent-driven sampling to recruit in-country migrant workers in China: a methodological assessment. *Scandinavian journal of public health*. 2012 Feb; 40(1):92–101. [PubMed: 21948979]
18. Johnston LG, Thurman TR, Mock N, Nano L, Carcani V. Respondent-driven sampling: A new method for studying street children with findings from Albania. *Vulnerable Children and Youth Studies*. 2010; 5(1):1–11.
19. Gwadz MV, Cleland CM, Quiles R, et al. CDC HIV testing guidelines and the rapid and conventional testing practices of homeless youth. *AIDS education and prevention : official publication of the International Society for AIDS Education*. 2010 Aug; 22(4):312–327. [PubMed: 20707692]
20. Mmari K, Blum R, Sonenstein F, et al. Adolescents' perceptions of health from disadvantaged urban communities: Findings from the WAVE study. *Social science & medicine (1982)*. 2014 Mar; 104:124–132. [PubMed: 24581070]
21. Johnston LG, Malekinejad M, Kendall C, Iuppa IM, Rutherford GW. Implementation challenges to using respondent-driven sampling methodology for HIV biological and behavioral surveillance: field experiences in international settings. *AIDS and behavior*. 2008 Jul; 12(4 Suppl):S131–S141. [PubMed: 18535901]
22. Johnston LG, Whitehead S, Simic-Lawson M, Kendall C. Formative research to optimize respondent-driven sampling surveys among hard-to-reach populations in HIV behavioral and biological surveillance: lessons learned from four case studies. *AIDS care*. 2010 Jun; 22(6):784–792. [PubMed: 20467937]
23. Ghanem KG, Hutton HE, Zenilman JM, Zimba R, Erbeling EJ. Audio computer assisted self interview and face to face interview modes in assessing response bias among STD clinic patients. *Sexually transmitted infections*. 2005 Oct; 81(5):421–425. [PubMed: 16199744]
24. White RG, Lansky A, Goel S, et al. Respondent driven sampling--where we are and where should we be going? *Sexually transmitted infections*. 2012 Oct; 88(6):397–399. [PubMed: 23012492]
25. Volz E, Heckathorn D. Probability based estimation theory for respondent driven sampling. *Journal of Official Statistics*. 2008; 24(1):79–97.
26. Robinson WT, Risser JM, McGoy S, et al. Recruiting injection drug users: a three-site comparison of results and experiences with respondent-driven and targeted sampling procedures. *Journal of urban health : bulletin of the New York Academy of Medicine*. 2006 Nov; 83(6 Suppl):i29–i38. [PubMed: 16933101]
27. Yeka W, Maibani-Michie G, Prybylski D, Colby D. Application of respondent driven sampling to collect baseline data on FSWs and MSM for HIV risk reduction interventions in two urban centres in Papua New Guinea. *Journal of urban health : bulletin of the New York Academy of Medicine*. 2006 Nov; 83(6 Suppl):i60–i72. [PubMed: 17066328]

28. Jaya, Hindin MJ, Ahmed S. Differences in young people's reports of sexual behaviors according to interview methodology: a randomized trial in India. *American journal of public health*. 2008 Jan; 98(1):169–174. [PubMed: 18160677]
29. Mensch BS, Hewett PC, Erulkar AS. The reporting of sensitive behavior by adolescents: a methodological experiment in Kenya. *Demography*. 2003 May; 40(2):247–268. [PubMed: 12846131]
30. Rathod SD, Minnis AM, Subbiah K, Krishnan S. ACASI and face-to-face interviews yield inconsistent estimates of domestic violence among women in India: The Samata Health Study 2005–2009. *Journal of interpersonal violence*. 2011 Aug; 26(12):2437–2456. [PubMed: 21282116]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figures 1-5.

Daily recruitment by gender and site.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

RDS recruitment summary by site

	Baltimore	Delhi	Ibadan	Johannesburg	Shanghai	Total
Catchment area	Five zip codes of East Baltimore	Badarpur area of Delhi	Ibadan North Local Government Area	Inner city Johannesburg	Taapu Community of the Putuo District of Shanghai (migrant youth only)	
a. Total # seeds	8	7	10	14	5	44
b. Total # coupons given	1158	657	847	1347	1166	5175
c. Total # coupons returned (sum d,e,f)	488	493	458	483	469	2391
d. Of returned coupons, # ineligible	20	0	3	0	16	39
e. Of returned coupons, # non participating	0	0	0	0	3	3
f. Of returned coupons, # eligible and participating	468	493	455	483	450	2349
g. Total number of participants (sum a and f)	476	500	465	497	455	2393
h. Participants providing incomplete or nonsensical data	15	0	5	1	17	38
i. Duplicate participants dropped	1	0	0	0	0	1
j. Lost or unsaved interviews	4	0	11	0	0	15
k. Final analytic sample	456	500	449	496	438	2339
l. Number of recruits by seed (range)	1–369	4–212	2–228	1–424	2–329	-
m. Number of recruitment waves (range)	1–14	1–8	1–16	1–15	1–20	-
n. Primary incentive	20 USD	None	N500	R100	100 CNY	-
o. Secondary incentive	10 USD per recruit, up to 3 recruits	None	N250 per recruit, up to 3 recruits	R10 per recruit, up to 3 recruits	50 CNY per recruit, up to 3 recruits	-
p. Location of data collection	Johns Hopkins School of Public Health	Population Council offices	University College Hospital; Lifebuilders	Wits Reproductive Health and HIV Institute offices	Taapu family planning service station	-

	Baltimore	Delhi	Ibadan	Johannesburg	Shanghai	Total
q. Language(s) of survey data collection	English	English Hindi	English Yoruba	English isiZulu Sesotho	English Mandarin	-
r. Data collection dates	April-May 2013	August-September 2013	September-October 2013	August-September 2013	May-July 2013	-
s. Elements of variation for seed matrix and seed characteristics	<ul style="list-style-type: none"> • School status • Employment status • Housing status • Drug involvement • Race 	<ul style="list-style-type: none"> • School status • Employment status (males) • Marital status (females) 	<ul style="list-style-type: none"> • School status • Employment status • Parental contact and living arrangements 	<ul style="list-style-type: none"> • School status • Employment status • Engagement in community activities • Living arrangements • Migrant status 	<ul style="list-style-type: none"> • School status • Employment status • Living arrangements 	

Table 2

Challenges in RDS implementation across sites

Challenges	Baltimore	Delhi	Ibadan	Johannesburg	Shanghai
Sample imbalance (eg. gender, student status)	X		X	X	X
Ineligible individuals trying to gain entry to study	X	X		X	X
Attempts at repeat enrollment	X			X	X
Unbalanced recruitment chains	X		X		X
Participants not taking survey seriously	X		X	X	X
Non-productive seeds	X			X	X
Recruitment pace	X	X	X	X	X

Table 3

Participation of high mobility and out-of--school youth, by site

	Baltimore	Delhi	Ibadan	Johannesburg	Shanghai	All Sites
	Weighted % N=475	Weighted % N=500	Weighted % N=465	Weighted % N=497	Weighted % N=455	Weighted % N=2388
Housing status **						
Stably housed	95.85	71.66	78.41	78.04	88.74	82.44
Unstably housed	4.15	28.34	21.59	21.96	11.26	17.56
School enrollment by age group						
Ages 15–16	N=252	N=284	N=267	N=123	N=127	N=1086
In school	97.80	92.63	93.57	90.58	59.43	87.53
Out of school	2.20	7.37	6.43	9.42	40.57	12.47
Ages 17–19	N=203	N=216	N=181	N=371	N=311	N=1302
In school	68.10	73.69	69.51	77.92	21.13	61.50
Out of school	31.90	26.31	30.49	22.08	78.87	38.50
Past month school absenteeism ***	N=388	N=428	N=364	N=406	N=125	N=1745
0 days	19.41	4.77	12.08	16.96	69.48	18.89
1–2 days	23.24	27.34	11.38	36.41	27.03	25.20
3–5 days	35.6	33.42	11.77	34.55	2.65	26.51
6+ days	21.76	34.48	64.77	12.09	0.84	29.40

* Weighted percentage adjusted for RDS and age

** Does not have a regular place to stay or has a regular place to stay but: 1) stayed there, on average, less than 4 nights per week in the past 30 days and 2) stayed in 2 or more different places in the past week

*** Number of days of school missed in the past 30 days among those currently in school