Published in final edited form as:

Biol Psychiatry. 2015 June 15; 77(12): 1079–1088. doi:10.1016/j.biopsych.2015.03.013.

Excitation-inhibition discoordination in rodent models of mental disorders

André A. Fenton

Neurobiology of Cognition laboratory, Center for Neural Science, New York University, 4 Washington Place, NYC, NY 10012

Abstract

Animal models of mental illness provide a foundation for evaluating hypotheses for the mechanistic causes of mental illness. Neurophysiological investigations of neural network activity in rodent models of mental dysfunction are reviewed from the conceptual framework of the discoordination hypothesis, which asserts that failures of neural coordination cause cognitive deficits in the judicious processing and use of information. Abnormal dynamic coordination of excitatory and inhibitory neural discharge in pharmacological and genetic rodent models support the discoordination hypothesis. These observations suggest excitation-inhibition discoordination and aberrant neural circuit dynamics as causes of cognitive impairment as well as therapeutic targets for cognition-promoting treatments.

Keywords

discoordination; neural synchrony; neural coordination; excitation-inhibition coupling; oscillations; neural ensemble

INTRODUCTION

In *Siddhartha*, Hermann Hesse penned an astute exchange between Kamaswami, the rich merchant, and the hero Siddhartha, who at this stage of his journey is without possessions and appears estitute (1).

K: "What is it that you've learned, what you're able to do?" S: "I can think. I can wait. I can fast."

K: "That's everything?"

S: "I believe, that's everything!"

© 2015 Published by Society of Biological Psychiatry

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

FINANCIAL DISCLOSURES

Dr. Fenton reports an equity interest in Bio-Signal Group Corp., a company he founded, US patent 7767195, and US patents pending 11/694,855, 11/694,816, 13/61554743, 13/284,886.

Siddhartha explains that he has full control of his mind; he is not distracted by impatience or bodily needs. With his singular ability, Siddhartha becomes rich and powerful.

This is an unusual beginning for a paper on rodent models of mental disorders, but this dialog makes the starting point of the present work: cognitive abilities are extremely valuable resources with major economic and social impacts beyond individual well being. The term “mental capital” expresses this notion that the mental capacities of individuals and the groups that they form are determinants of individual and national wealth and prosperity (2). Research with rodent models of mental disorders aims to improve mental capacities by improving understanding of mental function and dysfunction.

Importance of Animal Models

The importance of animal research that translates basic science to understanding mental disorders like schizophrenia has become increasingly apparent: knowledge of basic mechanisms grew enormously while treatment options only expanded slightly (3, 4). Understanding such disorders is impeded because animal research traditionally avoids the mental domain where mental illness is prominent. One difficulty in developing more sophisticated approaches to treating psychiatric illness is the gulf between the behavioral/mental spheres in which mental disorders manifest, and the biochemical/developmental domains where therapies and interventions are implemented. This is the problem of the missing middle (5). Tools are optimal for the microscopic (genomics and proteomics) and the large ensemble levels (fMRI and EEG). The middle level is difficult to access, representing analysis of, for example, temporally-organized discharge within neural ensembles or temporally-activated synapses called synapsensembles (6). Yet this middle level is needed to connect the nuts and bolts of receptors and transmitters with the level of clinical observables (7). A major challenge is to study normal and abnormal mental phenomena at the middle level of neural ensembles. This level of investigation is currently only practical in animal studies and is especially developed for rodents.

Patterns of neural circuit activity in rodent models of mental dysfunction are the focus of this review. I argue that the neurophysiology literature on animal models of mental dysfunction is converging on a specific form of neural discoordination as the basis for impaired cognition: when cognitive deficits manifest, the culprit is likely to be inappropriately coordinated dynamic interactions between excitatory and inhibitory neural discharge within and between neural networks.

The utility of animal models

What is an “animal model of a mental disorder” and how might it be useful? The phrase implies the animal mimics a patient with the disorder being modeled, raising issues of validity (8). However, the notion of a mimic is problematic for mental illness. The mental phenomena that are the foundation of a clinical diagnosis are rarely applicable to animals because it is unclear that animals have corresponding mental capacities, and if they do, they are unlikely to manifest like in people. Consider ‘psychosis’, a symptom that involves a distorted sense of objective reality, and profound alterations of personality. What would that look like in a non-human? Furthermore, many diagnoses of mental dysfunction are open

constructs, in that definitive criteria that define the disorder and differentiate it from another are unknown (9). How can one model something that is poorly defined? How can one judge if a model is a valid mimic? Psychosis is a symptom of schizophrenia, schizoaffective and personality disorders, as well as depression and bipolar disorders; if we could agree on what psychosis looked like in a rodent, which disorder would we have modeled? In this regard, the effort to base clinical diagnosis on objective biological criteria, the Research Domain Criteria (RDoC) project (10) may prove invaluable, and the identification of biomarkers for mental disorders, such as genetic variants and mutations, holds substantial promise. Even this can be ethologically problematic because pleiotropy can have species-specific outcomes, so that depending on the species, a gene may confer rather different phenotypes, making it challenging to interpret how a human genetic alteration might be mimicked in a model organism. The penetrance of mental disorder-related genetic variants poses a further complication for animal models because penetrance is typically just a few percent (11), meaning that the probability of a genetic modification leading to an abnormal phenotype can be difficult to detect in laboratory studies (12). There is even substantial overlap in the genetic abnormalities that associate with disorders as diverse as autism spectrum disorder, attention deficit-hyperactivity disorder, bipolar disorder, major depressive disorder, and schizophrenia (13, 14). These issues have been reviewed as they pertain to animal models that are relevant to schizophrenia (15), but the same issues are central to any attempts to model a mental disorder.

It may be ill advised to consider animal models of mental disorders as mimics of the target disorder (16). Rather, animal models are amongst our most powerful tools to test hypotheses about the disorder (15). We do not consider any of the rodent models reviewed here models of disease *per se*, rather they are tools, a reagent that when used effectively can evaluate key hypotheses to drive understanding of relevant mental disorders.

Neural coordination: network patterns of electrical activity and neural codes

Much of the research on neural network patterns in rodent models is conducted within the hypothesis that discoordinated patterns of electrical neural activity are a core deficit in a variety of mental disorders (17-22). According to the hypothesis, cognitive deficits arise because of inappropriately coordinated neural electrical activity within and between neural networks (23). This causes neural information processing failures that preferentially manifest when there are competing sources of information. These coordinating processes are best studied as the temporal discharge relationship between two or more neurons (24), and/or their relationship to the local field potential (LFP) that arises from the spatiotemporal patterns of synaptic currents (6).

The coordinating processes are thought to be distinct from the more unitary processes that determine spiking characteristics, such as the firing rate and the tuning curves of individual neurons (25), and the frequency of oscillations in the LFP. Coordinating activity can also impact these unitary properties and can manifest as failures to sufficiently amplify neural representations of relevant information and sufficiently suppress the representations of irrelevant information, without explicit discoordination of temporal discharge (26). It may be that such abnormalities in unitary processes are secondary to failures of coordination

between excitation and inhibition (26). Nonetheless, the discoordination hypothesis predicts that basic properties and neural network functions, such as responses to stimuli and memory, can maintain under simple conditions. In contrast, functional and discoordination abnormalities will manifest under complex conditions that require using relevant information and ignoring irrelevant information to meet competing demands as in tasks like the Stroop test for people (27), and for rodents, tasks that require contextual modulation of responses like the two-frame place avoidance task, set-shifting tasks, and other tests of cognitive flexibility (28-30), especially those that require the subject to selectively use the information that is inherently not preferred, which may be the case for the Room+Arena-variant of the place avoidance task (29).

The discoordination hypothesis emerges from a concept of how neurons represent information, which remains unknown. At the core of the dedicated-coding hypothesis is the notion of 'cardinal cells.' Analogous to how the red light in a traffic signal unambiguously means 'stop' (Fig. 1A), these are neurons dedicated to signaling high-order stimuli and concepts, like 'face' or 'grandmother.' Examples include single cell firing tuned to faces (31), celebrities, which was recorded from people (32) and hippocampus place cells (Fig. 1C). A place cell tends to discharge at a single location in standard experimental environments (33). Although the capacity of a dedicated code is limited, how brains read the information in the firing of such cardinal cells seems straightforward and isomorphic with perception (34).

The discoordination hypothesis is founded in the ensemble-coding hypothesis. Information is represented by patterns of activity across many cells in an ensemble code, analogous to a jumbotron display that uses many lights to signal a message (Fig. 1B). No particular bulb is essential for a message, many more messages can be encoded than there are bulbs, and the same bulbs can only represent one message at a time. Such properties require temporal coordination amongst neurons to represent, and transmit information, perhaps at multiple time scales (35, 36). Ensemble codes must avoid simultaneously representing multiple items with the same cells (37). Just as a jumbotron cannot simultaneously display two messages using many of the same lights, cell assemblies with many cells in common cannot coactivate if they share cells because the cells will merge into a unique, coactivity-defined cell assembly. Without effective neural coordination, multiple assemblies will coactivate merging into one assembly with catastrophic information loss (37). Hebb's cell assembly postulate is an ensemble-coding scheme (38) in which a subset of linked cells fire together to represent information by the pattern of active and inactive cells. Place cells that appear to signal single locations in small environments, discharge in multiple locations in larger environments (39, 40) indicating they are fundamentally an ensemble place code (Fig. 1C). Unlike dedicated codes, it is unclear how the brain might read an ensemble code (6).

There are multiple coordinating mechanisms, but this review focuses on the dynamic interactions between populations of excitatory and inhibitory cells, which can be measured directly from ensemble recordings of unitary spike trains as well as recordings of frequency-specific oscillations in the LFP. The oscillatory dynamics provide an electrophysiological infrastructure within which spiking activity is embedded and organized (6, 41, 42).

Figure 2 describes a winner-take-all configured competitive neural network architecture in which the interactions between excitatory and inhibitory neurons is explicit and illustrative of how optimal and aberrant excitation-inhibition interactions underlie neural coordination (Fig. 2A-C) and discoordination (Fig. 2D-F), respectively. Such networks are proposed to explain phenomena like cognitive control and attention (43) as well as the tuning of spatially selective neurons like place cells, head-direction cells and grid cells (44-46). The architecture is such that for any excitation, a subset of similarly tuned excitatory neurons will become active and suppress the remaining neurons. This is because of mutual excitation amongst subsets of the similarly tuned excitatory cells and because the excitatory activity activates interneurons that produce widespread inhibition. From the ensemble coding perspective, this network describes how subsets of cells are recruited to represent distinct classes of information by their coactivity. Disturbing the excitation-inhibition interactions can dramatically alter the network's ability to represent multiple classes of information, resulting in the superposition catastrophe (Fig. 2D-F). LFPs arise from the synchronous occurrence of postsynaptic potentials at many neighboring cells, resulting in rhythmic activity (47). These oscillations are typically temporally correlated with discharge of a small proportion of neurons due to strong feedback and feed-forward inhibition (39, 40). The correlation does not mean the oscillations are due to the spiking, quite the contrary, the synaptic activity underlying the oscillations forms a neural infrastructure that organizes temporal opportunities for spiking due to the coupling of excitation and inhibition (48, 49).

Gamma (30-100 Hz) is probably the best understood oscillation. Gamma tends not to synchronize across distant sites as these oscillations arise from the coincident local release of GABA because GABA_A receptor-mediated inhibition has a ~25 ms time constant (50-52). Tonic excitation of inhibitory neurons generates gamma oscillations by synchronous release of GABA that limits the possibility of excitation to the next gamma cycle. This organizes principal cell spiking in cortical networks (53, 54). Greater excitation drives faster oscillations because GABA inhibition can be overcome sooner (53). This interaction between excitation and inhibition illustrates how excitation and inhibition are dynamically coupled. As reviewed below, the exact timing between excitatory and inhibitory discharge within identified subsets of excitatory and inhibitory neurons provides a rich palette of excitation-inhibition coordination phenomena. Additional mechanisms provide for neural coordination and these include adjustments of EPSP-spike (E-S) coupling, which has an inhibitory component (55), as well as intrinsic rhythmic and bursting properties such as what is discussed next for theta oscillations (56, 57).

Theta oscillations synchronize neural activity on a ~140 ms time scale such that the sequence of theta phases is associated with the alternation between reduced and enhanced discharge probability (58, 59). In hippocampus this is largely due to rhythmic inhibition that waxes and wanes with corresponding membrane potential fluctuations. This temporal organization is also spatially organized; theta oscillations propagate as a travelling wave along the septo-temporal (dorsoventral) hippocampal axis (60, 61). Unlike gamma oscillations, which are generated locally and so do not synchronize across long ranges, theta oscillations can synchronize between distant sites, such as the hippocampus and prefrontal cortices (62-64) and is more synchronous between prefrontal cortex and ventral than dorsal hippocampus, consistent with the direct and indirect anatomical connectivity, respectively

(65-67). Spike-field coherence can be observed across distant sites; LFPs and single neuron spiking in hippocampus and neocortex can be synchronized at a variety of oscillatory frequencies during both waking and sleep behaviors (68-71).

Neural discoordination in pharmacological models of mental dysfunction

Under urethane anesthesia, exemplary aberrant neural coordination caused by excitation-inhibition discoordination is observed in one hippocampus after inactivating the other hippocampus by tetrodotoxin (26). Inactivation disinhibits the uninjected hippocampus (56, 72), transiently increasing excitatory cell discharge for ~15 minutes (Fig. 3A), and uncoupling excitatory and inhibitory cells (Fig. 3B), despite the inactivation lasting ~10 hours. This resulted in a persistent discoordination amongst principal cells an hour after tetrodotoxin, when baseline rates had restored. Excitatory cell pairs that had previously been weakly coactive were now strongly coactive; initially strongly coactive pairs were unchanged, demonstrating aberrant circuit function (Fig. 3C). This discoordination of cell pair spike timing was selectively associated with impaired cognitive control in the Room +Arena- variant of the place avoidance task, but not control task variants requiring limited cognitive control (73), as predicted by the discoordination hypothesis (Fig. 3D).

The discoordination hypothesis emerged from animal studies that used pharmacological agents that produce altered mental states in people, in addition to the theoretical considerations of how neurons code information. Cannabinoids like delta(9)-tetrahydrocannabinol cause acute, reversible memory impairments. Cannabinoids slightly decrease the overall firing of hippocampal principal cells but not interneurons. Nor do cannabinoids disrupt the spatial discharge of hippocampus place cells, demonstrating maintained single cell discharge properties (74).

In contrast, cannabinoids discoordinate the discharge of place cells in relation to each other and in relation to LFPs, at multiple timescales (75). At theta timescales and faster, cannabinoids reduce cell pair coactivity, possibly disrupting the gathering of principal cells into coactivity-defined cell assemblies (24) that carry place information (30, 76). Cannabinoids disrupt the relationship between the theta phase difference at which a pair of place cells fires and the distance between their place fields, a representation of place sequences (77, 78). At timescales of a few seconds, cannabinoids increase coactivity, potentially causing excessive formation of cell assemblies that signal distinctive contexts or tasks (35, 36). These forms of discoordination are accompanied by reduced theta, gamma, and sharp wave associated ripple (100-200 Hz) oscillations in the hippocampus LFP, and the theta reduction indexes impairment in a task that requires segregation of memories of more and less recent places (75). This pattern of preserved discharge in individual cells with disorganized inter-cell activity resembles the effects of unilateral tetrodotoxin and is predicted by the discoordination hypothesis.

Cannabinoids also discoordinate activity between brain areas. While hippocampal theta is reduced by cannabinoids, theta is preserved in prefrontal areas during stillness and rest, whereas gamma is decreased (79). Cannabinoids disinhibit by activating CB1 receptors on cholecystokinin-positive interneurons to decrease GABA release. CB1 receptors are also on excitatory terminals so that instead of increasing firing rates, the disinhibition primarily

disturbs spike timing (80-83). Cannabinoids reduce correlated <10 ms burst spiking in hippocampus, whereas the effect is to reduce prefrontal coactivity over a wider range of time scales (10-100 ms), again with minimal effects on overall firing rates. Although the cannabinoid effects coincide with impaired spatial choices in normal animals (79), the reductions in coordinated activity have beneficial potential when baseline activity is hypersynchronized, as in pathological conditions. Resting gamma hypersynchrony is observed in schizophrenia as well as some experimental conditions that selectively cause cognitive coordination deficits (26, 84, 85).

Different forms of neural discoordination are caused by psychotomimetic uncompetitive antagonists of N-methyl-D-aspartate receptors (NMDAR) like phencyclidine (PCP), MK801 and the non-competitive antagonist ketamine (86, 87). Psychotomimetics produce psychotic symptoms in healthy people (88) and exacerbate symptoms in schizophrenia patients (89) making them once the most widely used animal models in antipsychotic drug development (90). Use of PCP models is dwindling, perhaps due to an overemphasis on the PCP-induced sensorimotor deficits as a model for positive symptoms (90) as well as the recognition that cognitive impairments are a core deficit of schizophrenia (91, 92; see the MATRICS: <http://www.matrics.ucla.edu/> and CNTRICS: <http://cntrics.ucdavis.edu/> initiatives of NIMH). Because cognitive symptoms are found in nearly all patients (93, 94) and are amongst the strongest predictors of clinical outcome (91, 95), emphasis has shifted to develop cognition-promoting antipsychotic treatments (91, 92). Although it has not been sufficiently emphasized, psychotomimetics discoordinate neural activity and impair cognition (18), consistent with the discoordination hypothesis.

Psychotomimetics activate prefrontal (96, 97) and cingulate cortical areas (98, 99) in human neuroimaging studies, which seems paradoxical because the drugs reduce excitatory currents through NMDARs. In rat prefrontal areas, MK801 decreases interneuron firing. The drug increases principal cell firing rates in ~40% of these cells but decreases burst discharge (100, 101), and firing rates in a small fraction (102). This suggests MK801 disinhibits cortical networks, potentially resolving the paradox. Consistent with the reduced inhibitory control of principal cell spiking, psychotomimetics enhance action potential associated high frequency oscillations (103-105) in motor cortex concurrent with hyperlocomotion (106). MK801 also disrupts the coupling of spiking to gamma oscillations in the LFP (102).

Ketamine also discoordinates hippocampus discharge by interfering with the coupling of inhibitory and excitatory cells, producing abnormal coordination in a variety of ways. Ketamine weakly decreases pyramidal cell firing, increases and decreases firing rates in distinct inhibitory neuron subclasses, producing no net change (107). CA1 cells usually discharge asynchronously, but synchronize under ketamine (107). Ketamine causes both (excitatory) pyramidal cells and (inhibitory) parvalbumin-positive basket cells to increase near-synchronous discharge, such that they were inhibited from firing together for a few hundred milliseconds before and after coactive discharge. Ketamine changes the phase of theta at which putative basket cells discharge (107). Whereas pyramidal cells preferentially discharge at the trough of local theta oscillations, basket cells preferentially discharge just earlier, on the descending theta phase. Another class of interneuron (possibly bistratified cells) prefers to fire at the theta trough, when pyramidal cells discharge, but a third class,

possibly chandelier cells, prefers to fire at the peak of theta (108-110). Ketamine also changed the theta-related firing probabilities of these cell classes, which is how network discharge can discoordinate without changing overall activity. These observations demonstrate that over-simple notions, like more or less neural activity in an inhibitory or excitatory population, are poor descriptors and predictors of discoordination. Because distinct interneuron classes target distinct input-specific perisomatic and dendritic compartments of hippocampus and neocortex (111-113), these findings emphasize the importance of measuring and manipulating the subcellular and circuit-specific features of neural networks rather than standard analyses that are prone to over-averaging across heterogeneous subpopulations of signals and cell types.

Neural discoordination in genetic models of mental disorders

Human genomic investigations identify genetic alterations that are associated with mental disorders and animal models are valuable for identifying the functional consequences. The 22q11.2 microdeletion increases risk of schizophrenia and autism (114-117) but it was not obvious why. Abnormal neural synchrony between the dorsal hippocampus and prefrontal cortex was identified in a mouse model that mimics the deletion and expresses working memory deficits. During working memory tasks, rodents increase the synchrony of hippocampal – prefrontal theta and phase locking of prefrontal discharge to theta (63, 118). This coordination is reduced in the mutant and the reduction predicts the working memory deficit magnitude (118).

The coupling of theta between the prefrontal cortex and ventral hippocampus increases during anxiety-related behavior, as does the theta-phase synchrony of discharge that distinguishes between parts of an environment with different anxiogenic potential (119). The theta-synchrony increase is exaggerated in serotonin 1A receptor knockout mice, a model of increased anxiety-related behavior (67).

There is substantial evidence of impaired NMDAR function of cortical interneurons in schizophrenia (120, 121), the functional consequences of which were investigated by knocking out the NR1 receptor subunit in mouse interneurons either early or after adolescence (122). In addition to a variety of schizophrenia-relevant behavioral and histological abnormalities, the discharge of S1 neurons in the early knockouts was faster but poorly correlated. Although baseline oscillatory power in the auditory cortical LFP was greater in the mutant, evoked 20-Hz and 40-Hz oscillatory activity was reduced and short lived in response to steady-state click stimulus trains (123). The abnormalities were not observed if the gene was deleted after adolescence, indicating a neurodevelopmental consequence of NMDAR loss of function in interneurons (122).

Schizophrenia risk is associated with abnormality at a gene locus that codes for the phosphatase calcineurin (124). Calcineurin knockout mice revealed that loss of the phosphatase severely impairs LTD, slightly increases LTP, and causes schizophrenia-related behavioral abnormalities (125, 126). Hippocampal LFPs were normal, except ripples were more frequent in the knockout, indicating a specific form of increased excitability (127-129) that is sensitive to long-term synaptic plasticity changes (127). During exploration, individual hippocampal place cells appear normal in the knockout, but discharge timing is

abnormal during ripple-associated replay of the discharge sequences that were observed during exploration. This discoordination is specific to quiet rest when neural activity during recent experience is re-expressed, presumably to organize and store memory (129-134).

Summary: the discoordination hypothesis and beyond

Processing information and generating knowledge is valuable, making mental illness devastating and costly. Mental disorders are being reconceptualized and analysed as genetically and environmentally driven developmental alterations in specific neural circuit operations (135). Typically, these circuits are robust and adaptive, partly because of multiple synaptic, structural and functional plasticity mechanisms for maintaining function. Unfortunate and inconveniently timed genetic and environmental influences can alter neural circuits in myriad ways with consequences that depend on the exact genetic, environmental and developmental interactions. Accordingly, a good bet to promote understanding and effective treatment outcomes is to look (and hope) for disruption in a limited set of common pathways that mediate cognition and behavior (15). This is the inspiration for the discoordination hypothesis, which focuses on neural network function in hopes of affecting outcomes rather than correcting the causes.

The discoordination hypothesis does not eschew genetic or other etiological accounts for mental disorders, it just does not focus on or aim to correct them. The hypothesis embraces data suggesting that diverse etiology converges to change neural circuit function, often by altering synaptic structure and function (136). The hypothesis supposes that dysfunction emanates from distributed disturbances and secondary adaptations within synaptic networks, suggesting that pathology is difficult to localize in space and time. The patterns of neural activation and inactivation captured by the patterns and interactions of neural oscillations reviewed here can only be fully understood in the context of their use and time evolution. These dynamics may at once be the precious information that the brain processes and generates as well as the substrate for processing that information (56). The relationship between the information and the functional infrastructure for it is analogous to the relationship between water and the river that it at once forms and is guided by. Tools and concepts are developing to identify and precisely measure these dynamics. We even see that appropriately curated experience can itself repair and improve outcomes (137-139). The neural patterns themselves are the therapeutic target (92, 95, 140, 141).

ACKNOWLEDGEMENTS

I thank Kally O'Reilly, Fraser Sparks and Zoe Talbot for comments on the manuscript. Supported by NIH grants R01MH084038, R01MH099128, and R01AG043688.

REFERENCES

1. Hesse, H. Siddhartha - An Indic Poem. transl. 2007. Lesser, R., editor. Barnes and Noble Books; New York: 1922.
2. Beddington J, Cooper CL, Field J, Goswami U, Huppert FA, Jenkins R, et al. The mental wealth of nations. *Nature*. 2008; 455:1057–1060. [PubMed: 18948946]
3. Leucht S, Cipriani A, Geddes JR, Salanti G, Davis JM. Efficacy of antipsychotic drugs for schizophrenia - Authors' reply. *Lancet*. 2013; 382:1874–1875. [PubMed: 24315170]

4. Lieberman JA, Stroup TS, McEvoy JP, Swartz MS, Rosenheck RA, Perkins DO, et al. Effectiveness of antipsychotic drugs in patients with chronic schizophrenia. *N Engl J Med*. 2005; 353:1209–1223. [PubMed: 16172203]
5. Laughlin RB, Pines D, Schmalian J, Stojkovic BP, Wolynes P. The middle way. *Proc Natl Acad Sci U S A*. 2000; 97:32–37. [PubMed: 10618366]
6. Buzsaki G. Neural syntax: cell assemblies, synapsembles, and readers. *Neuron*. 2010; 68:362–385. [PubMed: 21040841]
7. Goldman-Rakic PS. The physiological approach: functional architecture of working memory and disordered cognition in schizophrenia. *Biol Psychiatry*. 1999; 46:650–661. [PubMed: 10472417]
8. Lipska BK, Weinberger DR. To model a psychiatric disorder in animals: schizophrenia as a reality test. *Neuropsychopharmacology*. 2000; 23:223–239. [PubMed: 10942847]
9. Corvin A, Morris DW. Genome-wide association studies: findings at the major histocompatibility complex locus in psychosis. *Biol Psychiatry*. 2013; 75:276–283. [PubMed: 24199664]
10. Cuthbert BN, Insel TR. Toward new approaches to psychotic disorders: the NIMH Research Domain Criteria project. *Schizophr Bull*. 2010; 36:1061–1062. [PubMed: 20929969]
11. Vassos E, Collier DA, Holden S, Patch C, Rujescu D, St Clair D, et al. Penetrance for copy number variants associated with schizophrenia. *Hum Mol Genet*. 2010; 19:3477–3481. [PubMed: 20587603]
12. Weatherall DJ. Phenotype-genotype relationships in monogenic disease: lessons from the thalassaemias. *Nat Rev Genet*. 2001; 2:245–255. [PubMed: 11283697]
13. Smoller, JWC-DGotPGC. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet*. 2013; 381:1371–1379. [PubMed: 23453885]
14. Lee SH, Ripke S, Neale BM, Faraone SV, Purcell SM, Perlis RH, et al. Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nat Genet*. 2013; 45:984–994. [PubMed: 23933821]
15. Mitchell, KJ.; O'Donnell, P.; Durstewitz, D.; Fenton, AA.; Gingrich, JA.; Gordon, JA., et al. A Framework for the Use of Models in Schizophrenia. In: Silverstein, SM.; Moghaddam, B.; Wykes, T., editors. *Schizophrenia: Evolution and Synthesis*. MIT Press; Cambridge, MA: 2013. p. 212-226.
16. O'Donnell P. Of mice and men: what physiological correlates of cognitive deficits in a mouse model of schizophrenia tell us about psychiatric disease. *Neuron*. 2013; 80:265–266. [PubMed: 24139031]
17. Tononi G, Edelman GM. Schizophrenia and the mechanisms of conscious integration. *Brain Res Brain Res Rev*. 2000; 31:391–400. [PubMed: 10719167]
18. Phillips WA, Silverstein SM. Convergence of biological and psychological perspectives on cognitive coordination in schizophrenia. *Behav Brain Sci*. 2003; 26:65–82. discussion 82-137. [PubMed: 14598440]
19. Lisman JE, Coyle JT, Green RW, Javitt DC, Benes FM, Heckers S, et al. Circuit-based framework for understanding neurotransmitter and risk gene interactions in schizophrenia. *Trends Neurosci*. 2008
20. Uhlhaas PJ, Singer W. Neural synchrony in brain disorders: relevance for cognitive dysfunctions and pathophysiology. *Neuron*. 2006; 52:155–168. [PubMed: 17015233]
21. Jones MW. Errant ensembles: dysfunctional neuronal network dynamics in schizophrenia. *Biochem Soc Trans*. 2010; 38:516–521. [PubMed: 20298213]
22. Fenton, AA. Neural coordination and psychotic disorganization. In: Holscher, C.; Munk, MH., editors. *Information Processing by Neuronal Populations*. Cambridge University Press; London: 2008. p. 387-408.
23. Moghaddam B, Wood J. Teamwork matters: coordinated neuronal activity in brain systems relevant to psychiatric disorders. *JAMA Psychiatry*. 2014; 71:197–199. [PubMed: 24305975]
24. Schneidman E, Berry MJ 2nd, Segev R, Bialek W. Weak pairwise correlations imply strongly correlated network states in a neural population. *Nature*. 2006; 440:1007–1012. [PubMed: 16625187]
25. Phillips WA, Singer W. In search of common foundations for cortical computation. *Behav Brain Sci*. 1997; 20:657–683. discussion 683-722. [PubMed: 10097008]

26. Olypher AV, Klement D, Fenton AA. Cognitive disorganization in hippocampus: a physiological model of the disorganization in psychosis. *J Neurosci*. 2006; 26:158–168. [PubMed: 16399683]
27. Stroop JR. Studies of interference in serial verbal reactions. *Journal of Experimental Psychology*. 1935; 18:643–662.
28. Robbins TW. The 5-choice serial reaction time task: behavioural pharmacology and functional neurochemistry. *Psychopharmacology (Berl)*. 2002; 163:362–380. [PubMed: 12373437]
29. Fenton AA, Wesierska M, Kaminsky Y, Bures J. Both here and there: simultaneous expression of autonomous spatial memories in rats. *Proc Natl Acad Sci U S A*. 1998; 95:11493–11498. [PubMed: 9736765]
30. Kelemen E, Fenton AA. Dynamic grouping of hippocampal neural activity during cognitive control of two spatial frames. *PLoS Biol*. 2010; 8:e1000403. [PubMed: 20585373]
31. Bruce C, Desimone R, Gross CG. Visual properties of neurons in a polysensory area in superior temporal sulcus of the macaque. *J Neurophysiol*. 1981; 46:369–384. [PubMed: 6267219]
32. Quiroga RQ, Reddy L, Kreiman G, Koch C, Fried I. Invariant visual representation by single neurons in the human brain. *Nature*. 2005; 435:1102–1107. [PubMed: 15973409]
33. Muller R. A quarter of a century of place cells. *Neuron*. 1996; 17:813–822. [PubMed: 8938115]
34. Cerf M, Thiruvengadam N, Mormann F, Kraskov A, Quiroga RQ, Koch C, et al. On-line, voluntary control of human temporal lobe neurons. *Nature*. 2010; 467:1104–1108. [PubMed: 20981100]
35. Kelemen, E.; Fenton, AA. The organization of neuronal discharge on timescales of milliseconds and seconds is related to the spatial response properties of hippocampal neurons. 1. Springer-Verlag; New York: 2012.
36. Kelemen E, Fenton AA. Key features of human episodic recollection in the cross-episode retrieval of rat hippocampus representations of space. *PLoS Biol*. 2013; 11:e1001607. [PubMed: 23874154]
37. von der Malsburg, C. The correlation theory of brain function (MPI Biophysical Chemistry, internal report 81-82). In: Domany, E.; van Hemmen, JL.; Schulten, K., editors. Reprinted in *Models of neural networks II*. Springer; Berlin: 1981. 1994.
38. Hebb, DO. The organization of behavior, a neuropsychological theory. Wiley; New York: 1949.
39. Park E, Dvorak D, Fenton AA. Ensemble Place Codes in Hippocampus: CA1, CA3, and Dentate Gyrus Place Cells Have Multiple Place Fields in Large Environments. *PLoS ONE*. 2011; 6:e22349. [PubMed: 21789250]
40. Fenton AA, Kao H-Y, Neymotin SA, Olypher AV, Vayntrub Y, Lytton WW, et al. Unmasking the CA1 ensemble place code by exposures to small and large environments: more place cells and multiple, irregularly-arranged, and expanded place fields in the larger space. *J Neurosci*. 2008; 28:11250–11262. [PubMed: 18971467]
41. Panagiotaropoulos TI, Deco G, Kapoor V, Logothetis NK. Neuronal discharges and gamma oscillations explicitly reflect visual consciousness in the lateral prefrontal cortex. *Neuron*. 2012; 74:924–935. [PubMed: 22681695]
42. Fukushima M, Saunders RC, Leopold DA, Mishkin M, Averbach BB. Spontaneous high-gamma band activity reflects functional organization of auditory cortex in the awake macaque. *Neuron*. 2012; 74:899–910. [PubMed: 22681693]
43. Reynolds JH, Chelazzi L, Desimone R. Competitive mechanisms subserve attention in macaque areas V2 and V4. *J Neurosci*. 1999; 19:1736–1753. [PubMed: 10024360]
44. Zhang K. Representation of spatial orientation by the intrinsic dynamics of the head-direction cell ensemble: a theory. *J Neurosci*. 1996; 16:2112–2126. [PubMed: 8604055]
45. Samsonovich A, McNaughton BL. Path integration and cognitive mapping in a continuous attractor neural network model. *J Neurosci*. 1997; 17:5900–5920. [PubMed: 9221787]
46. McNaughton BL, Battaglia FP, Jensen O, Moser EI, Moser MB. Path integration and the neural basis of the 'cognitive map'. *Nat Rev Neurosci*. 2006; 7:663–678. [PubMed: 16858394]
47. Buzsaki G, Anastassiou CA, Koch C. The origin of extracellular fields and currents--EEG, ECoG, LFP and spikes. *Nat Rev Neurosci*. 2012; 13:407–420. [PubMed: 22595786]
48. Traub RD, Miles R, Wong RK. Model of the origin of rhythmic population oscillations in the hippocampal slice. *Science*. 1989; 243:1319–1325. [PubMed: 2646715]

49. Brazhnik E, Borgnis R, Muller RU, Fox SE. The effects on place cells of local scopolamine dialysis are mimicked by a mixture of two specific muscarinic antagonists. *J Neurosci.* 2004; 24:9313–9323. [PubMed: 15496667]
50. Antkowiak B, Hentschke H. Cellular mechanisms of gamma rhythms in rat neocortical brain slices probed by the volatile anaesthetic isoflurane. *Neurosci Lett.* 1997; 231:87–90. [PubMed: 9291147]
51. Jefferys JG, Traub RD, Whittington MA. Neuronal networks for induced '40 Hz' rhythms. *Trends Neurosci.* 1996; 19:202–208. [PubMed: 8723208]
52. Traub RD, Whittington MA, Colling SB, Buzsaki G, Jefferys JG. Analysis of gamma rhythms in the rat hippocampus in vitro and in vivo. *J Physiol.* 1996; 493:471–484. Pt 2. [PubMed: 8782110]
53. Whittington MA, Traub RD, Jefferys JG. Synchronized oscillations in interneuron networks driven by metabotropic glutamate receptor activation. *Nature.* 1995; 373:612–615. [PubMed: 7854418]
54. Cardin JA, Carlen M, Meletis K, Knoblich U, Zhang F, Deisseroth K, et al. Driving fast-spiking cells induces gamma rhythm and controls sensory responses. *Nature.* 2009; 459:663–667. [PubMed: 19396156]
55. Daoudal G, Hanada Y, Debanne D. Bidirectional plasticity of excitatory postsynaptic potential (EPSP)-spike coupling in CA1 hippocampal pyramidal neurons. *Proc Natl Acad Sci U S A.* 2002; 99:14512–14517. [PubMed: 12391303]
56. Buzsaki G. Feed-forward inhibition in the hippocampal formation. *Prog Neurobiol.* 1984; 22:131–153. [PubMed: 6433403]
57. Buzsaki G, Eidelberg E. Phase relations of hippocampal projection cells and interneurons to theta activity in the anesthetized rat. *Brain Res.* 1983; 266:334–339. [PubMed: 6191827]
58. Fox SE. Membrane potential and impedance changes in hippocampal pyramidal cells during theta rhythm. *Exp Brain Res.* 1989; 77:283–294. [PubMed: 2792276]
59. Ylinen A, Soltesz I, Bragin A, Penttonen M, Sik A, Buzsaki G. Intracellular correlates of hippocampal theta rhythm in identified pyramidal cells, granule cells, and basket cells. *Hippocampus.* 1995; 5:78–90. [PubMed: 7787949]
60. Lubenov EV, Siapas AG. Hippocampal theta oscillations are travelling waves. *Nature.* 2009; 459:534–539. [PubMed: 19489117]
61. Agarwal G, Stevenson IH, Berenyi A, Mizuseki K, Buzsaki G, Sommer FT. Spatially distributed local fields in the hippocampus encode rat position. *Science.* 2014; 344:626–630. [PubMed: 24812401]
62. Siapas AG, Lubenov EV, Wilson MA. Prefrontal phase locking to hippocampal theta oscillations. *Neuron.* 2005; 46:141–151. [PubMed: 15820700]
63. Jones MW, Wilson MA. Theta rhythms coordinate hippocampal-prefrontal interactions in a spatial memory task. *PLoS Biol.* 2005; 3:e402. [PubMed: 16279838]
64. Uhlhaas PJ, Linden DE, Singer W, Haenschel C, Lindner M, Maurer K, et al. Dysfunctional long-range coordination of neural activity during Gestalt perception in schizophrenia. *J Neurosci.* 2006; 26:8168–8175. [PubMed: 16885230]
65. Verwer RW, Meijer RJ, Van Uum HF, Witter MP. Collateral projections from the rat hippocampal formation to the lateral and medial prefrontal cortex. *Hippocampus.* 1997; 7:397–402. [PubMed: 9287079]
66. Burwell, RD.; Witter, MP. Basic anatomy of the parahippocampal region in monkeys and rats. Oxford University Press; New York: 2002.
67. Adhikari A, Topiwala MA, Gordon JA. Synchronized activity between the ventral hippocampus and the medial prefrontal cortex during anxiety. *Neuron.* 2010; 65:257–269. [PubMed: 20152131]
68. Lubenov EV, Siapas AG. Decoupling through synchrony in neuronal circuits with propagation delays. *Neuron.* 2008; 58:118–131. [PubMed: 18400168]
69. Siapas AG, Wilson MA. Coordinated interactions between hippocampal ripples and cortical spindles during slow-wave sleep. *Neuron.* 1998; 21:1123–1128. [PubMed: 9856467]
70. Schroeder CE, Lakatos P. The gamma oscillation: master or slave? *Brain Topogr.* 2009; 22:24–26. [PubMed: 19205863]
71. Jones MW, Wilson MA. Phase precession of medial prefrontal cortical activity relative to the hippocampal theta rhythm. *Hippocampus.* 2005; 15:867–873. [PubMed: 16149084]

72. Buzsaki G, Eidelberg E. Convergence of associational and commissural pathways on CA1 pyramidal cells of the rat hippocampus. *Brain Res.* 1982; 237:283–295. [PubMed: 7082996]
73. Wesierska M, Dockery C, Fenton AA. Beyond memory, navigation, and inhibition: behavioral evidence for hippocampus-dependent cognitive coordination in the rat. *J Neurosci.* 2005; 25:2413–2419. [PubMed: 15745968]
74. Robbe D, Buzsaki G. Alteration of theta timescale dynamics of hippocampal place cells by a cannabinoid is associated with memory impairment. *J Neurosci.* 2009; 29:12597–12605. [PubMed: 19812334]
75. Robbe D, Montgomery SM, Thome A, Rueda-Orozco PE, McNaughton BL, Buzsaki G. Cannabinoids reveal importance of spike timing coordination in hippocampal function. *Nat Neurosci.* 2006; 9:1526–1533. [PubMed: 17115043]
76. Harris KD, Csicsvari J, Hirase H, Dragoi G, Buzsaki G. Organization of cell assemblies in the hippocampus. *Nature.* 2003; 424:552–556. [PubMed: 12891358]
77. Skaggs WE, McNaughton BL, Wilson MA, Barnes CA. Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus.* 1996; 6:149–172. [PubMed: 8797016]
78. Dragoi G, Buzsaki G. Temporal encoding of place sequences by hippocampal cell assemblies. *Neuron.* 2006; 50:145–157. [PubMed: 16600862]
79. Kucewicz MT, Tricklebank MD, Bogacz R, Jones MW. Dysfunctional prefrontal cortical network activity and interactions following cannabinoid receptor activation. *J Neurosci.* 2011; 31:15560–15568. [PubMed: 22031901]
80. Katona I, Sperlagh B, Sik A, Kafalvi A, Vizi ES, Mackie K, et al. Presynaptically located CB1 cannabinoid receptors regulate GABA release from axon terminals of specific hippocampal interneurons. *J Neurosci.* 1999; 19:4544–4558. [PubMed: 10341254]
81. Katona I, Rancz EA, Acsady L, Ledent C, Mackie K, Hajos N, et al. Distribution of CB1 cannabinoid receptors in the amygdala and their role in the control of GABAergic transmission. *J Neurosci.* 2001; 21:9506–9518. [PubMed: 11717385]
82. Karson MA, Tang AH, Milner TA, Alger BE. Synaptic cross talk between perisomatic-targeting interneuron classes expressing cholecystokinin and parvalbumin in hippocampus. *J Neurosci.* 2009; 29:4140–4154. [PubMed: 19339609]
83. Reich CG, Karson MA, Karnup SV, Jones LM, Alger BE. Regulation of IPSP theta rhythm by muscarinic receptors and endocannabinoids in hippocampus. *J Neurophysiol.* 2005; 94:4290–4299. [PubMed: 16093334]
84. Katona I, Freund TF. Endocannabinoid signaling as a synaptic circuit breaker in neurological disease. *Nat Med.* 2008; 14:923–930. [PubMed: 18776886]
85. Gandal MJ, Edgar JC, Klook K, Siegel SJ. Gamma synchrony: Towards a translational biomarker for the treatment-resistant symptoms of schizophrenia. *Neuropharmacology.* 2011
86. Javitt DC, Jotkowitz A, Sircar R, Zukin SR. Non-competitive regulation of phencyclidine/sigma-receptors by the N-methyl-D-aspartate receptor antagonist D-(-)-2- amino-5-phosphonovaleric acid. *Neurosci Lett.* 1987; 78:193–198. [PubMed: 2888059]
87. Anis NA, Berry SC, Burton NR, Lodge D. The dissociative anaesthetics, ketamine and phencyclidine, selectively reduce excitation of central mammalian neurones by N-methyl-aspartate. *Br J Pharmacol.* 1983; 79:565–575. [PubMed: 6317114]
88. Cohen BD, Rosenbaum G, Luby ED, Gottlieb JS. Comparison of phencyclidine hydrochloride (Sernyl) with other drugs. Simulation of schizophrenic performance with phencyclidine hydrochloride (Sernyl), lysergic acid diethylamide (LSD-25), and amobarbital (Amytal) sodium; II. Symbolic and sequential thinking. *Arch Gen Psychiatry.* 1962; 6:395–401. [PubMed: 13880223]
89. Itil T, Keskiner A, Kiremitci N, Holden JM. Effect of phencyclidine in chronic schizophrenics. *Can Psychiatr Assoc J.* 1967; 12:209–212. [PubMed: 6040448]
90. Abbott A. Schizophrenia: The drug deadlock. *Nature.* 2010; 468:158–159. [PubMed: 21068804]
91. Carter CS, Barch DM. Cognitive neuroscience-based approaches to measuring and improving treatment effects on cognition in schizophrenia: the CNTRICS initiative. *Schizophr Bull.* 2007; 33:1131–1137. [PubMed: 17630405]

92. Lakatos P, Schroeder CE, Leitman DI, Javitt DC. Predictive suppression of cortical excitability and its deficit in schizophrenia. *J Neurosci*. 2013; 33:11692–11702. [PubMed: 23843536]
93. Palmer BW, Heaton RK, Paulsen JS, Kuck J, Braff D, Harris MJ, et al. Is it possible to be schizophrenic yet neuropsychologically normal? *Neuropsychology*. 1997; 11:437–446. [PubMed: 9223148]
94. Morice R, Delahunty A. Frontal/executive impairments in schizophrenia. *Schizophr Bull*. 1996; 22:125–137. [PubMed: 8685655]
95. Lewis DA, Gonzalez-Burgos G. Pathophysiologically based treatment interventions in schizophrenia. *Nat Med*. 2006; 12:1016–1022. [PubMed: 16960576]
96. Breier A, Malhotra AK, Pinals DA, Weisenfeld NI, Pickar D. Association of ketamine-induced psychosis with focal activation of the prefrontal cortex in healthy volunteers. *Am J Psychiatry*. 1997; 154:805–811. [PubMed: 9167508]
97. Vollenweider FX, Leenders KL, Scharfetter C, Antonini A, Maguire P, Missimer J, et al. Metabolic hyperfrontality and psychopathology in the ketamine model of psychosis using positron emission tomography (PET) and [18F]fluorodeoxyglucose (FDG). *Eur Neuropsychopharmacol*. 1997; 7:9–24. [PubMed: 9088881]
98. Holcomb HH, Lahti AC, Medoff DR, Cullen T, Tamminga CA. Effects of noncompetitive NMDA receptor blockade on anterior cingulate cerebral blood flow in volunteers with schizophrenia. *Neuropsychopharmacology*. 2005; 30:2275–2282. [PubMed: 16034443]
99. Rowland LM, Bustillo JR, Mullins PG, Jung RE, Lenroot R, Landgraf E, et al. Effects of ketamine on anterior cingulate glutamate metabolism in healthy humans: a 4-T proton MRS study. *Am J Psychiatry*. 2005; 162:394–396. [PubMed: 15677610]
100. Homayoun H, Moghaddam B. NMDA receptor hypofunction produces opposite effects on prefrontal cortex interneurons and pyramidal neurons. *J Neurosci*. 2007; 27:11496–11500. [PubMed: 17959792]
101. Jackson ME, Homayoun H, Moghaddam B. NMDA receptor hypofunction produces concomitant firing rate potentiation and burst activity reduction in the prefrontal cortex. *Proc Natl Acad Sci U S A*. 2004; 101:8467–8472. [PubMed: 15159546]
102. Wood J, Kim Y, Moghaddam B. Disruption of prefrontal cortex large scale neuronal activity by different classes of psychotomimetic drugs. *J Neurosci*. 2012; 32:3022–3031. [PubMed: 22378875]
103. Dvorak D, Fenton AA. Toward a proper estimation of phase-amplitude coupling in neural oscillations. *J Neurosci Methods*. 2014; 225C:42–56. [PubMed: 24447842]
104. Belluscio MA, Mizuseki K, Schmidt R, Kempter R, Buzsaki G. Cross-frequency phase-phase coupling between theta and gamma oscillations in the hippocampus. *J Neurosci*. 2012; 32:423–435. [PubMed: 22238079]
105. Ray S, Maunsell JH. Different origins of gamma rhythm and high-gamma activity in macaque visual cortex. *PLoS Biol*. 2011; 9:e1000610. [PubMed: 21532743]
106. Phillips KG, Cotel MC, McCarthy AP, Edgar DM, Tricklebank M, O'Neill MJ, et al. Differential effects of NMDA antagonists on high frequency and gamma EEG oscillations in a neurodevelopmental model of schizophrenia. *Neuropharmacology*. 2011; 62:1359–1370. [PubMed: 21521646]
107. Kuang H, Lin L, Tsien JZ. Temporal Dynamics of Distinct CA1 Cell Populations during Unconscious State Induced by Ketamine. *PLoS One*. 2010; 5:e15209. [PubMed: 21165147]
108. Klausberger T, Somogyi P. Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science*. 2008; 321:53–57. [PubMed: 18599766]
109. Somogyi P, Klausberger T. Defined types of cortical interneurone structure space and spike timing in the hippocampus. *J Physiol*. 2005; 562:9–26. [PubMed: 15539390]
110. Klausberger T, Magill PJ, Marton LF, Roberts JD, Cobden PM, Buzsaki G, et al. Brain-state- and cell-type-specific firing of hippocampal interneurons in vivo. *Nature*. 2003; 421:844–848. [PubMed: 12594513]
111. Laszotz B, Klausberger T. Layer-Specific GABAergic Control of Distinct Gamma Oscillations in the CA1 Hippocampus. *Neuron*. 2014; 81:1126–1139. [PubMed: 24607232]

112. Klausberger T. GABAergic interneurons targeting dendrites of pyramidal cells in the CA1 area of the hippocampus. *Eur J Neurosci.* 2009; 30:947–957. [PubMed: 19735288]
113. Hartwich K, Pollak T, Klausberger T. Distinct firing patterns of identified basket and dendrite-targeting interneurons in the prefrontal cortex during hippocampal theta and local spindle oscillations. *J Neurosci.* 2009; 29:9563–9574. [PubMed: 19641119]
114. Karayiorgou M, Gogos JA, Galke BL, Jeffery JA, Nestadt G, Wolynec PS, et al. Genotype and phenotype analysis at the 22q11 schizophrenia susceptibility locus. *Cold Spring Harb Symp Quant Biol.* 1996; 61:835–843. [PubMed: 9246508]
115. Antshel KM, Aneja A, Strunge L, Peebles J, Fremont WP, Stallone K, et al. Autistic spectrum disorders in velo-cardio facial syndrome (22q11.2 deletion). *J Autism Dev Disord.* 2007; 37:1776–1786. [PubMed: 17180713]
116. Vorstman JA, Morcus ME, Duijff SN, Klaassen PW, Heineman-de Boer JA, Beemer FA, et al. The 22q11.2 deletion in children: high rate of autistic disorders and early onset of psychotic symptoms. *J Am Acad Child Adolesc Psychiatry.* 2006; 45:1104–1113. [PubMed: 16926618]
117. Karayiorgou M, Gogos JA. The molecular genetics of the 22q11-associated schizophrenia. *Brain Res Mol Brain Res.* 2004; 132:95–104. [PubMed: 15582150]
118. Sigurdsson T, Stark KL, Karayiorgou M, Gogos JA, Gordon JA. Impaired hippocampal-prefrontal synchrony in a genetic mouse model of schizophrenia. *Nature.* 2010; 464:763–767. [PubMed: 20360742]
119. Adhikari A, Topiwala MA, Gordon JA. Single units in the medial prefrontal cortex with anxiety-related firing patterns are preferentially influenced by ventral hippocampal activity. *Neuron.* 2011; 71:898–910. [PubMed: 21903082]
120. Gonzalez-Burgos G, Hashimoto T, Lewis DA. Alterations of cortical GABA neurons and network oscillations in schizophrenia. *Curr Psychiatry Rep.* 2010; 12:335–344. [PubMed: 20556669]
121. Lewis DA, Moghaddam B. Cognitive dysfunction in schizophrenia: convergence of gamma-aminobutyric acid and glutamate alterations. *Arch Neurol.* 2006; 63:1372–1376. [PubMed: 17030651]
122. Belforte JE, Zsiros V, Sklar ER, Jiang Z, Yu G, Li Y, et al. Postnatal NMDA receptor ablation in corticolimbic interneurons confers schizophrenia-like phenotypes. *Nat Neurosci.* 2010; 13:76–83. [PubMed: 19915563]
123. Nakao K, Nakazawa K. Brain state-dependent abnormal LFP activity in the auditory cortex of a schizophrenia mouse model. *Front Neurosci.* 2014; 8:168. [PubMed: 25018691]
124. Gerber DJ, Hall D, Miyakawa T, Demars S, Gogos JA, Karayiorgou M, et al. Evidence for association of schizophrenia with genetic variation in the 8p21.3 gene, PPP3CC, encoding the calcineurin gamma subunit. *Proc Natl Acad Sci U S A.* 2003; 100:8993–8998. [PubMed: 12851458]
125. Zeng H, Chattarji S, Barbarosie M, Rondi-Reig L, Philpot BD, Miyakawa T, et al. Forebrain-specific calcineurin knockout selectively impairs bidirectional synaptic plasticity and working/episodic-like memory. *Cell.* 2001; 107:617–629. [PubMed: 11733061]
126. Miyakawa T, Leiter LM, Gerber DJ, Gainetdinov RR, Sotnikova TD, Zeng H, et al. Conditional calcineurin knockout mice exhibit multiple abnormal behaviors related to schizophrenia. *Proc Natl Acad Sci U S A.* 2003; 100:8987–8992. [PubMed: 12851457]
127. Behrens CJ, van den Boom LP, de Hoz L, Friedman A, Heinemann U. Induction of sharp wave-ripple complexes in vitro and reorganization of hippocampal networks. *Nat Neurosci.* 2005; 8:1560–1567. [PubMed: 16222227]
128. Suh J, Foster DJ, Davoudi H, Wilson MA, Tonegawa S. Impaired hippocampal ripple-associated replay in a mouse model of schizophrenia. *Neuron.* 2013; 80:484–493. [PubMed: 24139046]
129. Gardner RJ, Kersante F, Jones MW, Bartsch U. Neural oscillations during non-rapid eye movement sleep as biomarkers of circuit dysfunction in schizophrenia. *Eur J Neurosci.* 2014; 39:1091–1106. [PubMed: 24712989]
130. Csicsvari J, O'Neill J, Allen K, Senior T. Place-selective firing contributes to the reverse-order reactivation of CA1 pyramidal cells during sharp waves in open-field exploration. *Eur J Neurosci.* 2007; 26:704–716. [PubMed: 17651429]

131. Foster DJ, Wilson MA. Reverse replay of behavioural sequences in hippocampal place cells during the awake state. *Nature*. 2006; 440:680–683. [PubMed: 16474382]
132. Buzsaki G. Memory consolidation during sleep: a neurophysiological perspective. *J Sleep Res*. 1998; 7(Suppl 1):17–23. [PubMed: 9682189]
133. Singer AC, Carr MF, Karlsson MP, Frank LM. Hippocampal SWR activity predicts correct decisions during the initial learning of an alternation task. *Neuron*. 2013; 77:1163–1173. [PubMed: 23522050]
134. Jadhav SP, Kemere C, German PW, Frank LM. Awake hippocampal sharp-wave ripples support spatial memory. *Science*. 2012; 336:1454–1458. [PubMed: 22555434]
135. Insel TR. Rethinking schizophrenia. *Nature*. 2010; 468:187–193. [PubMed: 21068826]
136. Lewis DA, Gonzalez-Burgos G. Neuroplasticity of neocortical circuits in schizophrenia. *Neuropsychopharmacology*. 2008; 33:141–165. [PubMed: 17805309]
137. Subramaniam K, Luks TL, Fisher M, Simpson GV, Nagarajan S, Vinogradov S. Computerized cognitive training restores neural activity within the reality monitoring network in schizophrenia. *Neuron*. 2012; 73:842–853. [PubMed: 22365555]
138. Lee H, Dvorak D, Kao HY, Duffy AM, Scharfman HE, Fenton AA. Early cognitive experience prevents adult deficits in a neurodevelopmental schizophrenia model. *Neuron*. 2012; 75:714–724. [PubMed: 22920261]
139. Anguera JA, Boccanfuso J, Rintoul JL, Al-Hashimi O, Faraji F, Janowich J, et al. Video game training enhances cognitive control in older adults. *Nature*. 2013; 501:97–101. [PubMed: 24005416]
140. Javitt DC, Spencer KM, Thaker GK, Winterer G, Hajos M. Neurophysiological biomarkers for drug development in schizophrenia. *Nat Rev Drug Discov*. 2008; 7:68–83. [PubMed: 18064038]
141. Luck SJ, Mathalon DH, O'Donnell BF, Hamalainen MS, Spencer KM, Javitt DC, et al. A roadmap for the development and validation of event-related potential biomarkers in schizophrenia research. *Biol Psychiatry*. 2010; 70:28–34. [PubMed: 21111401]

Figure 1.

Two classes of neural code. A) Dedicated codes represent information in a form that is analogous to a traffic light. Like the red like signals ‘stop’, whenever a neuron discharges it unambiguously represents the same information. B) Ensemble codes represent information analogous to a jumbotron, which represents information in the pattern of it many on and off lights. C) The discharge of hippocampus place cells is localized to single places in a familiar 70-cm cylinder (top and bottom rows) consistent with a dedicated code. However, in a large 1.4×1.5 m environment, the same cells now discharge in multiple locations. The color-coded firing rate maps range from blue = 0 AP/s to red. The red rate is given for each map. Adapted from reference (40).

Figure 2.

Winner-take-all competitive networks illustrate how excitation-inhibition interactions can organize neural network activity into coordinated network patterns. A) The network is configured such that there is relatively strong mutual excitation amongst subgroups of pyramidal cells and these subgroups drive feed forward inhibitory neurons. Because of this configuration, the subgroup that receives more excitation will out compete the other subgroup, causing it to suppress. B,C) Network simulations illustrate the network behavior. Each blue-to-red color-coded circle represents the collective activity of a 632 place cell network divided into two subnets, one for room locations (upper row) the other for arena (lower row) locations. The cells representing nearby locations were mutually exciting and cells representing far apart locations were less mutually exciting. Each cell generates global inhibition proportional to its current activity. The initial activity of all cells is low (blue) until a stimulus corresponding to a location in the room or arena occurs during the time of the second plot. Network activity persists after the stimulus is removed. B) The network first represents a room location then switches to an arena location in correspondence with the stimulation (square and circle) C) The network is simultaneously presented with separate room and arena stimulus locations, with the room input slightly stronger. Both the room and arena subnets initially activate but the room subnet predominates and the other is suppressed. D) The same network is corrupted by randomly increasing connections 0-25%. The most significant impact is to increase the connections between the weakly connected cells. After the network corruption E) responses to the initial stimulus location are normal, but the network fails to switch to represent an arena location. The corrupted response is to represent both the room and arena locations, amounting to a nonsense representation and an example of the superposition catastrophe. F) Similarly, the corrupted network fails to represent either of the room or arena locations when the two inputs are simultaneous. Adapted from reference (26).

Figure 3.

Neural discoordination following disinhibition of one hippocampus by unilateral injection of tetrodotoxin (TTX) into the other hippocampus. A) Rasters of an ensemble recording and B) average firing rate changes before and after TTX injection into the contralateral hippocampus. The injection transiently increased pyramidal (pyr) cell discharge without changing inhibitory interneuron (inh) discharge, consistent with disinhibition. C) Excitation-inhibition coupling was transiently reduced by the TTX injection when pyramidal cell rates increased. Coactivity of pyramidal cell – interneuron pairs was selectively reduced if the pairs were strongly correlated before the contralateral TTX injection. D) Cross-correlograms of individual cell pairs illustrating that 1 h after TTX injection, the coactivity of pyramidal cell pairs (left) that had previously discharged together did not change (right) whereas coactivity increased if the pair had previously only rarely discharged together. E) Summary of how initially weakly and strongly correlated cell pairs changed before and after TTX. F) 1 h after TTX injection, (left) Room+Arena– place avoidance with high cognitive control demand was impaired but not (right) Room+ place avoidance with low cognitive control demand. In both task variants the arena rotates at 1 rpm and the shock zone is defined in stationary room coordinates. The dry area has salient but irrelevant rotating olfactory cues to actively ignore in the Room+Arena– task variant, whereas the irrelevant cues are attenuated by shallow water in the Room+ task variant. Adapted from references (26, 73).