MEETING ABSTRACT

Open Access

Arthroscopic resection of wrist ganglia

Christophe Mathoulin

From 10th Congress of the Asia-Pacific Federation of Societies of Surgery for the Hand and the 6th Congress of Asia-Pacific Federation of Societies of Hand Therapists
Kuala Lumpur, Malaysia. 2-4 October 2014

Introduction

The purpose of this study is to analyse the rate of recurrences of the dorsal ganglion resected by trans-cystic arthroscopic technique. All patients were operated on by a single surgeon with a minimum follow-up of 2 years.

Material and methods

We report a series of 114 patients with dorsal ganglia (87 women and 27 men). The average age was 33.2 years (range: 12 to 63 years). All the patients had had an unsuccessful preliminary treatment. Although a minority of patients (32 cases) complained of pain, aesthetic motivation comprised the majority of cases (82 patients).

All the patients were operated under local-regional anaesthesia on an outpatient basis. The shaver was placed directly in trans-cystic, mostly into the midcarpal joint, to perform the capsulectomy and resection of the wall of cyst. The patients were allowed to mobilise their wrist freely immediately post-operatively.

Results

Our average follow-up was 42.3 months (between 24 and 84 months). The functional clinical data (mobility, pain and strength) were improved in a statistically significant way. We had 14 recurrences (12 %) which appeared after an average period of 16.8 months (between 2 and 35 months). The cyst was again removed by wrist arthroscopy in 11 cases at the request of the patients without any particular problem. We had no other complications

Conclusion

The arthroscopic resection of dorsal ganglia of the wrist is a technique which allows a satisfactory resection of the cyst and the adjacent articular capsule. The rate of recurrence appears significant initially but it is within comparable levels to the best series of classical open

techniques. The long follow-up of this series addresses the notion of true recurrence or return of a new cyst. Nevertheless the rate of satisfaction of the patients is very high particularly because there are no requirements for cutaneous suture or immobilization of the wrist.

Published: 19 May 2015

doi:10.1186/1753-6561-9-S3-A31

Cite this article as: Mathoulin: Arthroscopic resection of wrist ganglia. *BMC Proceedings* 2015 **9**(Suppl 3):A31.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

Institut de la Main, 75016, Paris, France

© 2015 Mathoulin; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.