Published in final edited form as:

Org Lett. 2013 August 16; 15(16): 4070–4073. doi:10.1021/ol401600j.

Atom Economical, One-Pot, Three Reaction Cascade to Novel Tricyclic 2,4-dihydro-1*H*-benzo[*f*]isochromenes

Robert J. Hinkle and Shane E. Lewis

Department of Chemistry, The College of William & Mary, P.O. Box 8795, Williamsburg, VA 23187-8795

Robert J. Hinkle: rjhink@wm.edu

Abstract

Reaction of 6-methyl-1-phenylhept-3-yn-2,6-diol with various aldehydes under Lewis acid conditions provides an atom economical, two-component cascade reaction sequence to novel 2,4-dihydro-1*H*-benzo[*f*]isochromene compounds. Aliphatic aldehydes as well as electron-deficient and electron-rich aromatic aldehydes can be used.

In efforts to increase atom economy,¹ reduce harmful environmental effects, and increase overall efficiency of experimental sequences, one-pot multi-component reactions² and cascade³ or domino reactions⁴ have become much more common in organic synthesis.^{5,6} We have recently described a three-component addition-cyclization protocol⁷ as well as a mechanistic investigation of a Bi(III)-initiated silyl-Prins cyclization toward dihydropyrans.⁸ Herein, we report a cascade sequence involving an alkynyl-Prins cyclization, Friedel-Crafts arylation, and dehydration/aromatization to afford unique tricyclic benzo[*f*]isochromene derivatives⁹ in which only H₂O is generated as a byproduct from the two reaction components. A minimum of 72% average yields are observed for each of the reactions involved in the cascade.

The Prins cyclization^{10,11} and related reactions^{12,13} are versatile methods for the synthesis of dihydropyran and tetrahydropyran rings as well as numerous derivatives. Recently, a tandem intermolecular Prins/Friedel-Crafts reaction was reported to furnish 4-aryldihydropyrans from homopropargylic alcohols and 4-aryltetrahydropyrans from homoallylic alcohols (Scheme 1) when the aromatic compound was used as solvent.¹⁴ An

Correspondence to: Robert J. Hinkle, rjhink@wm.edu.

Supporting Information Available X-ray crystallographic tables of lactone **4** and characterization data, including ¹H and ¹³C APT, and ¹³C DEPT NMR spectra for **2a-2c** and **3a-3j**; GC MS traces also included for **3a-3j**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

intramolecular variation of this reaction with homoallylic alcohols provided a number of tricyclic compounds via an intermediate secondary alkyl carbocation.¹⁵ While a number of fused polycyclic pyran derivatives exist in the literature, this represented a novel approach to their construction, as many other examples use substrates with a preformed ring system.¹⁶ In contrast to these results, the reaction we describe occurs via a putative, high-energy alkenyl cation and includes a subsequent dehydrative aromatization to afford 2,4-dihydro-1*H*-benzo[*f*]isochromenes, which also represent a novel class of heterotricycles.

Based on our previous work, we envisioned that a tricyclic 2,4,5,6-tetrahydro-1*H*-benzo[*f*]isochromen-5-ol product might be accessible utilizing a cascade sequence involving an alkynyl-Prins reaction^{17,18} and Friedel-Crafts cyclization¹⁹ from an unsymmetrical alkynediol and an aldehyde (*R*-CHO) under Lewis acid mediated conditions (Scheme 2).

The requisite unprotected alkynediol **2a** was synthesized from the known propargylic alcohol **1a**²⁰ by conversion to the dianion with *n*-Bu-Li followed by addition to isobutylene oxide in the presence of BF₃•OEt₂ (eq 1).²¹

(1)

Several different Lewis acid conditions were examined with alkynediol **2a** and propionaldehyde or isobutyraldehyde as electrophiles (Table 1). When using propionaldehyde (entry 1) relatively low yields were obtained with 1.1 equiv of BF₃•OEt₂ (average of 72% per reaction). Reactions with the silylated derivative **2b** (eq 1) and stoichiometric Bi(OTf)₃ made isolation of **3a** more difficult (entry 2). We also attempted cyclizations with the *mono*-TBDPS protected analog **2c**, but recovered starting materials. Reactions using isobutyraldehyde were more efficient (entries 3–6), although stoichiometric amounts of BiBr₃ and Bi(OTf)₃ provided **3b** in lower yields (entries 4 and 5). Halving the concentration of the reaction had no effect on isolated yield (entry 6).

As a result of our success with catalytic quantities of Bi(III) compounds in the synthesis of dihydropyrans, catalytic quantities (10–20%) of BiBr₃ and Bi(OTf)₃ were screened as potential initiators,^{7,8} but starting materials were largely recovered. Although the reaction did proceed with stoichiometric amounts of both reagents (entries 2, 4 and 5), the high molecular weight and associated cost limited the appeal of each. Other, less expensive Lewis acids such as TiCl₄ and SnCl₄ were examined in catalytic and stoichiometric quantities, but each was more hazardous to dispense and led to more complex mixtures as well as decomposition of starting materials. Overall, a 10 mol% excess of BF₃•OEt₂ over diol **2a** provided the least complex reaction mixtures (entries 1 and 3).

Alkynediol **2a** was then reacted with various other aldehydes in the presence of 1.1 equiv of $\text{BF}_3 \cdot \text{OEt}_2$ at room-temperature in CH_2Cl_2 (Table 2). Compounds **3a-3j** were all isolated as extremely viscous oils after column chromatography and were characterized by IR, ^1H , ^{13}C APT, and ^{13}C DEPT NMR spectroscopy as well as combustion or mass spectral analysis.

Alkyl-substituted products, **3a-3e**, were generally less stable than the corresponding aryl-substituted analogs, **3f-3j**, and decomposed at room temperature when exposed to air. As reported by Guiso et al., analogs with only one aromatic ring (isochromans) are easily oxidized to lactones.²² Based on X-ray crystallographic analysis of a sample left at rt in air, the alkyl 2,4-dihydro-1*H*-benzo[*f*]isochromenes **3a-3f** herein appear to be prone to oxidation via a similar pathway (Scheme 3).²³

Reactions using benzaldehyde and substituted benzaldehydes afforded benzo[*f*]isochromenes (**3f-3j**) containing both electron-withdrawing and electron-donating substituents. These aryl-substituted benzo[*f*]isochromenes were significantly more stable than the corresponding alkyl-substituted products **3a-3e**. This trend was also verified by GC MS analysis in which the molecular ions of the aryl-substituted substrates **3f-3j** were much more evident relative to fragments.

We believe that the reaction progresses through a sequence of steps outlined in Scheme 4. Either of the two hydroxyls in **2a** could attack the putative Lewis acid-activated aldehyde generating intermediate **I** or **IV** via *Path A* or *Path B*, respectively (Scheme 4). Subsequent formation of oxocarbenium ions **II** or **V** then allows for progression of the cascade. Intramolecular 6-*endo* attack of the alkyne via *Path A* would generate **III**, the alkenyl cation of a dihydropyran, whereas intramolecular attack of the alkyne via *Path B* would generate **VI**, the vinyl cation of a dihydrofuran. Classic studies conducted by Mayr et al. established that cyclic, five-membered alkenyl cations are approximately 10 kcal/mol higher in energy than cyclic, six-membered analogs, which are in turn similar to open-chain alkenyl cations.²⁴ In accord with this difference in energies, solvolyses of cyclopentenyl nonaflates occur by S-O bond cleavage rather than alkenyl cation formation.²⁵ As predicted by these energetic differences, we do not observe products from cyclization by the higher energy manifold (*Path B*); however, some complex polymerization products are generated, and we cannot completely discount this intermediate in formation of those materials. In *Path A*, a subsequent intramolecular Friedel-Crafts reaction would occur at the *ortho*- position of the pendant aromatic ring. After re-aromatization of the original phenyl moiety, the central ring is formed and acid-catalyzed dehydration/aromatization²⁶ occurs to generate the observed benzo[*f*]isochromenes **3a-3j**. In the case of protected alkynediol **2c** (vide supra), the large TBDPS group likely prevented cyclization due to severe steric interactions, and the initial intermolecular addition to **I** is reversible (Scheme 4, *Path A*).

It is somewhat surprising that tetrahydrofuran products are not observed. Rychnovsky and co-workers²⁷ as well as Cho and co-workers²⁸ reported that internal alkynes afford 5-*exo* products rather than the 6-*endo* adducts we observe. However, a 5-*exo* reaction of intermediate **II** would produce **VII** (Scheme 4, lower right), which would contain an alkenyl cation that would be destabilized by the adjacent hydroxyl group. Therefore, the products formed proceed through *Path A*.

In summary, 2,4-dihydro-1*H*-benzo[*f*]isochromenes **3a-3j** are novel tricyclic compounds that are obtained in four chemical steps from commercially-available trimethylsilylacetylene. The only ring present in the initial compound is the phenyl moiety shown in compounds **1a-1c** and **2a-2c**. Conversion of **2a** to the tricyclic products is achieved by an atom economical cascade involving an alkynyl-Prins cyclization, Friedel-Crafts arylation, and dehydrative aromatization. Although overall yields appear modest, three separate reactions are part of the cascade, and the average yields for each of these reactions involved in the cascade are a minimum of 72%. Further studies involving a wider variety of substrates and further optimization are underway and will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the AREA Program of the National Institutes of Health (RGM072525-01A2) as well as the National Science Foundation (CHE-1012305) for support of this work. We also thank Robert D. Pike, PhD, for X-ray structure analysis of lactone **4**.

References

1. Sheldon RA. Chem Soc Rev. 2012; 41:1437–1451. [PubMed: 22033698] Trost BM. Handbook of Green Chemistry. 2012; 7:1–33. Trost BM. Angew Chem Int Ed, Engl. 1995; 34:259–281.
2. Touré BB, Hall DG. Chem Rev. 2009; 109:4439–4486. [PubMed: 19480390]
3. Kleinke AS, Webb D, Jamison TF. Tetrahedron. 2012; 68:6999–7018. Nicolaou KC, Chen JS. Chem Soc Rev. 2009; 38:2993–3009. [PubMed: 19847336]
4. Tietze, L-F. Domino Reactions in Organic Synthesis. Wiley-VCH; Weinheim: 2006.
5. For reactions described in the literature as cascade reactions, see: Horning BD, MacMillan DWC. J Am Chem Soc. 2013; 135:6442–6445. [PubMed: 23586842] Ohno H. Asian J Org Chem. 2013; 2:18–28. Pintado-Sierra M, Rasero-Almansa AM, Corma A, Iglesias M, Sánchez F. J Cat. 2013; 299:137–145. Surendra K, Qiu W, Corey EJ. J Am Chem Soc. 2011; 133:9724–9726. [PubMed: 21644499] Li H, Loh TP. Org Lett. 2010; 12:2679–2681. [PubMed: 20481623] Dalgard JE, Rychnovsky SD. Org Lett. 2005; 7:1589–1591. [PubMed: 15816759]
6. For reactions described as domino sequences, see: Liu XL, Han WY, Zhang XM, Yuan WC. Org Lett. 2013; 15:1246–1249. [PubMed: 23452297] Long Y, She Z, Liu X, Chen Y. J Org Chem. 2013; 78:2579–2588. [PubMed: 23438089] Lu BZ, Wei HX, Zhang Y, Zhao W, Dufour M, Li G, Farina V, Senanayake CH. J Org Chem. 2013; 78:4558–4562. [PubMed: 23544431]
7. Hinkle RJ, Lian Y, Speight LC, Stevenson HE, Sprachman MM, Katkish LA, Mattern MC. Tetrahedron. 2009; 65:6834–6839. [PubMed: 20161364]
8. Lambert RF, Hinkle RJ, Ammann SE, Lian Y, Liu J, Lewis SE, Pike RD. J Org Chem. 2011; 76:9269–9277. [PubMed: 21916500]
9. Fused pyrans have been investigated as antioxidants and for anti-platelet and antiproliferative properties: Kishore KRKR, Longato GB, de CJE, Ruiz ALTG, Silva LF Jr. Molecules. 2012; 17:9621–9630. [PubMed: 23442985] Rafinejad A, Fallah-Tafti A, Tiwari R, Shirazi AN, Mandal D, Shafiee A, Parang K, Foroumadi A, Akbarzadeh T. Daru, J Pharm Sci. 2012; 20:100. Guiso M, Marra C, Arcos RR. Nat Prod Res. 2008; 22:1403–1409. [PubMed: 19023801] Lorenz P, Zeh M, Martens-Lobenhoffer J, Schmidt H, Wolf G, Horn TFW. Free Rad Res. 2005; 39:535–545.
10. For recent reviews on the Prins reaction and Prins cyclization in organic synthesis, see: Han X, Peh G, Floreancig PE. Eur J Org Chem. 2013:1193–1208. Pastor IM, Yus M. Curr Org Chem. 2012; 16:1277–1312.

11. For mechanistic studies on the Prins reaction, see: Ogawa Y, Painter PP, Tantillo DJ, Wender PA. *J Org Chem.* 2013; 78:104–115. [PubMed: 23121542] Zheng K, Liu X, Qin S, Xie M, Lin L, Hu C, Feng X. *J Am Chem Soc.* 2012; 134:17564–17573. [PubMed: 23030638] Chio FK, Warne J, Gough D, Penny M, Green S, Coles SJ, Hursthouse MB, Jones P, Hassall L, McGuire TM, Dobbs AP. *Tetrahedron.* 2011; 67:5107–5124. Gesinski MR, Tadpetch K, Rychnovsky SD. *Org Lett.* 2009; 11:5342–5345. [PubMed: 19873984] Gesinski MR, Van OLJ, Rychnovsky SD. *Synlett.* 2008:363–366. [PubMed: 20936058] Tadpetch K, Rychnovsky SD. *Org Lett.* 2008; 10:4839–4842. [PubMed: 18816133] Jasti R, Rychnovsky SD. *J Am Chem Soc.* 2006; 128:13640–13648. [PubMed: 17031979] Jasti R, Rychnovsky SD. *Org Lett.* 2006; 8:2175–2178. [PubMed: 16671810] Jasti R, Anderson CD, Rychnovsky SD. *J Am Chem Soc.* 2005; 127:9939–9945. [PubMed: 15998101]
12. The “silyl-Prins” reaction has been studied extensively: Hemery T, Wibbeling B, Froehlich R, Hoppe D. *Synthesis.* 2010:329–342. Meilert K, Brimble MA. *Org Biomol Chem.* 2006; 4:2184–2192. [PubMed: 16729128] Dobbs AP, Guesne SJJ. *Synlett.* 2005:2101–2103. Semeyn C, Blaaup RH, Hiemstra H, Speckamp WN. *J Org Chem.* 1997; 62:3426–3427.
13. The intramolecular silyl-mediated Sakurai reaction (ISMS) involves the same intermediate cation as the silyl-Prins reaction: Markó IE, Plancher JM. *Tetrahedron Lett.* 1999; 40:5259–5262. Markó IE, Bayston DJ. *Tetrahedron.* 1994; 50:7141–7156.
14. Reddy UC, Saikia AK. *Synlett.* 2010:1027–1032.
15. Subba Reddy BV, Kumar H, Borkar P, Yadav JS, Sridhar B. *Eur J Org Chem.* 2013:1993–1999. Reddy BVS, Jalal S, Borkar P, Yadav JS, Reddy PP, Kunwar AC, Sridhar B. *Org Biomol Chem.* 2012; 10:6562–6568. [PubMed: 22763607]
16. Guo K, Chen Y. *Mater Chem Phys.* 2013; 137:1062–1066. Schevenels F, Tinant B, Declercq JP, Markó IE. *Chem Eur J.* 2013; 19:4335–4343. [PubMed: 23335128] Yuan FQ, Han FS. *Adv Synth Catal.* 2013; 355:537–547. Shekhar AC, Kumar AR, Sathaiiah G, Raju K, Rao PS, Sridhar M, Narsaiah B, Srinivas PVSS, Sridhar B. *Helv Chim Acta.* 2012; 95:502–508. Sousa CM, Berthet J, Delbaere S, Coelho PJ. *J Org Chem.* 2012; 77:3959–3968. [PubMed: 22459032] Thirunavukkarasu VS, Donati M, Ackermann L. *Org Lett.* 2012; 14:3416–3419. [PubMed: 22691029] Silva LF Jr, Quintiliano SA. *Tetrahedron Lett.* 2009; 50:2256–2260.
17. Gharpure SJ, Prasath V. *J Chem Sci.* 2011; 123:943–949. Zheng DP, Gong WC, Ma ZD, Ma BZ, Zhao XL, Xie ZX, Li Y. *Tetrahedron Lett.* 2011; 52:314–317. Kim C, Bae HJ, Lee JH, Jeong W, Kim H, Sampath V, Rhee YH. *J Am Chem Soc.* 2009; 131:14660–14661. [PubMed: 19785427] Miranda PO, Carballo RnM, Martín VcS, Padrón JI. *Org Lett.* 2008; 11:357–360. [PubMed: 19093859]
18. The descriptor “Prins cyclization” has been used inconsistently in the literature. However, many authors now interpret the typical Prins cyclization to involve formation of two bonds in the tetrahydropyran product, and we are using this definition. See ref. 10b for a discussion of the Prins cyclization.
19. Reddy BVS, Reddy PS, Yadav JS, Sridhar B. *Tetrahedron Lett.* 2013; 54:1392–1396. Suzuki Y, Niwa T, Yasui E, Mizukami M, Miyashita M, Nagumo S. *Tetrahedron Lett.* 2012; 53:1337–1340. Basavaiah D, Reddy KR. *Org Lett.* 2006; 9:57–60. [PubMed: 17192084]
20. Xu CF, Xu M, Yang LQ, Li CY. *J Org Chem.* 2012; 77:3010–3016. [PubMed: 22356316]
21. Yamaguchi M, Hirao I. *Tetrahedron Lett.* 1983; 24:391–394.
22. Guiso M, Marra C, Piccioni F. *Nat Prod Res.* 2010; 24:331–340. [PubMed: 20221939]
23. See the Supporting Information for details of the X-ray analysis.
24. Mayr H, Schneider R, Wilhelm D, Schleyer PVR. *J Org Chem.* 1981; 46:5336–5340.
25. Apeloig, Y.; Muller, T. Theory and calculations for vinyl cations. In: Rappoport, Z.; Stang, PJ., editors. *Dicoordinated Carbocations*. Wiley; 1997. p. 9-104.
26. For an example of a Friedel-Crafts reaction followed by a dehydration/aromatization, see: Kuninobu Y, Tatsuzaki T, Matsuki T, Takai K. *J Org Chem.* 2011; 76:7005–7009. [PubMed: 21761938]
27. Jaber JJ, Mitsui K, Rychnovsky SD. *J Org Chem.* 2001; 66:4679–4686. [PubMed: 11421792]
28. Chavre SN, Choo H, Cha JH, Pae AN, Choi KI, Cho YS. *Org Lett.* 2006; 8:3617–3619. [PubMed: 16869674]

Scheme 1.
Intermolecular Friedel-Crafts.

Scheme 2.
Retrosynthetic Analysis.

Scheme 3.
Air Oxidation of **3a** and ORTEP of Lactone **4**.

Scheme 4.
Mechanistic Rationale for Product Formation.

Table 1

Brief Survey of Lewis Acid Activators.

entry	aldehyde	product	Lewis acid	yield (%) ^{a,b}
1			BF ₃ •OEt ₂	38
2		3a	Bi(OTf) ₃	19 ^c
3			BF ₃ •OEt ₂	47
4			Bi(OTf) ₃	42
5			BiBr ₃	36
6		3b	BF ₃ •OEt ₂	47 ^d

^aReactions were run under argon at ~0.25 M in **2a** using 1.2 equiv of aldehyde and 1.1 equiv of corresponding Lewis acid.^bIsolated yields of compounds after silica gel chromatography.^cThe *bis*-TES protected analog **2b** was used.^dReaction conducted at ~0.125 M.

Table 2One-Pot Cascade Synthesis of Benzo[*f*]isochromenes.

entry	product	aldehyde (R)	yield (%) ^a
1	3a	Et–	38
2	3b	<i>i</i> -Pr–	47
3	3c	<i>sec</i> -Bu–	52
4	3d	pentan-3-yl–	44
5	3e	PhCH ₂ CH ₂ –	43
6	3f	Ph–	58
7	3g	<i>p</i> -Br-Ph–	58
8	3h	<i>p</i> -CF ₃ -Ph–	43
9	3i	<i>p</i> -NO ₂ -Ph–	46
10	3j	<i>p</i> -MeO-Ph–	58

^a Isolated yields of compounds after silica gel column chromatography.