Anna Ardisson, MD*
Giuseppe Piscosquito,
MD*

Andrea Legati, PhD
Tiziana Langella, MSc
Eleonora Lamantea, MSc
Barbara Garavaglia, PhD
Ettore Salsano, MD
Laura Farina, MD
Isabella Moroni, MD
Davide Pareyson, MD
Daniele Ghezzi, PhD

A SLOWLY PROGRESSIVE MITOCHONDRIAL ENCEPHALOMYOPATHY WIDENS THE SPECTRUM OF *AIFM1* DISORDERS

To date, 3 *AIFM1* (apoptosis inducing factor mitochondrial 1, located on *Xq26.1*) mutations have been reported: 2 missense changes (c.923G>A/p.Gly308Glu; c.1478A>T/p.Glu493Val) and a 3-basepair deletion (c.601delAGA/p.Arg201del). Two mutations have been described in early-onset severe mitochondrial encephalomyopathy related to impaired oxidative phosphorylation.^{1,2} A third mutation is associated with Cowchock syndrome, or Charcot-Marie-Tooth X4 (CMTX4), a slowly progressive disorder characterized by axonal neuropathy, hearing loss, and mental retardation.^{3,4}

Case history. We report a 39-year-old man with an early-onset, slowly progressive mitochondrial disease associated with a novel *AIFM1* mutation, thus widening the clinical spectrum of *AIFM1*-related disorders.

The patient's family history was unremarkable. Psychomotor development was normal until age 1 year, when he started to show walking difficulties. During childhood, he developed gait and limb ataxia (sensory and cerebellar), hearing loss, and cognitive impairment. Visual deficit and distal muscle wasting and weakness became evident at puberty, with scanned speech, head and trunk titubation, and intention tremor, progressively more marked at repeated neurologic examinations. He became chair-bound at age 20 years, needed acoustic aids at age 22 years, and developed progressive sleep apnea syndrome since age 23 years. At age 30 years, visual loss had progressed to 1-2/10 bilaterally. Only one (13 years) of several assays showed elevated lactate and pyruvate serum levels (3,051 $\mu\text{mol/L}$, normal value 580–2,100; 146 $\mu\text{mol/L}$, normal value 40–140 $\mu\text{mol/L}$, respectively). Brain CT scan and MRI were normal at age 9 years, subsequently showing cerebellar atrophy (19 years) and mild cortical and thalamic atrophy (30 years). At last MRI (35 years), further involvement of bulbar olives, dentate nuclei, and spinal cord (slight cervical atrophy with T2 hyperintensity in posterior columns) were observed (figure, A–C). Brain spectroscopy was normal. Nerve conduction study showed a moderate-to-severe

axonal, more sensory than motor, neuropathy. EEG was normal. Optic atrophy and retinopathy were documented by visual evoked potentials (since his teens) and electroretinogram (25 years).

Results. Informed consent was obtained for biochemical and genetic analyses. Muscle biopsy (13 years) showed evidence of chronic denervation, ragged-red fibers, and fibers hyporeactive to cytochrome *c* oxidase (complex IV [cIV]) histochemical staining. Biochemical analysis revealed markedly reduced cIV activity, with partial reduction of other mitochondrial electron transfer chain (ETC) complex activities and increase of citrate synthase (figure, D). Mitochondrial DNA Southern blot and screening for *POLG1*, *SURF1*, *SCO1*, and *COX15* mutations were normal. Coenzyme Q10 and riboflavin supplementation did not modify clinical course. Recently, sequencing of a panel containing more than 100 nuclear genes associated with mitochondrialriopathies revealed a hemizygous c.784G>A/p.Gly262Ser mutation in *AIFM1* (NM_004208) (figure, E), the gene encoding apoptosis-inducing factor (AIF). This variant, inherited from the healthy mother (figure, F), is not reported in public databases, is predicted as probably pathogenic (table e-1 on the *Neurology*® Web site at Neurology.org), and affects a conserved amino acid residue (table e-2). Western blot analysis on patient fibroblasts showed strong reduction in AIF amount (figure, G), indicating instability of the mutant protein and confirming a deleterious outcome of the identified variant.

Discussion. AIF is a mitochondrial flavoprotein, attached by an N-terminal transmembrane domain to the inner mitochondrial membrane, with a poorly defined (FAD-dependent) NADH oxidase activity and a role in caspase-independent cell death.⁵ AIF defects can result in loss of its pro-survival function related to assembly/stabilization of ETC, or in increased pro-death activity, facilitating AIF translocation from mitochondria to the nucleus and eventually leading to DNA fragmentation.⁶ The few cases with *AIFM1* mutations previously reported are summarized in table e-3. Patients with a severe outcome included 2 cousins with early-onset, severe psychomotor delay, extrapyramidal signs, epilepsy, and axonal sensory-motor neuropathy associated with bilateral striatal necrosis¹; and 2 siblings with prenatal ventriculomegaly.² Recently, *AIFM1*

Supplemental data
at Neurology.org

(A) Sagittal T2-weighted image. (B) Coronal fluid-attenuated inversion recovery image. (C) Axial proton density-weighted image. (A–C) Diffuse cerebellar atrophy. Hyperintensity of dentate (B, arrow on the right) and olivary (C, arrow on the right) nuclei are also present. (D) Activities of mitochondrial respiratory chain complexes (cl, cl + III, cII, cII + III, cIV) in patient muscle homogenate, reported as percentages of the controls mean. The specific activities were normalized for citrate synthase (CS) activity. (E) Snapshot from IGV software of the region containing the mutation identified in the patient. *AIFM1* is on the reverse strand, hence the highlighted nucleotide (T) corresponds to the complementary of mutant c.784A. (F) Family pedigree shows the segregation of the c.784G>A mutation. The mother, heterozygous for the mutation, was asymptomatic; none of her 6 siblings has ever presented signs of mitochondrial disorder. No neurophysiologic testing was performed. (G) Western blot analysis of fibroblasts from patient (Pt) and controls (Ct1, Ct2, Ct3) using antibodies against AIFM1 and GAPDH, the latter used as loading control. The reported percentages correspond to the values of AIFM1/GAPDH signals obtained by densitometric analysis. One hundred percent corresponds to the mean value from controls.

mutation was found in a family with CMTX4.^{3,4} Mitochondrial encephalomyopathy with moderate clinical severity, slow progressive course despite early onset, and cerebellar involvement was not reported until now. Notwithstanding common clinical and instrumental features, the cases reported to date are heterogeneous. Even the biochemical profile is inconsistent, with multiple ETC deficiency reported in muscle for the most severe cases, mainly cIV deficiency found in the present patient, and normal activities described in CMTX4 individuals. These observations are partly in contrast with data obtained from mouse models suggesting that knockdown or knockout of AIF led to complex I deficiency, particularly in brain.⁷ However, biochemical defects seem to reflect the

clinical severity in patients, confirming the importance of AIF for proper ETC functioning and indicating that this AIF role is fundamental to reduce neuronal damage. Moreover, this case confirms the usefulness of broad genetic screening, for instance targeted sequencing, in highly heterogeneous syndromes resembling mitochondrial disorders.

*These authors contributed equally to the manuscript.

From the Foundation Institute of Neurology "Carlo Besta," Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS), Milan, Italy.

Author contributions: A.A., G.P., E.S., I.M., and D.P. evaluated the patients and wrote the case report. A.L. and T.L. performed genetic screening and analyzed exome-sequencing data. E.L. performed biochemical analyses under the supervision of B.G., D.G. monitored genetic/protein analyses. L.F. evaluated the MRIs. G.P., A.A., and D.G. wrote the manuscript. I.M. and D.P. critically revised the

manuscript for important intellectual content. All authors read and approved the manuscript.

Study funding: Supported by Fondazione Telethon (GGP11011), the Italian Ministry of Health (GR2010–2316392), CARIPLO grant 2011/0526, the Mariani Foundation (CM23), and the Italian Association of Mitochondrial Disease Patients and Families (Mitocon).

Disclosure: A. Ardisson reports no disclosures relevant to the manuscript. G. Piscosquito received funded travels from Kedrion SpA and Pfizer Italia srl. A. Legati, T. Langella, and E. Lamantea report no disclosures relevant to the manuscript. B. Garavaglia received funding and grants for research from Fondazione Mariani (CM23), Telethon Network of Genetics Biobank (GTB12001). E. Salsano received funded travels from Actelion Pharma Schweiz AG. L. Farina reports no disclosures relevant to the manuscript. I. Moroni received funding and grants for research from Telethon Italy (GUP11001, GUP13006). D. Pareyson received funding and grants for research from Telethon Italy (GUP09013, GUP10008, GUP10010, GUP13006), Regione Lombardia (grant on rare diseases), ACMT-Rete (funding for a CMT Registry), and Pfizer Italia srl (donations for research activities) and funded travels from Kedrion SpA and Pfizer Italia srl. He serves as member of the editorial boards of Journal of Peripheral Nervous System, Neurological Sciences, and Journal of Neuromuscular Disorders. D. Ghezzi received funding and grants for research from the Italian Ministry of Health, Telethon Italy (GGP11011), and CARIPLO Foundation. Go to Neurology.org for full disclosures.

This is an open access article distributed under the Creative Commons Attribution-Noncommercial License, which permits downloading, sharing, and reproducing the work in any medium, provided it is properly cited. The work cannot be used commercially.

Received November 7, 2014. Accepted in final form January 21, 2015.

Correspondence to Dr. Ghezzi: dghezzi@istituto-besta.it or Dr. Pareyson: davide.pareyson@istituto-besta.it

© 2015 American Academy of Neurology

1. Ghezzi D, Sevioukova I, Invernizzi F, et al. Severe X-linked mitochondrial encephalomyopathy associated with a mutation in apoptosis-inducing factor. *Am J Hum Genet* 2010; 86:639–649.
2. Berger I, Ben-Neriah Z, Dor-Wolman T, et al. Early prenatal ventriculomegaly due to an *AIFM1* mutation identified by linkage analysis and whole exome sequencing. *Mol Genet Metab* 2011;104:517–520.
3. Cowchock FS, Duckett SW, Streletz LJ, Graziani LJ, Jackson LG. X-linked motor-sensory neuropathy type-II with deafness and mental retardation: a new disorder. *Am J Med Genet* 1985;20:307–315.
4. Rinaldi C, Grunseich C, Sevioukova IF, et al. Cowchock syndrome is associated with a mutation in apoptosis-inducing factor. *Am J Hum Genet* 2012;91:1095–1102.
5. Sevioukova IF. Apoptosis-inducing factor: structure, function, and redox regulation. *Antioxid Redox Signal* 2011;14: 2545–2579.
6. Polster BM. AIF, reactive oxygen species, and neurodegeneration: a “complex” problem. *Neurochem Int* 2013;62: 695–702.
7. Vahsen N, Cande C, Briere JJ, et al. AIF deficiency compromises oxidative phosphorylation. *EMBO J* 2004;23: 4679–4689.

Davide Tonduti, MD*
Imen Dorboz, PhD*
Florence Renaldo, MD
Julien Masliah-Planchon,
PharmD, PhD
Monique Elmaleh-Bergès,
MD
Hélène Dalens, MD
Diana Rodriguez, MD,
PhD
Odile Boespflug-Tanguy,
MD, PhD

CYSTIC LEUKOENCEPHALOPATHY WITH CORTICAL DYSPLASIA RELATED TO *LAMB1* MUTATIONS

Several diseases related to anomalies of basement membrane components are known.^{1,2} Among these disorders, cerebral white matter involvement is well-documented with respect to mutations in *COL4A1-2*, coding for the $\alpha 1$ and $\alpha 2$ chains of collagen IV, as well as mutations in *LAMA2*, coding for laminin $\alpha 2$.² Recently, mutations in another laminin-encoding gene, *LAMB1*, have been reported in patients presenting a severe malformative encephalopathy.³ We present 2 siblings from a nonconsanguineous family who extend the phenotype of the *LAMB1*-related disorder.

Case reports. Patient 1 presented with infantile spasms associated with a congenital static right hemiplegia, and was diagnosed with cerebral palsy due to the presence of a left porencephalic cavity on CT. From the age of 12 years, she had a resurgence of epilepsy and progressively deteriorated. Bilateral anterior subcapsular lens opacification was observed. At 19 years of age, she lost the ability to walk; a tendency to macrocephaly, spastic tetraplegia predominating on the right side, optic atrophy, and mental deficiency were noted. At 23 years of age, mild bilateral deafness was detected. She is now 30 years old, and

the clinical picture has evolved in terms of a slowly progressive mental deterioration with stable epilepsy (1 seizure per month).

The older brother of patient 1 presented with a progressive isolated postnatal macrocephaly. Infantile depression was diagnosed when he presented well-controlled absence seizures at 7 years. At age 16 years, after a fall, he presented with generalized pharmacoresistant seizures and gait difficulties. Subsequently, neurocognitive and motor deterioration became evident, whereas seizures stabilized. At 24 years of age, he lost the ability to walk and presented relapsing and long-lasting (3 months) episodes of pain in the right arm related to a right axonal brachial plexopathy on EMG. Neurologic examination revealed spastic paraplegia with cerebellar signs and frontal cognitive dysfunction. Ophthalmologic evaluation revealed the presence of retinal vascular tortuosities (figure, K) and bilateral subcapsular lens opacification. During the last 10 years, he presented with 3 acute relapsing episodes of unexplained stupor coma induced by stress. Now 34 years old, he is totally dependent.

In both patients, the supratentorial white matter showed a diffuse abnormal MRI signal with anterior subcortical temporal cysts and a vacuolating aspect of the deep white matter. Agyria/lissencephaly type