Case Report

Success of Minimally Invasive Transumbilical Single-Incision Laparoscopic Surgery (T-SILS) Plus Double-Balloon Endoscopy (DBE) for Pediatric Intestinal Angiodysplasia: A Case Report

Koichiro Mori, Yuhki Koike, Mikihiro Inoue, Kohei Ohtake, Koji Tanaka, Keichi Uchida, Yasuhiko Mohri, Masato Kusunoki

Departments of Gastrointestinal and Pediatric Surgery, Mie University Graduate School of Medicine, Tsu, Japan

We describe here a child with angiodysplasia of the small intestine, diagnosed by double-balloon endoscopy (DBE), who was treated with transumbilical single-incision laparoscopic surgery (T-SILS). A 9-year-old boy presented to another hospital with intermittent fresh melena of a duration of 5 months and 4 days. Anoscopy and gastric mucosal membrane scintigraphy were unsuccessful diagnostically, and he was referred to our hospital for further examination and treatment. Under general anesthesia, we performed DBE by an anal route, with the ileum assessed for a distance of about 150 cm from the ileocecal valve. Although no bleeding lesion was found in the colon, a flat elevated venous lake approximately 20 mm in size was observed 20 cm from the ileocecal valve, suggesting angiodysplasia of the small intestine. A tattoo was made under DBE, and wedge resection by T-SILS was performed 1 month later under general anesthesia. The excised specimen appeared as a flat, elevated venous lake approximately 20 mm in size. Histopathologic analysis revealed several dilated and distorted veins within the submucosa of the small intestine. The patient was diagnosed with angiodysplasia of the small intestine. His postoperative course was uneventful, and he remained free of recurrent gastrointestinal bleeding 18 months after the operation. This is the first case report showing the diagnosis, localization, and tattooing of an obscure gastrointestinal bleeding lesion of the small intestine by preoperative DBE and removal of the lesion by subsequent T-SILS in children.

Corresponding author: Masato Kusunoki, Department of Gastrointestinal and Pediatric Surgery, Mie University Graduate School of Medicine, 2–174 Edobashi, Tsu, Mie 514-8507, Japan.

Tel.: +81 59 232 1111 (ext. 5645); Fax: +81 59 232 6968; E-mail: koichiromori@clin.medic.mie-u.ac.jp

Key words: Transumbilical single-incision laparoscopic surgery (T-SILS) – Double-balloon endoscopy (DBE) – Angiodysplasia – Small intestine

Approximately 1% to 5% of patients with gastrointestinal bleeding can be classified as having obscure gastrointestinal bleeding (OGIB), defined as bleeding of unknown origin that persists or recurs after a negative initial or primary colonoscopy and/or upper endoscopy.^{1–3} Small intestinal angiodysplasias are found in 30% to 40% of patients with OGIB.^{4,5} Small intestinal bleeding can be detected, diagnosed, and localized by double-balloon endoscopy (DBE).⁶ No consensus has been reached regarding the most effective therapeutic strategy for angiodysplasia of the small intestine. We describe here a pediatric patient with angiodysplasia of the small intestine who was successfully treated by DBE combined with transumbilical single-incision laparoscopic surgery (T-SILS).

Case Report

A 9-year-old boy presented to another hospital with intermittent fresh melena 5 months and 4 days in duration. Anoscopy and gastric mucosal membrane scintigraphy were unsuccessful in diagnosing his condition, and he was transferred to our hospital for further examination. The patient had symptomatic epilepsy, cerebral palsy, and mental retardation, but no obvious abnormal physical findings on admission, including blood biochemistry. His symptoms suggested lower gastrointestinal tract bleeding, perhaps small intestinal bleeding. DBE (Fujinon EN-450p5/20, Fujinon Inc, Saitama, Japan) was performed under general anesthesia. Although no bleeding lesion was found in either the total colon or the ileum (150 cm from the ileocecal valve), a flat, elevated venous lake approximately 20 mm in size was found 20 cm from the ileocecal valve (Fig. 1A). We suspected angiodysplasia of the small intestine. The area around the lesion was tattooed endoscopically for subsequent laparoscopic surgery. Because of the possibility of repeated future gastrointestinal bleeding, surgery was performed 1 month later. The patient was placed under general anesthesia, and T-SILS was performed with the patient in a supine position. A reverse Ω -shaped incision was made at the umbilicus, and a multichannel EZ access (Covidien, Tokyo, Japan) was inserted as a platform under direct vision. A 5-mm flexible scope (Olympus, Tokyo, Japan) and 2 noncrushing graspers were

introduced into the EZ access. The previously tattooed small intestine lesion was immediately identified (Fig. 1B) and extracted from the abdominal cavity through the umbilical incision. A wedge resection was made around the tattooed site. The excised specimen appeared as a flat, elevated venous lake approximately 20 mm in size (Fig. 1C). Histopathologic analysis showed several dilated and distorted veins within the submucosa of the small intestine (Fig. 1D), resulting in a diagnosis of angiodysplasia of the small intestine. The patient's postoperative course was uneventful, and he was discharged 7 days after the operation. The patient remains free of recurrence 18 months after the operation.

Discussion

Angiodysplasia of the small intestine may be treated endoscopically, pharmacologically, surgically, or by interventional radiology. No consensus has yet been reached about the most effective therapeutic strategy for bleeding angiodysplasia of the small intestine. Although approximately 90% of hemorrhagic angiodysplasias spontaneously cease bleeding,⁷ the risk of recurrent bleeding is high. For example, the cumulative incidences of overt rebleeding at 1 and 3 years were 26% and 45%, respectively, and the rates of occult rebleeding at 1 and 3 years were 46% and 64%, respectively.^{8,9} A previous episode of rebleeding is an independent risk factor for further rebleeding.¹⁰ Up to 12% of these patients require surgical treatment,¹¹ and the angiodysplasia related mortality rate is around 2%.¹² Curative surgical resection is recommended for patients with gastrointestinal angiodysplasia. Lesions too small and too soft to be palpated may be difficult to localize intraoperatively, even during laparotomy. Thus, marking a lesion before surgery is important.

The clinical introduction of DBE markedly changed the diagnosis and treatment of small intestinal bleeding.¹³ DBE allows endoscopic imaging of the entire small intestine, an organ inaccessible using conventional procedures, allowing the bleeding site to be diagnosed and localized. DBE was able to diagnose 64% of patients with small intestinal OGIB who could not be given a diagnosis

Fig. 1 Histologic appearance of the lesion in our patient. (A) A flat, elevated venous lake approximately 20 mm in size and located 10 cm from the ileocecal valve. (B) Laparoscopic view of the tattoo of the lesion in the ileum, enabling easy identification of the lesion during subsequent laparoscopic surgery. (C) Photograph of the excised specimen, showing lifting of a 20-mm area of the mucous membrane. (D) Several dilated and distorted veins present within the submucosa of the small intestine. Hematoxylin-eosin ($\times 40$).

by conventional approaches.¹⁴ We found that DBE could successfully give a diagnosis of OGIB and localize its site of origin in our patient. Labeling the lesion allowed its later surgical removal.

Surgical methods for small intestinal diseases have changed since the introduction of DBE. As in our patient, an integrated approach consisting of DBE and laparoscopic bowel surgery (LBS) was used to manage small intestinal diseases manifesting as OGIB and vague abdominal pain when nonoperative methods failed or were not appropriate. LBS plus DBE has advantages over laparotomy without DBE. First, the surgical incision is smaller in LBS than in laparotomy, reducing postoperative pain. Moreover, LBS is faster, safer, has better cosmetic outcomes, and is more cost-effective than laparotomy. Most laparoscopically assisted intestinal resections can be completed in 1 hour, with no mortality and minimal morbidity.¹⁵ Second, the use of LBS in a targeted fashion diminished the need for thorough exploration and manipulation of the entire intestine. In most cases, intestinal targets were easily identified at initial laparoscopic examination based on the “tattoo” procedure. Third, by exfoliating the adhesion laparoscopically, it is easy to displace the organ outside the abdominal cavity to remove any physiologic adhesion, such as in the ascending colon

or the ligament of Treitz. Exteriorization of the diseased intestinal loop at the nearest port site allowed any surgical procedure to be easily performed.

Resumption of bowel movement, surgical morbidity and mortality rates, and hospital stay of patients managed using LBS plus DBE compared favorably with those of patients treated with exploratory laparotomy and intraoperative endoscopy.^{2,16,17} T-SILS was minimally invasive, allowing good outcomes in our patient.

To date, 26 children with small intestinal disease have been managed by SILS.^{18–23} Many diseases are not accompanied by a specific lesion site, such as Meckel diverticulum and Crohn disease, making preoperative identification of any lesion important. To our knowledge, this is the first case report showing the diagnosis, localization, and tattooing of an OGIB lesion of the small intestine by preoperative DBE, and removal of the lesion by subsequent T-SILS.

In conclusion, DBE is useful in the diagnosis and localization of lesions in the small intestine. DBE plus T-SILS enabled the successful management of this OGIB in a child using minimally invasive laparoscopic surgery. The combination of DBE plus

T-SILS may be a standard approach to OGIB lesions in the small intestine.

References

1. Zuckerman GR, Prakash C, Askin MP, Lewis BS. AGA technical review on the evaluation and management of occult and obscure gastrointestinal bleeding. *Gastroenterology* 2000; **118**(1):201–221
2. Lau WY, Fan ST, Wong SH, Wong KP, Poon GP, Chu KW *et al.* Preoperative and intraoperative localisation of gastrointestinal bleeding of obscure origin. *Gut* 1987; **28**(7):869–877
3. Lewis BS. Small intestinal bleeding. *Gastroenterol Clin North Am* 1994; **23**(1):67–91
4. Foutch PG. Angiodysplasia of the gastrointestinal tract. *Am J Gastroenterol* 1993; **88**(6):807–818
5. Schmit A, Gay F, Adler M, Cremer M, Van Gossum A. Diagnostic efficacy of push-enteroscopy and long-term follow-up of patients with small bowel angiodysplasias. *Dig Dis Sci* 1996; **41**(12):2348–2352
6. Kaffes AJ, Siah C, Koo JH. Clinical outcomes after double-balloon enteroscopy in patients with obscure GI bleeding and a positive capsule endoscopy. *Gastrointest Endosc* 2007; **66**(2):304–309
7. Duchini A, Godino J, Wong P, Patti M, Talavera F, Achord JL *et al.* Angiodysplasia of colon. Available at: http://www.emedicine.com/med/topic_134.htm. Accessed July 17, 2008
8. Richter JM, Christensen MR, Colditz GA, Nishioka NS. Angiodysplasia: natural history and efficacy of therapeutic interventions. *Dig Dis Sci* 1989; **34**(10):1542–1546
9. Junquera F, Feu F, Papo M, Videla S, Armengol JR, Bordas JM *et al.* A multicenter, randomized, clinical trial of hormonal therapy in the prevention of rebleeding from gastrointestinal angiodysplasia. *Gastroenterology* 2001; **121**(5):1073–1079
10. Junquera F, Saperas E, Videla S, Feu F, Vilaseca J, Armengol JR *et al.* Long-term efficacy of octreotide in the prevention of recurrent bleeding from gastrointestinal angiodysplasia. *Am J Gastroenterol* 2007; **102**(2):254–260
11. Gupta N, Longo WE, Vernava AM III. Angiodysplasia of the lower gastrointestinal tract: an entity readily diagnosed by colonoscopy and primarily managed nonoperatively. *Dis Colon Rectum* 1995; **38**(9):979–982
12. Cappell MS, Gupta A. Changing epidemiology of gastrointestinal angiodysplasia with increasing recognition of clinically milder cases: angiodysplasia tend to produce mild chronic gastrointestinal bleeding in a study of 47 consecutive patients admitted from 1980–1989. *Am J Gastroenterol* 1992; **87**(2):201–206
13. Yamamoto H, Kita H, Sunada K, Hayashi Y, Sato H, Yano T *et al.* Clinical outcomes of double-balloon endoscopy for the diagnosis and treatment of small-intestinal diseases. *Clin Gastroenterol Hepatol* 2004; **2**(11):1010–1016
14. Arakawa D, Ohmiya N, Nakamura M, Honda W, Shirai O, Itoh A *et al.* Outcome after enteroscopy for patients with obscure GI bleeding: diagnostic comparison between double-balloon endoscopy and videocapsule endoscopy. *Gastrointest Endosc* 2009; **69**(4):866–874
15. Yeh TS, Liu KH, Su MY, Lin CH, Chiu CT, Tseng JH. Laparoscopically assisted bowel surgery in an era of double-balloon enteroscopy: from inside to outside. *Surg Endosc* 2009; **23**(4):739–744
16. Ress AM, Benacci JC, Sarr MG. Efficacy of intraoperative enteroscopy in diagnosis and prevention of recurrent, occult gastrointestinal bleeding. *Am J Surg* 1992; **163**(1):94–98
17. Cave DR, Cooley JS. Intraoperative enteroscopy: indications and techniques. *Gastrointest Endosc Clin N Am* 1996; **6**(4):793–802
18. Tam YH, Lee KH, Sihoe JD, Chan KW, Cheung ST, Pang KK. Initial experience in children using conventional laparoscopic instruments in single-incision laparoscopic surgery. *J Pediatr Surg* 2010; **45**(12):2381–2385
19. St Peter SD. Single-site umbilical laparoscopic segmental small bowel resection. *Semin Pediatr Surg* 2011; **20**(4):219–223
20. Garey CL, Laituri CA, Ostlie DJ, Snyder CL, Andrews WS, Holcomb GW III *et al.* Single-incision laparoscopic surgery in children: initial single-center experience. *J Pediatr Surg* 2011; **46**(5):904–907
21. Laituri CA, Fraser JD, Garey CL, Aguayo P, Sharp SW, Ostlie DJ *et al.* Laparoscopic ileocectomy in pediatric patients with Crohn's disease. *J Laparoendosc Adv Surg Tech A* 2011; **21**(2):193–195
22. Hattori K, Sato M, Miyauchi Y. Single-incision laparoscopic surgery for small bowel resection with an intraumbilical arcuate incision in children. *Asian J Endosc Surg* 2012; **5**(4):172–175
23. Inoue M, Uchida K, Otake K, Okigami M, Okita Y, Hiro J *et al.* Transumbilical single-incision laparoscopic ileocecal resection for Crohn's disease in a child. *Asian J Endosc Surg* 2013; **6**(1):48–51