Published in final edited form as:

Dev Psychopathol. 2014 November ; 26(4 0 2): 1277–1287. doi:10.1017/S0954579414001023.

***IL1B* Gene Variation and Internalizing Symptoms in Maltreated Preschoolers**

Kathryn K. Ridout^{a,b}, Stephanie H. Parade^{b,c}, Ronald Seifer^{b,c}, Lawrence H. Price^{a,b}, Joel Gelernter^d, Paloma Feliz^c, and Audrey R. Tyrka^{a,b,*}

^a Mood Disorders Research Program and Laboratory for Clinical and Translational Neuroscience, Butler Hospital, Providence, RI, USA

^b Department of Psychiatry and Human Behavior, Alpert Medical School of Brown University

^c Bradley/Hasbro Children's Research Center, E. P. Bradley Hospital, East Providence, RI, USA

^d Yale University School of Medicine, Departments of Psychiatry, Genetics, and Neurobiology and the VA CT Healthcare Center

Abstract

Evidence now implicates inflammatory proteins in the neurobiology of internalizing disorders. Genetic factors may influence individual responses to maltreatment; however, little work has examined inflammatory genetic variants in adults and none in children. The present study examined the role of an *IL1B* variant in preschoolers exposed to maltreatment and other forms of adversity in internalizing symptom development. One hundred ninety-eight families were enrolled, with one child (age 3-5 years) from each family. Adversity measures included child protective service documentation of moderate-severe maltreatment in the last 6 months and interview-assessed contextual stressors. Internalizing symptoms were measured using the Child Behavior Checklist (CBCL) and the Diagnostic Infant and Preschool Assessment (DIPA). Maltreated children had higher MDD and PTSD symptoms and marginally higher internalizing symptoms on the CBCL. Controlling for age, sex and race, *IL1B* genotype was associated with MDD symptoms ($p = .002$). Contextual stressors were significantly associated with MDD and PTSD and marginally with internalizing symptoms. The *IL1B* genotype interacted with contextual stress such that children homozygous for the minor allele had more MDD symptoms ($p = .045$). These results suggest that genetic variants of *IL1B* may modulate the development of internalizing symptoms in the face of childhood adversity.

Keywords

IL1B; cytokine; variant; maltreatment; children

* Address correspondence to Audrey Tyrka, M.D., Ph.D., 345 Blackstone Boulevard, Providence RI, 02906. Tel: 401-455-6520 Fax: 401-455-6534 Audrey_Tyrka@Brown.edu.

Introduction

Longitudinal investigations have demonstrated the persistence of social-emotional and behavioral problems in the internalizing domains into later life, even when they are identified as early as 12-48 months of age; such symptoms are strongly correlated with the development of significant psychopathology (Briggs-Gowan, Carter, Bosson-Heenan, Guyer, & Horwitz, 2006; Mesman & Koot, 2001; Roza, Hofstra, van der Ende, & Verhulst, 2003). Childhood maltreatment and adversity are established risk factors for internalizing symptoms (Carr, Martins, Stingel, Lemgruber, & Juruena, 2013; Hickman et al., 2013; Kim-Spoon, Cicchetti, & Rogosch, 2013; R. Mills et al., 2013; Norman et al., 2012) and associated conditions, such as depression, post-traumatic stress disorder (PTSD), and anxiety, both later in childhood and into adulthood (Benjet, Borges, & Medina-Mora, 2010; Green et al., 2010; Schilling, Aseltine, & Gore, 2007; Scott, Smith, & Ellis, 2010; Widom, 1999; Widom, DuMont, & Czaja, 2007). Individual responses to adversity and the subsequent development and severity of internalizing symptoms or disorders vary considerably, suggesting that there are underlying factors modifying an individual's risk, although little is known regarding what these factors may be, especially in very young children exposed to maltreatment (Lupien, McEwen, Gunnar, & Heim, 2009; Teicher & Samson, 2013). Identifying these moderating factors is critical for efforts to prevent or alter the long-term sequelae of early-onset internalizing symptoms.

Childhood adversity may have the greatest effects in individuals with genetic variants that influence the neural systems dysregulated in internalizing conditions (Heim & Binder, 2012; Hornung & Heim, 2014). In addition, some genes may influence response to a broad range of environmental conditions, with alleles that confer risk in the face of environmental stressors but provide benefit under positive conditions (Belsky et al., 2009; Belsky & Pluess, 2009). Variability in the outcome of some gene $\times$ environment (G $\times$ E) studies is likely due in part to difficulties inherent to designing such studies, which rely on retrospective accounts of adversity in adults or are limited in young children who cannot report their adversity history. The timing and type of exposure to maltreatment and other adversities is likely critical and introduces another source of variability between G $\times$ E studies (Pietrek, Elbert, Weierstall, Muller, & Rockstroh, 2013). This is likely in part due to developmental changes in attachment relationships and the development of neural systems that underlie the regulation of affect and behavior (Cicchetti & Toth, 2005), suggesting the importance of studying different types of adverse exposures at different developmental periods in elucidating G $\times$ E interactions. Additionally, families who may be most at risk are relatively inaccessible to research participation due to the effects of concurrent poverty and environmental stress exposure.

G $\times$ E interactions in maltreated children, psychopathology, and internalizing symptoms

Several studies in youth have examined how maltreatment exposure interacts with functional genetic variants that are thought to influence neurobiological systems relevant to internalizing disorders (Bellani, Nobile, Bianchi, van Os, & Brambilla, 2012; Nugent, Tyrka, Carpenter, & Price, 2011; Teicher & Samson, 2013). For example, the serotonin system is important to the development of internalizing disorders such as depression and

anxiety and thought to play a central role in stress-induced psychiatric disorders (Shikanai, Kimura, & Togashi, 2013). Numerous studies have focused on genetic variants in the serotonin-transporter-linked polymorphic region (*5-HTTLPR*) of the serotonin transporter protein (*SLC6A4*) gene, which appears to regulate serotonin uptake from the synaptic cleft (Lesch et al., 1996). Consistent with adult studies (Caspi et al., 2003), Kaufman et al. reported a significant association of *5-HTTLPR* genotype with increased depressive symptoms in maltreated children compared to maltreated children with other genotypes or non-maltreated children with the same genotype (Kaufman et al., 2004). This result has been replicated in other studies in youth and appears potentiated by factors such as low social supports (Aslund et al., 2009; Banny, Cicchetti, Rogosch, Oshri, & Crick, 2013; Kaufman et al., 2004). Other studies have demonstrated G×E interactions in the development of internalizing disorders involving genetic variants in the serotonergic, dopaminergic, noradrenergic, glutamatergic and GABAergic systems, other monoamine enzymes, cannabinoids, neuroendocrine, pro-survival factors and inflammatory mediators (for review of mechanisms see: (Mandelli & Serretti, 2013; Nugent et al., 2011).

Because these neurobiological systems are closely interconnected, a number of studies have tested for gene × gene interactions between regulatory genes in these systems (Masten & Cicchetti, 2010). Interactions of variants of *5-HTTLPR* and brain-derived neurotrophic factor (BDNF) genes were associated with higher depressive symptoms in maltreated but not in non-maltreated youth (Kaufman et al., 2006). Similar results for these genes were found in other studies of adolescents and young adults (Aguilera et al., 2009; Comasco, Aslund, Orelund, & Nilsson, 2013; Nederhof, Bouma, Oldehinkel, & Ormel, 2010). Other work has documented gene × gene interactions of *5-HTTLPR* with an *MAOA* gene variant in sexually-abused youth (Cicchetti, Rogosch, & Sturge-Apple, 2007), with a variant of the corticotropin releasing hormone receptor 1 (*CRHR1*) gene in maltreated children, and with the dopamine receptor D4 (*DRD4*) gene (Cicchetti, Rogosch, & Oshri, 2011; Cicchetti, Rogosch, & Toth, 2011). In work aiming to predict resilience in adaptive functioning among 595 maltreated and nonmaltreated low-income children, genetic variants in the *5-HTTLPR*, *CRHR1*, *DRD4*, and oxytocin receptor gene each gave a relative advantage in resilient functioning in nonmaltreated compared to maltreated children (Cicchetti & Rogosch, 2012). Taken together, these results show that neurobiologically-guided targets for examining G×E interactions has yielded some potential models by which adversity may modify individual risk of psychopathology. However, these models have been criticized on methodological grounds in some cases and more complete models advocated (Keller, 2014).

In addition to maltreatment, other forms of childhood adversity, such as parental loss, socioeconomic adversity, and poor social supports have been linked to the development of internalizing problems (Dunn et al., 2011; Manly, Kim, Rogosch, & Cicchetti, 2001). While some G×E interaction studies have focused on direct childhood maltreatment such as abuse and neglect, others have included more broadly experienced events, such as parental arguing/divorce, poverty, or parent education levels. Evidence supports the existence of differences in the impact of type of ELS (Cicchetti et al., 2007; Sjöberg et al., 2006). Poor social support predisposes to internalizing symptoms, and moderates the effects of childhood maltreatment on internalizing symptoms (Kaufman et al., 2006; Kaufman et al.,

2004). In a study examining childhood risk factors for adult psychopathology, participants with childhood parental separation, desertion or death were significantly more likely to report the subsequent onset of symptoms of a depressive or anxiety disorder. These effects were not fully explained by parental relationships or history of childhood maltreatment and a family history of depressive or anxiety disorders accounted for the effect of parental separation, suggesting a genetic predisposition contributing to the development of psychopathology after this exposure (Tyrka, Wier, Price, Ross, & Carpenter, 2008). Thus, in addition to maltreatment, exposure to other contextual stressors contributes to risk of developing internalizing symptoms or disorders in genetically-predisposed individuals.

Few studies on this topic have included young children and none have included children younger than age 5, so little is known regarding the effects of genetic variants that may potentiate responses to early adversity. While the cumulative effects of stress over time are likely to produce the most robust effects, evidence shows that even very young children develop internalizing symptoms and disorders (Luby, Gaffrey, Tillman, April, & Belden, 2014; Scheeringa & Haslett, 2010; Zeanah, Boris, & Scheeringa, 1997). Studies of young children are critical because early attachment plays an important role in the development of internalizing symptoms, and the neural systems that underlie internalizing symptoms undergo major developmental change during this time (Carrion & Wong, 2012; Rincon-Cortes & Sullivan, 2014).

Inflammatory cytokines and internalizing symptoms

Despite a wealth of evidence implicating inflammation in association with adversity and in the development of depressive and anxiety disorders, almost no work has examined genes that regulate inflammatory proteins in relation to internalizing symptoms or adverse childhood experiences (Mandelli & Serretti, 2013). The inflammatory response is a reaction to acute stress or injury involving release of inflammatory mediators, cellular recruitment, and release of pro-inflammatory cytokines. Cytokines are a diverse group of messenger proteins that act through intercellular signaling to regulate immune responses in response to injuries; cytokines are also activated in response to psychological stress (Lacy & Stow, 2011; Robles, Glaser, & Kiecolt-Glaser, 2005). C-reactive protein (CRP) is an acute-phase protein produced in response to acute infection or injury that is a marker of chronic, low-grade inflammation (Ferri et al., 2007; Lavie, Milani, Verma, & O'Keefe, 2009). While this inflammatory response is part of an adaptive response to minimize injury and promote healing in response to acute injuries or threats, excessive inflammation with prolonged and persistent elevation of pro-inflammatory cytokine levels is associated with a number of psychiatric disorders, including major depressive disorder (MDD) and PTSD (Mills, Scott, Wray, Cohen-Woods, & Baune, 2013; Rohleder, 2014).

Adults with MDD have elevated inflammatory responses and peripheral concentrations of some cytokines, including interleukin (IL)-6, IL-1 β , and tumor necrosis factor (TNF)- α (Dowlati et al., 2010; Hiles, Baker, de Malmanche, & Attia, 2012; Liu, Ho, & Mak, 2012), and PTSD in adults is associated with inflammation (Baker, Nievergelt, & O'Connor, 2012; Gola et al., 2013; O'Donovan et al., 2011; Pace et al., 2012). Several studies have demonstrated elevations of inflammatory markers in adults with a history of childhood

adversity (Coelho, Viola, Walss-Bass, Brietzke, & Grassi-Oliveira, 2014). In youth, there is now a wealth of evidence showing that adversity is associated with elevations of CRP (Appleton et al., 2012; Danese et al., 2008; Danese, Pariante, Caspi, Taylor, & Poulton, 2007; Low, Matthews, & Hall, 2013; Slopen, Kubzansky, McLaughlin, & Koenen, 2013), and a few studies have also shown higher levels of pro-inflammatory cytokines (Bertone-Johnson, Whitcomb, Missmer, Karlson, & Rich-Edwards, 2012; Rooks, Veledar, Goldberg, Bremner, & Vaccarino, 2012; Slopen et al., 2013). Recently, we found that salivary IL-1 β was positively associated with childhood adversity in preschool-aged children (Tyrka et al., under review). Links between inflammatory proteins and internalizing symptoms have also been documented in some, but not all, studies of youth (Mills et al., 2013; Slopen et al., 2013).

To our knowledge, only one study examined genetic determinants of the association between internalizing, adversity, and inflammatory proteins. In adults previously exposed to stressful life events, two promoter variants in the gene for *IL18* were associated with onset of depression (Haastrup et al., 2012). There is evidence that genes regulating cytokine expression have an important role in the development of inflammatory conditions in association with depression or adversity in adults (Cole et al., 2010; Kim et al., 2013) and adolescents (Cole et al., 2011). Prior studies have not examined inflammatory genetic variants in relation to internalizing behavior or adversity in children.

The role of IL-1 β in stress neurobiology

As noted above, studies have shown elevated IL-1 β in individuals with internalizing disorders associated with childhood maltreatment, suggesting this cytokine may have a role in the development of these disorders (Mitchell & Goldstein, 2014). Genes that regulate IL-1 β may be particularly important in this context because IL-1 β plays an important role in stress neurobiology (Dantzer, 2009). Animal models of stress exposure show increases in IL-1 β (Bailey, Kinsey, Padgett, Sheridan, & Leblebicioglu, 2009; Caso, Moro, Lorenzo, Lizasoain, & Leza, 2007; Nguyen et al., 1998; Porterfield, Gabella, Simmons, & Johnson, 2012; You et al., 2011), and in humans, IL-1 β increases acutely in response to a variety of stress challenges, including cognitive, social, and sleep-deprivation paradigms (Brydon et al., 2005; Mastroiardo, Alicino, Zefferino, Pasquini, & Picardi, 2007; Steptoe, Hamer, & Chida, 2007; Yamakawa et al., 2009). In animal models, central administration of IL-1 β activates the hypothalamic-pituitary-adrenal (HPA) axis, reduces hippocampal BDNF, and impairs hippocampal-dependent learning (Koo & Duman, 2008), important processes in the pathophysiology of internalizing disorders (Felger & Lotrich, 2013; N. T. Mills et al., 2013; von Kanel et al., 2007). Furthermore, there is evidence that activation of the IL-1 β receptor is necessary for stress to impair neurogenesis (Koo & Duman, 2008).

The Present Study

To summarize, a large body of evidence now implicates inflammatory proteins in the neurobiology of depressive and anxiety disorders in adults. Most of the work in children has focused on a general marker of inflammation, CRP, with little examination of cytokines, such as IL-1 β that are known to be important mediators of the neural response to stress. Almost no work has examined genes that regulate inflammation in relation to stress and

internalizing disorders and the single existing study was in adults. Although the onset of internalizing disorders often occurs later in childhood and adulthood, young children develop internalizing symptoms following adverse experiences and elucidating these processes during early development may be critical to an understanding of these disorders. Given the documented involvement of IL-1 β in neuronal function, stress responses, and MDD, the gene coding for this pro-inflammatory cytokine is a good candidate for evaluation in relation to internalizing symptoms and exposure to adversity.

The aim of the present study was to examine the role of a genetic variant in the *IL1B* gene in preschoolers exposed to maltreatment and other forms of adversity in the development of internalizing symptoms including symptoms of PTSD and MDD assessed using a structured diagnostic interview. This sample is a unique group of healthy but impoverished preschool-aged children. Children with a recent episode of maltreatment identified through the local child welfare agency and an emergency maltreatment assessment service are compared with children with no indicated case of maltreatment, and effects of additional stressors are assessed.

Methods

One-hundred and ninety-eight families enrolled in this study. One child from each family was included. Children ranged in age from 3 to 5 years ($M = 51.6$ months; $SD = 9$ months), 108 were female and 90 were male. The sample was racially and ethnically diverse, with 49 White non-Hispanic, 93 Hispanic, 30 Black, 26 other races. Most caregivers ($n = 184$) were biological mothers. Forty-three caregivers had less than a high school degree, 69 completed high school, 68 had some post-secondary education, 17 had a bachelor's degree, and one did not provide education information. One-hundred and thirteen caregivers were unemployed and 180 of the families qualified for public assistance. Ninety-five children (48%) had substantiated cases of moderate to severe child maltreatment within the past 6 months as described below.

Procedure

Families with a maltreated child were identified from the local child welfare agency and an emergency maltreatment assessment service via record review. Families of children with no indicated case of maltreatment within the past 6 months were recruited at a pediatric medical clinic during a well-child visit as well as at childcare centers. Based on review of available medical records and parent report, children with a chronic illness, medication use, obesity, and failure-to-thrive were excluded. Those with acute illness or medication use were included no less than 2 weeks following resolution of illness and discontinuation of medication. Families completed two home visits and questionnaires between the visits. The current report focuses on the first home visit, during which caregivers completed interviews on child stress exposure and a saliva sample for DNA isolation was collected from the children.

Measures

Child maltreatment status—All families consented to examination of child welfare records to determine maltreatment status. Trained research staff coded the records using the System for Coding Subtype and Severity of Maltreatment in Child Protective Records (Barnett, Manly, & Cicchetti, 1993). Five maltreatment subtypes and severity scores ranging from 1 (least severe) to 5 (most severe) were derived. Children with an episode that met the criteria for moderate to severe maltreatment (score of 3-5) within the last 6 months were included in the maltreated group ($n = 95$). Sixteen children had substantiated cases of physical abuse, 24 sexual abuse, 11 physical neglect/failure to provide, 25 physical neglect/lack of supervision, and 60 emotional maltreatment (emotional maltreatment typically included witnessing domestic violence). The comparison group included children who had never had a substantiated case of maltreatment. Seven children had a remote episode of maltreatment ($M = 33$ months prior to enrollment, $SD = 15$ months prior to enrollment). Results were consistent whether these children were in the maltreatment or comparison group; because they did not meet the criteria for the maltreatment group they were included in the comparison group.

Contextual stress interview—Caregivers completed a semi-structured interview developed in our laboratory to assess the child's experience of contextual stressors in the past 6 months. Categories were: death of a caregiver, separation from a caregiver, frequent change of residence or homelessness, inadequate food or clothing, and other events including witnessing neighborhood violence or parental arrest. Each domain was scored positive if at least one episode occurred. Possible scores ranged from 0 (no stressors) to 5 (stressors in all five domains).

Internalizing Behavior Problems—Caregivers complete the Child Behavior Checklist for Ages 1.5 to 5 (CBCL; (Achenbach & Ruffle, 2000) to assess internalizing behavior problems. For each of the 100 behaviors, parents assessed their children on a 3-point scale from 0 (Not True) to 2 (Very True). T scores were used for data analysis.

PTSD and MDD symptoms—The Diagnostic Infant and Preschool Assessment (DIPA; (Scheeringa & Haslett, 2010) interview was conducted with caregivers to assess symptoms of PTSD and MDD. Interviews were conducted by trained clinical social workers and a Ph.D.-level psychologist, reviewed in a group supervision format, and scored based upon group consensus. Consistent with the low prevalence of MDD and PTSD diagnoses reported in other studies of children under the age of six recruited from low-income mental health clinics (Bufferd, Dougherty, Carlson, & Klein, 2011; Scheeringa & Haslett, 2010), none of the children met full DSM-IV criteria for MDD and only one met criteria for PTSD; therefore the variables for analyses were the number of symptoms of MDD and of PTSD experienced within the past month.

Genotyping—Oragene kits (DNA Genotek, Ontario, Canada) were used to collect saliva, and DNA was extracted according to the manufacturer's protocol. The *IL1B* G⁺⁵⁸¹⁰A (rs1143633) single nucleotide polymorphism (SNP) was genotyped by the TaqMan method using an ABI PRISM 7900 Sequence Detection System (ABI, Foster City, CA, USA). This

SNP was selected because it has previously been associated with inflammatory phenotypes (Hahn, Cho, Kim, Kim, & Kang, 2009; Moxley, Han, Stern, & Riley, 2007; Solovieva et al., 2009), cognitive function (Sasayama, Hori, Teraishi, Hattori, Ota, Matsuo, et al., 2011), schizophrenia (Sasayama, Hori, Teraishi, Hattori, Ota, Iijima, et al., 2011), and with cortisol response to dexamethasone administration (Sasayama, Hori, Iijima, et al., 2011).

Statistical analysis

Mean differences in demographic characteristics and stress variables were examined using *t*-tests and Chi-square. General linear modeling was used for hypothesis testing. Child age, sex, and race (Black vs. all others and Hispanic vs. all others) were included as covariates in all models. Unadjusted means and standard deviations are provided in the text for each group. First, main and interactive effects of child maltreatment and *IL1B* genotype on child behavioral outcomes were tested. Second, main and interactive effects of contextual stress and *IL1B* genotype on child behavioral outcomes were tested. Significant interaction effects were probed by examining group differences in child behavioral outcomes.

Results

Sample Characteristics

The minor allele frequency (MAF) of the *IL1B* allelic variant in the sample was .32 for our total population, and the distribution conformed to the Hardy-Weinberg equilibrium ($X^2 = 0.05$, $p = .82$). Our overall sample MAF was not statistically different from the National Center for Biotechnology Information (NCBI) SNP database (dbSNP)-reported MAF of .35 ($p = .65$). Table 1 shows sample characteristics in relation to *IL1B* genotype; age, sex, and race did not differ according to genotype. Calculated MAF for Whites, Blacks, and Hispanics were .39, .27, and .32, respectively. The distribution of these alleles conformed to Hardy-Weinberg equilibrium ($X^2 = 0.05$, $p = .82$ for Whites, $X^2 = 3.0$, $p = .081$ for Blacks, and $X^2 = 3.1$, $p = .081$ for Hispanics) and did not significantly differ from MAF reported in the NCBI dbSNP for these populations of .40, .23, and .35 ($p = .89$, $p = .51$, $p = .65$, respectively). Low levels of symptoms were identified on the DIPA, with a range of 0-6 for MDD (mean = 0.45, $SD = 0.95$) and 0-15 for PTSD ($M = 1.4$, $SD = 2.6$), with similarly low levels of internalizing symptoms on the CBCL ($M = 52.7$, $SD = 8.8$). Sex was not associated with MDD, PTSD or internalizing symptoms. There were no differences in internalizing symptoms on the CBCL or PTSD or MDD symptoms based on race. Age was not associated with MDD symptoms or internalizing behaviors, but there was a trend association with PTSD symptoms ($p = .06$). Age, sex, and race were controlled in the models as described above.

IL1B genotype, maltreatment and G×E effects on child behavioral outcomes

As displayed in the main effects models in Table 2, maltreatment was associated with both PTSD and MDD symptoms, after controlling for race, sex, and age. In comparison with the non-maltreated children ($N = 103$), the maltreated group ($N = 95$) had more symptoms of PTSD ($M = 2.35$, $SD = 3.10$ versus $M = .54$, $SD = 1.5$) and MDD ($M = 0.62$, $SD = 1.09$ versus $M = 0.30$, $SD = 0.78$). Maltreatment also was associated with internalizing behaviors

at trend level with more internalizing behaviors in the maltreated group ($M = 53.90$, $SD = 7.87$) than non-maltreated children ($M = 51.61$, $SD = 9.39$).

IL1B genotype was associated with MDD symptoms. As illustrated in Figure 1, homozygotes for the A allele had more MDD symptoms ($M = 1.15$, $SD = 1.78$) than carriers of the G allele ($M = 0.40$, $SD = 0.82$ for G homozygotes and $M = 0.35$, $SD = 0.76$ for GA heterozygotes). In contrast, *IL1B* was not associated with internalizing behaviors or PTSD symptoms.

In moderation effects models (Table 2), the interaction of maltreatment and *IL1B* was not a significant predictor of internalizing behaviors, PTSD symptoms, or MDD symptoms.

***IL1B* genotype, contextual stress and G×E effects on child behavioral outcomes**

As displayed in the main effects models in Table 3, the number of contextual stressors was associated with both PTSD and MDD symptoms. Children exposed to more contextual stressors had greater PTSD and MDD symptoms. Contextual stress also was positively associated with internalizing behaviors at trend level.

Consistent with the effect of *IL1B* in the models with maltreatment, *IL1B* was associated with MDD symptoms. Homozygotes for the A allele had greater MDD symptoms than carriers of the G allele. In models with contextual stress, *IL1B* was also positively associated with PTSD symptoms at trend level. A homozygotes had more PTSD symptoms ($M = 2.35$, $SD = 3.51$) than GA heterozygotes ($p = .027$; $M = 1.20$, $SD = 2.12$), with G homozygotes at trend level ($p = .10$; $M = 1.39$, $SD = 2.69$).

Moderation effect models are also displayed in Table 3. The interaction of contextual stress and *IL1B* was significantly associated with MDD symptoms ($p = .045$). As illustrated in Figure 2, children with contextual stress who had the AA genotype had more MDD symptoms than children who were carriers of the G allele and those who had no contextual stress. Following recommendations of Keller (2014), the significant interaction effect was also tested controlling for the interaction of each covariate and *IL1B*, and the interaction of each covariate and contextual stress. None of the eight covariate interactions were significant, and the interaction effect was essentially unchanged ($F = 2.06$, $p = .051$). The interaction of contextual stress and *IL1B* was not associated with internalizing behaviors or PTSD.

Discussion

To our knowledge, this is the first study to examine the role of the *IL1B* gene in the development of internalizing symptoms in children. We found that maltreatment and other contextual stressors are associated with symptoms of PTSD and MDD and that a polymorphism in the *IL1B* gene is associated with depressive symptoms. In addition, the effect of contextual stress was accounted for by an interaction with *IL1B* genotype, such that only those with the AA genotype had higher depressive symptoms in the context of these stressors.

A growing body of research has established that pro-inflammatory cytokines have systemic effects far beyond the canonical immune response. As discussed above, evidence from animal and human studies support a role of inflammatory cytokines in the development of MDD (Groer & Morgan, 2007; Koo & Duman, 2008; O'Brien, Scott, & Dinan, 2004; Pace, Hu, & Miller, 2007) and anxiety disorders (Bauer, Wieck, Lopes, Teixeira, & Grassi-Oliveira, 2010; Hoge et al., 2009; von Kanel et al., 2007) as well as in the broad-ranging adverse health effects of stress-induced inflammation. IL-1 β has been linked to MDD in adults and plays a key role in the neural response to stress, but few studies have examined this cytokine in relation to stress exposure or psychiatric symptoms. A study of adolescent suicides found increased activity of IL-1 β , IL-6, and TNF- α in the prefrontal cortex (Pandey et al., 2012). In a sub-sample of 40 children from our parent study, we recently found that contextual stressors and traumatic life events were associated with saliva concentrations of IL-1 β (Tyrka et al., under review). Only seven subjects in that sub-sample had the AA genotype; these subjects had numerically higher saliva IL-1 β concentrations than G allele carriers but this did not reach significance in the small sample ($p = .11$). Genetic variants in *IL1B* may help define a subset of children who, when exposed to maltreatment, have differential IL-1 β responses and higher internalizing symptoms.

The rs1143633 SNP is an intronic variant in the *IL1B* gene. Introns can contribute to the enhancement of translation of gene products, possibly due to mechanisms including transcription initiation, termination, and regulation (Shabalina et al., 2010). A prior report found that this *IL1B* polymorphism was associated with cortisol response to dexamethasone challenge in a sample of 179 adults (Sasayama, Hori, Iijima, et al., 2011). Abnormal function of the hypothalamic-pituitary-adrenal (HPA) axis is well established in association with early stress exposure and internalizing symptoms and disorders (Carpenter et al., 2007; Carpenter, Shattuck, Tyrka, Geraciotti, & Price, 2011; Hartman, Hermanns, de Jong, & Ormel, 2013; Lupien et al., 2009; Morris, Compas, & Garber, 2012; Stetler & Miller, 2011). The HPA and inflammatory systems are tightly co-regulated, and an effect of this gene in HPA axis activation could be involved in the association between stress exposure and internalizing symptoms and disorders. This SNP has also been associated with decrements in verbal cognitive function in older females (Sasayama, Hori, Teraishi, Hattori, Ota, Matsuo, et al., 2011) and with schizophrenia (Sasayama, Hori, Teraishi, Hattori, Ota, Iijima, et al., 2011), as well as inflammatory phenotypes including arthritis and nephropathies (Hahn et al., 2009; Moxley et al., 2007; Solovieva et al., 2009). Thus, this SNP may be functional or linked with a region that has a functional effect on *IL1B* gene expression.

Our significant effects were seen with symptoms on a clinician-rated interview, but not on the internalizing symptom scale of the parent-report CBCL. Prior work suggests that parent reports may include a degree of bias (Saudino, Wertz, Gagne, & Chawla, 2004; Seifer, 2004), which may have contributed to less robust effects in models predicting parent reports of internalizing symptoms than clinician-rated PTSD and MDD symptoms. It is important to note that in the present study, maltreatment and contextual stressors usually directly involved the parent who completed the CBCL, which may have been another source of bias. Consistent with this, CBCL scores were in the normative range, despite the fact that most of the children experienced some form of adversity and would be expected to have higher

scores. Caregivers also provided the information in the diagnostic interview, but as the assessments were made by trained clinicians, they may be less subject to bias. Internalizing symptoms and clinician-rated MDD symptoms were correlated, $r = 0.22$, $p = .002$, indicating overlap but also substantial unique variance.

The *IL1B* genotype effect in this study was on depressive symptoms, but not symptoms of PTSD. Most of the literature on cytokine associations with psychopathology is on MDD, but there is also evidence of a link with PTSD for several cytokines. PTSD symptoms are tied to the experience of traumatic events, and assessment of symptoms requires knowledge of these events, which is limited in studies of young children.

It is of interest that the interaction of *IL1B* genotype with stress exposure on MDD symptoms was for contextual stressors, which related to separation, loss, or instability of the child's primary supports or neglect of basic needs. Previous work has shown that the lack of social supports predisposes to the development of internalizing symptoms and that genetic variants can moderate this effect (Dunn et al., 2011). That the G×E effect was seen for contextual stress but not maltreatment suggests that maltreatment had a robust effect on symptoms, irrespective of genotype, whereas only children who were at-risk on the basis of *IL1B* genotype had depressive symptoms following the more commonly-encountered types of stressors assessed with this measure.

Strengths of this study include the careful measurement of adverse experiences including documented evidence of maltreatment from child welfare records, in-home assessments with both parent- and clinician-rated assessments, and the involvement of an at-risk, impoverished sample. Limitations include the inclusion of only a single SNP (albeit one with numerous previous association reports) rather than a set that tags the entire locus, the relatively small number of children who were homozygous for the minor allele, and the smaller numbers involved in examining G×E effects. However, the sample size is similar to that of other risk allele groups previously published reporting G×E effects (Cicchetti, Rogosch, & Oshri, 2011). The lack of genetic characterization of ancestry is a further limitation, and this will be addressed in future studies. Children who were on medications or had any chronic medical conditions were excluded from the study, and the sample as a whole had few internalizing symptoms. However, the effect sizes seen were robust, and it is possible that adversity and *IL1B* genotype may predict the development of greater numbers of symptoms and internalizing disorders over time. These findings should be considered preliminary until replicated.

In summary, the current study is the first to demonstrate that a polymorphism in the *IL1B* gene is associated with depressive symptoms in children. Additionally, we found that the effect of contextual stress was accounted for by an interaction with *IL1B* genotype, such that only those with the AA genotype had higher depressive symptoms in the context of these stressors. The results of this preliminary study will be helpful in elucidating the effects of genetic variation in the *IL1B* and other inflammatory cytokine genes on internalizing symptoms and as a moderator of the effects of childhood adversity.

Acknowledgments

This research was supported by grant R01 MH083704 awarded to the last author from the National Institute of Mental Health. The content is solely the responsibility of the authors and does not necessarily reflect the official views of the NIMH.

References

- Achenbach TM, Ruffle TM. The Child Behavior Checklist and related forms for assessing behavioral/emotional problems and competencies. *Pediatrics in review / American Academy of Pediatrics*. 2000; 21(8):265–271. [PubMed: 10922023]
- Aguilera M, Arias B, Wichers M, Barrantes-Vidal N, Moya J, Villa H, Fananas L. Early adversity and 5-HTT/BDNF genes: new evidence of gene-environment interactions on depressive symptoms in a general population. *Psychological medicine*. 2009; 39(9):1425–1432. [PubMed: 19215635]
- Appleton AA, Buka SL, McCormick MC, Koenen KC, Loucks EB, Kubzansky LD. The association between childhood emotional functioning and adulthood inflammation is modified by early-life socioeconomic status. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association*. 2012; 31(4):413–422.
- Aslund C, Leppert J, Comasco E, Nordquist N, Orelund L, Nilsson KW. Impact of the interaction between the 5HTTLPR polymorphism and maltreatment on adolescent depression. A population-based study. *Behavior genetics*. 2009; 39(5):524–531.
- Bailey MT, Kinsey SG, Padgett DA, Sheridan JF, Leblebicioglu B. Social stress enhances IL-1beta and TNF-alpha production by Porphyromonas gingivalis lipopolysaccharide-stimulated CD11b+ cells. *Physiology & behavior*. 2009; 98(3):351–358. [PubMed: 19560480]
- Baker DG, Nievergelt CM, O'Connor DT. Biomarkers of PTSD: neuropeptides and immune signaling. *Neuropharmacology*. 2012; 62(2):663–673. [PubMed: 21392516]
- Banny AM, Cicchetti D, Rogosch FA, Oshri A, Crick NR. Vulnerability to depression: a moderated mediation model of the roles of child maltreatment, peer victimization, and serotonin transporter linked polymorphic region genetic variation among children from low socioeconomic status backgrounds. *Development and psychopathology*. 2013; 25(3):599–614. [PubMed: 23880379]
- Barnett, D.; Manly, JT.; Cicchetti, D. Defining child maltreatment: The interface between policy and research. In: Cicchetti, D.; Toth, SL., editors. *Child abuse, child development, and social policy*. Ablex Publishing Corp; Norwood, NJ: 1993. p. 7-73.
- Bauer ME, Wieck A, Lopes RP, Teixeira AL, Grassi-Oliveira R. Interplay between neuroimmunoendocrine systems during post-traumatic stress disorder: a minireview. *Neuroimmunomodulation*. 2010; 17(3):192–195. [PubMed: 20134200]
- Bellani M, Nobile M, Bianchi V, van Os J, Brambilla P. G × E interaction and neurodevelopment I. Focus on maltreatment. *Epidemiology and psychiatric sciences*. 2012; 21(4):347–351.
- Belsky J, Jonassaint C, Pluess M, Stanton M, Brummett B, Williams R. Vulnerability genes or plasticity genes? *Molecular psychiatry*. 2009; 14(8):746–754. [PubMed: 19455150]
- Belsky J, Pluess M. Beyond diathesis stress: differential susceptibility to environmental influences. *Psychological bulletin*. 2009; 135(6):885–908. [PubMed: 19883141]
- Benjet C, Borges G, Medina-Mora ME. Chronic childhood adversity and onset of psychopathology during three life stages: childhood, adolescence and adulthood. *Journal of psychiatric research*. 2010; 44(11):732–740. [PubMed: 20144464]
- Bertone-Johnson ER, Whitcomb BW, Missmer SA, Karlson EW, Rich-Edwards JW. Inflammation and early-life abuse in women. *American journal of preventive medicine*. 2012; 43(6):611–620. [PubMed: 23159256]
- Briggs-Gowan MJ, Carter AS, Bosson-Heenan J, Guyer AE, Horwitz SM. Are infant-toddler social-emotional and behavioral problems transient? *Journal of the American Academy of Child and Adolescent Psychiatry*. 2006; 45(7):849–858. [PubMed: 16832322]
- Brydon L, Edwards S, Jia H, Mohamed-Ali V, Zachary I, Martin JF, Steptoe A. Psychological stress activates interleukin-1beta gene expression in human mononuclear cells. *Brain, behavior, and immunity*. 2005; 19(6):540–546.

- Bufferd SJ, Dougherty LR, Carlson GA, Klein DN. Parent-reported mental health in preschoolers: findings using a diagnostic interview. *Comprehensive psychiatry*. 2011; 52(4):359–369. [PubMed: 21683173]
- Carpenter LL, Carvalho JP, Tyrka AR, Wier LM, Mello AF, Mello MF, Price LH. Decreased adrenocorticotrophic hormone and cortisol responses to stress in healthy adults reporting significant childhood maltreatment. *Biological psychiatry*. 2007; 62(10):1080–1087. [PubMed: 17662255]
- Carpenter LL, Shattuck TT, Tyrka AR, Geraciotti TD, Price LH. Effect of childhood physical abuse on cortisol stress response. *Psychopharmacology*. 2011; 214(1):367–375. [PubMed: 20838776]
- Carr CP, Martins CM, Stingel AM, Lemgruber VB, Juruena MF. The role of early life stress in adult psychiatric disorders: a systematic review according to childhood trauma subtypes. *The Journal of nervous and mental disease*. 2013; 201(12):1007–1020. [PubMed: 24284634]
- Carrion VG, Wong SS. Can traumatic stress alter the brain? Understanding the implications of early trauma on brain development and learning. *The Journal of adolescent health : official publication of the Society for Adolescent Medicine*. 2012; 51(2 Suppl):S23–28. [PubMed: 22794529]
- Caso JR, Moro MA, Lorenzo P, Lizasoain I, Leza JC. Involvement of IL-1 β in acute stress-induced worsening of cerebral ischaemia in rats. 2007; 17(9):600–607.
- Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, Poulton R. Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science*. 2003; 301(5631):386–389. [PubMed: 12869766]
- Cicchetti D, Rogosch FA. Gene $\times$ Environment interaction and resilience: effects of child maltreatment and serotonin, corticotropin releasing hormone, dopamine, and oxytocin genes. *Development and psychopathology*. 2012; 24(2):411–427. [PubMed: 22559122]
- Cicchetti D, Rogosch FA, Oshri A. Interactive effects of corticotropin releasing hormone receptor 1, serotonin transporter linked polymorphic region, and child maltreatment on diurnal cortisol regulation and internalizing symptomatology. *Development and psychopathology*. 2011; 23(4):1125–1138. [PubMed: 22018085]
- Cicchetti D, Rogosch FA, Sturge-Apple ML. Interactions of child maltreatment and serotonin transporter and monoamine oxidase A polymorphisms: depressive symptomatology among adolescents from low socioeconomic status backgrounds. *Development and psychopathology*. 2007; 19(4):1161–1180. [PubMed: 17931441]
- Cicchetti D, Rogosch FA, Toth SL. The effects of child maltreatment and polymorphisms of the serotonin transporter and dopamine D4 receptor genes on infant attachment and intervention efficacy. *Development and psychopathology*. 2011; 23(2):357–372. [PubMed: 23786683]
- Cicchetti D, Toth SL. Child maltreatment. *Annual review of clinical psychology*. 2005; 1:409–438.
- Coelho R, Viola TW, Walss-Bass C, Brietzke E, Grassi-Oliveira R. Childhood maltreatment and inflammatory markers: a systematic review. *Acta psychiatrica Scandinavica*. 2014; 129(3):180–192. [PubMed: 24205846]
- Cole SW, Arevalo JM, Manu K, Telzer EH, Kiang L, Bower JE, Fuligni AJ. Antagonistic pleiotropy at the human IL6 promoter confers genetic resilience to the pro-inflammatory effects of adverse social conditions in adolescence. *Developmental psychology*. 2011; 47(4):1173–1180. [PubMed: 21639625]
- Cole SW, Arevalo JM, Takahashi R, Sloan EK, Lutgendorf SK, Sood AK, Seeman TE. Computational identification of gene-social environment interaction at the human IL6 locus. *Proceedings of the National Academy of Sciences of the United States of America*. 2010; 107(12):5681–5686. [PubMed: 20176930]
- Comasco E, Aslund C, Orelund L, Nilsson KW. Three-way interaction effect of 5-HTTLPR, BDNF Val66Met, and childhood adversity on depression: a replication study. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology*. 2013; 23(10):1300–1306. [PubMed: 23481907]
- Danese A, Moffitt TE, Pariante CM, Ambler A, Poulton R, Caspi A. Elevated inflammation levels in depressed adults with a history of childhood maltreatment. *Archives of general psychiatry*. 2008; 65(4):409–415. [PubMed: 18391129]

- Danese A, Pariante CM, Caspi A, Taylor A, Poulton R. Childhood maltreatment predicts adult inflammation in a life-course study. *Proceedings of the National Academy of Sciences of the United States of America*. 2007; 104(4):1319–1324. [PubMed: 17229839]
- Dantzer R. Cytokine, sickness behavior, and depression. *Immunology and allergy clinics of North America*. 2009; 29(2):247–264. [PubMed: 19389580]
- Dowlati Y, Herrmann N, Swardfager W, Liu H, Sham L, Reim EK, Lancot KL. A meta-analysis of cytokines in major depression. *Biological psychiatry*. 2010; 67(5):446–457. [PubMed: 20015486]
- Dunn EC, Uddin M, Subramanian SV, Smoller JW, Galea S, Koenen KC. Research review: gene-environment interaction research in youth depression - a systematic review with recommendations for future research. *Journal of child psychology and psychiatry, and allied disciplines*. 2011; 52(12):1223–1238.
- Felger JC, Lotrich FE. Inflammatory cytokines in depression: neurobiological mechanisms and therapeutic implications. *Neuroscience*. 2013; 246:199–229. [PubMed: 23644052]
- Ferri C, Croce G, Cofini V, De Berardinis G, Grassi D, Casale R, Desideri G. C-reactive protein: interaction with the vascular endothelium and possible role in human atherosclerosis. *Current pharmaceutical design*. 2007; 13(16):1631–1645. [PubMed: 17584094]
- Gola H, Engler H, Sommershof A, Adenauer H, Kolassa S, Schedlowski M, Kolassa IT. Posttraumatic stress disorder is associated with an enhanced spontaneous production of pro-inflammatory cytokines by peripheral blood mononuclear cells. *BMC psychiatry*. 2013; 13:40. [PubMed: 23360282]
- Green JG, McLaughlin KA, Berglund PA, Gruber MJ, Sampson NA, Zaslavsky AM, Kessler RC. Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication I: associations with first onset of DSM-IV disorders. *Archives of general psychiatry*. 2010; 67(2):113–123. [PubMed: 20124111]
- Groer MW, Morgan K. Immune, health and endocrine characteristics of depressed postpartum mothers. *Psychoneuroendocrinology*. 2007; 32(2):133–139. [PubMed: 17207585]
- Haastrup E, Bukh JD, Bock C, Vinberg M, Thorner LW, Hansen T, Ullum H. Promoter variants in IL18 are associated with onset of depression in patients previously exposed to stressful-life events. *Journal of affective disorders*. 2012; 136(1-2):134–138. [PubMed: 21962565]
- Hahn WH, Cho BS, Kim SD, Kim SK, Kang S. Interleukin-1 cluster gene polymorphisms in childhood IgA nephropathy. *Pediatric nephrology*. 2009; 24(7):1329–1336. [PubMed: 19280228]
- Hartman CA, Hermanns VW, de Jong PJ, Ormel J. Self- or parent report of (co-occurring) internalizing and externalizing problems, and basal or reactivity measures of HPA-axis functioning: a systematic evaluation of the internalizing-hyperresponsivity versus externalizing-hyporesponsivity HPA-axis hypothesis. *Biological psychology*. 2013; 94(1):175–184. [PubMed: 23735709]
- Heim C, Binder EB. Current research trends in early life stress and depression: review of human studies on sensitive periods, gene-environment interactions, and epigenetics. *Experimental neurology*. 2012; 233(1):102–111. [PubMed: 22101006]
- Hickman LJ, Jaycox LH, Setodji CM, Kofner A, Schultz D, Barnes-Proby D, Harris R. How much does "how much" matter? Assessing the relationship between children's lifetime exposure to violence and trauma symptoms, behavior problems, and parenting stress. *Journal of interpersonal violence*. 2013; 28(6):1338–1362. [PubMed: 23266994]
- Hiles SA, Baker AL, de Malmanche T, Attia J. A meta-analysis of differences in IL-6 and IL-10 between people with and without depression: exploring the causes of heterogeneity. *Brain, behavior, and immunity*. 2012; 26(7):1180–1188.
- Hoge EA, Brandstetter K, Moshier S, Pollack MH, Wong KK, Simon NM. Broad spectrum of cytokine abnormalities in panic disorder and posttraumatic stress disorder. *Depression and anxiety*. 2009; 26(5):447–455. [PubMed: 19319993]
- Hornung OP, Heim CM. Gene-Environment Interactions and Intermediate Phenotypes: Early Trauma and Depression. *Frontiers in endocrinology*. 2014; 5:14. [PubMed: 24596569]
- Kaufman J, Yang BZ, Douglas-Palumberi H, Grasso D, Lipschitz D, Houshyar S, Gelernter J. Brain-derived neurotrophic factor-5-HTTLPR gene interactions and environmental modifiers of depression in children. *Biological psychiatry*. 2006; 59(8):673–680. [PubMed: 16458264]

- Kaufman J, Yang BZ, Douglas-Palumberi H, Houshyar S, Lipschitz D, Krystal JH, Gelernter J. Social supports and serotonin transporter gene moderate depression in maltreated children. *Proceedings of the National Academy of Sciences of the United States of America*. 2004; 101(49):17316–17321. [PubMed: 15563601]
- Keller MC. Gene $\times$ environment interaction studies have not properly controlled for potential confounders: the problem and the (simple) solution. *Biological psychiatry*. 2014; 75(1):18–24. [PubMed: 24135711]
- Kim-Spoon J, Cicchetti D, Rogosch FA. A longitudinal study of emotion regulation, emotion lability-negativity, and internalizing symptomatology in maltreated and nonmaltreated children. *Child development*. 2013; 84(2):512–527. [PubMed: 23034132]
- Kim JM, Stewart R, Kim SW, Kim SY, Bae KY, Kang HJ, Yoon JS. Physical health and incident late-life depression: modification by cytokine genes. *Neurobiology of aging*. 2013; 34(1):e351–359. 356.
- Koo JW, Duman RS. IL-1beta is an essential mediator of the antineurogenic and anhedonic effects of stress. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105(2):751–756. [PubMed: 18178625]
- Lacy P, Stow JL. Cytokine release from innate immune cells: association with diverse membrane trafficking pathways. *Blood*. 2011; 118(1):9–18. [PubMed: 21562044]
- Lavie CJ, Milani RV, Verma A, O'Keefe JH. C-reactive protein and cardiovascular diseases--is it ready for primetime? *The American journal of the medical sciences*. 2009; 338(6):486–492. [PubMed: 20010156]
- Lesch KP, Bengel D, Heils A, Sabol SZ, Greenberg BD, Petri S, Murphy DL. Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science*. 1996; 274(5292):1527–1531. [PubMed: 8929413]
- Liu Y, Ho RC, Mak A. Interleukin (IL)-6, tumour necrosis factor alpha (TNF-alpha) and soluble interleukin-2 receptors (sIL-2R) are elevated in patients with major depressive disorder: a meta-analysis and meta-regression. *Journal of affective disorders*. 2012; 139(3):230–239. [PubMed: 21872339]
- Low CA, Matthews KA, Hall M. Elevated C-reactive protein in adolescents: roles of stress and coping. *Psychosomatic medicine*. 2013; 75(5):449–452. [PubMed: 23576771]
- Luby JL, Gaffrey MS, Tillman R, April LM, Belden AC. Trajectories of Preschool Disorders to Full DSM Depression at School Age and Early Adolescence: Continuity of Preschool Depression. *The American journal of psychiatry*. 2014
- Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature reviews. Neuroscience*. 2009; 10(6):434–445. [PubMed: 19401723]
- Mandelli L, Serretti A. Gene environment interaction studies in depression and suicidal behavior: An update. *Neuroscience and biobehavioral reviews*. 2013; 37:2375–2397. 10 Pt 1. [PubMed: 23886513]
- Manly JT, Kim JE, Rogosch FA, Cicchetti D. Dimensions of child maltreatment and children's adjustment: contributions of developmental timing and subtype. *Development and psychopathology*. 2001; 13(4):759–782. [PubMed: 11771907]
- Masten AS, Cicchetti D. Developmental cascades. *Development and psychopathology*. 2010; 22(3):491–495. [PubMed: 20576173]
- Mastrolonardo M, Alicino D, Zefferino R, Pasquini P, Picardi A. Effect of psychological stress on salivary interleukin-1beta in psoriasis. *Archives of medical research*. 2007; 38(2):206–211. [PubMed: 17227730]
- Mesman J, Koot HM. Early preschool predictors of preadolescent internalizing and externalizing DSM-IV diagnoses. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001; 40(9):1029–1036. [PubMed: 11556626]
- Mills, Scott J, Alati R, O'Callaghan M, Najman JM, Strathearn L. Child maltreatment and adolescent mental health problems in a large birth cohort. *Child abuse & neglect*. 2013; 37(5):292–302. [PubMed: 23380430]

- Mills NT, Scott JG, Wray NR, Cohen-Woods S, Baune BT. Research review: the role of cytokines in depression in adolescents: a systematic review. *Journal of child psychology and psychiatry, and allied disciplines*. 2013; 54(8):816–835.
- Mills R, Scott J, Alati R, O'Callaghan M, Najman JM, Strathearn L. Child maltreatment and adolescent mental health problems in a large birth cohort. *Child abuse & neglect*. 2013; 37(5):292–302. [PubMed: 23380430]
- Mitchell RH, Goldstein BI. Inflammation in children and adolescents with neuropsychiatric disorders: a systematic review. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2014; 53(3):274–296. [PubMed: 24565356]
- Morris MC, Compas BE, Garber J. Relations among posttraumatic stress disorder, comorbid major depression, and HPA function: a systematic review and meta-analysis. *Clinical psychology review*. 2012; 32(4):301–315. [PubMed: 22459791]
- Moxley G, Han J, Stern AG, Riley BP. Potential influence of IL1B haplotype and IL1A-IL1B-IL1RN extended haplotype on hand osteoarthritis risk. *Osteoarthritis and cartilage / OARS, Osteoarthritis Research Society*. 2007; 15(10):1106–1112.
- Nederhof E, Bouma EM, Oldehinkel AJ, Ormel J. Interaction between childhood adversity, brain-derived neurotrophic factor val/met and serotonin transporter promoter polymorphism on depression: the TRAILS study. *Biological psychiatry*. 2010; 68(2):209–212. [PubMed: 20553751]
- Nguyen KT, Deak T, Owens SM, Kohno T, Fleshner M, Watkins LR, Maier SF. Exposure to acute stress induces brain interleukin-1beta protein in the rat. 1998; 18(6):2239–2246.
- Norman RE, Byambaa M, De R, Butchart A, Scott J, Vos T. The long-term health consequences of child physical abuse, emotional abuse, and neglect: a systematic review and meta-analysis. *PLoS medicine*. 2012; 9(11):e1001349. [PubMed: 23209385]
- Nugent NR, Tyrka AR, Carpenter LL, Price LH. Gene-environment interactions: early life stress and risk for depressive and anxiety disorders. *Psychopharmacology*. 2011; 214(1):175–196. [PubMed: 21225419]
- O'Brien SM, Scott LV, Dinan TG. Cytokines: abnormalities in major depression and implications for pharmacological treatment. *Human psychopharmacology*. 2004; 19(6):397–403. [PubMed: 15303243]
- O'Donovan A, Sun B, Cole S, Rempel H, Lenoci M, Pulliam L, Neylan T. Transcriptional control of monocyte gene expression in post-traumatic stress disorder. *Disease markers*. 2011; 30(2-3):123–132. [PubMed: 21508516]
- Pace TW, Hu F, Miller AH. Cytokine-effects on glucocorticoid receptor function: relevance to glucocorticoid resistance and the pathophysiology and treatment of major depression. *Brain, behavior, and immunity*. 2007; 21(1):9–19.
- Pace TW, Wingenfeld K, Schmidt I, Meinlschmidt G, Hellhammer DH, Heim CM. Increased peripheral NF-kappaB pathway activity in women with childhood abuse-related posttraumatic stress disorder. *Brain, behavior, and immunity*. 2012; 26(1):13–17.
- Pandey GN, Rizavi HS, Ren X, Fareed J, Hoppensteadt DA, Roberts RC, Dwivedi Y. Proinflammatory cytokines in the prefrontal cortex of teenage suicide victims. *Journal of psychiatric research*. 2012; 46(1):57–63. [PubMed: 21906753]
- Pietrek C, Elbert T, Weierstall R, Muller O, Rockstroh B. Childhood adversities in relation to psychiatric disorders. *Psychiatry research*. 2013; 206(1):103–110. [PubMed: 23261184]
- Porterfield VM, Gabella KM, Simmons MA, Johnson JD. Repeated stressor exposure regionally enhances beta-adrenergic receptor-mediated brain IL-1beta production. *Brain, behavior, and immunity*. 2012; 26(8):1249–1255.
- Rincon-Cortes M, Sullivan RM. Early Life Trauma and Attachment: Immediate and Enduring Effects on Neurobehavioral and Stress Axis Development. *Frontiers in endocrinology*. 2014; 5:33. [PubMed: 24711804]
- Robles TF, Glaser R, Kiecolt-Glaser JK. Out of Balance: A New Look at Chronic Stress, Depression, and Immunity. 2005; 14(2):111–115.
- Rohleder N. Stimulation of systemic low-grade inflammation by psychosocial stress. *Psychosomatic medicine*. 2014; 76(3):181–189. [PubMed: 24608036]

- Rooks C, Veledar E, Goldberg J, Bremner JD, Vaccarino V. Early trauma and inflammation: role of familial factors in a study of twins. *Psychosomatic medicine*. 2012; 74(2):146–152. [PubMed: 22286843]
- Roza SJ, Hofstra MB, van der Ende J, Verhulst FC. Stable prediction of mood and anxiety disorders based on behavioral and emotional problems in childhood: a 14-year follow-up during childhood, adolescence, and young adulthood. *The American journal of psychiatry*. 2003; 160(12):2116–2121. [PubMed: 14638580]
- Sasayama D, Hori H, Iijima Y, Teraishi T, Hattori K, Ota M, Kunugi H. Modulation of cortisol responses to the DEX/CRH test by polymorphisms of the interleukin-1beta gene in healthy adults. *Behavioral and brain functions : BBF*. 2011; 7:23. [PubMed: 21726461]
- Sasayama D, Hori H, Teraishi T, Hattori K, Ota M, Iijima Y, Kunugi H. Possible association between interleukin-1beta gene and schizophrenia in a Japanese population. *Behavioral and brain functions : BBF*. 2011; 7:35. [PubMed: 21843369]
- Sasayama D, Hori H, Teraishi T, Hattori K, Ota M, Matsuo J, Kunugi H. Association of interleukin-1beta genetic polymorphisms with cognitive performance in elderly females without dementia. *Journal of human genetics*. 2011; 56(8):613–616. [PubMed: 21614008]
- Saudino KJ, Wertz AE, Gagne JR, Chawla S. Night and day: are siblings as different in temperament as parents say they are? *Journal of personality and social psychology*. 2004; 87(5):698–706. [PubMed: 15535780]
- Scheeringa MS, Haslett N. The reliability and criterion validity of the Diagnostic Infant and Preschool Assessment: a new diagnostic instrument for young children. *Child psychiatry and human development*. 2010; 41(3):299–312. [PubMed: 20052532]
- Schilling EA, Aseltine RH Jr, Gore S. Adverse childhood experiences and mental health in young adults: a longitudinal survey. *BMC public health*. 2007; 7:30. [PubMed: 17343754]
- Scott KM, Smith DR, Ellis PM. Prospectively ascertained child maltreatment and its association with DSM-IV mental disorders in young adults. *Archives of general psychiatry*. 2010; 67(7):712–719. [PubMed: 20603452]
- Seifer R, Sameroff A, Dickstein S, Schiller M, Hayden LC. Your own children are special: Clues to the sources of reporting bias in temperament assessments. 2004; 27(3):323–341.
- Shabalina SA, Ogurtsov AY, Spiridonov AN, Novichkov PS, Spiridonov NA, Koonin EV. Distinct patterns of expression and evolution of intronless and intron-containing mammalian genes. *Molecular biology and evolution*. 2010; 27(8):1745–1749. [PubMed: 20360214]
- Shikanai H, Kimura S, Togashi H. Early life stress affects the serotonergic system underlying emotional regulation. *Biological & pharmaceutical bulletin*. 2013; 36(9):1392–1395. [PubMed: 23995647]
- Sjoberg RL, Nilsson KW, Nordquist N, Ohrvik J, Leppert J, Lindstrom L, Orelund L. Development of depression: sex and the interaction between environment and a promoter polymorphism of the serotonin transporter gene. *The international journal of neuropsychopharmacology / official scientific journal of the Collegium Internationale Neuropsychopharmacologicum*. 2006; 9(4):443–449.
- Slopen N, Kubzansky LD, McLaughlin KA, Koenen KC. Childhood adversity and inflammatory processes in youth: a prospective study. *Psychoneuroendocrinology*. 2013; 38(2):188–200. [PubMed: 22727478]
- Solovieva S, Kamarainen OP, Hirvonen A, Hamalainen S, Laitala M, Vehmas T, Leino-Arjas P. Association between interleukin 1 gene cluster polymorphisms and bilateral distal interphalangeal osteoarthritis. *The Journal of rheumatology*. 2009; 36(9):1977–1986. [PubMed: 19684156]
- Steptoe A, Hamer M, Chida Y. The effects of acute psychological stress on circulating inflammatory factors in humans: a review and meta-analysis. *Brain, behavior, and immunity*. 2007; 21(7):901–912.
- Stetler C, Miller GE. Depression and hypothalamic-pituitary-adrenal activation: a quantitative summary of four decades of research. *Psychosomatic medicine*. 2011; 73(2):114–126. [PubMed: 21257974]

- Teicher MH, Samson JA. Childhood maltreatment and psychopathology: A case for ecophenotypic variants as clinically and neurobiologically distinct subtypes. *The American journal of psychiatry*. 2013; 170(10):1114–1133. [PubMed: 23982148]
- Tyrka AR, Wier L, Price LH, Ross NS, Carpenter LL. Childhood parental loss and adult psychopathology: effects of loss characteristics and contextual factors. *International journal of psychiatry in medicine*. 2008; 38(3):329–344. [PubMed: 19069576]
- von Kanel R, Hepp U, Kraemer B, Traber R, Keel M, Mica L, Schnyder U. Evidence for low-grade systemic proinflammatory activity in patients with posttraumatic stress disorder. *Journal of psychiatric research*. 2007; 41(9):744–752. [PubMed: 16901505]
- Widom CS. Posttraumatic stress disorder in abused and neglected children grown up. *The American journal of psychiatry*. 1999; 156(8):1223–1229. [PubMed: 10450264]
- Widom CS, DuMont K, Czaja SJ. A prospective investigation of major depressive disorder and comorbidity in abused and neglected children grown up. *Archives of general psychiatry*. 2007; 64(1):49–56. [PubMed: 17199054]
- Yamakawa K, Matsunaga M, Isowa T, Kimura K, Kasugai K, Yoneda M, Ohira H. Transient responses of inflammatory cytokines in acute stress. *Biological psychology*. 2009; 82(1):25–32. [PubMed: 19446599]
- You Z, Luo C, Zhang W, Chen Y, He J, Zhao Q, Wu Y. Pro- and anti-inflammatory cytokines expression in rat's brain and spleen exposed to chronic mild stress: involvement in depression. *Behavioural brain research*. 2011; 225(1):135–141. [PubMed: 21767575]
- Zeanah CH, Boris NW, Scheeringa MS. Psychopathology in infancy. *Journal of child psychology and psychiatry, and allied disciplines*. 1997; 38(1):81–99.

Figure 1.

Association of *IL1B* and MDD symptoms

Notes: ** Indicates AA homozygotes are significantly different from the two other groups at $p < .01$.

Figure 2.

Interaction of Adversity and *IL1B*

Note: *** Indicates AA homozygotes are significantly different from the five other groups at $p < .001$.

Table 1Descriptive statistics and mean differences by *IL1B* genotype

	GG (<i>n</i> = 90)	GA (<i>n</i> = 88)	AA (<i>n</i> = 20)	<i>p</i>
Sex, <i>N</i> (%) male	38 (42.2)	42 (47.7)	10 (50.0)	.69
Age, <i>M</i> (<i>SD</i>)	4.3 (0.7)	4.4 (0.8)	4.4 (0.8)	.49
Race, <i>N</i> (%)				.11
White	18 (20.0)	24 (27.3)	7 (35.0)	
Black	18 (20.0)	8 (9.1)	4 (20.0)	
Hispanic	39 (43.3)	48 (54.5)	6 (30.0)	
Maltreated Status, <i>N</i> (%)	40 (44.4)	45 (51.1)	10 (50.0)	.66
Number of Stressors, <i>M</i> (<i>SD</i>)	1.1 (1.1)	1.2 (1.1)	0.9 (1.1)	.55
Any Adversity, <i>N</i> (%)	59 (65.6)	68 (77.3)	14 (70.0)	.22

Note: *p* values indicate *t*-test or chi-square significance level.

Table 2Main and Interactive Effects of Child Maltreatment and *IL1B* Genotype

	Internalizing	PTSD symptoms	MDD symptoms
Main Effects Models			
Maltreatment	$F = 3.14, p = .078$	$F = 23.73, p < .001$	$F = 4.69, p = .032$
<i>IL1B</i>	$F = 1.87, p = .157$	$F = 1.87, p = .157$	$F = 6.21, p = .002$
Moderation Effect Models			
Maltreatment X <i>IL1B</i>	$F = 1.28, p = .281$	$F = 0.37, p = .688$	$F = 0.11, p = .899$

Notes: Moderation effect models included main effects but not shown for parsimony.

Child age, sex, and race (Hispanic vs. all others and Black vs. all others) were included as covariates in all models.

Table 3Main and Interactive Effects of Contextual Stress and *IL1B* Genotype

	Internalizing	PTSD symptoms	MDD symptoms
Main Effects Models			
Contextual Stress	$F = 2.14, p = .063$	$F = 12.66, p < .001$	$F = 9.46, p < .001$
<i>IL1B</i>	$F = 1.64, p = .197$	$F = 2.52, p = .083$	$F = 7.07, p = .001$
Moderation Effect Models			
Contextual Stress X <i>IL1B</i>	$F = 1.53, p = .162$	$F = 0.97, p = .460$	$F = 2.03, p = .045$

Notes: Moderation effect models included main effects but not shown for parsimony. Child age, sex, and race (Hispanic vs. all others and Black vs. all others) were included as covariates in all models.