Published in final edited form as:

Brain Inj. 2008 August ; 22(9): 649–656. doi:10.1080/02699050802227162.

Parent perceptions of school-based support for students with traumatic brain injuries

SUSAN DAVIES GFROERER¹, SHARI L. WADE^{2,3}, and MICHELLE WU²

¹The University of Dayton, Dayton, OH, USA

²Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

³University of Cincinnati, Cincinnati, OH, USA

Abstract

Primary objective—To determine whether parents believe schools provided necessary support to their children who sustained traumatic brain injuries.

Research design—Interview, to determine parent perceptions.

Methods and procedure—Sixty-six primary caregivers of school-age children who experienced a TBI within the previous 2 years were interviewed regarding what types of special support were needed by and provided for their children during the 3 months immediately following school re-entry. They then rated how difficult it was to obtain support or services from the school and how satisfied they were with the support or services.

Main outcomes and results—The majority of participants did not perceive the need for school-based services, even when the injury was severe. Almost all students whose parents perceived a need for an adjusted schedule were granted that accommodation, but few students received school-based counselling or behavioural support.

Conclusions—Results indicated that participants perceived relatively few school-based supports, particularly given the actual academic, behavioural and social challenges experienced by children who have sustained a TBI. Schools and hospitals must take steps to ensure appropriate post-head injury support services.

Keywords

Traumatic brain injury; parent perception; return to school; school services; school support

Introduction

In 1990, the government of the USA passed legislation requiring provision of special education services for students with traumatic brain injuries (TBI) that adversely affect

© 2008 Informa Healthcare USA, Inc.

Correspondence: Susan Davies Gfroerer, Department of Counselor Education and Human Services, University of Dayton, 300 College Park, Dayton, OH 45469-0530, USA. gfroersd@notes.udayton.edu.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

educational performance [1]. More than one million children in the US sustain TBI every year, 30 000 of whom experience long-term disabilities as a result of the injury [2]. With the advancements in medical technology, many children with severe TBI now survive [3]. However, these survivors often experience a myriad of physical, cognitive and social-emotional deficits [4]. Specifically, severe TBI can contribute to physical impairments (i.e. neuromotor difficulties), lowered cognitive and academic skills relative to age expectations or pre-injury estimates and problems in school performance, behaviour, socialization and adaptive functioning [5–10]. Beyond documented global reductions in ability as measured by IQ testing, mild-to-severe TBI may result in specific impairments in language, memory, problem-solving, perceptual-motor skills, attention and executive function (EF) [5, 6, 11–13]. Each of these sequelae might impede subsequent development and learning [14]. Additionally, problems with attention, memory and fatigue may be subtle and difficult to distinguish from low motivation or pre-morbid issues. Thus, increasing rates of trauma survival have required the development of programmes and interventions to meet the long-term needs of these individuals [15].

Despite the prevalence and adverse effects of TBIs, very few systematic studies have addressed the educational needs of students with TBI [16]. Studying children with brain injuries can be difficult because one must consider neurodevelopmental aspects of the paediatric population, including age of injury and developmental stage [15, 17]. Mounting evidence suggests that children injured at an early age may experience more significant long-term deficits, with emerging skills being most vulnerable to disruption [17–19]. Fletcher-Janzen and Kade [15] summarized research that indicated that although the young brain may allow short-term recovery by shifting functions to another area of the brain, later abilities might be compromised. Thus, a child may temporarily appear superficially intact and then ‘grow into a deficit’ when the developing brain structures, that normally control an activity, become more important for the execution of the behaviour [20]. For example, a child injured at the age of 3 might not show observable outcomes of the injury until first grade, when greater sustained attention to task is expected. As a result of the delayed onset of symptoms, the school problems experienced by children who have sustained a TBI may be misattributed to factors such as laziness, an oppositional nature or emotional instability, rather than to the injury that occurred several years before. The failure of the school to recognize the types and degree of support needed may, in turn, lead to further behavioural deterioration.

The transition from a hospital or rehabilitation setting to school can be quite difficult for children who have sustained a TBI [21]. Many children who have sustained a severe TBI miss school due to the acute hospitalization and inpatient and outpatient rehabilitation, while others are injured during the summer months when schools are not in session. When children who have sustained brain injuries return to school, their teachers may not be aware of the child’s injuries and subsequent educational needs, particularly if there is a lack of communication between hospital/rehabilitation, parents and school. Readjusting to the school environment depends largely upon the quality of the transition plan that is implemented by the student’s care team. The plan must specify how the learning environment needs to be adapted in order to accommodate the child’s physical and cognitive deficits. Further, many of the challenges associated with school re-entry are related to

behavioural rather than academic concerns, such as the child's need to develop or re-learn the social skills enabling him or her to function within the school environment.

Despite the research supporting specialized service delivery for students with TBI, relatively few students with head injuries are identified under the TBI label [22]. Thus, school personnel may lack the experience to fully understand the unique needs of students with head injuries. Failure to appropriately identify students' needs may reduce the likelihood of their educational needs being met. When Glang et al. [23] surveyed parents whose children have sustained a TBI throughout Oregon, they found that lack of school staff knowledge of TBI and its effects was the primary reason for dissatisfaction with their children's instructional services.

Few studies have examined parent perceptions of students' educational needs [24–26]. These studies identified parents as important informants regarding educational issues such as eligibility assessments for students with special needs, inclusion of students with disabilities and academic engagement of typically developing students. Given the intimate knowledge parents have of their children's strengths and weaknesses and the fact that parents are critical partners in school-based intervention assistance teams, it is important that their perceptions of their child's educational needs be carefully considered. The importance of parental perceptions is evidenced in the fact that multi-factored evaluations of student performance typically include parent interviews and/or the completion of standardized checklists (e.g. Child Behavior Checklist; parent versions of the Behavior Assessment Scale for Children (BASC) and so forth). Following a TBI, the parents are likely to be in the best position to evaluate certain aspects of the child's functioning, such as whether full school days are too exhausting, if there have been significant behaviour changes that are impacting the child's ability to socialize and learn or whether the child is spending inordinate amounts of time completing homework that previously would have been considered 'easy'.

The purpose of this study was to examine the needs of and services provided to children who have sustained a moderate-to-severe TBI. The overarching goal of this study was to inform school teams of best practices in meeting the needs of head-injured students who are transitioning from hospital or rehabilitation back to school.

Method

Participants

A list of potential participants was constructed from the Trauma Registry at Cincinnati Children's Hospital Medical Center. Identified students had a TBI within the previous 2 years (Glasgow Coma Scale Score <12 or 13–15 with evidence of brain injury on neuroimaging; see Table I). Children with penetrating injuries were not included because the pathophysiology and effects of penetrating vs. non-penetrating injuries are quite different [27]. Out of 155 possible recruits, 135 were contacted; attempts to reach the other possible recruits were unsuccessful. Sixty-six were successfully recruited for the study, all of whom agreed to participate in an intervention that was part of a treatment study.

Participants in this study were parents of children who met the study criteria. Some of the caregivers declined to participate in the study because they did not feel their child was experiencing problems as a result of the brain injury. Thus, the sample of participants likely represents a group of students with more significant problems than if all potential participants had been successfully recruited for the study. Children with a pre-morbid history of learning or behavioural difficulties were not excluded from the study; even if children were already receiving services, it was still deemed important to re-evaluate the correspondence between their services and needs.

Of the children whose parents participated in the interviews, 70% were classified as having moderate head injuries; 30% had severe injuries, as determined by the Glasgow Coma Scale (GCS 3–8 was classified as severe; GCS 9–12 or >12 with evidencing of trauma-related abnormalities on imaging were classified as moderate). More of the students were in middle and high school (62%) than in elementary school (38%). There were nearly three times as many injured males than females (see Table I). Consistent with demographic composition of the region, 86% of the injured students were White, 11% were Black and 3% were multiracial.

Instrument and procedure

The *Back to School* Interview [28] was part of a broader 1.5–2 hour assessment of child adjustment and parental and family functioning. Interviewers spent time initially with parents or caregivers, establishing rapport and obtaining information about the injury. Parents of the children with head injuries were interviewed by trained research assistants. To ensure quality control, each interviewer administered five practice interviews and had ongoing supervision throughout the study. The *Back to School* Interview asked each caregiver to identify areas of concern in school. Participants were then asked what types of special support were needed. For each area in which a need for supports was identified, the participating caregiver was asked whether the supports were provided to the child during the 3 months immediately following school re-entry. If the participant indicated that a support was needed, he or she was also asked to rate how difficult it was to obtain support or services from the school as well as the level of satisfaction with the support or service provided.

Because children are not likely to be strong informants regarding their educational accommodations, primary caregivers were interviewed for this study. Thus, although the children with head injuries assented to be part of the study and completed some measures, they did not respond to the school interview that was used for this part of the study.

Results

Of the caregivers interviewed, 38% indicated their children needed special education services following the TBI, 38% indicated a need for an instructional assistant and 36% reported a need for an adjusted schedule (see Table II). The need for class changes (e.g. to move down one level from an honours course) was reported by 25% of participants. Only 20% reported a need for counselling/behavioural support and 17% reported a need for non-

academic support (e.g. speech, occupational therapy, physical therapy). Thus, most caregivers did not perceive a need for special services.

According to caregiver reports, nearly all students who needed the accommodation of having an adjusted schedule received that accommodation (see Table II). Parents least frequently reported receiving needed services in the domain of counselling/behavioural support. Sixteen of the participants reported a need for class changes; 11 reported that this change was provided. Of the 24 participants who indicated a need for an instructional assistant (IA), 16 were provided the IA. The percentage of students receiving non-academic support services at school, such as speech therapy, occupational therapy and physical therapy, was relatively high. Sixty-six per cent of participants indicating that their child needed special education services did in fact receive them (see Table II).

It was expected that severely injured students would be more likely to have perceived needs and would be more likely to receive school-based services. However, when comparing students' levels of injury severity, there were few differences between the needs of the groups, based on parent perceptions (see Table III). Even in students with severe head injuries (GCS <9), only 57% of caregivers indicated a perceived need for an adjusted schedule, with smaller percentages desiring other accommodations or services.

There was, however, a significant difference between severe and non-severe groups in terms of the type of non-academic support (speech, OT/PT) that was perceived as needed. More caregivers of students with severe TBI said their child needed non-academic supports than did parents of students with moderate TBI, $\chi^2(1, n = 44) = 8.404, p = 0.004$. Only 7% of caregivers of students who sustained moderate TBI perceived this need compared to 43% of caregivers of children with severe TBI. In other areas, there was no significant relationship between the perceived need for services and the child's degree of injury severity. That is, parents of more severely injured students did not perceive their children as needing more services (see Table III).

The researchers also examined whether there were differences in perceived needs or unmet needs reported by parents of elementary school students vs secondary (middle and high school) students; however, no significant differences were found (see Table III). Although more than twice as many parents of secondary students (33%) than elementary students (14%) perceived a need for counselling/behavioural support, this difference was not statistically significant. Elementary school and secondary students did not differ in the likelihood of having services provided. Although *no* elementary students in the sample were provided these services, compared to 33% of secondary students, these differences were not statistically significant (see Table III).

Finally, it was hypothesized that students injured during the summer might be less likely to receive services upon their return, but that was not the case (see Table III). There was no significant difference in perceived needs or unmet needs reported by caregivers whose children were injured during the summer vs those whose children were injured during the school year, except in the area of class changes. This, of course, would be expected, given that all students coming back to school after the summer experience a new class placement.

Using a 4-point rating scale, caregivers rated how difficult it was to obtain needed services (very easy, easy, difficult or very difficult). The number of respondents to these questions was very low, as many participants in the study were not receiving school-based services; therefore, reliable conclusions could not be drawn from the data. However, of note, 93% of parents indicated that it was *easy* or *very easy* to obtain needed academic supports, while only 40% of parents indicated that it was *easy* or *very easy* to obtain needed social/behavioural support.

Discussion

Although not every student who experiences a head injury requires special school-based services, the children whose caregivers were surveyed in this study experienced injuries that typically result in at least short-term deficits in cognition, attention and memory that might benefit from school-based adaptations [4]. However, families of children who have sustained TBI in the current study reported fewer than expected academic and behavioural concerns during the initial 3-months following the child's return to school. The reasons for this discrepancy are unclear. This may be related to the fact that parents do not yet know about or understand the academic and behavioural issues that their child is experiencing. Parents initially may have difficulty recognizing emerging limitations and needs and may misattribute them to other factors (i.e. being spoiled as a result of the injury and hospitalization). Given what families have been through, most parents are anxious to see their child return to school and get 'back to normal' and thus are focused on signs of recovery rather than impairment. Nonetheless, the child's return to school is a critical time to ease the transition and put services into place. Delaying services until the child experiences academic failure or social rejection may lead to an exacerbation of difficulties as well as a mislabelling of the condition (severe behaviour handicap rather than TBI).

Parents cannot advocate for their child if they do not perceive a need. With the passage of time, parents often come to understand the long-term implications of their child's injuries; in the first 3 months after returning to school, they may still be coming to terms with the injury. However, if parents are told to 'just wait and see' how their child readjusts to school on his or her own, their child might miss out on a critical window for neural as well as cognitive and behavioural recovery. This is particularly true while children are experiencing post-concussive symptoms such as fatigue and concentration difficulties that can significantly affect classroom performance. Clearly there is a need for information and education, as well as formal and informal advocacy, for families of children who have sustained a TBI [29, 30].

This study found that the greatest unmet need was in the area of emotional and behavioural supports. These results are consistent with previous research indicating that professional educators would benefit from enhancing their knowledge about the psychosocial/behavioural aspects of TBI as well as the cognitive and academic domains [23, 31]. However, there were few differences in caregivers' perceptions of unmet needs as a function of injury severity, timing of the injury (i.e. during the school year vs during the summer) or the child's grade level in school (i.e. elementary vs. secondary)..

The current sample included children with pre-morbid behavioural and learning difficulties. Such pre-injury problems can pose difficulties in understanding the educational needs of a child who has experienced a head injury in that one's deficits post-injury may actually reflect pre-injury problems rather than being a direct result of a head injury. Pre-morbid conditions, such as impulsivity and hyperactivity, may lead to higher head injury rates [32]. Thus, it is important to take into consideration the child's status prior to the injury, as children with pre-injury behavioural difficulties are at a greater risk for post-injury difficulties [33]. Further, TBI may interact with ongoing difficulties, such as pre-existing psychiatric disorders; learning disabilities; health conditions; low intellectual capacity; a previous head injury; or significantly disrupted families [33]. Because TBI can exacerbate pre-morbid difficulties resulting in changes in the need for academic services, it is important for both parents and teachers to be educated about these issues.

Limitations

The sample of participants likely represented a group of students with more significant problems than if all potential participants had been successfully recruited for the study. Thus, the sample may have been more likely to report problems. Conversely, because the study focuses on the first 3 months after the child returned to school, both parents and school professionals may have been reluctant to identify problems given everything the child had been through. Thus, it may reflect a 'honeymoon' or grace period in that parents are not yet aware or ready to acknowledge the long-term issues that may be involved with the child's injury. Thus, a follow-up study to investigate whether perception of getting needs met changes over the course of the first year of school re-entry might be of interest. Further, the students being studied were potentially up to 2 years post-injury. This is not an ideal time frame because parents were being asked to recall services that were provided in the first 3 months after school re-entry.

Some of the interviews had missing data, which limited analyses that could be conducted. Further, the study relied on parent reports and it was assumed that they were knowledgeable about their child's medical history and educational services. However, information from the schools regarding the provision of services would have been useful to corroborate parental reports. Finally, some of the questions were subjective and parents may have different ideas about what constitutes a 'need' (e.g. a need for special education, a need for counselling and so forth). A future study including the teacher's as well as the child's perceptions of services that are needed might be a valuable complement to this study.

An additional limitation is the fact that some of the children whose parents were interviewed may have had learning or behaviour problems prior to their injury, thus making it more difficult to identify and address newly emerging problems. The sample was also heterogeneous in respect to age. School-age children from elementary school through high school were included in the study. This age range comprises a wide developmental spectrum. While the study found some differences between the needs of elementary and secondary student needs, a more detailed examination of needs of same-age students would be beneficial.

Implications

There are several steps schools and hospitals can take to ensure appropriate support following head injury. Assessments should begin prior to a student's return to the classroom and might involve early identification of skills and deficits, the teacher's management style and tolerance for inappropriate behaviour, the level of classroom structure, and the degree of classroom control. Means of assessment might include behavioural observations and interviews with the teacher and school personnel, problem-oriented evaluation scales (e.g. Glasgow Assessment Schedule), samples of student work, parent interviews, analogue methods and functional analysis. School-based academic and psychological assessments should go beyond traditional standard practices (e.g. traditional norm-referenced tests of intelligence) and be aimed at insuring a continuum of regular education services, determining the need for special education, maintaining suitable levels of performance throughout the academic programme and maintaining positive social relationships [14].

The degree of intervention needed depends upon the stage of recovery, specific aspects of the disability and type of environmental support provided. Despite widespread beliefs that behaviour cannot be changed after a brain injury because of physical damage to the brain, appropriate intervention can help children who have sustained a TBI learn to respond to environmental stimuli in a more controlled, socially appropriate manner [21]. Rather than merely reducing maladaptive behaviours, interventions can also increase prosocial behaviour and help students interact more effectively with peers and adults.

In order to facilitate a smooth reintegration to the school setting, other agencies must be aware of school practises and procedures. Conflicts between hospitals or rehabilitation settings and schools might result when treatment centre personnel take a militant advocacy role for children who have sustained a TBI without adequately understanding school policies (e.g. special education eligibility requirements). Suggested solutions for this potential conflict included communication and mutual respect, as well as the assignment of a case manager to coordinate data and communicate about the child to respective systems [33]. Entering into the special education system can be particularly overwhelming for parents of older students who were not previously identified with special education needs [30].

Although children with brain injuries have some similarities to students in traditional special education classifications, they often demonstrate greater discrepancies between abilities and more uneven progress patterns. Therefore, simply placing them in classrooms designed for students with other difficulties fails to recognize the need for adjusting strategies to accommodate their unique deficits [34]. The behaviour patterns for a child with TBI can be dynamic as recovery of function progresses. Thus, assessment must be ongoing and intervention must be flexible. Intervention programmes should help a child with TBI 'recognize, understand, and learn to compensate for his or her deficits' ([34], p. 142). School accommodations should also be flexible and should change as the natural recovery process takes place.

Acknowledgments

This research was supported by grant HD40942-02 from the National Council on Medical Rehabilitation Research, National Institutes of Health and grant R49 CCR523224 from the National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.

References

1. Individuals with Disabilities Education Act of 1990. Pub L. No. 101-476, Section 602(a)(19). Washington, DC: US Government Printing Office
2. National Information Center for Children and Youth with Disabilities. Fact sheet number 18 (FS18). Washington, DC: NICCYD; 2004. Traumatic brain injury.
3. Clark, E. Brain injury. In: Bear, GG.; Minke, KM.; Thomas, A., editors. Children's needs II. Washington, DC: National Association of School Psychologists; 1997. p. 815-826.
4. Yeates, KO. Closed head injury. In: Yeates, KO.; Ris, MD.; Taylor, HG., editors. Pediatric neuropsychology: Research, theory, and practice. New York: The Guilford Press; 2000. p. 92-116.
5. Anderson VA, Catroppa C, Dudgeon P, Morse SA, Haritou F, Rosenfeld JV. Understanding predictors of functional recovery and outcome 30 months following early childhood head injury. *Neuropsychology*. 2006; 20:42-57. [PubMed: 16460221]
6. Ewing-Cobbs L, Miner ME, Fletcher JM, Levin HS. Intellectual, motor, and language sequelae following closed head injury in infants and preschoolers. *Journal of Pediatric Psychology*. 1989; 14:531-547. [PubMed: 2607392]
7. Stancin T, Drotar D, Taylor HG, Yeates KO, Wade SL, Minich NM. Health-related quality of life of children and adolescents after traumatic brain injury. *Pediatrics*. 2002; 109 Available online at: <http://www.pediatrics.org/cgi/content/full/109/2/e34>.
8. Schwartz L, Taylor HG, Drotar D, Yeates KO, Wade SL, Stancin T. Long-term postinjury-onset behavior problems following pediatric traumatic brain injury: Incidence, predictors, and correlates. *Journal of Pediatric Psychology*. 2003; 28:251-263. [PubMed: 12730282]
9. Yeates, KO. Pediatric closed-head injury. In: Yeates, KO.; Ris, MD.; Taylor, HG., editors. Pediatric neuropsychology: Research, theory, and practice. New York: Guilford; 2000. p. 92-116.
10. Yeates KO, Taylor HG, Wade SL, Drotar D, Stancin T, Minich N. A prospective study of short- and long-term neuropsychological outcomes after traumatic brain injury in children. *Neuropsychology*. 2002; 16:514-523. [PubMed: 12382990]
11. Fay GC, Jaffe KM, Polissar ML, Liao S, Rivara JB, Martin KM. Outcome of pediatric traumatic brain injury at three years: A cohort study. *Archives of Physical Medicine and Rehabilitation*. 1994; 75:733-741. [PubMed: 8024416]
12. Levin, HS.; Ewing-Cobbs, L.; Eisenberg, HM. Neurobehavioral outcome of pediatric head injury. In: Broman, S.; Michel, ME., editors. Traumatic head injury in children. Oxford: Oxford University Press; 1995. p. 70-94.
13. Taylor HG, Yeates KO, Wade SL, Drotar D, Klein SK, Stancin T. Influences on first-year recovery from traumatic brain injury in children. *Neuropsychology*. 1999; 13:76-89. [PubMed: 10067779]
14. Clark E, Russman S, Orme S. Traumatic brain injury: Effects on school functioning and intervention strategies. *School Psychology Review*. 1992; 28:242-251.
15. Fletcher-Janzen, E.; Kade, HD. Pediatric brain injury rehabilitation in a neurodevelopmental milieu. In: Reynolds, CR.; Fletcher-Janzen, E., editors. Handbook of child clinical neuropsychology. 2. New York: Plenum Press; 1997. p. 452-481.
16. Ewing-Cobbs, L.; Fletcher, JM. Neuropsychological assessment of traumatic brain injury in children. In: Bigler, ED., editor. Traumatic brain injury: Mechanisms of damage, assessment, intervention, and outcome. Austin, TX: Pro-Ed; 1990. p. 107-128.
17. Tonks J, Williams WH, Frampton I, Yates P, Slater A. The neurological bases of emotional dysregulation arising from brain injury in childhood: A 'when and where' heuristic. *Brain Impairment*. 2007; 8:143-153.
18. Anderson V, Moore C. Age at injury as a predictor of outcome following pediatric head injury: A longitudinal perspective. *Child Neuropsychology*. 1995; 1:187-202.

19. Taylor HG, Alden J. Age-related differences in outcomes following childhood brain insults: An introduction and overview. *Journal of the International Neuropsychological Society*. 1997; 3:1–13. [PubMed: 9072958]
20. Teeter, PA.; Semrud-Clikeman, M. *Child neuropsychology: Assessment and interventions for neurodevelopmental disorders*. Boston: Allyn and Bacon; 1997.
21. Dykeman BF. School-based interventions for treating social adjustment difficulties in children with traumatic brain injury. *Journal of Instructional Psychology*. 2003; 30:225–231.
22. [accessed 5 April 2006] Individuals with Disabilities Education Act Data [Internet]. Available online at: <http://www.ideadata.org/>
23. Glang A, Tyler J, Pearson S, Todis B, Morvant M. Improving educational services for students with TBI through statewide consulting teams. *NeuroRehabilitation*. 2004; 19:219–231. [PubMed: 15502255]
24. Kim YM, Sugawara AI, Kim G. Parents' perception of and satisfaction with the eligibility assessment of their children with special needs. *Early Child Development and Care*. 2000; 160:133–142.
25. Peck CA, Staub D, Gallucci C, Schwartz I. Parent perception of the impacts of inclusion on their nondisabled child. *Research and Practice for Persons with Severe Disabilities*. 2004; 29:135–143.
26. Ravet J. Making sense of disengagement in the primary classroom: A study of pupil, teacher and parent perceptions. *Research Papers in Education*. 2007; 22:333–362.
27. Jellinger, K. The neuropathology of pediatric head injury. In: Shapiro, K., editor. *Pediatric head trauma*. Mounth Kisco, NY: Futura; 1983. p. 87–115.
28. Glang A, Dise-Lewis J, Tyler J, Denslow P. Identification and appropriate service delivery for children who have TBI; abstracts from the 2nd Federal Interagency Conference on traumatic brain injury. *Journal of Head Trauma Rehabilitation*. 2006; 21:408–436. [PubMed: 17243310]
29. Leith KH, Phillips L, Sample PL. Exploring the service needs and experiences of persons with TBI and their families: The South Carolina experience. *Brain Injury*. 2004; 18:1191–1208. [PubMed: 15666564]
30. Savage RC, Depompei R, Tyler J, Lash M. Paediatric traumatic brain injury: A review of pertinent issues. *Pediatric Rehabilitation*. 2005; 8:92–103. [PubMed: 16089249]
31. Mohr JD, Bullock LM. Traumatic brain injury: Perspectives from educational professionals. *Preventing School Failure*. 2005; 49:53–57.
32. Rutter M. Psychological sequelae of brain damage in children. *American Journal of Clinical Neuropsychology*. 1981; 138:1533–1544.
33. Telzrow C. Management of academic and educational problems in head injury. *Journal of Learning Disabilities*. 1987; 20:536–545. [PubMed: 3320243]
34. Boll, TJ.; Stanford, LD. Pediatric brain injury: Mechanisms and amelioration. In: Reynolds, CR.; Fletcher-Janzen, E., editors. *Handbook of child clinical neuropsychology*. 2. New York: Plenum Press; 1997. p. 140–156.

Table I

Demographic characteristics.

	<i>M</i> (SD)	<i>n</i>	Percentage
Injury severity (GCS score), <i>n</i> = 65	10.96 (4.7)	Moderate ★ (9–15) = 44	70%
		Severe (3–8) = 19	30%
Age, <i>n</i> = 66	11.74 (3.68)	Elementary school age (K–5th) = 25	38%
		Secondary school age (6th–12th) = 41	62%
Gender, <i>n</i> = 64		Male = 46	72%
		Female = 18	28%
Race, <i>n</i> = 64		White = 55	86%
		Black = 7	11%
		Multiracial = 2	3%

★ 'Moderate' includes children who had a GCS of 13–15 with positive imaging findings.

Table IIServices needed and provided, $n = 64$.

	% Needed	% Provided	If needed, % provided
Adjusted schedule	36%	34%	94%
Non-academic support services (Speech, OT/PT)	17%	14%	82%
Class changes	25%	17%	68%
Special education instruction	38%	25%	66%
Instructional assistant	38%	25%	66%
Counselling/behavioural support	20%	5%	25%

Table III

Services needed and (if needed) provided: moderate★ vs. severe injuries; elementary vs. secondary school age; summer injury vs. school-year injury.

	Moderate % needed	Moderate if needed, % provided	Severe % needed	Severe if needed, % provided	Elementary % needed	Elementary if needed, % provided	Secondary % needed	Secondary if needed, % provided	Summer injury if needed, % provided	School-year injury if needed, % provided
<i>n</i>	44	44	44	44	48	48	48	48	49	49
Adjusted schedule	38%	92%	57%	88%	39%	100%	44%	83%	67%	73%
Non-academic support services (Speech, OT/PT)	7%	50%	43%	83%	24%	80%	15%	75%	22%	41%
Class changes	33%	67%	21%	100%	24%	75%	33%	78%	0%	52%
Special education instruction	7%	50%	27%	100%	19%	75%	19%	80%	25%	38%
Instructional assistant	43%	62%	43%	67%	38%	63%	52%	64%	40%	55%
Counselling/behavioural support	27%	38%	29%	0%	14%	0%	33%	33%	0%	17%

★, 'Moderate' includes children who had a GCS of 13–15 with positive imaging findings.