Published in final edited form as:

Urol Oncol. 2015 July ; 33(7): 331.e17–331.e23. doi:10.1016/j.urolonc.2015.04.002.

Quality of Life and Symptom Assessment in Randomized Clinical Trials of Bladder Cancer: A Systematic Review

Michael A. Feuerstein^a, Marc Jacobs^b, Alfonso Piciocchi^c, Bernard Bochner^a, Andrea Pusic^a, Peter Fayers^d, Jane Blazeby^e, and Fabio Efficace^c On behalf of Patient Reported Outcome Measurements Over Time In Oncology-PROMOTION Registry

^aDepartment of Surgery, Memorial Sloan Kettering Cancer Center, New York, USA ^bDepartment of Medical Psychology, Academic Medical Center/University of Amsterdam, Amsterdam, The Netherlands ^cData Center and Health Outcomes Research Unit, Italian Group for Adult Hematologic Disease (GIMEMA), Rome, Italy ^dInstitute of Applied Health Sciences, University of Aberdeen, Aberdeen, UK and Department of Cancer Research and Molecular Medicine, Faculty of Medicine, Norwegian University of Technology and Science, Trondheim, Norway ^eCentre for Surgical Research, School of Social & Community Medicine, University of Bristol and University Hospitals Bristol NHS Foundation Trust, Bristol, UK

Abstract

Objectives—Patient-reported outcomes (PRO) help patients, caretakers, clinicians and policy makers make informed decisions regarding treatment effectiveness. Our objective was to assess the quality of PRO reporting and methodological strengths and weaknesses in randomized controlled trials (RCT) in bladder cancer.

Methods—A systematic literature search of bladder cancer RCT published between January 2004 and March 2014 was performed. Relevant studies were evaluated using a predetermined extraction form that included trial demographics, clinical and PRO characteristics, and standards of PRO reporting based on recommendations of the International Society for Quality of Life Research.

Results—Nine RCTs enrolling 1,237 patients were evaluated. All studies were in patients with non-metastatic disease. In 5 RCTs, a PRO was the primary endpoint. The majority of RCTs did not report the mode of administration of the PRO instrument or the methods of collecting data. No RCT addressed the statistical approaches for missing data.

© 2015 Published by Elsevier Inc.

Corresponding Author: Michael A. Feuerstein, Department of Urology, Lenox Hill Hospital, 186 East 76th Street, Floor 1, New York, NY, USA 10021, Phone +1 212 434 4650, Fax +1 212 434 3250, mfeuerste1@nshs.edu.

Financial Disclosures and Conflicts of Interest

None

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conclusions—We found that few RCTs in bladder cancer report PRO as an outcome. Efforts to expand PRO reporting to more RCTs and improve the quality of PRO reporting according to recognized standards are necessary for facilitating clinical decision-making.

Keywords

bladder cancer; patient-reported outcomes; clinical trials; quality of life; clinical decision-making

1. Introduction

Bladder cancer is the 7th most common cancer in men worldwide, with an estimated total of 429,793 new cases and 165,068 deaths in 2012[1]. Bladder cancer is more common in the western world, and is the 6th most common cancer in the United States, accounting for an estimated 74,690 new cases and 15,580 deaths in 2014[2].

Approximately 30% of newly diagnosed patients will have muscle-invasive bladder cancer (MIBC), for which neoadjuvant chemotherapy and radical cystectomy with urinary diversion are considered the standard-of-care[3]. A subset of patients with non-muscle invasive bladder cancer (NMIBC) will progress to invasive disease while many others will have a protracted disease course that may include invasive monitoring and intravesical treatments[4].

For any stage of bladder cancer, informed decision-making needs to consider objective outcome measures with a high level of evidence as well as the patient's values and experience[5]. It is increasingly recognized that patient-reported outcomes (PRO) help patients, caretakers, clinicians and policy makers make decisions regarding treatment effectiveness[6; 7]. However, previous systematic reviews have noted several weaknesses in PRO studies in bladder cancer, including retrospective study design and use of non-validated questionnaires[8; 9]. Further, reviews in other cancers have shown poor PRO reporting in randomized controlled trials (RCT)[10; 11]. Therefore, standards for reporting PRO in RCTs have recently been established[12; 13]. The objective of this review was to identify the number of RCTs in bladder cancer that have included PRO as an endpoint, and to assess the quality of PRO reporting from these studies.

2. Materials and Methods

2.1. Search strategy and identification of studies

An electronic, systematic literature search using Pubmed/Medline, the Cochrane Library, PsycINFO and PsychARTICLES was used to identify RCTs in bladder cancer with a PRO component from January 2004 to March 2014. Details of our search strategy used in other cancers have been previously described[10; 11; 14]. We limited our search to the last 10 years because a previous MEDLINE search of the literature from 1966 to January 2004 found no RCT evaluating PRO after radical cystectomy[9]. Relevant studies listed as references were also considered.

2.2. Selection criteria

English-language RCTs involving adult patients with bladder cancer were included regardless of disease stage. Studies had to enroll at least 50 patients to be included. Studies of patients undergoing screening or involving patients with benign disease were excluded. Conference abstracts were not included. Interventions included any RCT comparing conventional treatments. Studies considering psychological intervention or complementary or alternative medicine were excluded. Any studies evaluating a PRO either as a primary or secondary outcome were included. This included both multidimensional HRQOL outcomes and any other type of PRO measuring the impact of an intervention. Studies evaluating only treatment adherence or satisfaction were not included.

2.3. Data extraction and type of information analyzed

Data were gathered through the Patient Reported Outcome Measurements Over Time IN Oncology (PROMOTION) Registry (<http://promotion.gimema.it>) [14]. For the purpose of this review, two broad types of information were extracted: 1) basic trial demographics and clinical and PRO characteristics; and 2) elements of PRO reporting based on recommendations from the International Society for Quality of Life Research (ISOQOL) [12].

3. Results

The systematic literature search yielded 1,682 records (Figure 1). After screening records, 58 full-text articles were assessed for eligibility of which 48 articles were excluded for being non-randomized (n=27), not including PRO (n=14), mixed sample (n=1), screening study (n=4), and non-English language (n=2). The result was 10 articles on 9 RCTs that met our study criteria enrolling 1,237 patients [15–24]. A summary of the main clinical results and PRO findings are presented in Table 1.

All 9 RCTs were performed in patients with non-metastatic disease: 5 in patients with NMIBC and 4 in patients with MIBC. In 5 RCTs, a PRO was the primary endpoint. The level of PRO reporting based on ISOQOL recommendations is presented in the Supplement Table.

Although 8 RCTs identified PRO as an endpoint in the abstract, only 2 stated a PRO hypothesis in the introduction. Seven RCTs did not report the mode of PRO administration or the methods of collecting data. A major limitation was the handling of missing data. None of the studies addressed the statistical approaches for missing data and only 1 stated the extent of, or the reasons for, missing data.

Four studies used versions of the EORTC instruments, and 3 studies used a bladder-specific instrument. However, 4 studies did not provide the rationale for choice of PRO instrument or evidence of PRO instrument validity.

Of the 5 RCT that used PRO as the primary endpoint, 3 described the limitations, generalizability and clinical significance of the PRO. All 5 interpreted the PRO in the discussion and discussed PRO in the context of other trial endpoints.

Two trials in patients undergoing radical cystectomy reported PRO differences. In a comparison of general plus epidural anesthesia versus general anesthesia alone, general plus epidural anesthesia was associated with improved pain control at all measured time points within 24 hours after surgery[21]. In a comparison of an early recovery after surgery protocol versus a conservative recovery regimen, patients treated with the early recovery protocol reported more favorable functioning and symptom scores on several EORTC measures on post-operative days 3, 7 and at discharge[18].

One study examined radiotherapy with and without chemotherapy for bladder-sparing treatment of MIBC. While this study met our inclusion criteria as a RCT of an intervention measuring PRO, this study did not report the instrument used, method or schedule of data collection, or PRO results[17].

Three of the 5 trials in NMIBC patients reported PRO differences. During transurethral resection of bladder tumors, fewer patients receiving spinal anesthesia with bupivacaine reported moderate or severe pain compared to patients receiving sufentanil[20]. However, bupivacaine was associated with intense motor blockade and longer time to recovery room discharge. A trial of oral anticholinergic therapy versus placebo during induction bacillus Calmette-Guerin (BCG) therapy reported more urinary and non-urinary symptoms with anticholinergic therapy with no clinical benefit[23]. In a comparison of gemcitabine versus one-third dose BCG for maintenance therapy, gemcitabine was associated with better functioning and symptoms on univariate but not multivariate analysis[16].

4. Discussion

Although bladder cancer is one of the most common cancers worldwide, there is paucity of evidence-based PRO from clinical trials. In the last decade, fewer than 1,300 bladder cancer patients were enrolled in a RCT evaluating PRO. Notably, 2 of these 9 studies focused primarily on anesthesia and peri-operative pain control rather than bladder cancer outcomes.

In contrast, a large number of RCTs with a PRO component have been conducted in prostate and gynecological cancers during a similar time period[10; 11]. Whereas we found no RCT in bladder cancer with robust PRO, in prostate and gynecological cancers the rate of high-quality PRO reporting likely to impact clinical decision-making was found to be 20% and 32%, respectively.

Historically, there have been few validated instruments in NMIBC or MIBC[8]. The EORTC QLQ-NMIBC24 was recently validated for use in clinical trials for intermediate and high-risk NMIBC[25]. This instrument was used by 1 of the 5 RCT in NMIBC we evaluated, while the other RCT used either ad hoc or generic cancer questionnaires. In MIBC, new instruments for evaluating the effects of radical cystectomy have recently been validated, including the Bladder Cancer Index and the Functional Assessment of Cancer Therapy-Vanderbilt Cystectomy Index (FACT-VCI)[26; 27]. Only 1 MIBC study we evaluated used a bladder-specific questionnaire. Four of the 9 studies evaluated did not provide the rationale for PRO instrument used and 4 did not provide the validity or reliability of the instrument used. The purpose of disease-specific, validated questionnaires

is to capture PRO data meaningful to stakeholders for clinical decision-making. We believe that the dissemination of newer instruments will improve the quality of PRO reporting.

Much of the interest in PRO in bladder cancer has been comparing outcomes by urinary diversion type: ileal conduit, continent cutaneous or orthotopic neobladder. Understandably, there are no RCT comparing these diversions, and therefore, few RCT in MIBC. Prior to January 2004, there were no RCT evaluating PRO after radical cystectomy[9]. Of the 3 RCT we found in patients undergoing radical cystectomy, 1 examined techniques in orthotopic neobladder creation and 2 examined post-operative recovery and pain control.

Some of the weaknesses of PRO reporting in bladder cancer we found have previously been described[8; 9]. In the current review, only 3 studies reported baseline PRO data. Four studies did not provide the rationale for choice of the PRO instrument or the psychometric properties.

As in prostate and gynecological cancers, one of the main weaknesses of PRO reporting was in the handling of missing data.[10; 11] Missing data leads to reduced power and can be a significant source of bias. Missing data in PRO studies are often not random and can be associated with the outcome of interest. For example, patients completing a PRO survey after surgery may have considerable demographic and clinical differences compared to patients who chose not to complete the survey. The extent of missing data and the statistical handling of this data are, therefore, critical to understanding the generalizability of PRO findings.

This study highlights the current weaknesses of PRO reporting in bladder cancer RCTs. In contrast to previous systematic reviews, we used a formal, objective approach to evaluating PRO reporting in the bladder cancer literature. As has been demonstrated in similar reviews in prostate and gynecological cancers, high-quality PRO reporting can facilitate clinical decision-making and approval of beneficial interventions.[10; 11] When designing RCTs in bladder cancer, investigators should recognize the importance of including valid and reliable PROs in the trial protocol and the necessity for detailing the methodology of PRO assessment.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the EORTC Quality of Life Group for having supported the development of the PROMOTION Registry. We also acknowledge the essential contribution of Alessandro Perreca and Salvatore Soldati, from the GIMEMA Health Outcomes Research Unit, for data management. Jane Blazeby is supported by the MRC ConDuCT-II Hub. Michael A. Feuerstein was supported by the National Cancer Institute/National Institutes of Health under Ruth L. Kirschstein National Research Service Award Institutional Research Training Grant T32 CA082088. The contents of this manuscript are solely the responsibility of the authors and do not necessarily represent the official views of the NIH.

References

1. Ferley, J.; Soerjomataram, I.; Ervik, M.; Dikshit, R.; Eser, S.; Mathers, C.; Rebelo, M.; Parkin, DM.; Forman, D.; Bray, F. GLOBOCON 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet]. 2013. Retrieved July 22, 2014, 2014, from <http://globoon.iarc.fr>
2. Siegel R, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. *CA Cancer J Clin.* 2014; 64(1):9–29. [PubMed: 24399786]
3. Clark PE, Agarwal N, Biagioli MC, Eisenberger MA, Greenberg RE, Herr HW, Inman BA, Kuban DA, Kuzel TM, Lele SM, Michalski J, Pagliaro LC, Pal SK, Patterson A, Plimack ER, Pohar KS, Porter MP, Richie JP, Sexton WJ, Shipley WU, Small EJ, Spiess PE, Trump DL, Wile G, Wilson TG, Dwyer M, Ho M. National Comprehensive Cancer N. Bladder cancer. *J Natl Compr Canc Netw.* 2013; 11(4):446–475. [PubMed: 23584347]
4. Sylvester RJ, van der Meijden AP, Oosterlinck W, Witjes JA, Bouffieux C, Denis L, Newling DW, Kurth K. Predicting recurrence and progression in individual patients with stage Ta T1 bladder cancer using EORTC risk tables: a combined analysis of 2596 patients from seven EORTC trials. *Eur Urol.* 2006; 49(3):466–465. discussion 475–467. [PubMed: 16442208]
5. Perlis N, Krahn M, Alibhai S, Finelli A, Ritvo P, Bremner KE, Kulkarni G. Conceptualizing global health-related quality of life in bladder cancer. *Qual Life Res.* 2014; 23(8):2153–2167. [PubMed: 24729055]
6. Howie L, Hirsch B, Locklear T, Abernethy AP. Assessing the value of patient-generated data to comparative effectiveness research. *Health Aff (Millwood).* 2014; 33(7):1220–1228. [PubMed: 25006149]
7. Basch E, Abernethy AP, Mullins CD, Reeve BB, Smith ML, Coons SJ, Sloan J, Wenzel K, Chauhan C, Eppard W, Frank ES, Lipscomb J, Raymond SA, Spencer M, Tunis S. Recommendations for incorporating patient-reported outcomes into clinical comparative effectiveness research in adult oncology. *J Clin Oncol.* 2012; 30(34):4249–4255. [PubMed: 23071244]
8. Botteman MF, Pashos CL, Hauser RS, Laskin BL, Redaelli A. Quality of life aspects of bladder cancer: a review of the literature. *Qual Life Res.* 2003; 12(6):675–688. [PubMed: 14516177]
9. Porter MP, Penson DF. Health related quality of life after radical cystectomy and urinary diversion for bladder cancer: a systematic review and critical analysis of the literature. *J Urol.* 2005; 173(4): 1318–1322. [PubMed: 15758789]
10. Efficace F, Feuerstein M, Fayers P, Cafaro V, Eastham J, Pusic A, Blazeby J. Group, E. Q. o. L. Patient-reported Outcomes in Randomised Controlled Trials of Prostate Cancer: Methodological Quality and Impact on Clinical Decision Making. *Eur Urol.* 2013
11. Efficace F, Jacobs M, Pusic A, Greimel E, Piciocchi A, Kieffer JM, Gilbert A, Fayers P, Blazeby J. E. Q. o. L. G. P. R. O. M. O. T. I. O.-P Registry. Patient-reported outcomes in randomised controlled trials of gynaecological cancers: investigating methodological quality and impact on clinical decision-making. *Eur J Cancer.* 2014; 50(11):1925–1941. [PubMed: 24825114]
12. Brundage M, Blazeby J, Revicki D, Bass B, de Vet H, Duffy H, Efficace F, King M, Lam CL, Moher D, Scott J, Sloan J, Snyder C, Yount S, Calvert M. Patient-reported outcomes in randomized clinical trials: development of ISOQOL reporting standards. *Qual Life Res.* 2013; 22(6):1161–1175. [PubMed: 22987144]
13. Calvert M, Blazeby J, Altman DG, Revicki DA, Moher D, Brundage MD, Group CP. Reporting of patient-reported outcomes in randomized trials: the CONSORT PRO extension. *JAMA.* 2013; 309(8):814–822. [PubMed: 23443445]
14. Efficace F, Rees J, Fayers P, Pusic A, Taphoorn M, Greimel E, Reijneveld J, Whale K, Blazeby J. European Organization for R, Treatment of Cancer Quality of Life G. Overcoming barriers to the implementation of patient-reported outcomes in cancer clinical trials: the PROMOTION Registry. *Health Qual Life Outcomes.* 2014; 12:86. [PubMed: 24902767]
15. Ahmadi H, Skinner EC, Simma-Chiang V, Miranda G, Cai J, Penson DF, Daneshmand S. Urinary functional outcome following radical cystoprostatectomy and ileal neobladder reconstruction in male patients. *J Urol.* 2013; 189(5):1782–1788. [PubMed: 23159582]
16. Gontero P, Oderda M, Mehnert A, Gurioli A, Marson F, Lucca I, Rink M, Schmid M, Kluth LA, Pappagallo G, Sogni F, Sanguedolce F, Schiavina R, Martorana G, Shariat SF, Chun F. The impact

of intravesical gemcitabine and 1/3 dose Bacillus Calmette-Guerin instillation therapy on the quality of life in patients with nonmuscle invasive bladder cancer: results of a prospective, randomized, phase II trial. *J Urol*. 2013; 190(3):857–862. [PubMed: 23545101]

17. James ND, Hussain SA, Hall E, Jenkins P, Tremlett J, Rawlings C, Crundwell M, Sizer B, Sreenivasan T, Hendron C, Lewis R, Waters R, Huddart RA, Investigators BC. Radiotherapy with or without chemotherapy in muscle-invasive bladder cancer. *N Engl J Med*. 2012; 366(16):1477–1488. [PubMed: 22512481]
18. Karl A, Buchner A, Becker A, Staehler M, Seitz M, Khoder W, Schneevoigt B, Weninger E, Rittler P, Grimm T, Gratzke C, Stief C. A new concept for early recovery after surgery for patients undergoing radical cystectomy for bladder cancer: results of a prospective randomized study. *J Urol*. 2014; 191(2):335–340. [PubMed: 23968966]
19. Koga H, Ozono S, Tsushima T, Tomita K, Horiguchi Y, Usami M, Hirao Y, Akaza H, Naito S, Group BCGTSS. Maintenance intravesical bacillus Calmette-Guerin instillation for Ta, T1 cancer and carcinoma in situ of the bladder: randomized controlled trial by the BCG Tokyo Strain Study Group. *Int J Urol*. 2010; 17(9):759–766. [PubMed: 20604814]
20. Marandola M, Antonucci A, Tellan G, Fegiz A, Fazio R, Scicchitano S, Delogu G. Subarachnoid sufentanil as sole agent vs standard spinal bupivacaine in transurethral resection of the bladder. *Minerva Anesthesiol*. 2005; 71(3):83–91. [PubMed: 15714184]
21. Ozyuvaci E, Altan A, Karadeniz T, Topsakal M, Besisik A, Yucel M. General anesthesia versus epidural and general anesthesia in radical cystectomy. *Urol Int*. 2005; 74(1):62–67. [PubMed: 15711112]
22. Sabichi AL, Lee JJ, Grossman HB, Liu S, Richmond E, Czerniak BA, De la Cerdá J, Eagle C, Viner JL, Palmer JL, Lerner SP. A randomized controlled trial of celecoxib to prevent recurrence of nonmuscle-invasive bladder cancer. *Cancer Prev Res (Phila)*. 2011; 4(10):1580–1589. [PubMed: 21881030]
23. Johnson MH, Nepple KG, Peck V, Trinkaus K, Klim A, Sandhu GS, Kibel AS. Randomized controlled trial of oxybutynin extended release versus placebo for urinary symptoms during intravesical Bacillus Calmette-Guerin treatment. *J Urol*. 2013; 189(4):1268–1274. [PubMed: 23123375]
24. Skinner EC, Skinner DG. Does reflux in orthotopic diversion matter? A randomized prospective comparison of the Studer and T-pouch ileal neobladders. *World J Urol*. 2009; 27(1):51–55. [PubMed: 19002689]
25. Blazeby JM, Hall E, Aaronson NK, Lloyd L, Waters R, Kelly JD, Fayers P. Validation and Reliability Testing of the EORTC QLQ-NMIBC24 Questionnaire Module to Assess Patient-reported Outcomes in Non-Muscle-invasive Bladder Cancer. *Eur Urol*. 2014
26. Anderson CB, Feuer ID, Large MC, Steinberg GD, Barocas DA, Cookson MS, Penson DF. Psychometric characteristics of a condition-specific, health-related quality-of-life survey: the FACT-Vanderbilt Cystectomy Index. *Urology*. 2012; 80(1):77–83. [PubMed: 22608798]
27. Gilbert SM, Dunn RL, Hollenbeck BK, Montie JE, Lee CT, Wood DP, Wei JT. Development and validation of the Bladder Cancer Index: a comprehensive, disease specific measure of health related quality of life in patients with localized bladder cancer. *J Urol*. 2010; 183(5):1764–1769. [PubMed: 20299056]

Figure 1.
Schematic breakdown of literature search results of bladder randomized controlled trials (PRO= patient-reported outcomes).

Table 1

Randomized controlled trials with PRO design: basic study characteristics.

Study*	Overall study sample size	Baseline PRO sample size	PRO instruments used	Primary endpoint	Treatment outline	Summary of main clinical results	Summary of PRO results/PRO treatment recommendations
Gontero P et al. J Urol. 2013;190: 857–862.	120	118	EORTC QLQ-C30 and EORTC QLQ-BLS24	PRO (including QoL or symptoms relief)	Intravesical gencitabine, 2000mg/50cc saline weekly for 6 weeks, maintenance once monthly for 1 year versus intravesical Connaught strain BCG 27mg/50cc saline (1/3 dose) weekly for 6 weeks, maintenance of 3 weekly treatments	Local and systemic side effects were more frequent in the BCG arm (56% vs 36%, p=0.03). The incidence of hematuria and fever were more common with BCG at T1 (post-induction), and dysuria was more common with BCG at T2	On univariate analysis, at T1, gencitabine arm had better cognitive and emotional functioning and urinary symptom distress. At T2, gencitabine arm had better cognitive functioning and less nausea and vomiting symptom distress.
James ND et al. N Engl J Med. 2012; 366:1477–88.	360	360	Not reported	Locoregional disease-free survival	55Gy or 64Gy radiotherapy to the bladder. Patients randomized to receive no additional therapy versus fluorouracil 500mg/m2 during fractions 1–5 and 16–20 plus mitomycin C 12mg/m2 on day 1	Radiotherapy with chemotherapy improved locoregional disease-free survival versus radiotherapy alone with no difference in adverse events or overall survival	Not reported
Ahmadi H et al. J Urol. 2013;189:1782–8 Skinner EC, Skinner DG. World J Urol. 2009;27, 51–55.	295	179	Modified version of the Bladder Cancer Index	Renal function and anatomy at 3 years following surgery	Urinary diversion using Studer versus T-pouch ileal orthotopic neobladder	Not reported	No differences
Sabichi, AL et al. Cancer Prev Res. 2011;4,1580–1589.	146	146	EORTC QLQ C-30 (abbreviated form)	Time to recurrence	Celecoxib 200mg versus placebo twice daily for at least 12 months for a maximum of 24 months or until recurrence	No differences	No differences
Karl A et al. J Urol. 2014; 191:335–340.	101	101	EORTC QLQ-C30	PRO (including QOL or symptoms relief)	Following radical cystectomy, early recovery after surgery (ERAS) versus conservative post-operative treatment regimen6	ERAS associated with lower rates of wound healing disorders, DVT and fever. ERAS group walked greater distances on Day 3. ERAS group spent less time in the	On Day 3, ERAS patients reporting better physical, emotional functioning and less constipation. On Day 7, ERAS patients reported better role, emotional, cognitive and social functioning and less