Published in final edited form as:

Econ Bot. 2014 March 1; 68(1): 1–15. doi:10.1007/s12231-014-9257-8.

Are ecologically important tree species the most useful? A case study from indigenous people in the Bolivian Amazon

Maximilien Guèze¹, Ana Catarina Luz¹, Jaime Paneque-Gálvez¹, Manuel J. Macía², Martí Orta-Martínez¹, Joan Pino³, and Victoria Reyes-García^{4,*}

¹Institut de Ciència i Tecnologia Ambientals (ICTA), Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

²Departamento de Biología, Área de Botánica, Universidad Autónoma de Madrid, Calle Darwin 2, 28049 Madrid, Spain

³Centre de Recerca Ecològica i Aplicacions Forestals (CREAF), Universitat Autònoma de Barcelona, 08193 Barcelona, Spain

⁴ICREA at the Institut de Ciència i Tecnologia Ambientals (ICTA), Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

Abstract

Researchers have argued that indigenous peoples preferably use the most apparent plant species, particularly for medicinal uses. However, the association between the ecological importance of a species and its usefulness remains unclear. In this paper we quantify such association for six use categories (firewood, construction, materials, food, medicines and other uses). We collected data on the uses of 58 tree species, as reported by 93 informants in 22 villages in the Tsimane' territory (Bolivian Amazon). We calculated the ecological importance of the same species by deriving their importance value index (IVI) in 48 0.1-ha old-growth forest plots. Matching both data sets, we found a positive relation between the IVI of a species and its overall use value (UV) as well as with its UV for construction and materials. We found a negative relation between IVI and UV for species that were reportedly used for medicine and food uses, and no clear pattern for the other categories. We hypothesize that species used for construction or crafting purposes because of their physical properties are more easily substitutable than species used for medicinal or edible purposes because of their chemical properties.

Keywords

ethnobotany; importance value index (IVI); species saliency; species usefulness; use value (UV); tropical forest; Tsimane' indigenous people

*Corresponding author; victoria.reyes@uab.cat.

Introduction

The ethnobotanical knowledge of people living in tropical forests has been increasingly well-documented in the last decades (Bennett 1992, Albuquerque and Lucena 2005). However, little is known about why people use some species more than others and particularly whether spatial distribution patterns of species relate to patterns of use. Ethnobotanists have addressed this question through the “ecological apparency” hypothesis (Phillips and Gentry 1993a, 1993b; Lucena et al. 2007). The hypothesis, first developed to explain plant-herbivores relations, considers that the visibility (or apparency) of the plant, together with its chemical composition, will influence herbivores’ behavior. When applied to human uses of plant species, researchers have hypothesized that most apparent species (i.e., most visible) have more known uses, and therefore a greater use value (UV) (*sensu* Phillips and Gentry 1993a), because they are readily available.

Some studies have tested this hypothesis mostly by relating the number of uses of species to their apparency, proxied by different ecological measures. For example, some authors have related the overall UV of a species with its density, dominance, and frequency (Phillips and Gentry 1993b, Galeano 2000, Lawrence et al. 2005) or with its importance value index (IVI) (Torre-Cuadros and Islebe 2003, Lucena et al. 2007, Thomas et al. 2009a). Particularly, the IVI of a species seems to be a good indicator of apparency because it provides information on basal area, frequency, and abundance, and therefore on the saliency of the species (Thomas et al. 2009a). Another study has related the UV to the accessibility of plant communities, proxied by the travel time to those sites (Thomas et al. 2009b). In all these studies the visibility of a species tends to be positively associated to its number of uses, thus giving support to the ecological apparency hypothesis.

Focusing on medicinal uses and based on the frequency with which a species had been collected by botanists, Berlin (2003) found that, among Highland Mayas in Mexico, common species are more commonly used as medicines than rare species. Stepp and Moerman (2001) have stressed that weeds, i.e., common species, are the most frequently found life form in the pharmacopoeia of the Mayas, probably because people seek highly bioactive compounds, which are commonly found in weeds. Voeks (2004) also reports the importance of secondary vegetation in traditional medicine. Findings from these studies are important in ethnopharmacological research because they challenge the traditional understanding that people obtain medicinal species from “unique and inaccessible habitats” (Berlin 2003, p. 2). However, a major setback in this body of research is that none of these studies has been based on ecological inventories. Results from studies measuring ecological apparency based on forest inventories have yielded equivocal results. For example, a study among rural people in areas dominated by *caatinga* vegetation (Brazil) found a negative association between medicinal importance (an index related to UV) and the density and frequency of woody species (Silva and Albuquerque 2005), whereas another study in the same environment revealed a positive association between the medicinal UV and woody species frequency (Lucena et al. 2007). Similarly, a study among indigenous groups of Bolivia found a positive correlation between the medicinal UV and the IVI of woody species (Thomas et al. 2009a), whereas a study among *mestizos* in the Peruvian Amazon reported no

association between the medicinal UV and the density, dominance, or frequency of woody species (Phillips and Gentry 1993b).

Within this context, the overall objective of this study is to analyze the relation between tree species' usefulness and their ecological importance in the territory of the Tsimane', a lowland Bolivian indigenous group. Specifically, our goals are to 1) test whether there is an association between the usefulness of a tree species, proxied by its UV, and its IVI, and 2) analyze different patterns of association between IVI and UV across different categories of use. If the ecological importance of a species drives its number of uses because people have more opportunities to learn about salient species, as predicted by the ecological apparency hypothesis, then we should find that the higher the IVI of a species, the larger its number of uses known (and therefore the larger its UV).

The Tsimane': lands and knowledge

The Tsimane' are one of the largest ethnic groups in the lowlands of Bolivia (Reyes-García et al. under review). Their territory lies between the foothills of the Andes and part of the Moxos savannas (Fig. 1). Annual mean temperature is 25.8°C (Navarro and Maldonado 2002) and annual mean precipitation is 1743mm (Godoy et al. 2008), with important inter-annual climatic variation. Climate is markedly seasonal with four months with less than 100mm rainfall and episodic southern cold winds. Soils are quaternary alluvial sediments of fluvial origin, mostly acrisols and ferralsols (Navarro and Maldonado 2002).

The territory of the Tsimane' is situated at the interface between three biogeographic regions: Amazonia, Andes, and Brazilian-Paraná. Most of the territory is covered with well-drained upland *terra firme* rainforest (Guèze et al. 2013). In some areas, different types of flooded forests occur according to the history of inundation. The northeastern part of the territory coincides with the edge of the Moxos savannas, where gallery forests and forest islands that seem to have an anthropogenic origin are found (Lombardo et al. 2011). In the southwestern part of the territory the hills form a transition with the Andean sub-montane forests.

Traditionally, the Tsimane' have been semi-nomadic hunters-gatherers and small-scale horticulturalists. They used to live in nuclear families scattered along rivers and streams. Nowadays they live clustered in permanent villages close to communication axes (main rivers and forest roads). The Tsimane' hold a deep knowledge of wild plants; Reyes-García et al. (2006) reported uses of 410 plant species. Traditional knowledge of wild plants is strongly shared by the whole ethnic group (Reyes-García et al. 2003), and has important consequences for Tsimane' life, as higher levels of knowledge have been found to be associated with less deforestation and more diversity in agricultural fields (Reyes-García et al. 2007a, 2008) and a better health (McDade et al. 2007). Nowadays, although the Tsimane' still strongly rely on mostly unfragmented forests for their livelihood, they experience different levels of encroachment upon their territory, integration into the market economy, and cultural changes, which in turn affect their knowledge and use of the forest (Reyes-García et al. 2007b). For example, some Tsimane' still practice subsistence slash-and-burn agriculture while others produce cash crops. It has also been shown that activities that drive

the Tsimane' to leave their villages are associated with lower levels of traditional knowledge whereas activities that keep them in their homelands contribute to the maintenance of this form of knowledge (Reyes-García et al. 2007b).

Materials and methods

Data collection

Since our study focuses on the relation between ecological importance and species usefulness, our data collection strategy includes methods from ecology and methods from socio-cultural anthropology. Our analysis focuses on trees because they are well-known by most Tsimane' and some species are known by a single local name (see below, personal observation). Palms were also included in the analyses because they are among the most useful plant species (Macía et al. 2011), and because –ecologically–palms resemble trees.

Ecological data—To obtain the IVI of tree species we established 48 0.1-ha plots in the territory of six Tsimane' villages (eight plots per village, hereafter called “villages with plots”) (Fig. 1). We selected villages on the basis of homogeneity in 1) forest types and topography, through the analysis of Landsat satellite images (Paneque-Gálvez et al. 2012) and 2) village characteristics, such as the number of households, based on a previous census (Reyes-García et al. 2012). Within each village we established plots in old-growth *terra firme* forest with no apparent sign of recent human activity (at a minimum distance of 500m from any agricultural field or fallow, and without large canopy gaps). In each plot we inventoried all trees with a diameter at breast height (dbh) ≥ 2.5 cm. We counted all stems rooting within the plot limits. Multiple stems were considered as one individual. We measured the dbh of each individual at 1.30m from the ground. We collected voucher specimens for all individuals that could not be identified in the field. Duplicates of the vouchers are deposited in LPB (Herbario Nacional de Bolivia, La Paz, Bolivia) and MA (Real Jardín Botánico, Madrid, Spain) and unicates are kept in LPB.

We selected 58 useful species, among all the tree species inventoried, following these criteria: 1) usefulness of the plant reported by at least three local guides while measuring them in plots and 2) unique correspondence between the vernacular name and the scientific name of the species, based both on the names reported by local guides and on two databases on Tsimane' ethnobotany (Huanca 1999, Reyes-García 2001).

Ethnobotanical data—After plots were inventoried, we collected ethnobotanical data on tree species uses through interviews based on vernacular names. Specifically, we gathered ethnobotanical information of the 58 selected tree species in 22 Tsimane' villages (hereafter called “villages with ethnobotanical prospection”) among which we included the 6 villages with plots. In each village, we asked participants to identify the most knowledgeable people about plant uses to interview them. Our initial target was to interview two men and two women per village, but the gender and the actual number of people interviewed in a village finally varied depending on the number of experts available and willing to participate. We interviewed a total number of 93 people.

To keep the interviews within a reasonable duration, we randomly split the list of 58 species into two lists of the same length. In each village we used one list to interview half of the informants (one man and one woman whenever possible) and the other list to interview the other informants. Interviews were conducted in Tsimane' language and with the help of translators. We asked each informant "Do you know the species X (vernacular name of the species)?" "Can you tell me all its possible uses?" We wrote down verbatim all the uses reported by the informants.

Data analysis

To analyze the relation between usefulness and apparency we used a multivariate model designed to explain the UV of a species as a function of its IVI. To do so, our first step was to match the ethnobotanical data collected in 22 villages and the ecological data collected in 6 villages. For most of their daily subsistence activities (hunting, fishing, and non-timber forest products collection), the Tsimane' use neighboring forest areas as well as a large common territory, often incurring on the territory of neighboring villages (Cruz-Burga et al. 2013). Therefore, we assumed that respondents' information on a species would be drawn from observations at the landscape level, not only from observations of areas around their own village. Consequently, a species' ecological importance in each of the six villages with plots would be associated with the uses reported by people from nearby villages with ethnobotanical prospection. We then defined "areas of influence" around the six villages with plots. These areas of influence comprised the closest villages with ethnobotanical prospection to the villages with plots (3-5 villages per area of influence; Fig. 1), and represented the area where people would normally carry out their subsistence and foraging activities. The distance between the centre of any village (its school) and any plot in its area of influence ranged between 4.9 (± 3.4) and 8.0 (± 3.4) km. We calculated the IVI of a species in the inventoried plots per area of influence as the sum of its relative density, relative dominance, and relative frequency (Curtis and Macintosh 1951).

We calculated 1) the total UV of the species as the average number of use-reports per species (Phillips and Gentry 1993a) and 2) the categorical UV of the species as the average number of use-reports per species within each of six categories. We followed Thomas et al. (2009a) to define five categories (food, firewood, construction, materials, and medicines). However, we grouped "environmental uses", "social uses" and "poison" in the category "other uses" since each of these categories had few use-reports for a small number of species. The category *construction* included house and canoe building and timber, and the category *materials* included tools, handicrafts, and toys. To calculate the categorical UV we counted each single use-report, even if there were several use-reports in the same category. For example, we counted all medicinal use-reports of a species mentioned, because the fact that a species has more medicinal use-reports is an indicator of its usefulness and could be related to its density or frequency. Our units of analysis are tree species per area of influence; therefore, our analysis relates the average UV of one tree species in one area of influence to the ecological importance of the species in the same area. We selected this approach as for each tree species the relation between usefulness and apparency might be specific for the area of influence (Macía 2008). Moreover, the approach also increases the number of observations for multivariate analysis (6 observations for each of the 58 species).

We studied the relation between both the total and the categorical UV and the IVI of the species per area of influence using scatter plots and Pearson's correlations, as well as ordinary least-square regressions. All regression models were run with robust standard errors and clustered by area of influence (Woolridge 2003). We included two controls in the models: one set of dummy variables for areas of influence and another set of dummy variables that capture the life form of the species (large tree, small tree, or palm). The dummies used for buffer areas (not shown in the tables) are binary variables that capture fixed effects of the areas of influence not measured in this study (Suits 1957). Such fixed effects include proximity to towns, ecological characteristics, and the like. We attributed the life form of each species on the basis of the dbh in the plots, verifying reported life forms in Killeen et al (1993). Small trees, i.e., understory species, were defined as all the species that never reached 10cm dbh (Guèze et al. 2013).

Results

We surveyed an average of 3.2 people per village. Overall, 1.6 informants in each village, or 35.2 informants in all the dataset, provided ethnobotanical information for each of the 58 species studied.

Out of the 58 useful species studied, 33 were ecologically important with an IVI>1 (Table 1). The four most ecologically important species, *Iriartea deltoidea*, *Hura crepitans*, *Poulsenia armata*, and *Socratea exorrhiza* showed a high UV. The three species with the highest UV were *Attalea phalerata*, *Swietenia macrophylla* and *Clarisia biflora* but only *C. biflora* showed a high IVI. *A. phalerata* and *S. macrophylla* were among the less ecologically important species.

Most of the 58 species had at least one use-report in the categories firewood and construction (54 and 53 species with more than 1 use-report in those categories, respectively). However, in the other categories the number of species with more than one use-report was much lower. Some patterns arise when examining the association between IVI and UV by categories. The most useful species in the categories construction and materials tended to have a high IVI (e.g., *I. deltoidea*, *S. exorrhiza*). The species with higher UV in the medicine category had low IVI, suggesting that medicinal species in our list were rare species (e.g. *Aniba canelilla*, *Galipea longiflora*). This tendency was also observed, although to a lesser extent, for food uses. For firewood and other uses, no tendency was found.

We found a positive but weak correlation between the total UV of the species and their IVI (Pearson's $r=0.18$, $P=0.002$). However, a close visual analysis of this relation suggests that two species (*H. crepitans* and *I. deltoidea*) with high IVI values (>27) in two areas of influence averaged upward the regression line (Fig. 2).

Three findings from the regression analyses between the total UV of a species and its IVI deserve comment (Table 2). First, a bivariate regression without controls (model [1]) indicates a positive and statistically significant (although weak) association between total UV and IVI. The coefficient in model [1] implies that an increase of 1 unit in the IVI of a

species would be associated with an increase of 0.027 units in its total UV ($P=0.000$). Since the IVI ranges from 0.04 to 17.05 and the UV ranges from 0.73 to 3.52, this corresponds approximately to a 1% increase in UV for each 6% increase in IVI. Second, when we control for life form and for village fixed-effects (model [2]), the association between UV and IVI found in model [1] remained significant, but weakened. Third, the analysis segregating between “rare” species ($IVI \leq 1$, model [3]) and “common” species ($IVI > 1$, model [4]) yielded a significant positive association ($P=0.004$) between UV and IVI only for common species. The number of observations was only 293 because some species were not found in the plots of two areas of influence.

The association between the categorical UV of a species and its IVI showed particularities (Fig. 3). A species UV showed a positive linear relation with its IVI within the categories construction (Pearson's $r=0.36$, $P<0.001$) and materials (Pearson's $r=0.22$, $P<0.001$), indicating that most species used for construction and materials were common species. However, a species UV showed a negative, although weak, linear relation with its IVI within the category medicine (Pearson's $r=-0.19$, $P<0.001$) and an also negative tendency with its IVI within the category food, indicating that rare tree species had more use-reports as foods and medicines. Last, we did not find a linear relation between IVI and UV in the categories firewood or other uses.

In Table 3 we present the results of the bivariate and multivariate regressions using only the average number of uses in a given category. We highlight three main findings from this table. First, after controlling for village fixed-effects and life form of the species, we found a positive but weak association between IVI and UV in the categories construction (model [6]) and materials (model [8]). Second, when including the controls, we found a negative association between IVI and UV in the categories food (model [2]) and medicine (model [10]). Third, the UV for the categories firewood and other uses showed no association with the IVI in bivariate models (models [3] and [11]). Thus, the significant association observed for the category firewood in multivariate analysis (model [4]) is probably an interaction effect between explanatory and control variables.

Discussion

Use value and ecological importance of tree species

The main finding of this work is that, when considering its overall UV, the more ecologically important a tree species is, the more uses it has, although the correlation is weak. Our finding is consistent with other studies that have related species usefulness and IVI (Torre-Cuadros and Islebe 2003, Lucena et al. 2007, Thomas et al. 2009a) and with studies that have related species usefulness and apparency estimated with indicators other than IVI (Paz-y-Miño et al. 1995, Galeano 2000). Thus, our finding provides additional support to the hypothesis that people use more the most apparent species in the forest.

However, our analyses provide insightful nuances to this finding. First, our results suggest that, despite the significance of the association found between IVI and UV, the coefficients of the regressions are small. Second, much of the variation in a species UV is captured by its life form. This is consistent with studies that found that a species' family is more important

than its apparency when explaining its uses (Phillips et al. 1993b, Thomas et al. 2009a). Here, palms (Arecaceae) are the most useful life form for the Tsimane', as reported widely in other Amazonian studies (Macía 2004, Byg et al. 2006, Macía et al. 2011). A potential explanation for their common use is that palms are among the most easily identifiable species. For instance, although *A. phalerata* is rare in our study area, it has a high UV. Since *A. phalerata* is common (i.e., visible or apparent) around Tsimane' villages settled in pampas (personal observation) and because it is useful and easily identifiable, it is possible that people talk more about this species than about others. Third, the relation between IVI and UV tends to be negative when the species are "rare" (IVI < 1), because some of these rare species are useful. Rare species in our study (e.g., *S. macrophylla*, *Cedrela odorata*, medicinal tree species such as *Genipa americana*) are often planted in home gardens. Thus, the pattern observed for these species could be due to the fact that they are well-known among the Tsimane' regardless of their ecological characteristics in old-growth forests. Indeed, we proxied the usefulness of a species with its UV (i.e., knowledge), but other measures such as actual frequencies of use, might yield different results (Albuquerque 2006, Reyes-García et al. 2006, Albuquerque and Hanazaki 2009, Lucena et al. 2012).

Additionally, results from this study must be taken with caution at least for two reasons. First, part of the variation in the results is probably due to sampling methods. Ecological values are likely biased due to samples size and vary depending on the tree diameter classes (Macía 2008). Although our ecological sampling seems accurate (all trees > 2.5cm dbh), we inventoried 0.8 hectares of forest per area of influence, which may not fully capture the actual ecological features sought for the trees inventoried (relative density, relative dominance, and relative frequency). Second, the ecological apparency hypothesis implies causality: the resource is the driver of users' behavior. Researchers have often rejected the alternative hypothesis that management and use of the species influence their ecological importance (but see Thomas et al. 2009a). However, reverse causality cannot be discarded in our study area. On the one side, our study area seems to have been inhabited for a long time (Denevan 1966), so the actual floristic composition could just be the result of anthropogenic perturbation or management on composition and abundances, as previous research suggest that those perturbations might persist for centuries (Macía 2008). On the other side, some of the most useful tree species in the area have suffered intensive exploitation. For example, mahogany (*S. macrophylla*) and Spanish cedar (*C. odorata*) are some of the most useful species and have been selectively logged in the last decades (Gullison et al. 1996); *A. phalerata* has probably been overexploited until local extinction in forest areas. Thus, it is likely that the IVI of these species have decreased as a consequence of their own usefulness. A plausible scheme, as suggested by Lawrence et al. (2005), is a negative feedback between apparency and uses, i.e. the apparency of the species increases its use value, leading to a negative effect on its abundance.

Categorical use values and ecological importance of tree species

We found a positive relation between IVI and UV in the categories construction and materials. This is consistent with the positive association found in other studies between species relative dominance and usefulness for construction (Lucena et al. 2007, Thomas et al. 2009a) and with the results of Phillips and Gentry (1993b), which stress a positive

association between species' relative density and frequency and their UV for construction and technology (including materials).

We also found a negative association between the medicinal and food UV and the IVI of the species, consistent with some earlier studies (Silva and Albuquerque 2005), but not with others (Phillips and Gentry 1993b, Lucena et al. 2007, Thomas et al. 2009a). These inconsistencies might be in part due to the fact that researchers do not always include the same uses in the same categories, particularly when comparing studies that have been conducted among indigenous and among non-indigenous populations (Galeano 2000, Schwantes and Felfili 2001, Lawrence et al. 2005). For example, firewood is sometimes included in a "technology" category (Phillips and Gentry 1993a) and sometimes considered as a category on its own (Galeano 2000, Macía et al. 2011, this study). The association between usefulness and ecological importance may be strongly specific of the area and the people studied.

What would explain the differences in the associations between IVI and categorical UV? We hypothesize that the physical properties of the species, particularly meaningful for the use of the species in construction or materials, are more easily substitutable than their chemical properties, which are more meaningful in medicinal and food uses. On the one hand, physical properties (e.g., mechanical resistance or durability) are likely to be shared by many species, such as palms or common construction trees (e.g., Meliaceae, Euphorbiaceae). The Tsimane' have a large choice of species, which might explain the positive relation between IVI and UV in the categories house construction and materials: if many species have adequate physical properties for those uses, they might just tend to use the most apparent of all the available species. On the other hand, specific chemical properties, such as those responsible for edibility, taste, or chemical compounds used for medicinal purpose, are likely to be much harder to substitute. Since in our study site those properties tend to be more often found in "rare" species, this would explain the negative association between IVI and UV in the categories medicine and food. Stepp (2004) argues that common weedy species invest in mobile compounds such as alkaloids, cardiac glycosides and terpenoids as a defense against herbivory. These compounds are valued as medicines, which might explain the role of weeds (i.e., common plants) as medicinal plants. Although we do not challenge this finding, we argue that the traits associated with the production of such compounds (short-lived leaves and fast-growing capacity) can also be found in old-growth forest tree species as many present in our study area, especially in understory species and deciduous large tree species. For example, deciduous canopy trees such as *Calycophyllum spruceanum* have been reported to have medicinal uses in Brazil (Costa et al. 2011). Moreover, leaves are not always the most useful part of medicinal trees, but rather people use other parts such as barks or roots.

In this study we do not control for cultural and social factors, which also explain preferences regarding the use of plants (Thomas 2012). For example, the palm *Iriartea deltoidea* has no medicinal use among the Tsimane' although it does for a neighboring group, the Yuracaré (Thomas et al. 2009a). Therefore, cultural uses might impose important limitations to the role of inherent physical or chemical properties of the plants, and require further study.

In sum, the overall results of this work suggest that the ecological apparency hypothesis has to be taken with caution when applied to humans. While herbivores use plants only as food, humans maintain a more complex relationship with plant species, which make the association between the ecological importance and the usefulness of vegetal species – in domains as diverse as medicinal uses and tool-making – far more unpredictable.

Acknowledgements

We thank the Gran Consejo Tsimane' and the Dirección General de la Biodiversidad (Bolivia) for research permits, as well as all the Tsimane' for their interest and cooperation in the field. We also thank P. Pache, E. Tayo, F. Sarabia and D. Durvano for help with the interviews, M. Lero, E. Huasnay and D. Cari for help with the inventories, T. Huanca and E. Conde for logistic support, and botanists from the LPB for help with plant identifications. M. Pardo de Santayana, R. Godoy and TAPS researchers kindly provided comments on earlier drafts. Research was funded by a grant from the BBVA Foundation (BIOCON_06_106-07) and written under the framework of an ERC-FP7 project under grant agreement n°261971 (LEK).

References

- Albuquerque UP. Re-examining hypotheses concerning the use and knowledge of medicinal plants: a study in the Caatinga vegetation of NE Brazil. *Journal of Ethnobiology and Ethnomedicine*. 2006; 2 [online] URL: <http://www.ethnobiomed.com/content/2/1/30>.
- Albuquerque UP, Lucena RFP. Can apparency affect the use of plants by people in tropical forests? *Interciencia*. 2005; 30:506–511.
- Albuquerque UP, Hanazaki N. Five problems in current ethnobotanical research - and some suggestions for strengthening them. *Human Ecology*. 2009; 37:653–661.
- Bennett BC. Plants and people of the Amazonian rainforests. *Biogeosciences*. 1992; 42:599–607.
- Berlin, B. The common flora = the medicinal flora: Theoretical implications of a comparison of medical ethnobotanical and general floristic surveys in the Chiapas Highlands; Symposium “Ethnobotany of southern Mexico”; Society of Economic Botany. 2-5 June 2003;
- Byg A, Vormisto J, Balslev H. Using the useful: characteristics of used palms in south-eastern Ecuador. *Environment Development and Sustainability*. 2006; 8:495–506.
- Costa LM, Santos VA, Ohana DT, Lima ES, Pereira MM, Souza TP. Technological development of aqueous extracts from *Calycophyllum spruceanum* (mulateiro) using factorial design. *Brazilian Journal of Pharmacognosy*. 2011; 21:181–186.
- Cruz-Burga Z, Reyes-García V, Novoa JA, Paneque-Gálvez J, Luz AC. Uso de territorio e integración a la economía de mercado. Estudio de caso en la Amazonía Boliviana. *Natura@Economía*. 2013; 1:105–121.
- Curtis JT, McIntosh RP. An upland forest continuum in the prairie-forest border region of Wisconsin. *Ecology*. 1951; 32:476–496.
- Denevan, WM. The aboriginal cultural geography of the Llanos de Mojos of Bolivia. University of California Press; Berkeley and Los Angeles: 1966.
- Galeano G. Forest use at the Pacific coast of Chocó, Colombia: a quantitative approach. *Economic Botany*. 2000; 54:358–376.
- Godoy R, Tanner S, Reyes-García V, Leonard WR, McDade TW, Vento M, Broesch J, Fitzpatrick IC, Giovanni P, Huanca T, Jha N. The effect of rainfall during gestation and early childhood on adult height in a foraging and horticultural society of the Bolivian Amazon. *American Journal of Human Biology*. 2008; 20:23–34. [PubMed: 17941036]
- Guèze M, Paneque-Gálvez J, Luz AC, Pino J, Orta-Martínez M, Reyes-García V, Macía MJ. Determinants of tree species turnover in a southern Amazonian rain forest. *Journal of Vegetation Science*. 2013; 24:284–295.
- Gullison E, Panfil SN, Strouse J, Hubbell S. Ecology and management of mahogany (*Swietenia macrophylla* King) in the Chimanes Forest, Beni, Bolivia. *Botanical Journal of the Linnean Society*. 1996; 122:9–34.

- Huanca, T. Tsimane' indigenous knowledge, swidden fallow management, and conservation. University of Florida; 1999. Ph.D. Dissertation
- Killeen, TJ.; García, EE.; Beck, SG., editors. Guía de árboles de Bolivia. Herbario Nacional de Bolivia and Missouri Botanical Garden; La Paz, Bolivia: 1993.
- Lawrence A, Phillips OL, Reategui A, Lopez M, Rose S, Wood D, Farfan AJ. Local values for harvested forest plants in Madre de Dios, Peru: towards a more contextualised interpretation of quantitative ethnobotanical data. *Biodiversity and Conservation*. 2005; 14:45–79.
- Lombardo U, Canal-Beeby E, Fehr S, Veit H. Raised fields in the Bolivian Amazonia: a prehistoric green revolution or a flood risk mitigation strategy? *Journal of Archaeological Science*. 2011; 38:502–511.
- Lucena RFP, Araújo EL, Albuquerque UP. Does the local availability of woody caatinga plants (Northeastern Brazil) explain their use value? *Economic Botany*. 2007; 61:347–361.
- Lucena RFP, Medeiros PM, Araújo EL, Alves AGC, Albuquerque UP. The ecological apparency hypothesis and the importance of useful plants in rural communities from Northeastern Brazil: an assessment based on use value. *Journal of Environmental Management*. 2012; 96:106–115. [PubMed: 22208403]
- McDade TW, Reyes-García V, Blackinton P, Tanner S, Huanca T, Leonard WR. Ethnobotanical knowledge is associated with indices of child health in the Bolivian Amazon. *Proceedings of the National Academy of Science*. 2007; 104(15):6134–6139.
- Macía MJ. Multiplicity in palm uses by the Huaorani of Amazonian Ecuador. *Botanical Journal of the Linnean Society*. 2004; 144:149–159.
- Macía MJ. Woody plants diversity, floristic composition and land use history in the Amazonian rain forests of Madidi National Park, Bolivia. *Biodiversity and Conservation*. 2008; 17:2671–2690.
- Macía MJ, Armesilla PJ, Cámara-Leret R, Paniagua-Zambrana N, Villalba S, Balslev H, Pardo-de-Santayana M. Palm uses in northwestern South America: a quantitative review. *Botanical Review*. 2011; 77:462–570.
- Navarro, G.; Maldonado, M., editors. Geografía ecológica de Bolivia: vegetación y ambientes acuáticos. Centro de Ecología Simón I. Patiño; Cochabamba, Bolivia: 2002.
- Paneque-Gálvez J, Mas J-F, Moré G, Cristóbal J, Orta-Martínez M, Luz AC, Guèze M, Pino J, Macía MJ, Reyes-García V. Enhanced land use/cover classification in heterogeneous tropical landscapes using support vector machines and textural homogeneity. *International Journal of Applied Earth Observation and Geoinformation*. 2012 DOI: 10.1016/j.jag.2012.10.007.
- Paz-y-Miño G, Balslev H, Valencia R. Useful lianas of the Siona-Secoya Indians from Amazonian Ecuador. *Economic Botany*. 1995; 49:269–275.
- Phillips O, Gentry AH. The useful plants of Tambopata, Peru: I. Statistical hypotheses tests with a new quantitative technique. *Economic Botany*. 1993a; 47:15–32.
- Phillips O, Gentry AH. The useful plants of Tambopata, Peru: II. Additional hypothesis testing in quantitative ethnobotany. *Economic Botany*. 1993b; 47:33–43.
- Reyes-García, V. Indigenous people, ethnobotanical knowledge, and market economy. A case study of the Tsimane' Amerindians in Lowland Bolivia. University of Florida; 2001. PhD dissertation
- Reyes-García V, Godoy R, Vadez V, Apaza L, Byron E, Huanca T, Leonard WR, Perez E, Wilkie D. Ethnobotanical knowledge shared widely among Tsimane' Amerindians, Bolivia. *Science*. 2003; 299:1707. [PubMed: 12637738]
- Reyes-García V, Vadez V, Byron E, Apaza L, Leonard WR, Perez E, Wilkie D. Market economy and the loss of folk knowledge of plant uses: estimates from the Tsimane' of the Bolivian Amazon. *Current Anthropology*. 2005; 46:651–656.
- Reyes-García V, Huanca T, Vadez V, Leonard W, Wilkie D. Cultural, practical, and economic value of wild plants: a quantitative study in the Bolivian Amazon. *Economic Botany*. 2006; 60:62–74.
- Reyes-García V, Vadez V, Tanner S, Huanca T, Leonard WR, McDade T. Ethnobotanical skills and clearance of tropical rain forest for agriculture: a case study in the lowlands of Bolivia. *Ambio*. 2007a; 26:406–408. [PubMed: 17847806]
- Reyes-García V, Vadez V, Huanca T, Leonard WR, McDade T. Economic development and local ecological knowledge: A deadlock? Quantitative research from a native Amazonian society. *Human Ecology*. 2007b; 35:371–377.

- Reyes-García V, Vadez V, Marti N, Huanca T, Leonard WR, Tanner S. Ethnobotanical knowledge and crop diversity in swidden fields: A study in a native Amazonian society. *Human Ecology*. 2008; 36:569–580.
- Reyes-García V, Ledezma JC, Paneque-Gálvez J, Orta-Martínez M, Gueze M, Lobo A, Guinard D, Huanca T, Luz AC, TAPS Bolivia Study Team. Presence and purpose of non-indigenous peoples on indigenous lands. A descriptive account from the Bolivian Lowlands. *Society and Natural Resources*. 2012; 25(3):270–284.
- Reyes-García V, Paneque-Gálvez J, Bottazzi P, Luz AC, Guèze M, Macia M, Pacheco P. Indigenous land reconfiguration and fragmented institutions: A historical political ecology of the Tsimane' lands. *Journal of Rural Studies*. Accepted pending revisions.
- Schwantes MB, Felfili JM. Ethnobotanical comparison of “Pau Brasil” (*Brosimum rubescens* Taub.) forests in a Xavante Indian and non-Xavante community in eastern Mato Grosso state, Brazil. *Economic Botany*. 2001; 55:555–569.
- Silva ACO, Albuquerque UP. Woody medicinal plants of the caatinga in the state of Pernambuco (Northeast Brazil). *Acta Botanica Brasilica*. 2005; 19:17–26.
- Stepp JR. The role of weeds as sources of pharmaceuticals. *Journal of Ethnopharmacology*. 2004; 92:163–166. [PubMed: 15137997]
- Stepp JR, Moerman DE. The importance of weeds in ethnopharmacology. *Journal of Ethnopharmacology*. 2001; 75:19–23. [PubMed: 11282438]
- Suits DB. Use of dummy variables in regression equations. *Journal of the American Statistical Association*. 1957; 52:548–551.
- Thomas E. The impact of traditional lifestyle, provenance and contact history on plant use knowledge and management: a cross-cultural comparison of two small-scale societies from the Bolivian Amazon. *Human Ecology*. 2012; 40:355–368.
- Thomas E, Vanderbroek I, Van Damme P. Valuation of forests and plant species in Indigenous Territory and National Park Isiboro-Sécure, Bolivia. *Economic Botany*. 2009a; 63:229–241.
- Thomas E, Vandebroek I, Van Damme P, Goetghebeur P, Douterlungne D, Sanca S, Arrazola S. The relation between accessibility, diversity and indigenous valuation of vegetation in the Bolivian Andes. *Journal of Arid Environments*. 2009b; 73:854–861.
- Torre-Cuadros MA, Islebe GA. Traditional ecological knowledge and use of vegetation in southeastern Mexico: a case study from Solferino, Quintana Roo. *Biodiversity and Conservation*. 2003; 12:2455–2476.
- Voeks RA. Disturbance pharmacopoeias: medicine and myth from the humid tropics. *Annals of the Association of American Geographers*. 2004; 94:868–888.
- Woolridge JM. Cluster-sample methods in applied econometrics. *The American Economic Review*. 2003; 93:133–138.

Figure 1.

Map of the sampling area showing the distribution of 48 0.1-plots inventoried in the territory of 6 Tsimane' villages, Bolivian Amazon, along with areas of influence including 22 villages where data of plant uses were collected.

Figure 2.

Scatter plot of the average use value (UV) against importance value index (IVI) of 58 useful species in six Tsimane' areas, Bolivian Amazon. Each point represents the combination of UV and IVI of each species in one of the six surveyed areas (n=293).

Figure 3. Scatter plot of the average use value (UV) per ethnobotanical categories, against importance value index (IVI) of 58 tree species in six Tsimane' areas, Bolivian Amazon. Each point represents the combination of UV and IVI of each species in one of the six surveyed areas (n=293)

Table 1

Importance value index (IVI) and use value (UV) of 58 tree species used in Tsimane' villages, Bolivian Amazon, ranked by decreasing average IVI. The UV of the five species with highest UV in each of the use categories are bolded.

Scientific name	Family	Tsimane' name	Mean IVI	Mean UV					Total
				Food	Firewood	Construction	Materials	Medicine	Others
Number of species				33	54	53	46	43	29
All species			2.44	0.43	0.39	0.36	0.34	0.28	0.09
<i>Iriarteella deltoidea</i> Ruiz & Pav.	Arecaceae	ojdo'	17.05	0.46	0.04	1.29	1.03	0	0
<i>Hura crepitans</i> L.	Euphorbiaceae	conofoto	13.80	0	0.28	0.90	0.08	0.27	1.10
<i>Poulsenia armata</i> (Miq.) Standl.	Moraceae	asha'ba	10.65	0.73	0.46	0.07	0.99	0.01	0.01
<i>Socratea exorrhiza</i> (C. Martius) H.A. Wendl.	Arecaceae	vijiri	9.78	0	0.07	0.81	1.43	0.01	0.03
<i>Leonia crassa</i> L.B. Sm. & A. Fernánde	Violaceae	rojro	6.51	0.07	0.61	0.04	0	0.13	0.17
<i>Otoba parvifolia</i> (Markgraf) A. Gentry	Myristicaceae	cam	5.35	0	0.53	0.86	0.04	0.16	0.50
<i>Astrocaryum murumuru</i> Mart.	Arecaceae	shibo'	4.90	0.49	0.09	1.09	0.61	0	0
<i>Unonopsis floribunda</i> Diels	Annonaceae	veya	4.84	0	0.65	0.67	0.43	0.05	0
<i>Celtis schippii</i> Standl.	Ulmaceae	ñove	4.72	0	0.89	0.23	0.04	0	0
<i>Clarisia racemosa</i> Ruiz & Pav.	Moraceae	väväij	4.66	0	0.34	0.93	1.40	0.06	0.03
<i>Iryanthera juruensis</i> Warburg	Myristicaceae	po'cocos	4.33	0	0.34	0.45	0.52	0.15	0
<i>Rinorea viridifolia</i> Rusby	Violaceae	shayimo	4.32	0	0.30	0.24	0.50	0.17	0
<i>Clarisia biflora</i> Ruiz & Pav.	Moraceae	mu'suru	2.91	0.88	0.67	0.17	1.16	0	0.04
<i>Duguetia spixiana</i> Mart.	Annonaceae	pi'serej	2.74	0.98	0.45	0.20	0.12	0.74	0
<i>Rheedia acuminata</i> Miers	Clusiaceae	tsocón	2.73	1.00	0.61	0.04	0	0	0
<i>Rheedia gardneriana</i> Miers	Clusiaceae	ibijqui	2.68	0	0.53	0.58	0.09	0.08	0
<i>Symphonia globulifera</i> L. f.	Clusiaceae	puñipuson	2.68	0.96	0.34	0.14	0	0.14	0
<i>Dipteryx odorata</i> (Aublet) Willd.	Fabaceae	cojma	2.65	1.00	0.42	0.34	0.38	0.05	0.20
<i>Spondias mombin</i> L.	Anacardiaceae	moco'	2.59	1.13	0.41	0.21	0.02	0.13	0.05
<i>Tetragastris altissima</i> (Aublet) Swart	Burseraceae	na'fa	2.42	0.98	0.47	0.23	0.78	0.30	0.10
<i>Sylogyne cauliflora</i> (C. Martius & Miq.) Mez	Myrsinaceae	viñaj	2.37	1.04	0.33	0.08	0	0.10	0
<i>Siparuna bifida</i> (Poeppig & Endl.) A. DC.	Siparunaceae	vatasan	2.12	0.03	0.19	0.05	0.06	0.41	0
<i>Guarea gomma</i> Pulle	Meliaceae	rosón	2.02	0	0.53	0.60	0.15	0	0.04

Scientific name	Family	Tsimane' name	Mean IVI	Mean UV						Total
				Food	Firewood	Construction	Materials	Medicine	Others	
<i>Terminalia oblonga</i> (Ruiz & Pav.) Steudel	Combretaceae	cotison	1.84	0	0.97	0.19	0.10	0	0	1.30
<i>Pourouma cecropiifolia</i> Mart.	Moraceae	moväij	1.79	1.00	0.71	0	0	0	0	1.75
<i>Cordia nodosa</i> Lam.	Boraginaceae	tayei	1.72	1.00	0.15	0	0	0.24	0	1.48
<i>Terminalia amazonia</i> (Gmelin) Exell	Combretaceae	cavaquis	1.68	0	0.57	0.67	0.13	0.04	0.08	1.54
<i>Sloanea obtusifolia</i> (Moric.) Schumann	Elaeocarpaceae	shesherena	1.64	0	0.81	0.29	0.65	0	0.15	1.94
<i>Abuta grandifolia</i> (C. Martius) Sandw.	Menispermaceae	odo'odo'	1.39	0.98	0.34	0.09	0	0.03	0	1.59
<i>Pourouma minor</i> Benoist	Moraceae	ajmo	1.23	0.78	0.38	0.24	0.04	0.03	0	1.54
<i>Euterpe precatoria</i> Mart.	Arecaceae	ma'ñerej	1.21	0.54	0	0.81	0.08	0.30	0.07	1.90
<i>Calyptanthus lanceolata</i> O. Berg	Myrtaceae	bunau'	1.11	0.95	0.34	0.12	0.05	0.07	0	1.55
<i>Triplaris poeppigiana</i> Wedd.	Polygonaceae	j'i'tyi	1.03	0	0.86	0.35	0.07	0.03	0.04	1.35
<i>Sloanea fragrans</i> Rusby	Elaeocarpaceae	copo'tare	0.99	0.03	0.81	0.27	0	0	0	1.19
<i>Attalea butyracea</i> (Mutis ex L. f.) Wess. Boer	Arecaceae	bitire'	0.94	1.09	0.01	0.58	0.86	0.14	0	2.82
<i>Alibertia pilosa</i> E.H. Krause	Rubiaceae	shishibutuj	0.71	1.00	0.54	0.04	0	0	0	1.58
<i>Aspidosperma rigidum</i> Rusby	Apocynaceae	vambason	0.61	0	0.53	0.03	0.97	0.89	0	2.41
<i>Gallisia integrifolia</i> (Sprengel) Harms	Phytolaccaceae	shepi'	0.60	0.02	0.29	0.03	0.39	1.48	0.05	2.26
<i>Pseudolmedia macrophylla</i> Trécul	Moraceae	pomo	0.59	1.03	0.37	0.19	0	0.16	0	1.79
<i>Bactris maraja</i> Mart.	Arecaceae	cocope'	0.47	0.83	0	0	1.04	0	0	1.91
<i>Copaifera reticulata</i> Ducke	Fabaceae	copaiva	0.40	0	0.23	0.32	0.02	1.62	0.16	2.36
<i>Myroxylon balsamum</i> (L.) Harms	Fabaceae	chuna	0.37	0	0.24	0.77	0.57	0.29	0.08	1.94
<i>Triplaris americana</i> L.	Polygonaceae	chij	0.33	0.05	0.81	0.05	0	1.09	0	1.99
<i>Bactris major</i> Jacq.	Arecaceae	chijchiva'	0.29	0.91	0	0	0.54	0	0.04	1.53
<i>Galipea longiflora</i> K. Krause	Rutaceae	tam'tac	0.24	0	0.16	0.01	0.05	2.07	0	2.30
<i>Borojoa claviflora</i> (K. Schum.) Cuatrec.	Rubiaceae	vuvujiri	0.22	1.05	0.32	0.06	0.05	0	0	1.57
<i>Cedrela odorata</i> L.	Meliaceae	siyamo	0.20	0	0.34	1.19	0.11	0.04	0.44	2.17
<i>Calophyllum brasiliense</i> Cambess.	Clusiaceae	yijdyä'dyä	0.20	0	0.40	0.99	0.24	0.37	0.52	2.51
<i>Calycophyllum spruceanum</i> (Benth.)	Rubiaceae	tunenes	0.19	0	0.82	0.20	0.10	0.11	0	1.23
<i>Swietenia macrophylla</i> King	Meliaceae	chura'	0.17	0	0.23	0.96	0.53	1.06	0.34	3.12
<i>Attalea phalerata</i> Mart. ex Spreng	Arecaceae	mana'i	0.13	1.19	0.01	0.85	0.93	0.45	0.03	3.52

Scientific name	Family	Tsimane' name	Mean IVI	Mean UV						Total
				Food	Firewood	Construction	Materials	Medicine	Others	
<i>Chrysophyllum venezuelanense</i> (Pierre) Penn.	Sapotaceae	ejtere'	0.12	0.36	0.20	0.12	0.06	0.03	0.02	0.97
<i>Helicarpus americanus</i> L.	Tiliaceae	mu'	0.11	0	0.38	0.26	1.07	0.17	0	1.87
<i>Carinitana estrellensis</i> (Raddi) Kuntze	Lecythidaceae	cocoma	0.08	0	0.37	0.35	0.16	0.08	0.08	1.04
<i>Protium aracouchini</i> (Aublet) Marchand	Burseraceae	vi'si	0.08	0	0.31	0.20	0.89	0.70	0.07	2.25
<i>Aniba canelilla</i> (H.B.K.) Mez	Lauraceae	chorecho	0.06	0.54	0.27	0.24	0	1.30	0.07	2.43
<i>Alphanes aculeata</i> Willd.	Arecaceae	cajna	0.04	1.05	0	0	0.11	0	0.05	1.26
<i>Genipa americana</i> L.	Rubiaceae	tyi'	0.04	1.00	0.46	0.05	0.05	0.58	0.64	2.87

Table 2

Results of ordinary least-square bivariate and multiple regressions of overall use value (UV) of 58 tree species against importance value index (IVI) with robust standard errors and clustering by area of influence.

	All species		Species with IVI 1	Species with IVI>1
	Model [1] Bivariate	Model [2] Multivariate	Model [3] Bivariate	Model [4] Bivariate
IVI	0.027 ***	0.019 ***	-0.328	0.040 **
Dummy for life form (palms: omitted category)				
Small trees	^	-0.825 ***	^	^
Trees	^	-0.367 *	^	^
Constant	1.92 ***	2.395 ***	2.056 ***	1.765 ***
N	293	293	161	132
R ²	0.032 ***	0.170	0.014	0.137 **

Notes: N=293. Cells are coefficients followed by a sign for statistical significance

In model [2] six dummy variables were included to control for village fixed-effects.

* P 0.05;

** P 0.01;

*** P 0.001.

^ = variable intentionally omitted.

Table 3

Results of bivariate and multiple ordinary least-square regressions of categorical use value (UV) of 58 tree species against their importance value index (IVI), with robust standard errors and clustering by area of influence.

	Food		Firewood		Construction		Materials		Medicine		Other uses	
	Model [1] Bivariate	Model [2] Multivariate	Model [3] Bivariate	Model [4] Multivariate	Model [5] Bivariate	Model [6] Multivariate	Model [7] Bivariate	Model [8] Multivariate	Model [9] Bivariate	Model [10] Multivariate	Model [11] Bivariate	Model [12] Multivariate
IVI	-0.0085**	-0.0119**	-0.0022	0.0024*	0.0289**	0.0241**	0.0202**	0.0141**	-0.0199****	-0.0184**	0.0086	0.0092
Life form (palms: omitted category)												
Small trees	^	-0.1375*	^	0.2573***	^	-0.5421***	^	-0.6484***	^	0.2827***	^	0.0015
Trees	^	-0.3838***	^	0.4437***	^	-0.2749*	^	-0.4239***	^	0.1956**	^	0.1281***
Constant	0.4596***	0.7602***	0.3817***	-0.0395	0.2987***	0.5700***	0.2982***	0.7091***	0.3665***	0.2744***	0.0752***	0.0198
R ²	0.0076	0.1089	0.0015	0.4381	0.1261	0.2744	0.0498	0.2294	0.0393	0.0831	0.0356	0.1117

Notes: N=293. Cells are coefficients followed by a sign for statistical significance

In multivariate models a set of dummy variables controlling for village fixed effect was included.

* P 0.05;

** P 0.01;

*** P 0.001.

^ = variable intentionally omitted