Published in final edited form as:

Tetrahedron Lett. 2015 June 3; 56(23): 3151–3154. doi:10.1016/j.tetlet.2014.12.071.

Studies toward the synthesis of cinachyramine. An efficient route to 1,5-diazabicyclo[4.4.0]dec-5-enes

Olga V. Barykina-Tassa and Barry B. Snider^{*,†}

Department of Chemistry MS 015, Brandeis University, Waltham MA 02454-9110, USA

Abstract

Hydrogenation (3 atm) of readily available pyrido[1,2-*a*]pyrimidines **10**, **14**, and **17** over 5% Rh/Al₂O₃ forms 1,5-diazabicyclo[4.4.0]dec-5-enes **9**, **15**, and **18** in > 95% yield, providing a general route to this little-studied class of compounds. All attempts to form the tetrahydro-1,2,4-triazine moiety of cinachyramine (**1**) by rearrangement of amidinium dimethylhydrazone **8** using the procedures developed by Kamatori to convert hydrazone **3a** to tetrahydro-1,2,4-triazine **4a** were unsuccessful.

Keywords

1,5-diazabicyclo[4.4.0]dec-5-enes; pyrido[1,2-*a*]pyrimidines; hydrogenation; hydrazones; 1,5-hydrogen shifts

Introduction

Cinachyramine (**1**), a novel alkaloid possessing a hydrazone and two amins, was isolated in 2006 from the Okinawan sponge *Cinachyrella* sp. (see Scheme 1).¹ The structure was assigned by spectroscopic analysis and degradation under acidic conditions to afford azoalkene **2**. Cinachyramine trifluoroacetate showed weak cytotoxic activity against HeLa S₃ cells with an IC₅₀ of 6.8 μg/mL. The structural novelty of cinachyramine and our continuing interest in amidine- and guanidine-containing natural products prompted us to attempt its synthesis.

Kamatori and coworkers reported an extensive series of studies on the rearrangements of the dimethylhydrazones of 1,1,1-trifluoro-2,3-diones such as **3a** (see Scheme 2).² Heating **3a** absorbed on silica gel with ammonium acetate (50 equiv) at 60 °C for 2 days afforded **4a** (36%) with the same tetrahydro-1,2,4-triazine ring as cinachyramine.^{2c} The mechanism

© 2014 Elsevier Ltd. All rights reserved.

^{*}Corresponding author. Tel.: +1-781-736-2550; fax: +1-781-736-2516; snider@brandeis.edu.

[†]Dedicated to the memory of Professor Harry H. Wasserman

Supplementary Material: Supplementary data (experimental details and spectra) associated with this article can be found in the online version, at <http://dx.doi.org/10.1016/j.tetlet...>

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

probably involves imine formation and protonation to give **5a**, which can also be drawn as the resonance structure **6a**. A 1,5-sigmatropic hydrogen shift will give iminium salt **7a**, which will cyclize to give **4a** after deprotonation.

We hoped that amidinium dimethylhydrazone **8** would undergo a similar 1,5-sigmatropic hydrogen shift followed by cyclization to give cinachyramine (**1**) (see Scheme 3). This route was particularly appealing because the conditions are mild enough for a similar sequence to occur in the biosynthesis of **1**. We expected that oxidation of the hydroxy group of **9** to a ketone and hydrazone formation would lead to **8**. We thought that hydroxy amidinium salt **9** should be available by partial hydrogenation of the readily available pyrido[1,2-*a*]pyrimidinium salt **10**.^{3,4}

Results and Discussion

Reaction of 2-amino-3-hydroxypyridine (**11**) with 1,1,3,3-tetraethoxypropane in 60% perchloric acid and ethanol at 80 °C by the literature procedure led to the formation of **10** which precipitated from solution and was isolated in pure form in 85% yield by filtration (see Scheme 4).⁴ We were delighted to find that hydrogenation⁵ of **10** under 3 atm of H₂ over 5% Rh/Al₂O₃ provided **9** as the perchlorate salt in 97% yield. Oxidation of the hydroxy group in **9** with Jones reagent gave oxo amidinium perchlorate salt **12** in 42% yield. Although the amidinium salt is quite stable, deprotonation with K₂CO₃ or NaOH led to partial hydrolysis of the unprotonated amidine. Consistent with this observation, the Swern oxidation and other neutral or basic oxidation procedures were unsuccessful. Therefore, Jones oxidation was the method of choice despite the moderate yield.

The structure of **12** was confirmed by X-ray crystal structure analysis (see Figure 1), which also established that the perchlorate ion was tightly bound and did not exchange with sulfate ion from the sulfuric acid in the Jones oxidation.⁶ It should also be noted that oxo amidinium cation **12** appeared to slowly equilibrate with the hemiketal during prolonged storage in CD₃OD solution.

Oxo amidinium perchlorate **12** was stirred in excess 1,1-dimethylhydrazine for 4 h to afford hydrazone salt **8** in quantitative yield. With the key intermediate **8** in hand we began to investigate the 1,5-sigmatropic hydrogen shift and cyclization needed to complete the synthesis of cinachyramine (**1**). Unfortunately, no reaction occurred on heating **8** in the presence of silica gel with or without NH₄OAc in an oil bath or in a MW oven at 70-100°C. No reaction occurred in TFA at 25 °C or on microwave heating with silica gel prior to decomposition at 250 °C.

To verify that we were carrying out the reaction properly, trifluoroacetyl hydrazone **3b** was prepared by the literature procedure.^{2a,b} Heating **3b** with NH₄OAc in the presence of silica in the MW oven for 1 hour cleanly afforded a 1.7:1 mixture of tetrahydro-1,2,4-triazine **4b** and the analogous tetrahydro-oxadiazine resulting from a 1,5-sigmatropic hydrogen shift prior to imine formation. Therefore the facile 1,5-sigmatropic hydrogen shift that occurs with the trifluoromethyliminium dimethylhydrazone cation **5** does not occur under the same

conditions with amidinium dimethylhydrazone **8**. Presumably the trifluoromethyliminium cation of **5** is much more acidic and therefore more reactive than the amidinium cation of **8**.

We briefly investigated other procedures for the conversion of **8** to cinachyramine (**1**). No reaction occurred on irradiation with 300 or 350 nm UV light in CD₃OD or D₂O. Treating **8** with a wide range of bases either gave recovered **8** or extensive decomposition without any evidence for the formation of cinachyramine (**1**).

Although this approach to cinachyramine was not successful, the hydrogenation of **10** provides a very simple and practical route to 7-hydroxy-1,5-diazabicyclo[4.4.0]dec-5-ene (**9**). 1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU) and 1,5-diazabicyclo[4.3.0]non-5-ene are readily available and widely used as bases,⁸ but the intermediate compound 1,5-diazabicyclo[4.4.0]dec-5-ene (DBD, **16**) has been reported only a few times.⁹ We therefore decided to explore the generality of this route to DBD derivatives. Condensation of 2-aminopyridine (**13**) with 1,1,3,3-tetraethoxypropane by the literature procedure¹⁰ in EtOH and 60% HClO₄ at 80 °C for 2 h provided **14** in 86% yield (see Scheme 5). Hydrogenation (3 atm) of a suspension of **14** in EtOH over 5% Rh/Al₂O₃ for 18 h afforded amidinium salt **15** in 95% yield. We did not determine the minimum amount of rhodium needed, but note that the hydrogenation was efficient with 0.5 % Rh (10 mg of 5% catalyst per mmol of **14**). Amidinium salts such as **15** are stable, but free amidines such as **16** hydrolyze very readily to the *N*-aminopropyl lactam as has also been noted for DBU.^{8b, 11, 12} Free amidine **16** was prepared by passing a solution of **15** in CH₂Cl₂ through DOWEX 550A (OH⁻ form) resin and concentration.¹² Evaporative distillation gave **16** in 90% yield and >95% purity.

Condensation of 2-aminopyridine (**13**) with 3,3-dimethoxybutan-2-one by the literature procedure^{10b,13} in MeOH and 70% HClO₄ at 25 °C for 24 h provided **17** in 61% yield (see Scheme 6). Hydrogenation (3 atm) of a suspension of **17** in EtOH over 5% Rh/Al₂O₃ for 18 h afforded methyl-substituted amidinium salt **18** in 95% yield.¹⁴

In conclusion, we were unable to convert amidinium dimethylhydrazone **8** to cinachyramine (**1**) using the procedures developed by Kamatori to convert hydrazone **3a** to tetrahydro-1,2,4-triazine **4a**. However, we have shown that hydrogenation (3 atm) of readily available pyrido[1,2-*a*]pyrimidines **10**, **14**, and **17** over 5% Rh/Al₂O₃ forms 1,5-diazabicyclo[4.4.0]dec-5-enes **9**, **15**, and **18** in > 95% yield. Since pyrido[1,2-*a*]pyrimidines can be prepared in a single high-yield step from substituted 2-aminopyridines and 1,3-dicarbonyl compounds,^{10b} hydrogenation provides a general route to a wide variety of novel substituted 1,5-diazabicyclo[4.4.0]dec-5-enes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the NIH (GM-50151) for partial support of this research.

References and notes

1. Shimogawa H, Kuribayashi S, Teruya T, Suenaga K, Kigoshi H. *Tetrahedron Lett.* 2006; 47:1409–1411.
2. (a) Kamitori Y, Hojo M, Masuda R, Fujitani T, Ohara S, Yokoyama T. *J Org Chem.* 1988; 53:129–135. (b) Kamitori Y, Hojo M, Masuda R, Yoshida T, Ohara S, Yamada K, Yoshikawa N. *J Org Chem.* 1988; 53:519–526. (c) Kamitori Y, Hojo M, Masuda R, Takahashi T, Wada M, Hiyama T, Mimura Y. *Heterocycles.* 1994; 38:803–809. (d) Kamitori Y, Sekiyama T. *Heterocycles.* 2005; 65:2139–2150. (e) Kamitori Y, Sekiyama T, Okada E. *Heterocycles.* 2007; 71:2219–2226.
3. For reviews of pyrido[1,2-*a*]pyrimidinium salts, see: Hermecz I, Mészáros Z. *Adv Heterocycl Chem.* 1983; 33:242–330. Hermecz I. *Adv Heterocycl Chem.* 2003; 85:173–285. Hajós G. *Curr Org Chem.* 2006; 10:319–322.
4. (a) Leon Palomino MI, Zaitsev BE, Gashev SB, Nikitin SV, Smirnov LD, Koval'chukova OV. *Chem Heterocycl Compd.* 1991; 27:1110–1115. *Chem Abstr.* 1992; 117:48471. (b) Dennin F, Blondeau D, Sliwa H. *Tetrahedron Lett.* 1991; 32:4307–4308.
5. Hydrogenation of 2-aminopyridines over Rh/Al₂O₃ affords amidines. See, for instance: Hansen DW Jr, Currie MG, Hallinan EA, Fok KF, Hagen TJ, Bergmanis AA, Kramer SW, Lee LF, Metz S, Moore WM, Peterson KB, Pitzele BS, Spangler DP, Webber RK, Toth MV, Trivedi M, Tjoeng FS. *U S Patent* 6,046,211. *Apr* 4. 2000 *Chem Abstr.* 2000; 132:265104.
6. Analysis of the X-ray crystal structure data suggests that the counterion is a perchlorate not a bisulfate: (a) There is no evidence for a longer S–O single bond corresponding to a S–O–H bisulfate group. (b) There is no evidence for any plausible O–H...O hydrogen bonds based on O...O contacts. (c) The Cl=O distances should be slightly shorter than S=O distances as is observed. (d) There are no likely H peaks near the perchlorate oxygens, but this last argument is weak owing to minor disorder in that region. Crystallographic data (excluding structure factors) for the structure in this Letter (**12**) have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no CCDC 1033656. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK, (fax: +44-(0)1223-336033 or e-mail: deposit@ccdc.cam.ac.uk).
7. The ¹H NMR spectrum of a freshly prepared solution of **12** shows a single compound. The spectrum becomes more complex over time. The ¹³C NMR spectrum shows a mixture of both the ketone and the hemi-ketal formed from CD₃OD. MS analysis of a methanolic solution of **12** shows cations with *m/z* of 153.1 (MH⁺) and 185.1 (MH⁺ + MeOH).
8. (a) Oediger H, Möller F, Eiter K. *Synthesis.* 1972:591–598. (b) Nakatani K, Hashimoto S. *Yuki Gosei Kagaku Kyokai.* 1975; 33:925–935. *Chem Abstr.* 1976; 84:164644. (c) Hermecz I. *Adv Heterocycl Chem.* 1987; 42:83–202.
9. (a) Möhrle H, Seidel CM. *Arch Pharm.* 1976; 309:471–479. (b) Rokach J, Hamel P, Hunter NR, Reader G, Rooney CS, Anderson PS, Cragoe EJ Jr, Mandel LR. *J Med Chem.* 1979; 22:237–247. [PubMed: 423205] (c) Birman VB, Li X, Han Z. *Org Lett.* 2007; 9:37–40. [PubMed: 17192079]
10. (a) Pollak A, Stanovnik B, Tišler M. *J Org Chem.* 1971; 36:2457–2462. (b) Sawyer JRH, Wibberly DG. *J Chem Soc, Perkin Trans.* 1973; 1:1138–1143. [PubMed: 4739012] (c) Fischer GW. *J Prakt Chem.* 1974; 316:474–484. (d) Tamura S, Ono M. *Chem Pharm Bull.* 1978; 26:3167–3177.
11. Heidelberger C, Guggisberg A, Stephanou E, Hesse M. *Helv Chim Acta.* 1981; 64:399–406.
12. Kumagai N, Matsunaga S, Shibasaki M. *Angew Chem, Int Ed.* 2004; 43:478–482.
13. (a) Nesmeyanov AN, Rybinskaya MI, Bel'skii NK. *Dokl Akad Nauk SSSR.* 1957; 113:343–346. *Chem Abstr.* 1957; 51:81441. (b) Nesmeyanov AN, Rybinskaya MI. *Dokl Akad Nauk SSSR.* 1958; 118:297–298. *Chem Abstr.* 1958; 52:55911.
14. Hydrogenation of **17** over platinum black in EtOH was reported to give the saturated aminal based on the absorption of 5 equivalents of hydrogen and elemental analysis for nitrogen.^{13a} However, we found that hydrogenation of **14** (1 mmol) over Pt₂O (10 mg) in EtOH under 3 atm gave only amidinium perchlorate **15** resulting from the absorption of 4 equivalents of hydrogen, although the hydrogenation was not as clean as that run over Rh/Al₂O₃.

Figure 1.
Crystal structure of oxo amidinium perchlorate **12**.

Scheme 1.

Structure and decomposition product of cinachyramine (**1**).

Scheme 2. Conversion of hydrazone **3** to tetrahydro-1,2,4-triazine **4**

Scheme 3.
Retrosynthesis of cinachyramine (**1**).

Scheme 4.
Synthesis of amidinium hydrazone **8**.

Scheme 5.
Synthesis of 1,5-diazabicyclo[4.4.0]dec-5-ene (DBD, **16**).

Scheme 6.

Synthesis of 2-methyl-1,5-diazabicyclo[4.4.0]dec-5-enium perchlorate.