Technical Section [TECHNICAL NOTES AND TIPS

BD Nokor™ admix needle for percutaneous needle fasciotomy: a novel use of a universal pharmacy needle

M Javed, T Wright, M Murison

Welsh Centre for Burns and Plastic Surgery, Morriston hospital, Swansea, UK

CORRESPONDENCE TO

Muhammad Javed, E: umair.dr@gmail.com

The authors declare they have no conflicts of interest.

Percutaneous needle fasciotomy is one of the less invasive surgical options for treating Dupuytren's contracture.¹ We describe the use of a ubiquitous pharmacy needle, the BD Nokor™ admix needle (Beckton, Dickinson and Company, Franklin Lakes, NJ, US), as an alternative for performing percutaneous fasciotomy.

The needle has a blade at its tip that can be used as a small scalpel for division of fibrous bands (Fig 1). The technique is similar to performing a fasciotomy with a scalpel blade or a conventional needle. In our experience, the BD Nokor™ admix needle can be used effectively for percutaneous fasciotomy in palmar lesions. We would not advocate its use in the digits.

ACKNOWLEDGEMENT

The authors are grateful to Steve Atherton (Medical Illustrations Department, Welsh Centre for Burns and Plastic Surgery) for the illustration in Figure 1.

The material in this technical tip was presented as a poster at the 24th Scandinavian Hand Society Meeting held in Copenhagen, 2012.

Figure 1 Illustration of target tissue (abnormal cords in palmar fascia) in needle fasciotomy and the scalpel tip of the BD Nokor™ admix needle

Reference

1. van Rijssen AL, Gerbrandy FS, Ter Linden H *et al.* A comparison of the direct outcomes of percutaneous needle fasciotomy and limited fasciectomy for Dupuytren's disease: a 6-week follow-up study. *J Hand Surg Am* 2006; **31**: 717–725.

The use of a hypodermic needle to pass intraosseous cerclage wire through bone

S Al-Benna

Ruhr University Bochum, Germany

CORRESPONDENCE TO

Sammy Al-Benna, E: sammy.al-benna@ruhr-uni-bochum.de

Passing a cerclage wire through drill holes can be difficult. This manoeuvre can be facilitated by cutting a 19G hypodermic needle from its base and using the bevelled end to drill a transverse hole on one side of the fracture line (Fig 1). Once the channel has been created, the tip of a 0.6mm cerclage wire is passed into the bevelled end of the needle and the needle is then withdrawn, pulling the cerclage wire through the drill holes (Fig 2).

Figure 1 The bevelled end of the needle is used to drill a transverse hole on one side of the fracture line

Figure 2 The wire is passed through the needle

Figure 1 Intraoperative photograph of incision site following localisation with sonography and preoperative marking of the lesion with the patient standing

Intraoperative ultrasonography localisation of intramuscular abdominal lipomas

GAC Wheble, K Asaad, M Turner, NJ Bennett
Portsmouth Hospitals NHS Trust, UK

CORRESPONDENCE TO

George Wheble, E: george.wheble@porthosp.nhs.uk

BACKGROUND

Ultrasonography is used frequently to determine the site and size of abdominal lipomas prior to surgery. These lesions are often easily palpable on standing. However, they may disappear when the patient lies supine.

TECHNIQUE

A portable ultrasonography scanner is used intraoperatively with the patient supine to confirm the location of the lesion. This often differs noticeably from the position marked preoperatively (Fig 1). Once the skin is incised, the probe can be reapplied, and the ultrasonography images are greatly improved without overlying skin and fat between probe and lesion (Fig 2). The lesion can then be clearly identified and excised.

DISCUSSION

It is common practice to mark patients standing up prior to surgery. From experience, we know these markings would be different if done with the patient supine. We also know that location of

pathology can vary with patient position, especially in patients of larger body habitus.

Intraoperative sonography has been described in other clinical settings such as detection and characterisation of focal liver lesions.¹ It has also been shown to aid localisation of non-palpable breast malignancies in breast conserving surgery.²

Portable ultrasonography machines are readily available in theatre for accurate placement of regional anaesthesia. We advocate the use of intraoperative ultrasonography guided localisation of such lesions; it ensures accurate visualisation of deeper lesions, and helps to minimise operative time and unnecessary dissection.

Figure 2 The lesion is visualised easily on intraoperative imaging