The use of FRED® to reduce intraoperative spectacle fogging

R Ricks¹, G Barrett²

¹Plymouth Hospitals NHS Trust, UK

²Royal Cornwall Hospitals NHS Trust, UK

CORRESPONDENCE TO

Rupert Ricks, E: rupertricks@nhs.net

Bespectacled surgeons are familiar with the annoyance of intraoperative fogging, reducing visibility and surgical effectiveness. Most surgeons combat the risk of fogging with surgical tape applied to the upper edge of their mask, sticking it to their skin, which can cause irritation. FRED® (Covidien, Dublin, Ireland) is a widely used antifog solution that prevents misting of endoscopes intraoperatively. There is often surplus solution, which can be used to prevent misting of surgeon's spectacles. A single application to both sides of spectacle lenses will last for an entire day, preventing the need for irritating surgical tape.

Figure 2 Ball is visible at the entry point

Calibration of the small bowel in stricture-forming small-bowel Crohn's disease

D Luke¹, D Bowley², N Mortensen³

¹Royal Shrewsbury Hospital,

²Royal Centre for Defence Medicine, and

³Oxford University Hospitals, UK

CORRESPONDENCE TO

David Luke, E: davidluke@btinternet.com

BACKGROUND

In 1932, Crohn, Ginzburg and Oppenheimer described a disease of the terminal ileum 'characterised by a process that frequently leads to

Figure 3 Ball is 'chased' along the lumen of the small bowel

Figure 1 Ball is introduced into the enterotomy

stenosis of the lumen of the intestine'.¹ Surgical treatment of Crohn's disease is now based on conservation (rather than resection) of the bowel.² Strictureplasty plays an important part in this strategy.³

TECHNIQUE

Several methods have been reported for the assessment of strictures. We advocate the use of a medical-grade 20-mm steel ball to calibrate the small bowel in Crohn's disease. This technique has been described previously,^{4,5} but not used widely.

An enterotomy is made at the site of an obvious stricture. The ball is inserted into the lumen of the small bowel (Figs 1 and 2). Soft bowel clamps are positioned proximally at the duodenal-jejunal flexure and distally on the small bowel. Then, the steel ball is 'run' along the length of the small bowel (Fig 3). The passage of the ball is impeded by the stricture. Strictureplasty can be carried out as necessary and the ball can continue to be run until all strictures have been identified and treated. The ball is then returned to the original enterotomy, extracted, and the final strictureplasty at the site of the enterotomy completed.