Figure 1 Use of blocking screws for intramedullary nail fixation may lead to overcorrection of axial deformity.

Figure 2 Fine-tuning of blocking screws in overcorrection of axial deformity

TECHNIQUE

If the blocking screw prevents reamer head or nail passage beyond it or overcorrects the deformity it is possible to back the blocking screw out so that it becomes unicortical (Fig 2). This reduces (but does not eliminate) the efficiency of the blocking screw prior to completing the procedure.

DISCUSSION

This fine-tuning technique is easily performed and more expedient than completely resiting the blocking screw. Once the nail has been locked proximally and distally, the screw can be removed if it is no longer required. Alternatively, if the surgeon wishes to retain it as an additional point of fixation for enhanced fracture stability, it may need to be exchanged for a shorter screw.

References

1. Krettek C, Stephan C, Schandelmaier P *et al*. The use of Poller screws as blocking screws in stabilising tibial fractures treated with small diameter intramedullary nails. *J Bone Joint Surg Br* 1999; **81**: 963–968.
2. Stedtfeld HW, Mittlmeier T, Landgraf P, Ewert A. The logic and clinical applications of blocking screws. *J Bone Joint Surg Am* 2004; **86**(Suppl 2): 17–25.

Combined endoscopic approach for patients with multiple bladder stones

M Darrad, M Collins, J Inglis

Royal Wolverhampton NHS Trust, UK

CORRESPONDENCE TO

Maitrey Darrad, E: mpsdarrad@gmail.com

BACKGROUND

Bladder stones account for 5% of all urinary tract stones and are caused by bladder outlet obstruction, neurogenic voiding dysfunction, infection or foreign bodies.¹ While small bladder stones can be managed effectively by transurethral methods and larger stones by open approaches, there is debate as to the best method for managing patients with multiple intermediate sized stones. An effective combined endoscopic approach for the treatment of such patients is described.

Figure 1 Computed tomography before the procedure with arrow showing bladder stones (top) and afterwards with arrow showing suprapubic catheter tip (bottom)

TECHNIQUE

With the patient in the lithotomy position, a 26Fr resectoscope is inserted transurethrally, followed by percutaneous suprapubic puncture. The suprapubic tract is dilated and a 30Fr Amplatz sheath inserted for nephroscope access. Through a 6Fr ureteric catheter, a 365µm laser fibre is passed via the nephroscope. Holmium yttrium aluminium garnet laser (2J and 5Hz) is used to break the larger stones into smaller fragments, which are removed by rigid graspers through the nephroscope. Residual debris is cleared by Ellick evacuation through the resectoscope. A 16Fr suprapubic and urethral catheter is inserted, and removed after 24 hours. Computed tomography prior to and following the procedure is shown in Figure 1. The total operative time is less than 60 minutes.

DISCUSSION

This approach has been described previously in the literature but was advocated for the treatment of large bladder stones with two

urologists working simultaneously with separate video monitors and equipment.² We believe laser lithotripsy is the most effective bladder stone fragmentation method because it enables stone fixation against the bladder wall. The combined endoscopic technique is safe, quick and effective for patients with multiple intermediate sized bladder stones.

References

1. Yoshida O, Okada Y. Epidemiology of urolithiasis in Japan: a chronological and geographical study. *Urol Int* 1990; **45**: 104–111.
2. Sofer M, Kaver I, Greenstein A *et al*. Refinements in treatment of large bladder calculi: simultaneous percutaneous suprapubic and transurethral cystolithotripsy. *Urology* 2004; **64**: 651–654.

Avoiding open reduction and internal fixation in the intramedullary nailing of subtrochanteric femoral fractures

SF Bellringer, C Gee, DGG Wilson, P Stott

Brighton and Sussex University Hospitals NHS Trust, UK

CORRESPONDENCE TO

Simon Bellringer, E: simon.bellringer@bsuh.nhs.uk

BACKGROUND

Subtrochanteric femoral fractures commonly present with significant displacement and are difficult to reduce (Fig 1). They can be pathological and are likely to increase in prevalence owing to bisphosphonate use and an ageing population.¹ Intramedullary nailing of subtrochanteric fractures is considered the gold standard of treatment and because of the deforming forces, many surgeons advocate open reduction for all displaced fractures.² Closed reduction would minimise soft tissue and fracture site disturbance, and we describe a technical tip for this.

Figure 1 Lateral (A) and anteroposterior (B) plain radiography of a pathological subtrochanteric fracture of the left femur