Published in final edited form as:

Metabolism. 2010 November ; 59(11): e18–e19. doi:10.1016/j.metabol.2010.07.017.

Author reply: Characterization of the metabolic and physiologic response to chromium supplementation in subjects with type 2 diabetes mellitus

Dr. William T. Cefalu, M.D.

Pennington Biomedical Research Center

Dear Dr. Field,

We are responding to the specific comments from Dr. Nanne Kleefstra as outlined in the Letter to the Editor in response to our recent article in your journal (1). First and foremost, Dr. Kleefstra stated concerns with the methodology and interpretation of our results in which we suggested that chromium supplements might be beneficial in a selected population of insulin resistant patients in poor glycemic control. Dr. Kleefstra stated they performed a trial in such a patient group, which showed that chromium supplementation did not result in any significant effect compared to placebo (2). With all due respect, we have carefully evaluated Dr. Kleefstra's study in comparison to our study to evaluate the reasons for the different outcomes and actually have reported the findings (3). In this regard, patient selection may be playing a major role. In the studies reported by Kleefstra et al (2), subjects were also from a Western population and were considered obese (BMI ranging 33-35). However, the subjects evaluated by Kleefstra et al(2) were more advanced in their disease process, e.g diabetes duration ranged 10.9 –18.4 years as compared to our recently reported study in subjects who were much earlier in their disease process, generally less than 5 years (1). In addition, by design, the cohort evaluated by Kleefstra et al (2) were taking high dose insulin (dose from 78 – 105 units/day) in addition to many subjects already taking metformin (2,3). Subjects from our recent study evaluating individuals with type 2 diabetes were excluded if they were on medications known to affect carbohydrate metabolism, including anti-diabetic medication. With this data, it remains unclear as to what are the specific differences that account for the conflicting results among carefully controlled trials. One consideration that appears very likely is that differences in subject characteristics may be one of the reasons that may explain the differences noted in efficacy as reported for various studies.

Our data suggesting that phenotype may be important when evaluating chromium response is supported by pre-clinical data (4,5) . For example, we evaluated the response to chromium in both lean and obese JCR rats, a model of obesity and insulin resistance and reported that response to Cr was dependent on baseline phenotype, In a pilot human study, we reported

that baseline patient characteristics can influence the response to Cr as the baseline insulin sensitivity was significantly associated with the clinical response to Cr ($p=0.0004$) and accounted for 40% (partial $R^2=0.4038$) of the variance in the clinical response (6). Finally, our conclusion that a clinical response to chromium is more likely in insulin resistant individuals with type 2 diabetes who have more elevated fasting glucose and A1c levels appears to agree with other prior reports (7).

Dr. Kleefstra commented that reporting the overall results, then dividing the intervention group in two separate cohorts of responders and non-responders seems stretching statistical possibilities. But, as pointed out in his letter, data management was indeed designed to not only look at the effect in the total cohort, but to specifically evaluate what characteristics define a response.

The final point made by Dr. Kleefstra was our definition of chromium status. However, we would like to point out that our study is the most comprehensive study to date for which an overall assessment of chromium status defined as multiple plasma sampling over time combined with urinary excretion. With this design, we demonstrated that chromium “status” in plasma and urine did not predict response. The point to be made is that we don't feel a clinical response to chromium is dictated by serum levels or urinary excretion. Clearly, other factors are involved that predict or are responsible for the effect. This information alone is novel and clarifies a major controversy in human chromium research.

The last point we would like to make is that our study represents the largest cohort to date that have evaluated chromium supplementation with use of “state of the art” metabolic techniques including use of euglycemic-hyperinsulinemic clamps for measuring insulin sensitivity, metabolic chambers for measuring energy expenditure, ingestive behavior labs for assessing dietary intake and use of MRS for assessing myocellular and hepatic fat. Thus, our completed study reported in the recent volume of *Metabolism* represents the most comprehensive, double-blinded, placebo controlled trial to date on chromium supplementation in subjects with Type 2 DM.

References

1. Cefalu WT, Rood J, Pinsonat P, Qin J, Sereda O, Levitan L, Anderson RA, Zhang XH, Martin JM, Martin CK, Wang ZQ, Newcomer B. Characterization of the metabolic and physiologic response to chromium supplementation in subjects with type 2 diabetes mellitus. *Metabolism*. 2010; 59:755–62. [PubMed: 20022616]
2. Kleefstra N, Houweling ST, Jansman FG, Groenier KH, Gans RO, Meyboom-de Jong B, Bakker SJ, Bilo HJ. Chromium treatment has no effect in patients with poorly controlled, insulin-treated type 2 diabetes in an obese Western population: a randomized, double-blind, placebo-controlled trial.
3. Wang ZQ, Cefalu WT. Current concepts about chromium supplementation in type 2 diabetes and insulin resistance. *Curr Diab Rep*. Apr. 2010; 10(2):145–51. [PubMed: 20425574]
4. Wang ZQ, Zhang XH, Russell JC, Hulver M, Cefalu WT. Chromium picolinate enhances skeletal muscle cellular insulin signaling in vivo in obese, insulin-resistant JCR:LA-cp rats. *J Nutr*. 2006; 136(2):415–20. [PubMed: 16424121]
5. Cefalu WT, Wang ZQ, Zhang XH, Baldor LC, Russell JC. Oral chromium picolinate improves carbohydrate and lipid metabolism and enhances skeletal muscle Glut-4 translocation in obese, hyperinsulinemic (JCR-LA corpulent) rats. *J Nutr*. Jun; 2002 132(6):1107–14. [PubMed: 12042418]

6. Wang ZQ, Qin J, Martin J, Zhang XH, Sereda O, Anderson RA, Pinsonat P, Cefalu WT. Phenotype of subjects with type 2 diabetes mellitus may determine clinical response to chromium supplementation. *Metabolism*. 2007; 56(12):1652–5. [PubMed: 17998017]
7. Albarracin CA, Fuqua BC, Evans JL, Goldfine ID. Chromium picolinate and biotin combination improves glucose metabolism in treated, uncontrolled overweight to obese patients with type 2 diabetes. *Diabetes Metab Res Rev*. Jan-Feb;2008 24(1):41–51. [PubMed: 17506119]