Published in final edited form as:

Org Lett. 2015 May 15; 17(10): 2378–2381. doi:10.1021/acs.orglett.5b00896.

Synthesis of Polycyclic Nitrogen Heterocycles via Cascade Pd-Catalyzed Alkene Carboamination/Diels-Alder Reactions

Derick R. White and John P. Wolfe*

Department of Chemistry, University of Michigan, 930 North University Avenue, Ann Arbor, Michigan, 48109-1055, United States

John P. Wolfe: jpwolfe@umich.edu

Abstract

Cascade Pd-catalyzed alkene carboamination/Diels-alder reactions between bromodienes and amines bearing two pendant alkenes are described. These transformations generate 4 bonds, three rings, and 3–5 stereocenters to afford polycyclic nitrogen heterocycles with high diastereoselectivity.

Polycyclic scaffolds that contain a nitrogen heterocycle fused to a six-membered carbocycle are common features of many biologically active molecules. These structures are displayed in natural products such as the yohimbine alkaloids (e.g., reserpine, **1**)¹ and molecules of potential pharmaceutical relevance such as thrombin receptor antagonist **2**.²

*Corresponding Author: jpwolfe@umich.edu.

The authors declare no competing financial interest.

Supporting Information

Experimental procedures, characterization data for all new compounds and copies of ¹H and ¹³C NMR spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Our group has previously illustrated that Pd-catalyzed alkene carboaminations between aryl or alkenyl halides or triflates and alkenes bearing pendant nitrogen nucleophiles are a concise, efficient, and stereoselective means of generating nitrogen heterocycles.^{3,4} We reasoned that coupling our Pd-catalyzed alkene carboamination method with a Diels-Alder cycloaddition could provide a rapid and efficient approach to the synthesis of polycyclic heterocycles with a high degree of stereocontrol. As shown in Scheme 1, a γ -aminoalkene derivative such as **3**, which bears a second pendant alkene unit, could be coupled with a bromodiene in the presence of a Pd-catalyst and a base to afford intermediate **4**. This intermediate could then undergo a Diels-Alder [4+2] cycloaddition⁵ to yield **5**. Our prior studies have illustrated that most Pd-catalyzed alkene carboamination reactions proceed with high diastereoselectivity, and examples of stereocontrolled intramolecular Diels-Alder reactions are well-established. In addition, a few prior examples of Pd-catalyzed cross-coupling/Diels-Alder reactions supported the overall feasibility of this transformation.⁶ Importantly, given the broad scope of the carboamination reactions, this strategy should not be limited to the synthesis of polycyclic pyrrolidine derivatives such as **5**, but rather a much broader array of heterocyclic architectures could be obtained by employing different alkene-tethered nucleophiles.

In our initial experiments we elected to employ 9-bromoanthracene as the bromodiene substrate, as this compound is commercially available, and would allow us to explore the diastereoselectivity (relative face selectivity) of the Diels-Alder reaction in a simple system with no possible stereoisomers resulting from *endo* vs. *exo*-cycloaddition. As shown in Scheme 2, after exploring a range of phosphine ligands, palladium precatalysts,⁷ and reaction conditions for the coupling of **3a** with 9-bromoanthracene we discovered that the best results were obtained using a catalyst composed of Pd(OAc)₂ and Dpe-Phos. Other phosphine ligands provided various amounts of Heck arylation side product **6**, or failed to efficiently promote the carboamination step.

Once suitable reaction conditions had been identified we began to explore the scope of this one-pot reaction sequence. Since chemical yields were modest in some instances, a few different phosphine ligands were examined for most substrate combinations. The Dpe-Phos

ligand provided satisfactory results in most cases, but for some substrates other ligands such as Xantphos, Cy₄Dpe-Phos, or RuPhos proved superior. As shown in Table 1, pentenylamine derivatives **3a–b** bearing a *N*-crotonyl group or an *N*-cinnamyl group were coupled with 9-bromoanthracene to afford polycyclic products **5a–b** with high diastereoselectivity.⁸ The presence of a phenyl substituent adjacent to the *N*-atom was tolerated in the reaction of **3c**, although the desired product **5c** was generated with lower, but still useful, diastereoselectivity (5:1 dr). A similar outcome was observed in the reaction of 3-phenylhydroxylamine derived substrate **7** (3.5:1 dr). In both of these cases the diminished stereocontrol occurred during the Diels-Alder step, as the diastomeric products differed in the configuration of the stereocenters generated in the cycloaddition relative to those formed in the carboamination step. Substrates **8a–b** obtained via acylation of 2-allylaniline also underwent the carboamination/Diels-Alder reaction sequence to afford **10a** and **10b** in moderate yield and high dr. In most instances satisfactory outcomes were achieved simply through extended heating of the reaction mixture at a constant temperature (usually 130 °C). However, for substrate **7** the best results were obtained when the carboamination step was conducted at 90 °C until the substrate had been completely consumed, at which point 1 equiv of LiOTf was added and the temperature was raised to facilitate the cycloaddition.⁹

Although the cascade reactions between 9-bromoanthracene and **3a–c**, **7**, and **8a–b** were reasonably efficient, efforts to extend these transformations to other bromodienes were less successful. For example, the Pd-catalyzed carboamination of **3a** with 2-bromofuran proceeded in good yield to afford **11** (Scheme 3, eq 1), but the Diels-Alder cycloaddition of this compound was not achieved even under forcing conditions (170 °C, microwave irradiation).¹⁰ Efforts to employ non-aromatic bromodienes such as 1-bromo-4-phenylbutadiene in the carboamination reactions of **3a–c**, **7**, or **8a–b** led to the formation of complex mixtures of products even at relatively low temperatures (Scheme 3, eq 2).

Given these limitations, we elected to explore the reactivity of urea or sulfamide derived substrates, as our prior studies had illustrated these types of substrates can be efficiently coupled with various alkenyl bromides. As shown in Table 2, *N,N'*-diallylureas and sulfamides proved to be viable substrates for the cascade Pd-catalyzed carboamination/Diels-Alder reactions, with broader substrate scope than was observed with **3a–c**, **7**, and **8a–b**. Reactions of **12a** or **12b** with 9-bromoanthracene or 2-bromofuran proceeded in moderate to good yield and high diastereoselectivity to afford polycyclic products **13a–d**. In some instances the best results were obtained when the carboamination step was conducted at relatively low temperature (70 °C) until the starting material had been consumed, at which time the temperature was raised to promote the Diels-Alder reaction.⁷ This protocol helped to minimize competing base-mediated alkene isomerization. Substrate **12b** was successfully coupled with non-aromatic bromodienes to yield **13e** and **13f**. However, in these cases diastereoselectivity was poor (ca. 1:1), presumably due to a low preference for *exo*- vs. *endo*-cycloaddition pathways. Finally, cyclohexadienyl triflate also proved to be a viable electrophile in a cascade reaction with **12b**; product **13g** was generated in 30% yield and 19:1 dr. In this case use of modified reaction conditions, in which *t*-BuOLi and PhCF₃ were employed as base and solvent, provided results superior to those obtained using standard conditions.

Finally, we briefly explored the formation of six-membered heterocycles using the cascade reaction sequence. Phenylenediamine derivatives **14a–b** were converted to polycyclic products **15a–c** in moderate to good yield with excellent diastereoselectivity (Scheme 4, eq 3–4). Unfortunately, efforts to employ non-aromatic bromodienes with these substrates did not provide satisfactory results.

In conclusion, we have developed a concise approach to the stereocontrolled synthesis of polycyclic heterocycles via a cascade Pd-catalyzed alkene carboamination/Diels-Alder reaction sequence. The transformations effect the formation of 4 bonds, 3 rings, and 3–5 stereocenters with high levels of diastereoselectivity in most cases examined. Future studies will be directed towards expanding the scope of these reactions to a broader array of diene electrophiles and further exploration of substrate scope.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank the NIH (GM 071650) for financial support of this work.

References

1. Chan F-E, Huang J. *Chem Rev.* 2005; 105:4671–4706. [PubMed: 16351058]
2. Chackalamannil, S.; Wang, Y. US Patent. Scifinder Scholar 20070149518A1. Jun 28. 2007 p. 118210
3. For reviews on Pd-catalyzed carboamination reactions, see: Wolfe JP. *Eur J Org Chem.* 2007:571–582. Wolfe JP. *Synlett.* 2008:2913–2937. Schultz DM, Wolfe JP. *Synthesis.* 2012; 44:351–361. [PubMed: 23243321] Wolfe JP. *Top Heterocycl Chem.* 2013; 32:1–38.
4. For examples of Pd-catalyzed carboamination reactions that afford N-acyl pyrrolidines, isoxazolidines, imidazolidin-2-ones, cyclic sulfamides, or dihydroquinoxalines, see: Bertrand MB, Wolfe JP. *Tetrahedron.* 2005; 61:6447–6459. Lemen GS, Giampietro NC, Hay MB, Wolfe JP. *J Org Chem.* 2009; 74:2533–2540. [PubMed: 19239234] Fritz JA, Nakhla JS, Wolfe JP. *Org Lett.* 2006; 8:2531–2534. [PubMed: 16737306] Fornwald RM, Fritz JA, Wolfe JP. *Chem Eur J.* 2014; 20:8782–8790. [PubMed: 24938206] Hopkins BA, Wolfe JP. *Chem Sci.* 2014; 5:4840–4844. [PubMed: 25431650]
5. For recent reviews on the intramolecular Diels-Alder reaction, see: Parvatker PT, Kadam HK, Tilve SG. *Tetrahedron.* 2014; 70:2857–2888. Juhl M, Tanner D. *Chem Soc Rev.* 2009; 38:2983–2992. [PubMed: 19847335] Takao K, Munakata R, Tadano K. *Chem Rev.* 2005; 105:4779–4807. [PubMed: 16351062]
6. (a) Marsault E, Deslongchamps P. *Org Lett.* 2000; 2:3317–3320. [PubMed: 11029199] (b) Brückner S, Abraham E, Klotz P, Suffert J. *Org Lett.* 2002; 4:3391–3393. [PubMed: 12323026] (c) de Meijere A, von Zezschwitz P, Bräse S. *Acc Chem Res.* 2005; 38:413–422. [PubMed: 15895979] (d) von Zezschwitz P, de Meijere A. *Top Organomet Chem.* 2006; 19:49–89. (e) D'Souza DM, Kiel A, Herten D-P, Rominger F, Mueller TJJ. *Chem Eur J.* 2008; 14:529–547. [PubMed: 17933002] (f) D'Souza DM, Rominger F, Mueller TJJ. *Angew Chem, Int Ed.* 2005; 44:153–158.
7. See the Supporting Information for a table describing details of optimization studies.
8. For examples of intramolecular Diels-Alder reactions of anthracene, see: Ciganek E. *J Org Chem.* 1980; 45:1497–1505.
9. For examples of lithium ion-accelerated Diels-Alder reactions, see: Grieco PA, Nunes JJ, Gaul MD. *J Am Chem Soc.* 1990; 112:4595–4596. Auge J, Gil R, Kalsey S, Lubin-Germain N. *Synlett.* 2000:877–879. Nair V, Maliakal D, Treesa PM, Rath NP, Eigendorf GK. *Synthesis.* 2000:850–856.

10. For examples of intramolecular Diels-Alder reactions of furan, see: Padwa A, Flick AC. *Adv Heterocycl Chem.* 2013; 110:1–41.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Scheme 1.
Cascade Carboamination/Diels Alder Reaction

Scheme 2.

Preliminary experiment – optimal conditions

Scheme 3.
Limitations

Scheme 4.
Transformations of phenylenediamine derivatives

Table 1Cascade Carboamination/Diels-Alder Reactions of Amides and Hydroxylamines^a

substrate	ligand	t (°C)	product	yield (%) ^c
3a	Dpe-Phos	105	5a	57
			dr ^b	>20:1

substrate	ligand	t (°C)	product	dr ^b	yield (%) ^c
3b	Xantphos	130	5b	>20:1	64
3c	Cy ₄ Dpe-Phos	130	5c	5:1	77

^a Conditions: 1.0 equiv substrate, 2.0 equiv 9-bromoanthracene, 2.0 equiv *t*-BuONa, 4 mol % Pd(OAc)₂ or 2 mol % Pd₂(dba)₃, 8 mol % ligand, xylenes (0.07 M), 24 h.

^b Diastereomeric ratio of isolated material. Diastereomeric ratios were similar for crude reaction mixtures unless otherwise noted.

^c Isolated yield (average of two experiments).

^d The reaction was conducted at 90 °C for 24 h until the carboamination step had proceeded to completion, then 1 equiv of LiOTf was added to the reaction mixture and the temperature was increased to 130 °C for an additional 24 h.

^e Diastereomeric ratio and yield for initial purification in which diastereomers were not separated. One pure diastereomer was isolated following a second chromatography

Table 2

Cascade Carboamination/Diels-Alder Reactions of Ureas and Sulfamides^a

substrate	diene	product	dr ^b	yield (%) ^c
12a		13a	>20:1	60 ^d
	13b	>20:1	40 ^d
12b		13c	>20:1	53
	13d	>20:1	60 ^d

substrate	diene	product	dr ^b	yield (%) ^c
			1:1	40 ^e
			1.4:1	49 ^f
			19:1	30 ^g

^a Conditions: 1.0 equiv substrate, 2.0 equiv bromodiene, 2.0 equiv *t*-BuONa, 2 mol % Pd₂(dba)₃, 8 mol % ligand, xylenes (0.07 M), 24 h.

^b Diastereomeric ratio of isolated material. Diastereomeric ratios were similar for crude reaction mixtures unless otherwise noted.

^c Isolated yield (average of two experiments).

^d The reaction was conducted at 70 °C for 24 h until the carboamination step had proceeded to completion, then the temperature was increased to 130 °C for an additional 24 h.

^e The reaction was conducted for 48 h.

^f The reaction was conducted using P(2-furyl)₃ as ligand.

^g The reaction was conducted using 4 mol % Pd(OAc)₂, 12 mol % CPhos, 2 equiv *t*-BuOLi, and trifluorotoluene (0.7 M).