Published in final edited form as:

Heart. 2015 February ; 101(4): 253–256. doi:10.1136/heartjnl-2014-306379.

Basic mechanisms for adverse cardiovascular events associated with air pollution

Michael T. Chin, M.D., Ph.D.

Division of Cardiology, Department of Medicine, University of Washington, Seattle, WA, USA

Abstract

Air pollution is a significant cause of cardiovascular morbidity and mortality worldwide. Although the epidemiologic association between air pollution exposures and exacerbation of cardiovascular disease is well established, the mechanisms by which these exposures promote cardiovascular disease are incompletely understood. In this review I will give an overview of the components of air pollution, an overview of the cardiovascular effects of air pollution exposure and a review of the basic mechanisms that are activated by exposure to promote cardiovascular disease.

Introduction

A wide variety of epidemiological studies conducted over the last decade have demonstrated a strong association between exposure to air pollution and cardiovascular disease (reviewed in ¹). Acute exposure has been associated with an increased incidence of myocardial infarction, arrhythmias, heart failure and all cause mortality. Chronic exposure also promotes cardiovascular disease and all cause mortality to a significantly greater degree than acute exposure. Although the predominant cardiovascular complication of exposure to air pollution is ischemic heart disease, statistically significant associations are also seen for arrhythmias, heart failure and cardiac arrest. While the association between air pollution and cardiovascular disease is now commonly accepted based on a large number of studies, the exact causal agents within the umbrella of airborne pollutants and the pathophysiological mechanisms by which they cause cardiovascular disease are incompletely understood. Many earlier studies linking air pollution and cardiovascular disease have been comprehensively reviewed ¹. Accordingly, this review will summarize briefly the components of air pollution that have been linked to cardiovascular disease and will focus on more recent mechanistic studies outlining how ambient particulate air pollution promotes adverse cardiovascular effects in both animal and controlled human exposure models.

There are six major air pollutants currently regulated by the United States Environmental Protection Agency (Table 1). The gaseous pollutants include carbon monoxide, nitrogen oxides, sulfur dioxide and ozone, while the ambient particulate matter (PM) components are

Correspondence: Michael T. Chin, M.D., Ph.D., Center for Cardiovascular Biology, UW Medicine at South Lake Union, Box 358050, 850 Republican Street, Brotman 353, Seattle, WA 98109, mtchin@u.washington.edu.

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in HEART editions and any other BMJPL products to exploit all subsidiary rights

generally subdivided according to size. These subdivisions reflect the distinct chemical and physical properties of different sized particles, most notably their ability to travel into different parts of the respiratory system. Thoracic particles (PM_{10}), consisting of particles smaller than $10\text{ }\mu\text{m}$, generally do not penetrate beyond the extrathoracic and upper bronchial regions. Fine particles ($PM_{2.5}$), consisting of particles smaller than $2.5\text{ }\mu\text{m}$, penetrate into the small airways and alveoli. Ultrafine particles ($PM_{0.1}$), consisting of particles smaller than $0.1\text{ }\mu\text{m}$, penetrate into the alveoli and possibly the systemic circulation. Coarse particles, in general, arise from natural sources, such as soil, agriculture, plants, volcanos and surface mining. Fine particles generally arise from combustion, smog, diesel and gasoline sources. Ultrafine particles exclusively arise from diesel and gasoline combustion, and are a direct consequence of manmade activity. Diesel exhaust is a prime contributor to air pollution, and consists of both particulate and gaseous components. The particles are composed primarily of elemental carbon, adsorbed organic compounds and small amounts of sulfate, nitrate, metals and other trace elements. Diesel particulates can account for up to approximately 90% of $PM_{2.5}$ in major cities ¹.

While each of the six regulated components of air pollution has been linked to cardiovascular diseases such as hypertension, myocardial infarction, stroke, heart failure and cardiac arrest (Table 1), the evidence for an association between PM exposure and cardiovascular disease far exceeds that for other components. Ambient particulate air pollution is the ninth leading cause of disability adjusted life years worldwide, the fourth leading cause in East Asia and was responsible for 3.2 million deaths worldwide in 2010 ². Interestingly, the effect estimates of air pollution on cardiovascular complications such as myocardial infarction are higher than the effects on lung disorders such as lung cancer and even on all cause mortality resulting in higher population attributable risk.

Although a significant proportion of ambient $PM_{2.5}$ is derived from diesel engines, the chemical composition of $PM_{2.5}$ can vary greatly depending on proximity to other sources such as power generation, industry, agriculture or aviation. To study the mechanisms by which airborne particulates promote cardiovascular disease, many animal and human studies use either diesel engine exhaust particulates (DEPs) or ambient particulate material concentrated directly from the atmosphere (CAPs). These particulates are introduced into a controlled exposure chamber at a defined dose for a defined length of time. CAPs consist of $PM_{2.5}$ from traffic related sources such as diesel exhaust, but also from aviation, agricultural and other intermittent industrial sources of $PM_{2.5}$ that may be located nearby. The advantage of using CAPs is that the $PM_{2.5}$ exposures closely match the human exposures in the regional population. The main disadvantage is that these exposures are difficult to compare when obtained from different regions because $PM_{2.5}$ composition will vary widely both regionally and temporally. In contrast, DEPs are well defined but relatively simple in composition. The inflammatory and toxic potential of these particles is influenced by their content of specific environmental components such as endotoxin and metals.

Both acute and chronic exposure to PM can lead to a variety of specific cardiovascular effects. Associations have been reported with exacerbation of ischemic heart disease ³, heart failure ⁴, cerebrovascular disease ⁵, deep venous thrombosis ⁶, hypertension ⁷ and cardiac arrhythmias ⁹, with varying degrees of evidence supporting these associations. Analysis of

the specific molecular and cellular mechanisms involved reveals diverse and overlapping effects in which systemic inflammation, oxidative stress, neurohumoral and epigenetic modifications contribute in a variety of ways. The specific mechanisms by which PM promotes cardiovascular disease are described below.

Mechanisms by which PM Exposure promotes cardiovascular disease

At present, there are three distinct hypotheses to explain the association between PM exposure and cardiovascular disease, with varying degrees of evidence and consensus ¹. The first hypothesis is best supported and asserts that PM entering the lungs provokes an inflammatory response that promotes oxidative stress and is sufficient to promote systemic oxidative stress and inflammation. This pro-inflammatory state is then thought to promote a variety of pathological processes related to cardiovascular disease, such as increased thrombosis, hypercoagulability, endothelial dysfunction, atherosclerosis progression, insulin resistance and dyslipidemia. The second hypothesis is somewhat supported and asserts that pulmonary exposure leads to activation of lung autonomic nervous system (ANS) arcs mediated by TRP channels that then cause ANS imbalance leading to pathological alterations in vasoconstriction, endothelial dysfunction, hypertension, platelet aggregation, tachycardia, increased heart rate variability and increased arrhythmia potential. The studies implicating TRP channels are largely done with inhibitors but await confirmation with knockout mice. The third hypothesis is supported by only a few studies and asserts that airborne particulates and/or their constituents inhaled through the lungs directly enter the circulation where they may directly interact with tissue components to promote vasoconstriction, endothelial dysfunction, atherosclerosis, hypertension, platelet aggregation, systemic oxidative stress and inflammation. Evidence has also been accumulating that PM exposure can lead to chemical modification of DNA, resulting in epigenetic dysregulation of gene expression. We will review the evidence in support of these hypotheses from exposure studies in both animal and human subjects (Table 2). A working model of how air pollution promotes cardiovascular disease is presented in Figure 1.

Systemic inflammation

Diesel exhaust particles have been shown to induce pulmonary inflammation, oxidative stress and pulmonary cell toxicity in both animal models and human subjects, releasing cytokines and other pro-inflammatory mediators into the circulation. In obese mice, CAPs exposure also promotes adipose tissue inflammation, which is thought to propagate systemic inflammation even further (reviewed in ¹). PM_{2.5} exposure also triggers the appearance of oxidized phospholipids in the lungs that then mediates a systemic inflammatory response mediated through TLR4/NADPH oxidase-dependent mechanisms ¹⁰. Short-term exposure studies in humans have also shown evidence of systemic inflammation as measured by levels of circulating cytokines such as IL-1 β , IL-6 and TNF- α ¹¹. Additional human studies have demonstrated changes in gene expression in peripheral blood mononuclear cells associated with inflammation, oxidative stress and coagulation ¹²¹³, suggesting that these mechanisms are clinically relevant. Other human studies have not shown changes in circulating inflammatory mediators, however, indicating that these studies are not always consistent ¹⁴.

Systemic Oxidative stress

Systemic oxidative stress can be induced by circulating cytokines and inflammatory cells that are activated by events induced in the lungs after exposure to PM, or by leaching of soluble components such as oxidized phospholipids ¹⁰, thereby promoting oxidative stress in the vasculature and other organs such as adipose tissue, the liver and the heart (reviewed in ¹). Exposure of pregnant mice to DEPs has been associated with placental vascular oxidative stress, implying that systemic oxidative stress is induced ¹⁵. Induction of systemic oxidative stress in human subjects exposed to PM under well-characterized conditions has been measured by urinary 8-hydroxy-2'-deoxyguanosine (8-OHdG), a marker of oxidative DNA damage. Urinary 8-OHdG levels correlate with reduction in heart rate variability and PM_{2.5} exposure, confirming the clinical plausibility of this biological mechanism ¹⁶. Generation of reactive oxygen species is a hallmark of activated leukocytes and underscores the close link between systemic oxidative stress and systemic inflammation.

Thrombosis and Coagulation

In the cardiovascular system, animal studies have suggested that airborne particles can promote hypercoagulability, from either direct translocation of particles into the blood or release of circulating factors, although early studies were done with either intratracheal administration of particulates or with PM₁₀ (reviewed in ¹). A more recent study showed that exposure of mice to inhaled CAPs leads to activation of the sympathetic nervous system and systemic catecholamine release, which in turn activates β 2-adrenergic receptor-dependent release of IL-6 from alveolar macrophages to promote a prothrombotic state ¹⁷. Controlled human exposure studies have suggested that brief exposure to DEPs promotes an increase in thrombus formation measured *ex vivo* in a Badimon chamber as well as an increase in platelet-neutrophil and platelet-monocyte aggregates, implying that DEPs induce platelet activation ¹⁸.

Hypertension

Effects of PM exposure on blood pressure have been variable, probably reflecting differences in methodology but also reflecting a relatively weak effect ¹. Long-term exposure to CAPs reportedly promotes vascular oxidative stress, increased sensitivity to angiotensin II, activation of Rho/ROCK, activation of the sympathetic nervous system, increased expression of endothelin, systemic inflammation, hypothalamic inflammation and systemic hypertension in mice ¹¹⁹. A controlled exposure study in human subjects indicated that acute exposure to CAPs in conjunction with ozone leads to transient diastolic hypertension, possibly due to autonomic imbalance ⁸. A more recent study demonstrated that short-term, controlled human exposure to PM_{2.5} CAPs resulted in an increase in systolic blood pressure that correlated with reduced DNA methylation in Alu repetitive elements in circulating leukocytes ⁷.

Vascular dysfunction and atherosclerosis

PM has been shown to have a multitude of vascular effects such as inhibition of endothelium dependent nitric oxide induced vasodilation, enhanced adrenergic induced vasoconstriction, induction of reactive oxygen species, vascular oxidative stress and vascular inflammation,

after both acute and chronic exposures in numerous animal studies (reviewed in ¹). Long-term exposure to CAPs has been shown to promote experimental atherosclerosis in two different mouse models through increased CD36 expression in plaque macrophages that facilitates greater uptake and accumulation of an oxidized form of cholesterol in atherosclerotic lesions ²⁰. Chronic CAP exposure also promotes inflammatory monocyte egress from bone marrow, production of reactive oxygen species and subsequent vascular dysfunction through TLR4 activation of NADPH oxidase in mice ¹⁰. Controlled exposure studies in human subjects have also been demonstrated to promote vascular dysfunction acutely through alteration of responsiveness to vasoactive mediators, to reduce endogenous fibrinolysis and also to enhance exercise induced myocardial ischemia in patients with coronary disease ²¹. Confirmation of effects of particulates on atherosclerosis in human patients through controlled exposures is unlikely to be done, given the decades long incubation period for human atherosclerosis.

Heart Rate Variability and arrhythmias

Exposure of rodents to PM has been shown to decrease heart rate variability and promote arrhythmias, in both acute and chronic settings. This effect correlates with the presence of iron and nickel within the particulates ²². Controlled exposure studies in humans have confirmed that acute exposure to CAPs can cause a temporary reduction in heart rate variability ²³, although a more recent study using diesel exhaust showed no effect ²⁴.

Myocardial fibrosis and heart failure

A 3-month exposure of mice to CAPs has been shown to exacerbate angiotensin II-induced cardiac hypertrophy in a Rho kinase dependent manner ²⁵. A 9-month exposure of mice to CAPs results in increased ventricular size along with systolic and diastolic dysfunction attributable to myocardial fibrosis ²⁶. Interestingly, exposure of pregnant mice to DEPs conveyed a long lasting susceptibility to myocardial fibrosis and heart failure in pups raised to adulthood ¹⁵, suggesting that mediators in the maternal circulation could cross the placenta and promote susceptibility in the developing embryo hearts, through a potential epigenetic mechanism. Although epidemiologic studies have linked air pollution exposure to heart failure ⁴, controlled human exposure studies have not yet demonstrated an effect of particulates on heart function, to the best of our knowledge.

Epigenetic changes and gene expression

Animals exposed to air from steel plants develop mutations in an expanded simple tandem repeat locus analyzed in sperm DNA. The sperm DNA was also found to be hypermethylated when compared to DNA from control mice breathing filtered air, and these changes persisted after the exposure was terminated ²⁷. Controlled human exposure studies showing induction of epigenetic changes in the heart, vasculature or immune cells that alter gene expression and mediate cardiovascular effects have been limited, but a recent controlled exposure study demonstrated an effect of short term particulate exposure on reduction of DNA methylation in Alu repetitive DNA elements in circulating leukocytes that correlated with reduction in blood pressure ⁷.

Clinical and scientific challenges for the future

The epidemiological link between air pollution exposure and cardiovascular disease is well established, and the global impact is clear. Proper environmental policy will be required to minimize unnecessary exposures and therefore limit the impact of air pollution on the incidence and prevalence of cardiovascular disease. Health care workers can advise patients about their risks to help minimize individual exposure. Future studies identifying molecular pathways involved such as IL-6 and Rho/ROCK may lead to the development of pharmacological agents that will mitigate the damage in individuals who cannot avoid exposure.

Acknowledgments

I apologize to the many investigators whose work could not be included in this manuscript due to space limitations. I thank Michael Rosenfeld and Terry Kavanagh for critical reading of the manuscript. MTC is supported by NIH grant ES023581.

References

1. Brook RD, Rajagopalan S, Pope CA 3rd, et al. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation*. 2010; 121(21):2331–78. [PubMed: 20458016]
2. Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012; 380(9859):2224–60. [PubMed: 23245609]
3. Gardner B, Ling F, Hopke PK, et al. Ambient fine particulate air pollution triggers ST-elevation myocardial infarction, but not non-ST elevation myocardial infarction: a case-crossover study. *Part Fibre Toxicol*. 2014; 11:1. [PubMed: 24382024]
4. Shah AS, Langrish JP, Nair H, et al. Global association of air pollution and heart failure: a systematic review and meta-analysis. *Lancet*. 2013; 382(9897):1039–48. [PubMed: 23849322]
5. Stafoggia M, Cesaroni G, Peters A, et al. Long-Term Exposure to Ambient Air Pollution and Incidence of Cerebrovascular Events: Results from 11 European Cohorts within the ESCAPE Project. *Environ Health Perspect*. 2014
6. Baccarelli A, Martinelli I, Zanobetti A, et al. Exposure to particulate air pollution and risk of deep vein thrombosis. *Arch Intern Med*. 2008; 168(9):920–7. [PubMed: 18474755]
7. Bellavia A, Urch B, Speck M, et al. DNA hypomethylation, ambient particulate matter, and increased blood pressure: findings from controlled human exposure experiments. *Journal of the American Heart Association*. 2013; 2(3):e000212. [PubMed: 23782920]
8. Urch B, Silverman F, Corey P, et al. Acute blood pressure responses in healthy adults during controlled air pollution exposures. *Environ Health Perspect*. 2005; 113(8):1052–5. [PubMed: 16079078]
9. Bartell SM, Longhurst J, Tjoa T, et al. Particulate air pollution, ambulatory heart rate variability, and cardiac arrhythmia in retirement community residents with coronary artery disease. *Environ Health Perspect*. 2013; 121(10):1135–41. [PubMed: 23838152]
10. Kampfrath T, Maiseyeu A, Ying Z, et al. Chronic fine particulate matter exposure induces systemic vascular dysfunction via NADPH oxidase and TLR4 pathways. *Circulation research*. 2011; 108(6):716–26. [PubMed: 21273555]
11. Tsai DH, Amyai N, Marques-Vidal P, et al. Effects of particulate matter on inflammatory markers in the general adult population. *Part Fibre Toxicol*. 2012; 9:24. [PubMed: 22769230]
12. Peretz A, Peck EC, Bammler TK, et al. Diesel exhaust inhalation and assessment of peripheral blood mononuclear cell gene transcription effects: an exploratory study of healthy human volunteers. *Inhal Toxicol*. 2007; 19(14):1107–19. [PubMed: 17987463]

13. Pettit AP, Brooks A, Laumbach R, et al. Alteration of peripheral blood monocyte gene expression in humans following diesel exhaust inhalation. *Inhal Toxicol.* 2012; 24(3):172–81. [PubMed: 22369193]
14. Mills NL, Robinson SD, Fokkens PH, et al. Exposure to concentrated ambient particles does not affect vascular function in patients with coronary heart disease. *Environ Health Perspect.* 2008; 116(6):709–15. [PubMed: 18560524]
15. Weldy CS, Liu Y, Liggitt HD, et al. In utero exposure to diesel exhaust air pollution promotes adverse intrauterine conditions, resulting in weight gain, altered blood pressure, and increased susceptibility to heart failure in adult mice. *PLoS One.* 2014; 9(2):e88582. [PubMed: 24533117]
16. Lee MS, Eum KD, Fang SC, et al. Oxidative stress and systemic inflammation as modifiers of cardiac autonomic responses to particulate air pollution. *Int J Cardiol.* 2014; 176(1):166–70. [PubMed: 25074558]
17. Chiarella SE, Soberanes S, Urich D, et al. beta(2)-Adrenergic agonists augment air pollution-induced IL-6 release and thrombosis. *J Clin Invest.* 2014; 124(7):2935–46. [PubMed: 24865431]
18. Lucking AJ, Lundback M, Mills NL, et al. Diesel exhaust inhalation increases thrombus formation in man. *European heart journal.* 2008; 29(24):3043–51. [PubMed: 18952612]
19. Ying Z, Xu X, Bai Y, et al. Long-term exposure to concentrated ambient PM_{2.5} increases mouse blood pressure through abnormal activation of the sympathetic nervous system: a role for hypothalamic inflammation. *Environ Health Perspect.* 2014; 122(1):79–86. [PubMed: 24240275]
20. Rao X, Zhong J, Maiseyeu A, et al. CD36-Dependent 7-Ketocholesterol Accumulation in Macrophages Mediates Progression of Atherosclerosis in Response to Chronic Air Pollution Exposure. *Circulation research.* 2014
21. Mills NL, Tornqvist H, Gonzalez MC, et al. Ischemic and thrombotic effects of dilute diesel-exhaust inhalation in men with coronary heart disease. *N Engl J Med.* 2007; 357(11):1075–82. [PubMed: 17855668]
22. Chang CC, Hwang JS, Chan CC, et al. Interaction effects of ultrafine carbon black with iron and nickel on heart rate variability in spontaneously hypertensive rats. *Environ Health Perspect.* 2007; 115(7):1012–7. [PubMed: 17637915]
23. Gong H Jr, Linn WS, Terrell SL, et al. Altered heart-rate variability in asthmatic and healthy volunteers exposed to concentrated ambient coarse particles. *Inhal Toxicol.* 2004; 16(6-7):335–43. [PubMed: 15204749]
24. Mills NL, Finlayson AE, Gonzalez MC, et al. Diesel exhaust inhalation does not affect heart rhythm or heart rate variability. *Heart.* 2011; 97(7):544–50. [PubMed: 20962342]
25. Ying Z, Yue P, Xu X, et al. Air pollution and cardiac remodeling: a role for RhoA/Rho-kinase. *Am J Physiol Heart Circ Physiol.* 2009; 296(5):H1540–50. [PubMed: 19286943]
26. Wold LE, Ying Z, Hutchinson KR, et al. Cardiovascular remodeling in response to long-term exposure to fine particulate matter air pollution. *Circ Heart Fail.* 2012; 5(4):452–61. [PubMed: 22661498]
27. Yauk C, Polyzos A, Rowan-Carroll A, et al. Germ-line mutations, DNA damage, and global hypermethylation in mice exposed to particulate air pollution in an urban/industrial location. *Proceedings of the National Academy of Sciences of the United States of America.* 2008; 105(2): 605–10. [PubMed: 18195365]
28. Xu X, Sun Y, Ha S, et al. Association between ozone exposure and onset of stroke in Allegheny County, Pennsylvania, USA, 1994–2000. *Neuroepidemiology.* 2013; 41(1):2–6. [PubMed: 23548644]
29. Ensor KB, Raun LH, Persse D. A case-crossover analysis of out-of-hospital cardiac arrest and air pollution. *Circulation.* 2013; 127(11):1192–9. [PubMed: 23406673]
30. Straney L, Finn J, Dennekamp M, et al. Evaluating the impact of air pollution on the incidence of out-of-hospital cardiac arrest in the Perth Metropolitan Region: 2000–2010. *J Epidemiol Community Health.* 2014; 68(1):6–12. [PubMed: 24046350]
31. Mustafic H, Jabre P, Caussin C, et al. Main air pollutants and myocardial infarction: a systematic review and meta-analysis. *Jama.* 2012; 307(7):713–21. [PubMed: 22337682]

32. Bedada GB, Smith CJ, Tyrrell PJ, et al. Short-term effects of ambient particulates and gaseous pollutants on the incidence of transient ischaemic attack and minor stroke: a case-crossover study. *Environ Health*. 2012; 11:77. [PubMed: 23067103]
33. Navas-Acien A, Guallar E, Silbergeld EK, et al. Lead exposure and cardiovascular disease--a systematic review. *Environ Health Perspect*. 2007; 115(3):472–82. [PubMed: 17431501]

Fig. 1.

A working model of how air pollution exposure promotes adverse cardiovascular effects.

Table 1
EPA Regulated Air Pollution Components and Cardiovascular Effects in Humans

Air Pollution Component	Cardiovascular Effect	Selected References
Ozone	Hypertension	8
	Stroke	28
	Out of hospital cardiac arrest	29
Particulate Matter	Ischemic heart disease	3
	Heart Failure	4
	Cerebrovascular disease	5
	Thrombosis	6
	Hypertension	78
	Arrhythmias	19
Carbon Monoxide	Heart failure	4
	Out of hospital cardiac arrest	30
	Increased MI risk	31
Nitrogen Oxides	Heart failure	4
	Transient ischemic attack and stroke	32
	Increased MI risk	31
Sulfur Dioxide	Heart failure	4
	Increased MI risk	31
Lead	Hypertension	33

Table 2
Basic Pathophysiological Mechanisms Activated by Exposure to Particulate Matter in
Animal and Human Subjects

Pathophysiological Mechanism	Selected References
Systemic inflammation	1011
Systemic oxidative stress	1016
Thrombosis and coagulation	1718
Hypertension	7819
Vascular dysfunction and atherosclerosis	102021
Heart rate variability, ECG changes and arrhythmias	22-24
Myocardial fibrosis and heart failure	152526
Epigenetic changes and gene expression	727