CASE REPORT

Spontaneous sphenopalatine pseudoaneurysm: recurrent epistaxis in a patient with cryptogenic vasculopathy

Samantha Goh,¹ Louisa Ritchie,¹ Theo Joseph,¹ Jocelyn Brookes²

¹Department of ENT, Royal National Throat, Nose and Ear Hospital, University College London Hospitals NHS Foundation Trust, London, UK

²Department of Interventional Radiology, University College London Hospitals NHS Foundation Trust, London, UK

Correspondence to

Samantha Goh,
jianlisamantha.goh@nhs.net

Accepted 3 June 2015

SUMMARY

We present a unique case of recurrent epistaxis secondary to a spontaneous sphenopalatine artery aneurysm in a young woman. This was investigated with an external carotid artery angiogram and subsequently treated successfully with endovascular embolisation of the aneurysm.

BACKGROUND

We present a unique case of a patient with recurrent episodes of epistaxis secondary to a spontaneous sphenopalatine artery (SPA) pseudoaneurysm. SPA pseudoaneurysm is a rare cause of epistaxis. In the literature, other cases have been associated with previous nasal or maxillofacial surgery.^{1–4} The gold standard for diagnosis is external carotid artery (ECA) angiography. Cases present in the literature report successful treatment of SPA pseudoaneurysms with microcatheter embolisation.^{2–5}

In the epistaxis management ladder, patients who fail non-surgical management may require endoscopic artery ligation or electrocauterisation.^{6–7} Some authors advocate earlier embolisation therapy, even before surgical arrest of epistaxis,^{7–8} as it is a well tolerated and effective procedure. However, this depends on the local expertise and availability of the interventional radiology service.

CASE PRESENTATION

We present a case of a 29-year-old woman with a complex medical history of cryptogenic vasculopathy; a prothrombotic state with multiple intracranial aneurysms. Since 2007, she suffered recurrent strokes, a combination of haemorrhagic and ischaemic, including thalamic and pontine infarcts. In 2010, she had a subarachnoid haemorrhage and was also found to have multiple inoperable cerebral aneurysms. The same year, she was further diagnosed with a large central pulmonary embolus, on a background of multiple smaller pulmonary emboli, and required an inferior vena cava filter insertion. She has ongoing multidisciplinary team input and remains on long-term daily dalteparin anticoagulation. In spite of her diagnosis of cerebral aneurysms, she does not have a personal or family history of connective tissue disorder. Tests for antiphospholipid syndrome and other thrombophilias were negative, and, to date, a unifying diagnosis is yet to be found for her vasculopathy.

The patient's first presentation to ear, nose and throat (ENT) was in 2013, with recurrent epistaxis, predominantly from the right nasal cavity. There was

no history of trauma or surgery to her nose. She had multiple visits to the emergency department and urgent access ENT clinics, and was managed initially with non-surgical intervention, nasal cavity tamponade and silver nitrate cautery to the anterior nasal septum. Over a period of 12 months, in addition to her emergency attendances, she had three further emergency admissions for recurrent epistaxis, which were managed non-surgically. In 2014, she once again was an inpatient for right-sided epistaxis that failed with nasal cautery and repeated nasal packing. However, due to her complex medical problems, she was a high-risk candidate for surgery. In light of this, a CT angiogram and input from interventional radiology was sought to gain control of the epistaxis. Selective right ECA revealed that the patient had a right SPA pseudoaneurysm (figure 1). Microcatheter embolisation was successfully carried out, preserving the rest of the nasal vasculature (figures 2 and 3). She was subsequently discharged home with no further episodes of epistaxis when reviewed after a 12-month interval.

DISCUSSION

Our case is the first reported example of a spontaneous SPA pseudoaneurysm. Other previously published cases have all been associated with nasal or maxillofacial surgery.^{2–5} Even reportedly atraumatic

Figure 1 Right external carotid artery (ECA) angiography pre-embolisation; fusiform dilation of microbranch of the ECA. Appearance suggestive of pseudoaneurysm, marked with an asterisk (*).

To cite: Goh S, Ritchie L, Joseph T, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-210159

Figure 2 Selective right external carotid artery angiography pre-embolisation; fusiform dilation of pseudoaneurysm identified as coming from a branch of sphenopalatine artery, marked with an asterisk (*).

aneurysms are associated with previous head injury in 90% of cases, usually affecting the internal carotid artery.⁷ Our patient's triad of spontaneous SPA pseudoaneurysm, cerebral aneurysms

Figure 3 Selective right external carotid artery angiography, postembolisation; preservation of normal vasculature. Sphenopalatine artery aneurysm successfully embolised.

and subarachnoid haemorrhages, is also the first to be described in the literature.

It is possible that recurrent nasal packing traumatised the SPA, leading to the formation of a pseudoaneurysm. However, considering the patient's complex vasculitic history, it is more conceivable that her SPA aneurysm was secondary to the same condition causing her intracerebral aneurysms, resulting in pre-existing fragility of the nasal vasculature. Angiography, in this case, allowed the exact position of the defect to be identified and selectively embolised, maximally preserving normal vasculature. The angiogram also excluded other vascular anomalies or malformations that may have complicated the patient's management.

Rudmik and Leung compared the cost-effectiveness of endoscopic SPA ligation with endovascular embolisation, based on healthcare costs in the USA. They concluded that embolisation is more effective but more costly than SPA ligation.⁹ Both treatment modalities are known to have excellent success rates, over 90%,^{6–9} therefore the management of choice will depend on local availability of surgical and interventional radiology expertise.

Learning points

- ▶ Sphenopalatine artery pseudoaneurysms can occur in patients with cerebral aneurysms and lead to intractable epistaxis.
- ▶ Angiography of the nasal vasculature not only enhances diagnostic sensitivity, but also offers a therapeutic opportunity to embolise and definitively manage the bleeding.
- ▶ This therapeutic modality should be considered in intractable epistaxis cases, especially in individuals with known vasculopathies and those unfit for surgery.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Kim YW, Baek MJ, Kim HD, *et al.* Massive epistaxis due to pseudoaneurysm of the sphenopalatine artery: a rare post-operative complication of orthognathic surgery. *J Laryngol Otol* 2013;127:610–13.
- 2 Campbell RG. Sphenopalatine artery pseudoaneurysm after endoscopic sinus surgery: a case report and literature review. *Ear Nose Throat J* 2012;91:E4–11.
- 3 Lanigan DT, Hey JH, West RA. Major vascular complications of orthognathic surgery: false aneurysms and arteriovenous fistulas following orthognathic surgery. *J Oral Maxillofac Surg* 1991;49:571–7.
- 4 Raymond J, Hardy J, Czepko R, *et al.* Arterial injuries in transsphenoidal surgery for pituitary adenoma: the role of angiography and endovascular treatment. *AJNR Am J Neuroradiol* 1997;18:655–65.
- 5 Gökdoğan O, Kizil Y, Aydil U, *et al.* Sphenopalatine artery pseudoaneurysm: a rare cause of intractable epistaxis after endoscopic sinus surgery. *J Craniofac Surg* 2014;25:539–41.
- 6 Abrich V, Brozek A, Boyle TR, *et al.* Risk factors for recurrent spontaneous epistaxis. *Mayo Clin Proc* 2014;89:1636–43.
- 7 Krajina A, Chrobok V. Radiological diagnosis and management of epistaxis. *Cardiovasc Intervent Radiol* 2014;37:26–36.
- 8 Seidel DU, Remmert S, Brassel F, *et al.* Superselective microcoil embolisation in severe intractable epistaxis: an analysis of 12 consecutive cases from an otorhinolaryngologic and interventional neuroradiologic point of view. *Eur Arch Otorhinolaryngol* 2014.
- 9 Rudmik L, Leung R. Cost-effectiveness analysis of endoscopic sphenopalatine artery ligation vs arterial embolization for intractable epistaxis. *JAMA Otolaryngol Head Neck Surg* 2014;140:802–8.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow