CORRESPONDENCE

Immunization Rates at the School Entry in 2012—Nationwide German Data With Regional and Local Breakdown

by Dr. med. Martin Weigel, PD Dr. med. habil. Roswitha Bruns, Kerstin Weitmann, Prof. Dr. med. Wolfgang Hoffmann, MPH in issue 46/2014

Unanswered Questions

I wish to thank the authors for summarizing immunization rates in Germany (1). Special recognition is due to the physicians in the new German states, especially pediatricians. I see the brief and exclusive explanation for the difference in vaccination rates between eastern and western Germany as a problem, however: “One possible explanation is the fact that immunization was compulsory in the former German Democratic Republic.” This implies that socialism still has (dictatorial) aftereffects, 25 years after the wall came down.

Perhaps the doctors who provide immunization have a particular awareness of, or pay particular attention to, achieving the best possible vaccination coverage.

Perhaps the idea of solidarity is greater among the parents of children, and they make decisions that go beyond their own individual situations.

Perhaps a certain egocentric attitude—if everyone has been immunized then I won’t need to worry about getting immunized myself—is less pronounced.

Perhaps the permanent fear that pharma-sponsored decision-making bodies—such as the Standing Committee on Vaccination (*Ständige Impfkommission*, STIKO) or vaccinating physicians are interested in nothing but money is not the main issue here.

Perhaps it would have been possible to evaluate immunization rates in eastern Germany in a more positive light.

In my practice and in the context of a 6-year evaluation interval, 97% of children had received the complete vaccination schedule at the preventive checkup on the occasion of their 2nd birthday.

DOI: 10.3238/arztebl.2015.0402a

REFERENCES

1. Weigel M, Bruns R, Weitmann K, Hoffmann W: Immunization rates at the school entry in 2012—nationwide German data with regional and local breakdown. *Dtsch Arztebl Int* 2014; 111: 788–94.

Dr. med. Steffen Lüder
Berlin, Germany
steffen_lueder@yahoo.de

Conflict of interest statement

Dr. Lüder has received honoraria for consultancy from Pfizer, Abbvie, and MSD. He has been reimbursed for travel expenses and has received speaking fees from Abbvie, Pfizer, and MSD.

Greater Differentiation Would Have Been Desirable

The authors write: “High rates of immunization in the population are desirable.” (1). I would have wished for a more differentiated approach from an institute that has made prevention and health promotion its mission. It should be possible to question each and every recommendation for immunization on any and every occa-

sion, even if it does not originate from a standing vaccination committee that is riven with conflicts of interests.

As a doctor who administers vaccines I ask myself about the risk to which I expose infants if I vaccinate them against hepatitis B, an infection that is primarily sexually transmitted. I would like to know more about the cost–benefit profile of the pertussis vaccine, which is known to be poorly tolerated and whose efficacy tends towards zero (2, 3). I am concerned about the mumps vaccine in girls, since contracting mumps would lower their risk for ovarian cancer. I would like to hear a plausible reason for why I should prevent chickenpox in childhood if it means more Herpes zoster infections in the current generation and greater numbers of cases of chickenpox in adults in future generations.

Vaccines are not “clinically tested before they are given license approval,” and neither is their safety subject to “continuing postmarketing surveillance afterwards.” Necessary hurdles before license approval are lacking—such as genuine placebo studies or surveillance of possible long-term adverse effects.

Parents are not reluctant because they “fear [...] unknown side effects,” as the product information leaflets list sufficient amounts of vaccination risks that anyone would gladly avoid. In a democracy, it will not do to deprive parents of their right to make their own decision by means of campaigns, but they should be enabled to make responsible decisions on behalf of their children and society. The lower immunization rates in southern Germany may thus be seen in a rather more positive light: they stand for a greater number of people who will not let themselves be patronized.

DOI: 10.3238/arztebl.2015.0402b

REFERENCES

1. Weigel M, Bruns R, Weitmann K, Hoffmann W: Immunization rates at the school entry in 2012—nationwide German data with regional and local breakdown. *Dtsch Arztebl Int* 2014; 111: 788–94.
2. Bertilone C, Wallace T, Selvey LA: Finding the ‘who’ in whooping cough: vaccinated siblings are important pertussis sources in infants 6 months of age and under. *Commun Dis Intell Q Rep* 2014; 38: e195–200.
3. van der Maas NA, Mooi FR, de Greeff SC, et al.: Pertussis in the Netherlands, is the current vaccination strategy sufficient to reduce disease burden in young infants? *Vaccine* 2013; 31: 4541–7.

Dr. med. Martin Hirte
Munich, Germany
martin.hirte@t-online.de

Conflict of interest statement

Dr. Hirte has received an author fee for a book publication on the topic of “Immunization: the pros and cons” from Droemer-Knaur publishers.

Motivation Is Crucial for Immunization Success

On reading the comprehensive study reported by Weigel et al. regarding nationwide immunization rates of children starting school in 2012 I could not help noticing that after Bavaria and Baden–Württemberg, North Rhine–Westphalia is among the federal states with the greatest variations for the individual immunizations (1). I am a pediatrician working in hospital in the Warendorf rural district, one of the two rural districts in North Rhine–Westphalia where immunization rates of more than 95% are the norm for all vaccines, and I wish to draw attention to

several factors that may be crucial for achieving these outstanding results and that may serve as a model for other regions.

In 2007 and 2008, immunization rates for poliomyelitis and hepatitis B in the Warendorf district with its primarily rural population were above 95%; the rates for measles were 92% and 93% (G. Fleissner-Busse, Gesundheitsamt [public health authority] Warendorf, personal communication). The latter had also risen to above 95% by 2012. These high immunization rates, which compare well in the intrastate and national comparisons, may be explained, among other factors, by the fact that all 17 pediatricians working in the rural district actively implement the STIKO recommendations. A further possible reason for the high vaccination rates in the Warendorf district may be the fact that, in a setting of 22–25% of children with a migration background, we succeeded in achieving identical rates to those in children without a migration background or even slightly higher rates (up to 2 percentage points higher). This is in stark contrast to the results of the KiGGS study [the German Health Interview and Examination Survey for Children and Adolescents], for example, in which the proportion of migrants was comparable but the immunization rates were 1–7% lower (2, 3). DOI: 10.3238/arztebl.2015.0402c

REFERENCES

1. Weigel M, Bruns R, Weitmann K, Hoffmann W: Immunization rates at the school entry in 2012—nationwide German data with regional and local breakdown. *Dtsch Arztebl Int* 2014; 111: 788–94.
2. Poethko-Müller C, Kuhnert R, Schlaud M: Durchimpfung und Determinanten des Impfstatus in Deutschland. Ergebnisse des Kinder- und Jugendgesundheits-surveys (KiGGS). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheits-schutz* 2007; 50: 851–62.
3. Schenk L, Ellert U, Neuhauser H: Kinder und Jugendliche mit Migrationshintergrund in Deutschland. Methodische Aspekte im Kinder- und Jugendgesundheits-surveys (KiGGS). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheits-schutz* 2007; 50: 590–9.

Dr. med. Carsten Krüger, MIH, FRCPCH

Klinik für Kinder und Jugendliche, St. Franziskus-Hospital, Ahlen, Germany
thea.carsten.krueger@web.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank our correspondents for their interest in our article on immunization rates at the regional level.

We agree with Dr. Lüder that the compulsory vaccinations in the former GDR are not the only reason for the differing immunization rates in the old and new German states. The alternative factors he lists are all plausible but were not investigated in our study owing to methodological constraints. We thank Dr. Krüger for pointing out the active commitment on pediatricians' part as a crucial success factor, which is supported by our data.

In response to Dr. Hirte's letter, we wish to point out that thus far no causal association has been identified between hepatitis B vaccination and autoimmune disorders. The frequency of visits to the doctor decreases the older children get. For this reason, a high immunization rate is more reliably delivered by means of the basic immunization program in infancy. The swap to an acellular pertussis vaccine (Pa) put an end to the adverse effects described for the whole-pathogen vaccine. Waning immunity over time after immunization remains a challenge; for this reason, booster vaccinations are required.

Unfortunately, vaccinated persons can still transmit the pertussis bacterium, which is an additional hindrance to limiting the spread of the disease. It is worth mentioning in this context that even the immunity conferred after the illness itself is limited to 15–20 years (1). The benefit of vaccination is a reduction in severe disease courses that may result in admission to hospital.

Studies have confirmed that contracting mumps lowers the rate of ovarian cancer in girls. However, most of the relevant studies were done at a time when the control group consisted mainly of unvaccinated women. The only study so far with a higher proportion of vaccinated women in the control group did not find any difference in the rate of ovarian cancers, although the manifestation of MUC1 antibodies, which has been credited with reducing the risk, does not occur in vaccinated subjects (2). Complications such as encephalitis, hearing loss, or oophoritis therefore constitute an indication for preventing infection, even in girls.

Since the varicella zoster vaccine was introduced the number of cases of infection necessitating admission to hospital has dropped significantly. The observed increase in cases of zoster infection is still not fully understood in terms of its etiology. However, one striking feature is an increase that happened years before the introduction of the varicella vaccine. Consequently, other risk factors have to be assumed to be possible causes. In spite of the waning immunity over time after vaccination, no increase in the incidence of varicella infections has been observed in adults thus far. This is also the case for the US, where the varicella vaccine has been available since 1995 (3).

The European Medicines Agency (EMA) is responsible for deciding on clinical trials before license approval is granted for medical drugs (4). According to Germany's Act on the Prevention and Control of Infectious Diseases in Man (Protection against Infection Act), possible complications and harms occurring in the context of immunization are notifiable. The Paul Ehrlich-Institute collects such reports and publishes them annually. DOI: 10.3238/arztebl.2015.0403

REFERENCES

1. RKI-Ratgeber Infektionskrankheiten: Merkblätter für Ärzte – Pertussis; *Epidemiologisches Bulletin* 2001; 43: 328–31.
2. Cramer DW, Vitonis AF, Pinheiro SP, et al.: Mumps and ovarian cancer: modern interpretation of an historic association. *Cancer Causes Control* 2010; 21: 1193–201.
3. Siedler A, Hecht J, Rieck T, Tolksdorf K, Hengel H: Die Varizellenimpfung in Deutschland. *Bundesgesundheitsblatt* 2013; 56: 1313–20.
4. European Medicines Agency, Committee for Human Medicinal Products: Note for guidance on the clinical evaluation of vaccines. 2005.
5. Weigel M, Bruns R, Weitmann K, Hoffmann W: Immunization rates at the school entry in 2012—nationwide German data with regional and local breakdown. *Dtsch Arztebl Int* 2014; 111: 788–94.

On behalf of the authors:

Dr. med. Martin Weigel

Abteilung Versorgungsepidemiologie und Community Health,
Institut für Community Medicine
Universitätsmedizin Greifswald
weigelm@uni-greifswald.de

Conflict of interest statement

Dr. Weigel has received study support (third-party funding) from Wyeth Pharma, Sanofi Pasteur MSD, and Novartis-Behring.

Dr. Bruns has received consultancy fees (advisory board) from Pfizer, AstraZeneca, and Sanofi Pasteur MSD. She has received conference delegate fees and travel expenses from Pfizer. She has also received honoraria for speaking from Pfizer, AstraZeneca, Sanofi Pasteur MSD, GlaxoSmith Kline, Novartis, and Abbvie.

Ms. Weitmann and Professor Hoffmann declare that no conflict of interest exists.