What is already known on this topic

Narrowing of the small blood vessels may have a role in the pathogenesis of hypertension, though few prospective trials support such a hypothesis

What this study adds

People with smaller retinal arteriolar diameters are more likely to develop hypertension than people with larger arteriolar diameters, independent of known risk factors for hypertension

This finding suggests that narrowed arterioles are associated with the development of hypertension and therefore that small vessel disease may be a target for antihypertensive treatment

attenuated. Strengths of our study include its population based nature, long follow up, and quantitative evaluation of retinal arteriolar diameters.

Clinical implications

Our results may have potential clinical implications. There is increasing recognition of the value of specifically targeting the microcirculation in the prevention and treatment of hypertension and its complications.^{19 20} For example, there is some evidence that angiotensin converting enzyme inhibitors have direct beneficial effects on microvascular structure and may have therapeutic value in preventing cardiovascular morbidity beyond that of lowering blood pressure.²¹ Our study indicates that further research to evaluate approaches to treatment focused on the microcirculation may be important.

These data were presented at the 44th Annual Conference on Cardiovascular Disease Epidemiology and Prevention, American Heart Association, San Francisco, March 2004.

Contributors: See bmj.com

Funding: American Diabetes Association (RK, AS), NIH grants EYO6594 (RK, BEKK) and HL66018 (RK, TYW), and the Singapore Biomedical Research Council Young Investigator Award (TYW).

Competing interests: None declared.

Ethical approval: All data were collected with Institutional Review Board approval in conformity with all country, federal, and state laws.

- 1 Beevers G, Lip GY, O'Brien E. ABC of hypertension: the pathophysiology of hypertension. *BMJ* 2001;322:912-6.
- 2 Folkow B. Physiological aspects of primary hypertension. *Physiol Rev* 1982;62:347-504.
- 3 Mulvany MJ, Aalkjaer C. Structure and function of small arteries. *Physiol Rev* 1990;70:921-61.
- 4 Rizzoni D, Castellano M, Porteri E, Bettoni G, Muiesan ML, Agabiti-Rosei E. Vascular structural and functional alterations before and after the development of hypertension in SHR. *Am J Hypertens* 1994;7:193-200.
- 5 Norrelund H, Christensen KL, Samani NJ, Kimber P, Mulvany MJ, Korsgaard N. Early narrowed afferent arteriole is a contributor to the development of hypertension. *Hypertension* 1994;24:301-8.
- 6 Mulvany MJ, Baumbach GL, Aalkjaer C, Heagerty AM, Korsgaard N, Schiffrin EL, et al. Vascular remodeling. *Hypertension* 1996;28:505-6.
- 7 Mulvany MJ. Are vascular abnormalities a primary cause or secondary consequence of hypertension? *Hypertension* 1991;18(3 suppl):52-7.
- 8 Hubbard LD, Brothers RJ, King WN, Clegg LX, Klein R, Cooper LS, et al. Methods for evaluation of retinal microvascular abnormalities associated with hypertension/sclerosis in the atherosclerosis risk in communities (ARIC) study. *Ophthalmology* 1999;106:2269-80.
- 9 Klein R, Klein BEK, Linton KLP, DeMets DL. The Beaver Dam eye study: visual acuity. *Ophthalmology* 1991;98:1310-5.
- 10 Klein R, Klein BEK, Lee KE. The changes in visual acuity in a population. The Beaver Dam eye study. *Ophthalmology* 1996;103:1169-78.
- 11 Klein R, Klein BE, Lee KE, Cruickshanks KJ, Chappell RJ. Changes in visual acuity in a population over a 10-year period: the Beaver Dam eye study. *Ophthalmology*. 2001;108:1757-66.
- 12 Wong TY, Klein R, Klein BEK, Meuer SM, Hubbard LD. Retinal vessel diameters and their associations with age and blood pressure. *Invest Ophthalmol Vis Sci* 2003;44:4644-50.
- 13 Knudtson MD, Lee KE, Hubbard LD, Wong TY, Klein R, Klein BEK. A revised method for summarizing retinal vessel diameters. *Curr Eye Res* 2003;27:143-9.
- 14 Klein R, Klein BEK. *Beaver Dam eye study. Manual of operations*. Springfield, VA: Department of Ophthalmology, University of Wisconsin-Madison, 1991. (NTIS Accession No. PB 91-149823.)
- 15 Hypertension Detection and Follow-up Program Cooperative Group. The hypertension detection and follow-up program. *Prev Med* 1976;5:207-15.
- 16 Tso MOM, Jampol LM. Pathophysiology of hypertensive retinopathy. *Ophthalmology* 1982;89:1132-45.
- 17 Klein R, Sharrett AR, Klein BE, Chambless LE, Cooper LS, Hubbard LD, et al. Are retinal arteriolar abnormalities related to atherosclerosis? The atherosclerosis risk in communities study. *Arterioscler Thromb Vasc Biol* 2000;20:1644-50.
- 18 Wong TY, Klein R, Sharrett AR, Duncan BB, Couper DJ, Klein BEK. Retinal arteriolar diameters and risk of hypertension. *Ann Intern Med* 2004;140:248-55.
- 19 Thybo NK, Stephens N, Cooper A, Aalkjaer C, Heagerty AM, Mulvany MJ, et al. Effect of antihypertensive treatment on small arteries of patients with previously untreated essential hypertension. *Hypertension* 1995;25:474-81.
- 20 Rizzoni D, Porteri E, Boari GE, De Ciuceis C, Sleiman I, Muiesan ML, et al. Prognostic significance of small-artery structure in hypertension. *Circulation* 2003;108:2230-5.
- 21 Levy BI, Ambrosio G, Pries AR, Struijker-Boudier HA. Microcirculation in hypertension: a new target for treatment? *Circulation* 2001;104:735-40. (Accepted 29 April 2004)

doi 10.1136/bmj.38124.682523.55

bmjlearning.com

Crunching the numbers

A 70 year old man with giant cell arteritis complains of pains in his shoulders. What percentage of patients with giant cell arteritis also have polymyalgia rheumatica?

- A: Less than 25%
- B: 25-40%
- C: Greater than 60%
- D: 80%

The use of imprecise terms in examination questions causes considerable confusion. A common error when writing such questions is to use inconsistent numeric data.¹ When you use numeric data, you should list the options in numeric order and also in a single format (that is, as single terms or as ranges).

In the example above, options A, B, and C are expressed as ranges whereas option D is a specific percentage. Mixing the formats like this is confusing, and readers could easily mistake option C (greater than 60%) with option D (80%).

BMJ Learning offers online learning modules, throughout which we ask questions, but we avoid imprecise and ambiguous questions. Our new learning module on giant cell arteritis outlines how to diagnose and treat this condition. It also gives an update on recent advances such as the role of colour flow Doppler in diagnosis and how best to avoid corticosteroid induced osteoporosis. You will also find out what percentage of patients with giant cell arteritis also have polymyalgia rheumatica.

Kieran Walsh *editorial registrar, BMJ Learning*
(bmjlearning@bmjgroup.com)

- 1 Case SM, Swanson DB. *Constructing written test questions for the basic and clinical sciences*. 3rd ed. Philadelphia PA: National Board of Medical Examiners, 2001.