Published in final edited form as:

Ann Rheum Dis. 2015 August ; 74(8): 1627–1629. doi:10.1136/annrheumdis-2015-207416.

***ERAP2* is associated with ankylosing spondylitis in *HLA-B27*-positive and *HLA-B27*-negative patients**

A full list of authors and affiliations appears at the end of the article.

The association of endoplasmic reticulum aminopeptidase 2 (*ERAP2*) with ankylosing spondylitis (AS) was recently described in the large International Genetics of AS Consortium Immunochip study.¹ Variants in *ERAP2* have also been associated with inflammatory bowel disease, psoriasis, acute anterior uveitis and birdshot chorioretinopathy.^{2–5} Subsequent investigation demonstrated an association of *ERAP2* with AS which was present when one conditioned on one of the two independent haplotypes of *ERAP1* associated with AS or when *HLA-B27*-negative patients were analysed separately.¹ These two analyses provide analogous evidence for the association of *ERAP2* with AS in *HLA-B27*-negative cases because of the genetic interaction between *HLA-B27* and the AS-associated *ERAP1* variants in AS cases. *ERAP1* and *ERAP2* are located on chromosome 5q15 in the opposite orientation. The locus is challenging to analyse because of the strong linkage disequilibrium (LD) across the locus and the epistasis between *ERAP1* and *HLA-B* alleles associated with AS. We therefore sought to investigate the association of *ERAP2* with AS in *HLA-B27*-positive patients.⁶ This is of clinical importance because functional studies have demonstrated that the strongly AS-protective variant rs2248374 causes a functional *ERAP2* protein knockout, because its G allele causes a loss of *ERAP2* protein expression.⁵⁷ There is also a variant of *ERAP2* which changes its enzyme catalytic activity and specificity (rs2549782, K392A⁸). Because this is in almost complete LD with rs2248374 (1000 Genomes $D'=1.00$, $r^2=0.90$), it is almost never translated in vivo. Further, the very strong LD between these markers means that analysis of rs2549782 for association would yield results almost identical to the results for rs2248374 presented below. Therefore, it is of relevance to determine whether the association of *ERAP2* with *HLA-B27*-negative disease is also found in *HLA-B27*-positive cases, since *ERAP* inhibition may offer a novel therapeutic for AS.⁹

Correspondence to Dr Matthew Brown, The University of Queensland Diamantina Institute, Translational Research Institute, Princess Alexandra Hospital, Woolloongabba, Brisbane QLD 4102, Australia; matt.brown@uq.edu.au.

Contributors PCR, MAB: conceived study; contributed patients/materials; contributed to or performed analysis. M-EC, PL, AC, DE: contributed to or performed analysis; wrote/reviewed/edited the paper. LAB, KH, SL, KBJ, S-CS, MW, MW, XZ, H-JG, GC, JN, BAL, ØF, JT, KL, LJ, YL, XW, DE, RB-V, LSG, SS, NH, JM, JLF-S, MAG-G, CL-L, PB, KG, JSHG, DDG, PR, WPM, HX, IEvdH-B, C-TC, RV-O, CR-S, IMH, FMP-S, RDI, JM, MB, JDR, T-HK, BPW: contributed patients/materials; wrote/reviewed/edited the paper.

Competing interests None declared.

Ethics approval UQ Human Ethics and Metro South Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The data that the study uses have been fully detailed elsewhere.¹

We studied European Immunochip *HLA-B*27*-positive patients (n=7772), *HLA-B*27*-positive controls (n=1204) and unselected European controls (n=13,578). *HLA-B*27*-positive status was taken as an imputed allele dosage of 0.6 or greater in the *HLA-B*27* tag SNP rs116488202 that has 99% sensitivity and 99% specificity for *HLA-B*27* in Europeans.¹ We used rs2248374 as a marker of the associated *ERAP2* haplotype.

We then constructed a range of logistic regression models that controlled for *ERAP1* haplotype association and included *HLA-B*27* controls and unselected controls and also included the first four eigenvectors to control for any potential population stratification. Analyses including only *HLA-B*27*-positive controls are somewhat underpowered relative to the unselected control cohort but were included for illustrative purposes. Analyses were performed using R and the glm() function.

The results show that when using only *HLA-B*27*-positive patients with AS and either *HLA-B*27*-positive or unselected controls there is significant association at the *ERAP2* locus (6.03×10^{-4} and 3.54×10^{-9} respectively; table 1 and figure 1). We also present the models excluding the two associated *ERAP1* haplotypes to demonstrate that the nearby influence of these associations masks the underlying *ERAP2* association.

The significance of this finding is substantial. First, the conditional association noted in all patients with AS previously could have been solely the result of association in *HLA-B*27*-negative patients in the dataset.¹ These data demonstrate that excluding *HLA-B*27*-negative patients still results in a robust level of association in logistic regression models that include only *HLA-B*27*-positive patients. The finding that *ERAP2* is associated with protection from AS can now be unequivocally extended from *HLA-B*27*-negative patients with AS to all patients with AS, meaning potentially 8–9 times more patients could benefit from *ERAP2* inhibition. Such aminopeptidase inhibitors are currently in development¹⁰ and have exciting therapeutic potential for AS and other immune-mediated diseases including inflammatory bowel disease, uveitis and psoriasis.

Authors

Philip C Robinson^{1,2}, Mary-Ellen Costello², Paul Leo², Linda A Bradbury², Kelly Hollis², Adrian Cortes², Seunghun Lee³, Kyung Bin Joo³, Seung-Cheol Shim⁴, Michael Weisman⁵, Michael Ward⁶, Xiaodong Zhou⁷, Henri-Jean Garchon^{8,9}, Gilles Chiochia⁸, Johannes Nossent^{10,11}, Benedicte A Lie^{12,13}, Øystein Førre¹⁴, Jaakko Tuomilehto^{15,16,17}, Kari Laiho¹⁸, Lei Jiang¹⁹, Yu Liu¹⁹, Xin Wu¹⁹, Dirk Elewaut^{20,21}, Ruben Burgos-Vargas²², Lianne S Gensler²³, Simon Stebbings²⁴, Nigil Haroon²⁵, Juan Mulero²⁶, Jose Luis Fernandez-Sueiro²⁷, Miguel A Gonzalez-Gay²⁸, Carlos Lopez-Larrea^{29,30}, Paul Bowness³¹, Karl Gafney³², John S Hill Gaston³³, Dafna D Gladman^{34,35,36}, Proton Rahman³⁷, Walter P Maksymowych³⁸, Huji Xu¹⁹, Irene E van der Horst-Bruinsma³⁹, Chung-Tei Chou^{40,41}, Raphael Valle-Oñate⁴², María Consuelo Romero-Sánchez⁴², Inger Myrnes Hansen⁴³, Fernando M Pimentel-Santos⁴⁴, Robert D Inman²⁵, Javier Martin⁴⁵, Maxime Breban^{8,46}, David Evans^{2,47,48}, John D Reveille⁷, Tae-Hwan Kim³, B Paul Wordsworth⁴⁹, and Matthew A Brown²

Affiliations

¹Centre for Neurogenetics and Statistical Genomics, Queensland Brain Institute, University of Queensland, Brisbane, Queensland, Australia ²The University of Queensland Diamantina Institute, Translational Research Institute, Princess Alexandra Hospital, Brisbane, Queensland, Australia ³Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases, Seoul, Republic of Korea ⁴Department of Medicine, Division of Rheumatology, Eulji University Hospital, Daejeon, Republic of Korea ⁵Department of Medicine/ Rheumatology, Cedars-Sinai Medical Center, Los Angeles, California, USA ⁶National Institute of Arthritis and Musculoskeletal and Skin Diseases, US National Institutes of Health, Bethesda, Maryland, USA ⁷Department of Rheumatology and Clinical Immunogenetics, University of Texas Health Science Center at Houston, Houston, Texas, USA ⁸INSERM U1173, UFR Simone Veil, Versailles-Saint-Quentin University, Laboratoire d'Excellence INFLAMEX, France ⁹Genetics Division, Ambroise Paré Hospital (AP-HP), Boulogne-Billancourt, Paris, France ¹⁰School of Medicine, University of Western Australia, Western Australia, Perth, Australia ¹¹Department of Rheumatology, Sir Charles Gairdner Hospital, Perth, Western Australia, Australia ¹²Department of Medical Genetics, University of Oslo and Oslo University Hospital, Oslo, Norway ¹³Department of Immunology, Oslo University Hospital, Oslo, Norway ¹⁴Department of Rheumatology, University Hospital Oslo, Oslo, Norway ¹⁵Centre for Vascular Prevention, Danube-University Krems, Krems, Austria. ¹⁶Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland ¹⁷King Abdulaziz University, Jeddah, Saudi Arabia ¹⁸Department of Medicine, Päijät-Häme Central Hospital, Lahti, Finland ¹⁹Department of Rheumatology and Immunology, Shanghai Changzheng Hospital, Second Military Medical University, Shanghai, China ²⁰Department of Rheumatology, Ghent University Hospital, Ghent, Belgium ²¹VIB Inflammation Research Center, Ghent, Belgium ²²Department of Rheumatology, Faculty of Medicine, Hospital General de México, Universidad Nacional Autónoma de México, Mexico City, México ²³University of California, San Francisco, California, USA ²⁴Department of Medicine, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand ²⁵Division of Rheumatology, Toronto Western Hospital, Toronto, Ontario, Canada ²⁶Rheumatology Department, Hospital Puerta de Hierro, Madrid, Spain ²⁷Rheumatology Department, Complejo Hospitalario La Coruña, Instituto de Investigación Biomédica A Coruña (INIBIC), La Coruña, Spain ²⁸Rheumatology Department, Hospital Marqués de Valdecilla, Instituto de Formación e Investigación Marqués de Valdecillas (IFIMAV), Santander, Spain ²⁹Department of Immunology, Asturias Central University Hospital, Oviedo, Spain ³⁰Fundación Renal Iñigo Alvarez de Toledo, Madrid, Spain ³¹National Institute for Health Research (NIHR) Oxford Musculoskeletal Biomedical Research Unit, Nuffield Orthopaedic Centre, Oxford, UK ³²Department of Rheumatology, Norfolk and Norwich University Hospital, Norwich, UK ³³Department of Medicine, University of Cambridge, Addenbrooke's Hospital, Cambridge, UK ³⁴Division of Rheumatology, University of Toronto, Toronto, Ontario, Canada ³⁵Toronto Western Research

Institute, Toronto, Ontario, Canada ³⁶Psoriatic Arthritis Program, University Health Network, Toronto, Ontario, Canada ³⁷Memorial University of Newfoundland, St. John's, Newfoundland, Canada ³⁸Department of Medicine, University of Alberta, Edmonton, Alberta, Canada ³⁹Department of Rheumatology, VU University Medical Center, Amsterdam, The Netherlands ⁴⁰Department of Medicine, Division of Allergy, Immunology, Rheumatology, Taipei Veterans General Hospital, Taipei, Taiwan ⁴¹School of Medicine, National Yang-Ming University, Taipei, Taiwan ⁴²SpA Group Hospital Militar, Universidad Militar Nueva Granada, Bogotá, Colombia ⁴³Helgelandssykehuset, Mo i Rana, Norway ⁴⁴Chronic Diseases Research Centre (CEDOC), Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Lisbon, Portugal ⁴⁵Instituto de Parasitología y Biomedicina López-Neyra, Consejo Superior de Investigaciones Científicas, Granada, Spain ⁴⁶Division of Rheumatology, Ambroise Paré Hospital, Assistance Publique-Hôpitaux de Paris, Boulogne-Billancourt, Paris, France ⁴⁷MRC Integrative Epidemiology Unit, University of Bristol, UK ⁴⁸School of Social and Community Medicine, University of Bristol, Bristol, UK ⁴⁹National Institute for Health Research (NIHR) Oxford Musculoskeletal Biomedical Research Unit, Nuffield Orthopaedic Centre, Oxford, UK

Acknowledgements

We wish to thank all the study participants who generously donated their DNA to the AS Immunochip study.

Funding PR is funded by the RACP-ARA-Starr Research Fellowship. MAB is a National Health and Medical Research Council Senior Principle Research Fellow. This work was in part funded by grants from Arthritis Research UK (19536 & 18797), the NIHR Oxford Comprehensive Biomedical Research Centre (immunity and inflammation theme A93081), NIHR Thames Valley collaborative research network and National Ankylosing Spondylitis Society (UK) and NIH/NIAMS—P01-AR052915.

REFERENCES

1. Cortes A, Hadler J, Pointon JP, et al. Identification of multiple risk variants for ankylosing spondylitis through high-density genotyping of immune-related loci. *Nat Genet.* 2013; 45:730–738. [PubMed: 23749187]
2. Robinson PC, Claushuis TA, Cortes A, et al. Genetic dissection of acute anterior uveitis reveals similarities and differences in associations observed with ankylosing spondylitis. *Arthritis Rheumatol.* 2015; 67:140–151. [PubMed: 25200001]
3. Tsoi LC, Spain SL, Knight J, et al. Identification of 15 new psoriasis susceptibility loci highlights the role of innate immunity. *Nat Genet.* 2012; 44:1341–1348. [PubMed: 23143594]
4. Franke A, McGovern DP, Barrett JC, et al. Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. *Nat Genet.* 2010; 42:1118–1125. [PubMed: 21102463]
5. Kuiper JJ, Van Setten J, Ripke S, et al. A genome-wide association study identifies a functional ERAP2 haplotype associated with birdshot chorioretinopathy. *Hum Mol Genet.* 2014; 23:6081–6087. [PubMed: 24957906]
6. Evans DM, Spencer CC, Pointon JJ, et al. Interaction between ERAP1 and HLA-B27 in ankylosing spondylitis implicates peptide handling in the mechanism for HLA-B27 in disease susceptibility. *Nat Genet.* 2011; 43:761–767. [PubMed: 21743469]
7. Andres AM, Dennis MY, Kretschmar WW, et al. Balancing selection maintains a form of ERAP2 that undergoes nonsense-mediated decay and affects antigen presentation. *PLoS Genet.* 2010; 6:e1001157. [PubMed: 20976248]

8. Evnouchidou I, Birtley J, Seregin S, et al. A common single nucleotide polymorphism in endoplasmic reticulum aminopeptidase 2 induces a specificity switch that leads to altered antigen processing. *J Immunol.* 2012; 189:2383–2392. [PubMed: 22837489]
9. Robinson PC, Brown MA. Genetics of ankylosing spondylitis. *Mol Immunol.* 2014; 57:2–11. [PubMed: 23916070]
10. Zervoudi E, Saridakis E, Birtley JR, et al. Rationally designed inhibitor targeting antigen-trimming aminopeptidases enhances antigen presentation and cytotoxic T-cell responses. *Proc Natl Acad Sci USA.* 2013; 110:19890–19895. [PubMed: 24248368]

Figure 1.

Local association plots of the *ERAP2* locus when the association model using *HLA-B27*-positive controls (upper) and unselected controls (lower) are used in logistic regression models that incorporate the two independent ankylosing spondylitis-associated *ERAP1* haplotypes, rs30187 and rs10050860, as covariates.

Logistic regression models of *ERAP2* (rs2248374) association controlling for the AS-associated *ERAP1* haplotypes (tagged by rs30187 and rs10050860) in HLA-B27-positive cases and either HLA-B27-positive or unselected controls

Table 1

Model	Controls	rs2248374 G allele β	SE	OR (95% CI)	p Value
rs2248374+rs30187+rs10050860+1st 4 eigenvectors	HLA-B27 positive	-0.16	0.047	0.85 (0.78 to 0.93)	6.03×10^{-4}
rs2248374+rs30187+rs10050860+1st 4 eigenvectors	Unselected	-0.13	0.021	0.88 (0.85 to 0.92)	3.54×10^{-9}
rs2248374+1st 4 eigenvectors	HLA-B27 positive	-0.041	0.044	0.96 (0.88 to 1.05)	0.36
rs2248374+1st 4 eigenvectors	Unselected	-0.028	0.020	0.97 (0.93 to 1.01)	0.16

AS, ankylosing spondylitis.