CASE REPORT

Aborted sudden cardiac death associated with an anomalous right coronary artery

Rienzi A Diaz,¹ Julio Valdés²

¹Departamento de Medicina Interna, Escuela de Medicina, Universidad de Valparaíso, Viña del Mar, Chile

²Unidad de Cuidados Intensivos, Clínica Reñaca, Viña del Mar, Chile

Correspondence to Professor Rienzi A Diaz, diaz.rienzi@gmail.com

Accepted 24 June 2015

SUMMARY

Coronary artery anomalies arising from the opposite sinus of Valsalva and having an interarterial course between the aorta (AO) and pulmonary artery (PA) are the second most common cause of sudden cardiac death among young athletes, after hypertrophic cardiomyopathy. The right coronary artery (RCA) originating from the AO above the left sinus of Valsalva (LSV) is an extremely rare anomaly. We report the first case of a RCA arising from the AO above the LSV that subsequently runs between the AO and the PA, discovered by a 64-slice multidetector coronary CT, in a patient who was successfully resuscitated from ventricular fibrillation (VF) cardiac arrest while running in a marathon race.

BACKGROUND

This case clearly illustrates how a coronary anomaly can lead to sudden cardiac death (SCD) due to ventricular fibrillation (VF) induced by exercise and that a multidetector coronary CT (MDCCT) should be considered to evaluate patients experiencing cardiovascular symptoms during exercise and routinely in competitive athletes.

CASE PRESENTATION

Background

Coronary artery anomalies are found in 0.6–1.3% of patients undergoing coronary angiography^{1 2} and represent the second most common cause of SCD in young athletes who died during or shortly after strenuous physical activity.^{3 4} Coronary arteries arising from the opposite sinus of Valsalva and having an interarterial course between the aorta (AO) and the pulmonary artery (PA) are considered to be life-threatening anomalies.^{5–7} A right coronary artery (RCA) originating from the AO above the left sinus of Valsalva (LSV) is an extremely rare anomaly, scarcely reported in the literature, and represents fortuitous findings at autopsy⁸ or during coronary angiography.^{9 10} We report the first case of a RCA arising from the AO above the LSV that subsequently runs between the AO and the PA, discovered by a 64-slice MDCCT in an athlete who had a VF cardiac arrest while competing in a marathon.

Case presentation

A 38-year-old woman, an athlete, was admitted to the intensive care unit at Clínica Reñaca after collapsing with loss of consciousness while competing in a marathon race. Two months before this event, she had begun to experience dyspnoea on exertion.

She underwent a medical check-up with an exercise ECG and a two-dimensional echocardiography because of the symptoms; the tests were normal except for the diagnosis of exercise-induced asthma. She was prescribed treatment in the form of a salbutamol inhaler.

At the scene of her collapse, the patient was immediately resuscitated with use of cardiopulmonary resuscitation, which was continued during ambulance transportation to our hospital. An automated external defibrillator was unavailable. On admission, the patient was in VF, which was successfully converted to sinus tachycardia with 120 J. After cardioversion, she regained consciousness and her blood pressure was 130/85 mm Hg. Pulmonary and cardiac auscultations were normal. A 12-lead ECG showed sinus rhythm without signs of ischaemia. Chest X-ray and a 2D echocardiogram were normal. Initial laboratory investigations showed arterial blood gases (using FiO₂ of 0.50): pH 7.1, PaO₂ 67.0 mm Hg, PaCO₂ 50.1 mm Hg, HCO₃ 15.2 (base excess –14.5), oxyhaemoglobin saturation 85.6%, arterial lactate 57.4 mg/dL, troponin <0.03 ng/mL, creatine kinase-myocardial band 2.8 ng/mL and blood urea nitrogen 32.1 pg/mL. We considered that further diagnostic evaluation was warranted. A MDCCT depicted an anomalous origin of the RCA originating from the left side of the AO anteriorly and above the LSV (figure 1), with an acute angle take-off along the aortic wall that subsequently courses between the AO and the PA, resulting in compression and narrowing of the proximal RCA lumen (figure 2). We felt that the RCA likely became compressed during the patient's vigorous activity and that she subsequently suffered VF arrest due to RCA-associated myocardial ischaemia.

She was counselled about this risk and discussions were held with her about options for treatment, including reimplantation of the RCA into its typical native location or implantation of an implantable cardioverter-defibrillator coupled with restrictions on exercise activity levels. She elected to consider these options and requested hospital dismissal. Our patient was discharged from hospital free of symptoms at day 4.

OUTCOME AND FOLLOW-UP

After our patient was discharged from hospital, we lost the follow-up.

DISCUSSION

The anomalous origin of a coronary artery from the opposite aortic sinus and coursing between the

To cite: Diaz RA, Valdés J. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-210850

Figure 1 (A) Three-dimensional volume rendered MDCCT of the coronary arteries, showing an anomalous RCA (yellow arrow) arising from the left side of the aorta anteriorly and superiorly of the (LMCA, white arrow) coming from the LSV. (B) Volume rendered tree view of the coronary arteries. This clearly depicts that the RCA arises ectopically from the left side of the aorta (yellow arrow) anteriorly and above the LSV (blue arrow) and the LMCA (white arrow; LAD, left anterior descending coronary artery; LCX, left circumflex artery; LMCA, left main coronary artery; LSV, left sinus of Valsalva; MDCCT, multidetector coronary CT; RCA, right coronary artery).

AO and PA is one of the most common causes of SCD in the young.^{3 4 11} Although SCD may be the first manifestation of this heart condition, almost half of these patients might experience some warning with cardiovascular symptoms such as palpitations, syncope, chest pain or dyspnoea on exertion,^{11–13} as occurred in our patient. Certainly, the diagnosis of this disease requires a high index of suspicion and access to more advance cardiovascular imaging modalities such as a MDCCT. The tests that were chosen to initially evaluate our patient were reasonable and appropriate from a cost-effective evaluation perspective. However, they were unable to detect her congenital aortic abnormality. These tests are frequently normal in this population of patients.^{11 13}

This case also highlights another unusual aspect of an anomalous origin of a RCA. Usually, the anomalous origin from the contralateral side of the AO is from within the sinus of Valsalva or from the proximal normal left coronary artery.^{6 14–17}

The RCA originating from the left side of the AO above the LSV is an extremely rare anomaly scarcely reported in the medical literature.^{8–10} Our patient is the first case of a RCA anomaly arising from the AO above the LSV with a subsequent course between the AO and the PA, detected by a MDCCT in a competitive athlete who developed a cardiac arrest from VF during a marathon race. This non-invasive imaging technology provides exceptional quality of the 3D anatomical coronary images allowing the visualisation of the

Figure 2 Axial images from a multidetector coronary CT coronary angiogram. (A) Multiplanar reconstruction at the LSV (blue arrow) showing the LMCA (white arrow) arising from it. Note the anomalous RCA (yellow arrow) located between the AO and the PA. (B) Multiplanar reconstruction through the tubular portion of the ascending AO, above the LSV, depicts the high origin of the anomalous RCA with an acute angle take-off along the aortic wall (yellow arrow), coursing anteriorly and to the right with an interarterial course between the AO and the PA. (C) Curved multiplanar reformation of the anomalous RCA depicts that the proximal portion of the RCA is narrowed and has an acute angle take-off from the AO, and follows an interarterial course between the AO and PA (AO, aorta; PA, pulmonary artery; LMCA, left main coronary artery; LSV, left sinus of Valsalva; RCA, right coronary artery).

origin, course and distal portion of these arteries.¹¹ Furthermore, we believe that MDCCT is likely the single best imaging technique today to evaluate patients presenting with cardiovascular symptoms during exercise and routinely in competitive athletes.^{13 18}

Conclusion

We suggest that MDCCT is the diagnostic method of choice when a clinician suspects an unusual mechanism for dyspnoea or the potential for an anomalous coronary artery, owing to the increasing availability of this technology.

Learning points

- ▶ Coronary artery anomalies are rare but represent the second most common cause of sudden cardiac death in young athletes.
- ▶ Sudden cardiac death may be the first manifestation of this heart condition; however, half of the patients may experience cardiovascular symptoms such as palpitations, syncope, chest pain or dyspnoea on exertion, but often routine cardiac tests such as exercising ECG and two-dimensional transthoracic echocardiography are normal.
- ▶ A multidetector coronary CT is currently the best imaging technique to evaluate patients with cardiovascular symptoms triggered by exercise and should be considered routinely in competitive athletes.

Contributors RAD contributed to authorship including the conception of this manuscript, the acquisition, analysis and interpretation of data, as well as drafting the work and giving final approval of the version to be sent for publication. JV contributed to authorship including analysis of data and revising the manuscript critically, as well as giving final approval of the version to be published.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Kardos A, Barbai L, Rudas L, *et al*. Epidemiology of congenital coronary artery anomalies: a coronary arteriography study on a Central European population. *Catheter Cardiovasc Diagn* 1997;42:270–5.
- 2 Yildiz A, Okun B, Peker T, *et al*. Prevalence of coronary artery anomalies in 12,457 adult patients who underwent coronary angiography. *Clin Cardiol* 2010;33:E60–4.
- 3 Van Camp SP, Bloor CM, Mueller FO, *et al*. Nontraumatic sports death in high school and college athletes. *Med Sci Sports Exerc* 1995;27:641–7.
- 4 Maron BJ, Thompson PD, Puffer JC, *et al*. Cardiovascular pre participation screening of competitive athletes: a statement for health professionals from the Sudden Death Committee (Clinical cardiology) and Congenital Cardiac Defects Committee (Cardiovascular disease in the young), American Heart Association. *Circulation* 1996;94:850–6.
- 5 Angelini P. Coronary artery anomalies—current clinical issues: definitions, classification, incidence, clinical relevance, and treatment guidelines. *Tex Heart Inst J* 2002;29:271–8.
- 6 Angelini P. Coronary artery anomalies: an entity in search of an identity. *Circulation* 2007;115:1296–305.
- 7 Sundaram B, Kreml R, Patel S. Imaging of coronary artery anomalies. *Radiol Clin North Am* 2010;48:711–27.
- 8 Alexander RW, Griffith GC. Anomalies of the coronary arteries and their clinical significance. *Circulation* 1956;14:800–5.
- 9 King BD, Ambrose JA, Stein JH, *et al*. Anomalous origin of the right coronary artery from the ascending aorta above the left coronary sinus. *Catheter Cardiovasc Diagn* 1982;8:277–80.
- 10 Young-Hyman PJ, Tommaso CL, Singleton RT. A new double coronary artery anomaly: the right coronary artery originating above the coronary sinus giving off the circumflex artery. *J Am Coll Cardiol* 1984;4:1329–31.
- 11 Angelini P. Novel imaging of coronary artery anomalies to assess their prevalence, the causes of clinical symptoms, and the risk of sudden cardiac death. *Circ Cardiovasc Imaging* 2014;7:747–54.
- 12 Basso C, Maron BJ, Corrado D, *et al*. Clinical profile of congenital coronary artery anomalies with origin from the wrong aortic sinus leading to sudden death in young competitive athletes. *J Am Coll Cardiol* 2000;35:1493–501.
- 13 Ishisone T, Satoh M, Okabayashi H, *et al*. Usefulness of multidetector CT angiography for anomalous origin of coronary artery. *BMJ Case Rep* 2014;2014:pii: bcr2014205180.
- 14 Chaitman BR, Lespérance J, Saltiel J, *et al*. Clinical, angiographic and hemodynamic findings in patients with anomalous origin of the coronary arteries. *Circulation* 1976;53:122–31.
- 15 Engel HJ, Torres C, Page HL Jr. Major variations in anatomical origin of the coronary arteries. *Catheter Cardiovasc Diagn* 1975;1:157–69.
- 16 Liberthson RR, Dinsmore RE, Bharati S, *et al*. Aberrant coronary artery origin from the aorta: diagnosis and clinical significance. *Circulation* 1974;50:774–9.
- 17 Erol C, Seker M. Coronary artery anomalies: the prevalence of origination, course, and termination anomalies of coronary arteries detected by 64-detector computed tomography coronary angiography. *J Comput Assist Tomogr* 2011;35:618–24.
- 18 Graham TP, Driscoll DJ, Gersony WM, *et al*. Task Force 2: congenital heart disease. *J Am Coll Cardiol* 2005;45:1326–33.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow