CORRESPONDENCE

Chronic and Treatment Resistant Depression— Diagnosis and Stepwise Therapy

by Prof. Dr. med. Tom Bschor, Prof. Dr. rer. nat. Dr. med. Dipl.-Biol. Michael Bauer,
PD Dr. med. Mazda Adli in issue 45/2014

Two Additional Aspects

In their article the authors provide a comprehensive and clear description of an evidenced-based stepwise therapy for unipolar depression (1). However, there are two further aspects that should be considered. First, the meta-analysis on antipsychotic augmentation (2) assumes comparable levels of efficacy for aripiprazole, olanzapine, quetiapine, and risperidone. That means that if, for example, the approved quetiapine cannot be used because of metabolic contraindications, aripiprazole would be available as a metabolic more neutral agent for evidence-based—albeit off-label—use. The drug has been approved in the United States for the indication of major depression. In addition, the substance will become available as a generic drug in 2015. The second aspects concerns the use of electroconvulsive therapy (ECT). In the light of the high relapse rates after successful acute treatment, the option of maintenance ECT should be mentioned as well. Although still limited, the available evidence supports a combined use of pharmacotherapy and ECT (3, 4) for relapse prevention; consequently, this approach has also been recommended as a treatment option in various guidelines (5).

DOI: 10.3238/arztebl.2015.0419a

REFERENCES

1. Bschor T, Bauer M, Adli M: Chronic and treatment resistant depression —diagnosis and stepwise therapy. Dtsch Arztebl Int 2014; 111: 766–76.
2. Nelson JC, Papakostas GI: Atypical antipsychotic augmentation in major depressive disorder: a meta-analysis of placebo-controlled randomized trials. Am J Psychiatry 2009; 166: 980–91.
3. Gagné GG Jr, Furman MJ, Carpenter LL, Price LH: Efficacy of continuation ECT and antidepressant drugs compared to long-term antidepressants alone in depressed patients. Am J Psychiatry 2000; 157: 1960–5.
4. Nordenskjöld A, Knorrung L von, Ljung T, et al.: Continuation electroconvulsive therapy with pharmacotherapy versus pharmacotherapy alone for prevention of relapse of depression: a randomized controlled trial. J ECT 2013; 29: 86–92.
5. Zilles D, Wolff-Menzler C, Wiltfang J: [Electroconvulsive therapy for the treatment of major depression]. Nervenarzt. 2015; 86: 549–56

Dr. med. David Zilles, MD
University Medical Center
Dpt. of Psychiatry und Psychotherapy
Göttingen
david.zilles@med.uni-goettingen.de

Conflict of interest statement

Dr. Zilles has received conference fees and reimbursement of travel expenses from Pfizer and Lundbeck.

Bio-psycho-social Aspects of an Overall Treatment Plan

We like to thank the authors for their article which discusses in great detail the pharmacotherapy, psychotherapy and electroconvulsive therapy (ECT) used to treat chronic and treatment-resistant depression (1). What we found missing is the bio-psycho-social aspect of an overall treatment plan. It is our understanding that pharmacotherapy, ECT, psychotherapy, and sociocotherapy should be well-orchestrated parts of an overall strategy, not just isolated measures that patients have to go through in a particular sequence. To achieve complete remission and to prevent chronification, the integrated use of all available resources is required, while taking into account the patients' wishes.

Apart from that we would like to point out that the "high rate of early recurrence" associated with ECT, can be deduced without difficulty from the abrupt termination of a highly effective intervention. In case of antidepressant therapy, successful treatment is continued over several months. Consequently, ECT is increasingly and successfully used as a maintenance therapy. Unfortunately, patients are frequently informed too late, not adequately or not at all because of irrational concerns and the treatment's alleged stigma (2). Since the efficacy of ECT declines with increasing duration and treatment resistance of the disease (3), not starting the treatment in time is detrimental to the patient. For this reason, the psychiatric societies of Germany (DGPPN), Austria (ÖGPP), Switzerland (SGPP), and South Tyrol (SIP) called for the timely and adequate use of ECT in a joint statement issued in 2012 (4). Earlier in 2003, the German Medical Association and, in 2004, the Austrian ÖGPP pleaded for the evidence-based use of ECT. These statements from German-speaking countries are in line with those from international psychiatric societies.

DOI: 10.3238/arztebl.2015.0419b

REFERENCES

1. Bschor T, Bauer M, Adli M: Chronic and treatment resistant depression—diagnosis and stepwise therapy. Dtsch Arztebl Int 2014; 111: 766–76.
2. Pfaff M, Seidl A, Angst K, et al.: [Electroconvulsive therapy as a „last resort“ in the treatment of depression?]. Psychiatr Prax 2013; 40: 385–90.
3. Kho KH, Zwinderman AH, Blansjaar BA: Predictors for the efficacy of electroconvulsive therapy: chart review of a naturalistic study. J Clin Psychiatry 2005; 66: 894–9.
4. Grözinger M, Conca A, DiPauli J, Ramseier F: Electroconvulsive therapy: Psychiatric associations in four countries recommend its timely and appropriate use. http://www.dgppn.de/fileadmin/user_upload/_medien/dokumente/referate/Klinische_Stimulationsverfahren/ECT-Statement_200712.pdf

Prof. Dr. med. Michael Grözinger
Klinik für Psychiatrie, Psychotherapie und Psychosomatik,
Universitätsklinikum Aachen, RWTH Aachen
mgroezinger@ukaachen.de

Prof. Dr. med. Andreas Conca
Psychiatrischer Dienst, Sanitätsbetrieb, Bozen, Italien

Conflict of interest statement

The authors declare that no conflict of interest exists.

List of Drugs That Can Cause Depression

As a neurologist working in a large hospital without a department of psychiatry, I am regularly asked questions about the treatment of depressive disorders when providing consultation services. Compared with reading through the current article (1), it is significantly more taxing to study a guideline of more than 250 pages or its summary with still more than 50 pages. I find the authors' clear statement with regard to the diagnosis and stepwise therapy of chronic and treatment-resistant depression plausible and it will become a clear guiding principle.

Only the list of drugs that can cause depression was somewhat disappointing. Initially, I was enthusiastic about it as it addresses a common problem. When the Department of Internal Medicine calls me for a consultation, I am frequently confronted with patients taking ten different medicines and I am asked whether these are responsible for the patient's depressive mood. This list, unfortunately, seems to have no sensible structure (item 1 would include items 2 to 4; are digitalis preparations no cardiac medications?) And there are mistakes (levodopamine is actually levodopa—by the way, it reportedly has an antidepressant effect, at least in patients with Parkinson's disease; or is the substance blamed for the depression often associated Parkinson's disease?). It would have been very helpful if the authors had highlighted substances from each of the various drug classes that have a comparatively low depression-promoting effect. In clinical practice, it is unfortunately unrealistic to simply discontinue the heart and anti-hypertensive medications. To compile such a table would have been much more difficult—from such an authoritative source, I would have wished to receive this kind of support.

Nevertheless, I am happy to add this article—especially in its electronic version—to my “consultation library”.

DOI: 10.3238/arztebl.2015.0420a

REFERENCES

1. Bschor T, Bauer M, Adli M: Chronic and treatment resistant depression—diagnosis and stepwise therapy. Dtsch Arztebl Int 2014; 111: 766–76.

Dr. med. Stephan Kinze

Klinik für Neurologie, Unfallkrankenhaus Berlin
stephan.kinze@ukb.de

Conflict of interest statement

Dr. Kinze is a lecturer at the Institut für Verhaltenstherapie (IVT) GmbH in Lübben, Germany.

Not Transferable to Outpatient Care

While being revealing and stimulating, unfortunately not every aspect of the article by Bschor et al. (1) can be applied to outpatient care, as our access to laboratory and ECG services is limited. Therefore, therapeutic drug monitoring (TDM) often remains an unfulfilled dream, as does lithium therapy. Ultimately, what is left—besides the early evaluation of an indication for psychotherapy—are the following steps:

Step 1: If patients are rather agitated and present with symptoms such as restlessness, anxiety, loss of appetite, weight loss, aggressive impulses, sleep-maintenance insomnia, and early waking, a presynaptic alpha-2 receptor antagonist (trazodone, mianserin or mirtazapine) should be prescribed; in all other cases, a non-sedating serotonin or norepinephrine reuptake inhibitor (SSRI, NSRI), both in standard doses. However, because of adverse reactions—not because of their efficacy—a change in medication is often required. Many patients treated with mirtazapine become significantly overweight, while those on reuptake inhibitors often develop sexual dysfunction. Only trazodone has neither of these two side effects; however, I think its effect is too weak to warrant its use as a monotherapy. In an outpatient setting, patients have very little tolerance of side effects, especially since they often continue to work or take care of children and the household. Finally, tricyclic antidepressants remain as a third-line treatment; however, they are considered problematic because of their many side effects. In this class, the most suitable agent may be nortriptyline which is only weakly sedative.

Step 2: A reuptake inhibitor should be combined with either quetiapine (beware: weight gain!) or with a presynaptic alpha-2 receptor antagonist (trazodone, mianserin or mirtazapine).

Step 3: I have seen astonishing improvements with tranylcypromine, but its use is challenging. It requires a diet low in tyramine, long safety intervals, and daily blood pressure monitoring; in addition, many medicines are contraindicated.

DOI: 10.3238/arztebl.2015.0420b

REFERENCES

1. Bschor T, Bauer M, Adli M: Chronic and treatment resistant depression—diagnosis and stepwise therapy. Dtsch Arztebl Int 2014; 111: 766–76.

Martin Sansoni

Praxis, Krefeld
sansoni@arcor.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Hormone Replacement in Patients With Depression

This chronic mood disorder is common among the elderly, but the significant increase in its incidence in postmenopausal women and men with the onset of hypogonadism is often overlooked. This stage of life is often associated with cognitive impairments and psychosocial withdrawal; therefore, it pays to address its hormonal aspects.

According to data from the Berlin age studies performed twelve years ago, only 10% of these patients received adequate treatment with antidepressants. In men, one likely reason for this is the development of more pronounced sexual dysfunction while taking antidepressants. Half of the patients taking selective serotonin reuptake inhibitors (SSRI) report a decrease

in libido and the same percentage complains about increased orgasm- and ejaculation-related problems. With this, the association of depression with hypogonadism should become apparent.

According to the Care Report 2013/14 on depression, neurologists observe the steepest rise in care among people aged between 40 and 60 years. This applies equally to both women and men, although the level among the latter is somewhat lower. In these two decades, gonadal function progressively declines.

In 2001, a study (1) demonstrated a significant correlation between androgen receptor structures and depression in men. Low testosterone levels were associated with a 5-fold increased risk of depression. Another study (Shores MM, et al.: The prevalence and incidence of depression in older hypogonadal men. Abstract, presented at the Annual meeting of the American Association of Geriatric Psychiatry 2001) found a 3-fold increase of the risk for depression associated with untreated hypogonadism. Both studies showed a very efficient elimination of symptoms of depression. As the reason for that, direct effects of testosterone on the central nervous system based on monoaminergic neurotransmission-related mechanisms were cited. These direct antidepressant effects were complemented by anabolic effects of testosterone on the energy level, resulting in greater stamina. This approach improved depressive symptoms independent of baseline serum testosterone levels close to the limit for hypogonadism.

A prospective long-term study on male hypogonadism confirmed that the depression rates without testosterone replacement are significantly higher (2).

Conclusion: More attention should be paid to testosterone deficiency in older men with depression.

DOI: 10.3238/arztebl.2015.0420c

REFERENCES

1. Seidman SN, Araujo AB, Roose SP, Mc Kinley JB: Testosterone level, androgen receptor polymorphism and depressive symptoms in middle-aged men. *Biol Psychiatry* 2001; 50: 371–6..
2. Burris AS, Banks SM, Carter CS, Davidson JM, Sherins RJ: A long-term, prospective study of the physiologic and behavioural effects of hormone replacement in untreated hypogonadal men. *J Androl* 1992; 13: 297–304.
3. Bschor T, Bauer M, Adli M: Chronic and treatment resistant depression—diagnosis and stepwise therapy. *Dtsch Arztebl Int* 2014; 111: 766–76.

Prof. Dr. med. Dipl. Psych. J. M. Wenderlein
Universität Ulm
wenderlein@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Repetitive Transcranial Magnetic Stimulation

We think that this review (1) is in actual fact very relevant, but somewhat incomplete as it does not include repetitive transcranial magnetic stimulation (rTMS) as a treatment option. This method, though not approved in Germany, has been cleared for use in the

United States and several European and non-European countries. In Germany, rTMS may be used within the legal framework of an individual treatment attempt (compassionate use). Scientific evidence of its efficacy in chronic and treatment-resistant depression is available, showing both statistically significant and clinically relevant benefits (2) and recently it has been recommended in the guidelines on the therapeutic use of repetitive transcranial magnetic stimulation with a high (A) level of evidence (3).

DOI: 10.3238/arztebl.2015.0421a

REFERENCES

1. Bschor T, Bauer M, Adli M: Chronic and treatment resistant depression—diagnosis and stepwise therapy. *Dtsch Arztebl Int* 2014; 111: 766–76.
2. George MS, Lisanby SH, Avery D, et al.: Daily left prefrontal transcranial magnetic stimulation therapy for major depressive disorder a sham-controlled randomized trial. *Arch Gen Psychiatry* 2010 ;67: 507–16.
3. Lefaucheur JP, André-Obadia N, Antal A, et al.: Guidelines. Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS). *Clin Neurophysiol* 2014; 125: 2150–206.

Dr. med. Stephan. G. Zipper

St. Marien-Krankenhaus Frankfurt/Main,
Frankfurt/Main
zipper@em.uni-frankfurt.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply

We have received an exceptionally large number of positive responses to our CME article (1). We also warmly welcome those critical comments.

We would like to thank Dr. Zilles and Prof. Grözing for their valuable additions with regard to anti-psychotic augmentation, the overall treatment plan, and especially maintenance ECT. We can agree with everything they have stated.

We fully understand Dr. Kinzes' disappointment about the list of drugs that can cause depression which was only published in the supplement to the online version of our article. We had submitted the manuscript without this list, but were then asked by the reviewers and editors to hand one in. Unfortunately, we could not compile a truly reliable list as there is a puzzling abundance of drugs that may be relevant combined with a general lack of data regarding their potential to cause depression. Therefore, as discussed in the article, we are using a different approach: The key indicators that a medicine may have caused depression are the temporal relationship between the start of a pharmacotherapy and the onset of depressive symptoms, which needs to be established as precisely as possible when the patient's history is taken, and, whenever possible, a treatment-free interval or a switch to another medicine that should be attempted. For most antihypertensive drugs, alternatives are available.

Mr. Sansoni sketches a sad picture of outpatient psychiatric management. The aim of our article was to

encourage at least the psychiatrists to use medicines supported by reliable evidence in patients with treatment-resistant depression even if that requires additional effort. In internal medicine, close monitoring of pharmacotherapy is not uncommon, for example in patients receiving anticoagulants or immunosuppressants. Specialists in internal medicine consider this naturally and with competence as their sphere. Likewise, psychiatrists should be proud of using their core competencies. These include lithium treatment, treatment with MAO inhibitors and treatment control based on therapeutic drug monitoring (TDM). Fortunately, we know from our daily clinical experiences that this is the case in the majority of practices. By the way, attempts to generate scientific evidence to support the usefulness of differential indications for pharmacotherapy based on subtypes of depression, as applied by Mr. Sanson, have been unsuccessful.

The sexual side effects of selective serotonin reuptake inhibitors (SSRIs) highlighted by Professor Wenderlein are indeed a common problem in clinical practice. However, these are not explained by hypogonadism, but by serotonergic stimulation of 5-HT₂ receptors. A particularly high prevalence of depression in a specific age group is, of course, not sufficient evidence to demonstrate its relationship with the decline in sex hormone levels. A temporal correlation is by no means a causal relationship, and there are numerous other biological and psychosocial changes that occur in people aged between 50 and 70 years. Naturally, hormone replacement therapy according to endocrinological guidelines has to be initiated in patients with hormone levels that are clearly too low (e.g. in case of overt hypothyroidism).

However, hormone replacement therapy is neither indicated in patients with hormone levels within the borderline range mentioned by Professor Wenderlein nor in patients with a typical age-related drop in hormone levels. It should be added that depression is usually treated by psychiatrists, not neurologists.

We would like to thank Dr. Zipper for highlighting the option of repetitive transcranial magnetic stimulation (rTMS). Its effects in the treatment of depression are, however, rather weak and significantly milder than those of ECT. Individual studies, such as the one cited by Dr. Zipper, are positive; older meta-analyses (2) arrive at very cautious positive conclusions; however, a current systematic review and meta-analysis (3) concludes: "rTMS generates very small improvements when used in the treatment of non-refractory depression. The results in treatment-resistant depression are somewhat better, but still reflect only minimal clinical

improvement." Consequently, the recommendations of the S3-Guideline/National Disease Management Guideline Unipolar Depression (4) are as follows: "The evidence on repetitive transcranial magnetic stimulation (rTMS) [...] currently available is not sufficient to allow recommendations for its general clinical utility and usability." DOI: 10.3238/arztebl.2015.0421b

REFERENCES

1. Bschor T, Bauer M, Adli M: Chronic and treatment resistant depression —diagnosis and stepwise therapy. *Dtsch Arztebl Int* 2014; 111: 766–76.
2. Lam RW, Chan P, Wilkins-Ho M, Yatham LN: Repetitive transcranial magnetic stimulation for treatment-resistant depression: a systematic review and metaanalysis. *Can J Psychiatry* 2008; 53: 621–31.
3. Lepping P, Schönfeldt-Lecuona C, Sambhi RS, et al.: A systematic review of the clinical relevance of repetitive transcranial magnetic stimulation. *Acta Psychiatr Scand* 2014; 130: 326–41.
4. DGPPN, BÄK, KBV, et al. (eds.) on behalf of the Guideline Group on Unipolar Depression: S3-Guideline/National Disease Management Guideline Unipolar Depression. Berlin, Düsseldorf: DGPPN, ÄZQ, AWMF 2009.

Prof. Dr. med. Tom Bschor

Schlosspark-Klinik, Abteilung Psychiatrie, Berlin

bschor@schlosspark-klinik.de

und

Klinik und Poliklinik für Psychiatrie und Psychotherapie, Universitätsklinikum

Carl Gustav Carus an der Technischen Universität Dresden

bschor@mailbox.tu-dresden.de

Prof. Dr. rer. nat. Dr. med. Dipl.-Biol. Michael Bauer

Klinik und Poliklinik für Psychiatrie und Psychotherapie, Universitätsklinikum

Carl Gustav Carus an der Technischen Universität Dresden

PD Dr. med. Mazda Adli

Fliegener Klinik Berlin und Klinik für Psychiatrie und Psychotherapie

Charité – Universitätsmedizin Berlin, Campus Charité Mitte, Berlin

Conflict of interest statement

Prof. Bschor has received reimbursement of meeting participation fees and of travel and accommodation expenses from Lundbeck and AstraZeneca. He has received honoraria for lectures at continuing medical education events from Lilly, BMS, esparma (Aristo), Servier, AstraZeneca, Sanofi, and Lundbeck. Prof. Bschor is a regular member of the Drug Commission of the German Medical Association and also the spokesman for the psychiatric working group within it. He is a board member of IGSLI and President of the Berlin Society for Psychiatry and Neurology.

Prof. Bauer has served as a paid consultant for Alkermes, AstraZeneca, BristolMyersSquibb, Ferrer Internacional, Janssen, Lilly, Lundbeck, Otsuka, Servier, Takeda, and Novartis. He has received honoraria from UNI MED Verlag, Elsevier, Kohlhammer, and Springer for authorship of a publication relating to the topic of this article. He has received reimbursement of meeting participation fees from Servier. He has received payment for the preparation of continuing medical education events from AstraZeneca, BristolMyers Squibb, Ferrer Internacional, Lilly, Lundbeck, Otsuka, Pfizer, and Servier. He has also received payment for carrying out clinical trials on behalf of Lilly, Servier, and Astra-Zeneca.

PD Adli has served as a paid consultant for Lundbeck, esparma (Aristo), and Merz and has received reimbursement of meeting participation fees and of travel and accommodation expenses from Lundbeck and Servier. He has received payment for the preparation of continuing medical education events from Lundbeck, Servier, and esparma (Aristo).