Published in final edited form as:

Neurosci Biobehav Rev. 2015 August ; 55: 36–52. doi:10.1016/j.neubiorev.2015.04.005.

Actions and interactions of estradiol and glucocorticoids in cognition and the brain: Implications for aging women.

Alexandra Ycaza Herrera^a and Mara Mather^a

^aUniversity of Southern California, 3715 McClintock Ave, Los Angeles, Ca 90089, United States

Abstract

Menopause involves dramatic declines in estradiol production and levels. Importantly, estradiol and the class of stress hormones known as glucocorticoids exert countervailing effects throughout the body, with estradiol exerting positive effects on the brain and cognition, glucocorticoids exerting negative effects on the brain and cognition, and estradiol able to mitigate negative effects of glucocorticoids. Although the effects of these hormones in isolation have been extensively studied, the effects of estradiol on the stress response and the neuroprotection offered against glucocorticoid exposure in humans are less well known. Here we review evidence suggesting that estradiol-related protection against glucocorticoids mitigates stress-induced interference with cognitive processes. Animal and human research indicates that estradiol-related mitigation of glucocorticoid damage and interference is one benefit of estradiol supplementation during perimenopause or soon after menopause. The evidence for estradiol-related protection against glucocorticoids suggests that maintaining estradiol levels in post-menopausal women could protect them from stress-induced declines in neural and cognitive integrity.

Keywords

Menopause; Glucocorticoids; Estradiol; Stress; Working Memory; Neurodegeneration; Neuroprotection; Hippocampus; Executive Function

1. Introduction

Women's transition into reproductive senescence is marked by reductions in ovarian function and output, referred to as menopause (Rannevik et al., 1986). However, the dramatic declines in estradiol production and estradiol levels occurring during menopause (Rannevik et al., 1986) may alter more than just reproductive capability. The systems governing stress and fluctuations in female reproductive hormones and the primary hormones of each system (glucocorticoids and estradiol, respectively) intimately interact and

Corresponding Author Ycaza@usc.edu, 213.821.5703.
Mara.mather@usc.edu

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

influence one another. Thus, as one system changes, the nature of the interactions between them may also change.

Research on how glucocorticoids and estradiol affect the brain, body, and cognition spans a number of species, ranging from rodents to humans. However, how glucocorticoids (GC) and estradiol interact to change the effect and action of the other on the brain, body, and cognition in human females is less studied. Due to the dramatic change in the production and release of estradiol in human females with age, understanding whether and how these systems interact to affect women is of importance.

In addition to interacting, glucocorticoids and estradiol also exert opposing effects on various bodily systems, in males and females. For instance, development of metabolic syndrome (Pasquali et al., 2006; for review see, Rosmond, 2005), unhealthy alterations in fat distribution (Rebuffe-Scrive et al., 1992), promotion of hyperglycemia and hyperinsulinemia (McGuinness et al., 1993; Rebuffe-Scrive et al., 1992), promotion of bone resorption (O'Brien et al., 2004), and maintenance of bone degrading osteoclasts (Jia et al., 2006) have been associated with long-term GC exposure. In contrast, opposite patterns of less unhealthy fat distribution (Green et al., 2004; Musatov et al., 2007), lesser occurrence of hyperglycemia and hyperinsulinemia (Krotkiewski et al., 1983; Musatov et al., 2007), and promotion and maintenance of bone mineral density (Delmas et al., 1997; Felson et al., 1993; Sowers et al., 1998), have been linked to estradiol, the primary estrogen in females (See Table 1).

Similar contrasting effects of GCs and estradiol occur in neural tissue and cognition, where chronic or extreme GC exposure leads to dendritic retraction, and neuropil and neuron damage or loss (Behl et al., 1997; Filipovic et al., 2013; Hiller et al., 2013; MacPherson et al., 2005; Magarinos et al., 2011; Stein-Behrens et al., 1992; for review see, Tata and Anderson, 2010; see also, Tombaugh et al., 1992; Tynan et al., 2013; Tynan et al., 2010; Woolley et al., 1990b), and impairments in cognitive performance (Alexander et al., 2007; Duncko et al., 2009; Elzinga and Roelofs, 2005; Luethi et al., 2009; Oei et al., 2006; Schoofs et al., 2009; Young et al., 1999). In contrast, estradiol promotes neural growth and protection (Brinton et al., 2000; Chen et al., 2006; Gerstner et al., 2007; Hao, 2006; Saravia et al., 2007) and improvement of cognitive function (Baker et al., 2012; Krug et al., 2006; Shaywitz et al., 2003; Velázquez-Zamora et al., 2012; Wolf et al., 1999)¹.

Due to the fluctuations and eventual loss of estradiol throughout the adult lifespan of human females, it is important to understand the implications of the opposing effects of stress and estradiol on the brain and cognition. Much of the literature reviewed here will span from basic research findings to findings in human research, and explore patterns of these hormone effects in males and females. We will focus on the negative effects of stress on neural tissue and cognition in both males and females, and discuss whether the opposing effects of estradiol in the same domains may offer some protection against these effects. We propose that estradiol loss after menopause leaves women more vulnerable to developing

¹GCs also are necessary for normal neuronal functioning (Nadeau and Rivest, 2003), and are beneficial for memory of emotional stimuli or events (e.g., Buchanan and Lovaglio, 2001). Similarly, estradiol can damage already compromised neural tissue (Chen et al., 2006), and is associated with worse emotional memory (for review see, Sakaki and Mather, 2012).

dysfunctional stress responses and therefore, larger effects of stress on cognition. We also argue that the loss of estradiol in midlife leaves peri- and post-menopausal women more vulnerable to the negative effects of stress hormone exposure, accelerating age-related declines in neural and cognitive integrity.

2. Interactions of the stress and estradiol systems

2.1 The “Stress System”

The hypothalamic-pituitary-adrenal (HPA) axis governs response to stressors. Emotional or physiological challenges cause the paraventricular nucleus of the hypothalamus to release corticotropin-releasing hormone. The portal system carries the neuropeptide to the anterior pituitary, causing release of adrenocorticotrophic hormone. Adrenocorticotrophic hormone travels via the bloodstream to the adrenal glands, causing release of GCs from the adrenal cortex (the outer layers of the adrenal gland) and the catecholamines epinephrine and norepinephrine from the adrenal medulla (for review see, Lupien et al., 2009). Due to our interest in the effects of GCs, and interactions between GCs and estradiol, we will focus on this portion of the stress response.

During the stress response, GCs assist in the fight-or-flight response by providing immediate energy to the body via assistance converting protein to glucose, fat to usable energy, and shunting blood flow from immediately nonessential systems, such as the gut, to skeletal muscles for energy. Cortisol, the primary GC in humans, also is responsible for terminating the HPA response to a stressor via negative feedback. Glucocorticoids initiate shutdown of the HPA response by exerting direct inhibition of the hypothalamus and pituitary gland, as well as recruiting additional brain regions, particularly the prefrontal cortex and hippocampus, to send inhibitory signals.

2.2 The “Estradiol System”

Unlike the HPA axis, the hypothalamic-pituitary-gonadal (HPG) axis terminates at different targets for males and females, the testes or ovaries, respectively. Here we will focus on the female HPG axis, as we are concerned with estradiol levels pre and post menopause in women.

Like the HPA axis, the HPG axis originates in the hypothalamus. Several subnuclei, including the arcuate nucleus and preoptic area, contain neurons that produce and release gonadotropin-releasing hormone. The portal system delivers gonadotropin-releasing hormone to the anterior pituitary gland, initiating the release of luteinizing hormone and follicle stimulating hormone. Luteinizing hormone and follicle stimulating hormone then act on the ovaries to cause release of estrogens and progestins. Of the three major forms of estrogens (estrone, estradiol, and estriol), estradiol (E2) is the estrogen predominantly responsible for estrogenic effects observed in the brain (McClure et al., 2013; for review see, Purves et al., 2004).

Estrogens both stimulate and inhibit the HPG axis (for reviews see, Herbison, 1998; Jones, 2009). Throughout most of the monthly menstrual cycle, E2 inhibits release of luteinizing hormone and follicle stimulating hormone by decreasing the sensitivity of neurons

regulating these hormones to the effects of gonadotropin releasing hormone. However, large increases in E2 lasting for 2 or more days, such as the increase observed just before ovulation, increases the sensitivity of neurons regulating luteinizing hormone and follicle stimulating hormone to the effects of gonadotropin releasing hormone, stimulating release of the two hormones. Reductions in E2 after this surge return gonadotropin-releasing hormone, luteinizing hormone, and follicle stimulating hormone to more common moderate-to-low levels (for review see, Jones, 2009).

2.3 Interactions of the Stress and Estradiol Systems

It has long been known that the stress and E2 systems influence each other's function (e.g., Selye, 1939) and the complex interactions of these two systems span beyond what is discussed in this brief section. For instance, there is a large literature examining how interactions between these systems likely affect the development of emotional pathologies, such as depression and post-traumatic stress disorder, predominately in women (for review see, Becker et al., 2007). Additionally, the interactions begin early in development (for reviews see, Dunn et al., 2011; Goel and Bale, 2009; see also, Mueller and Bale, 2007; Mueller and Bale, 2008) influencing the effects of stress throughout the lifespan. Although HPA and HPG interactions during development can alter stress responsiveness later in life, this review focuses on how normal age-related changes in stress-estradiol system interactions may affect women during and after the menopause transition.

One of the first interactions noted between the two systems was the ability of stress to interfere with normal reproductive function (Selye, 1939). Stress-induced interference of HPG function appears to rely on GCs (for review see, Breen and Karsch, 2006), which reduce sensitivity of the ovaries to luteinizing hormone, decrease aromatase activity in ovarian cells (Hsueh and Erickson, 1978), and block ovulation (for review see, Rivier and Rivest, 1991). Furthermore, HPA axis activation is capable of interfering with normal HPG axis activity at every level of the HPG axis, including the hypothalamus and pituitary (for review see, Tilbrook et al., 2000), and the ovaries (for review see, Tetsuka, 2007), where GCs can stop follicle maturation. How and at what level the HPA can disrupt normal HPG function depends on whether stress exposure is acute or chronic (for example, Gore et al., 2006).

On the other side of the interaction, E2 can both reduce and promote HPA function. The majority of circulating GC is often bound to corticosteroid binding globulin, making only 5-10% of the hormone available to enter and act on cells (Burke and Roulet, 1970). One way E2 reduces HPA function is by upregulating corticosteroid binding globulin which decreases exposure to released GCs (for review see, Brien, 1981). On the other hand, E2 upregulates corticotropin releasing hormone expression in the hypothalamus (Seale et al., 2004a; Seale et al., 2004b), which could result in increased HPA function by stimulating release of the hormone that initiates the HPA response. The bi-directionality of HPA-HPG interactions makes it important to understand whether, and how, these interactions may change when one system is dramatically altered, such as when women go through the menopause transition where E2 levels rapidly and severely decrease.

3. Effects of Glucocorticoids and Estradiol on Cognitive Processes

The following sections review effects of GC on cognition and then contrast those with E2 effects on cognition.

3.1 Glucocorticoids can Facilitate and Impair Memory Processes

Exposure to long-term or extreme stress can impair learning and memory. However, stress exposure does not always lead to cognitive impairment. Different factors contribute to whether stress will lead to impairment, such as the type of learning and memory being examined and the amount of stress applied. With regard to the amount of stress applied, effects of stress on learning and memory sometimes follow an inverted-U shaped curve, with too little or too much being related to poor performance and a moderate amount related to peak performance in a variety of tasks from shock avoidance to spatial learning and working memory (Mateo, 2008; Salehi et al., 2010; Yerkes and Dodson, 1908), an effect which also has been observed for measures of neural plasticity (Diamond et al., 1992). Further, what is considered an optimal amount of stress can shift depending on task difficulty (Anderson, 1994; Dodson, 1915; Salehi et al., 2010; for review see, Shors, 2004).

Stress can also be beneficial in learning about emotional stimuli. For instance, administration of hydrocortisone (Buchanan and Lovallo, 2001) and stress induction (Payne et al., 2007; Smeets et al., 2006) prior to encoding improved memory for emotional stimuli while having no effect or even impairing memory for neutral stimuli. However, memory for emotional items is not impervious to stress. Stress applied before retrieval, rather than before encoding, impaired recall of emotional and neutral words (Kuhlmann et al., 2005). Thus, while stress is beneficial to learning and memory in some instances, there are still limitations to when stress can be beneficial, even in these situations.

3.2 Glucocorticoids Impair Memory Processes in Animals

With regard to instances of chronic or excessive stress, the animal and human literature is rife with reports of GC- or stress-induced cognitive impairment (for review see, Holmes and Wellman, 2009). In male rodents, eight weeks of daily corticosterone injections resulted in more working memory errors during a Y-maze task (spending equal amounts of time in a familiar and novel arm, instead of spending more time exploring the novel arm) compared with animals receiving a sesame oil control injection (Coburn-Litvak et al., 2003), while nine weeks of corticosterone treatment induced more working memory errors during a radial-arm-maze task (reentering already traversed arms; Arbel et al., 1994), and three weeks of daily corticosterone injections impaired male rodents' reference memory on another measure of working memory, the Barnes maze (Coburn-Litvak et al., 2003).

Similar to effects of exogenous GC administration, cognitive impairments also are observed in animals repeatedly exposed to psychological and physical stressors. For instance, male rats exposed to a cat for six hours then housed with different combinations of 3-4 rats daily over a course of five weeks showed impaired working memory on a radial arm water maze (Park et al., 2001). Males exposed to an unfamiliar environment during a testing delay period for twenty-one days made more working memory errors in a radial arm maze

compared with animals housed in their home cages during the delay period (Diamond et al., 1996). Likewise, male rats exposed to restraint stress for 6 hours/day for 21 days exhibited worse working memory in a Y-maze (Kleen et al., 2006), compared with rats not exposed to restraint stress, with similar results observed in animals trained on a delayed alternation task in a T-maze (Lee and Goto, 2015; See Table 2 for effects of working memory). Shorter stress exposures also can induce cognitive impairments in animals. Four weeks of unpredictable stress resulted in impaired acquisition, working, and reference memory of a modified Morris water maze task and impaired reversal learning in male rodents. The study also found that shorter regimens of six days of unpredictable stress negatively impacted reference memory and reversal learning, while three days of unpredictable stress induced deficits in reference memory on the task (Cerqueira et al., 2007). Similarly, corticosterone administration a mere thirty minutes prior to testing resulted in impaired performance on a delayed alternation task in T-maze in male rodents (Roozendaal, 2004), while fifteen minutes of tail pinch stress increased error rates during a delayed win-shift task in a radial arm maze, an effect blocked in animals given a GC antagonist (Butts et al., 2011).

Interestingly, depending on the type of memory under investigation, these effects of stress on cognition appear more consistently in male animals than female animals (Beck and Luine, 2002; for review see, Beck and Luine, 2010; see also, Bowman, 2003; Bowman et al., 2009; for review see, Luine, 2002). Despite female rodents showing larger HPA responses to a stressor compared with males, E2 may be a possible protective agent in the face of chronic stress, as female animals² habituate to chronic stressors sooner than do males (Bowman, 2003; Bowman et al., 2002; Bowman et al., 2009).

3.3 Glucocorticoids Impair Working Memory and Executive Function in Humans

Although much of the animal literature focuses on the effects of chronic GC or stress exposure on cognitive processes, most laboratory research with humans focuses on the effects of *acute* stress on cognitive function. Nevertheless, the impairing effects observed in animals also are seen with acute pharmacological, psychological, and physical stress administration in humans. For instance, men treated with acute intravenous (Young et al., 1999) and chronic oral (Lupien et al., 1999) hydrocortisone showed greater working memory errors during a visuospatial sketchpad task (Young et al., 1999) and the Sternberg item recognition task (Lupien et al., 1999), as well as higher error rates in a paired associates task (Young et al., 1999).

With respect to laboratory stressors, one frequently used acute psychological stressor is the Trier Social Stress Test (TSST). The TSST reliably elevates participants' cortisol levels by requiring them to give a speech and perform mental arithmetic in front of an audience (Elzinga and Roelofs, 2005; Oei et al., 2006; Schoofs et al., 2008; Wolf et al., 2009). Completing the TSST resulted in increased reaction times and impairments on tests of working memory given 10-45 minutes later, such as the reading span task (Luethi et al., 2009), the Sternberg item recognition task (Oei et al., 2006), and the N-back task (Schoofs et

²“Female animals” here refers to intact females. In many studies, female animals are ovariectomized (ovaries removed) in order to halt endogenous production of sex hormones and allow for experimenters to control levels of ovarian hormones, such as estradiol. If female animals remain intact, then they are still endogenously producing sex hormones, such as estradiol.

al., 2008) in men, and the digit span task (Elzinga and Roelofs, 2005) in men and women. The TSST also impairs performance on tests of executive function, or cognitive flexibility. When tested during execution of the social stressor, men and women made fewer correct responses during the compound remote associates test and exhibited increased latencies to complete anagrams (Alexander et al., 2007), and men exhibited impaired performance on a go no-go task (Scholz et al., 2009). Impaired performance on a mental arithmetic task was also impaired in men exhibiting the highest cortisol responses during the task or after a giving a surprise speech (Al'Absi et al., 2002; See Table 3 for effects on executive function and other cognitive processes).

Another commonly employed acute laboratory stressor is the cold pressor task (CPT). The CPT is a physical stressor that reliably induces a stress response by requiring participants to hold one of their hands in ice water (Bullinger et al., 1984; Edelson and Robertson, 1986; Lighthall et al., 2009; Lighthall et al., 2011; Mather et al., 2010). Like the TSST, men showed impaired performance on working memory measures such as the operation span task and digit span backward (Schoofs et al., 2009), while a sample of men and women displayed impairment on the Sternberg item recognition task (Duncko et al., 2009) following exposure to the CPT.

The ecological validity of laboratory studies is difficult to assess, however, some work indicates that stressors encountered outside of the laboratory can interfere with cognition in both sexes. For instance, a meta-analysis revealed that intermediate intensity exercise resulted in declines in a range of working memory tasks (McMorris et al., 2011). In other examples, attention-shifting was impaired in students preparing for a major academic exam, and those students' reports of chronic stress on the Perceived Stress Scale predicted their performance on the attention-shift task (Liston et al., 2009). Those experiencing anticipatory stress after being told they would need to give a surprise speech exhibited impaired decision making on the game of dice task (Starcke et al., 2008), while another cohort with high perceived chronic stress also showed poorer working memory than those rating lower perceived chronic stress (Öhman et al., 2007). Even within a population of patients suffering from unipolar major depression, morning cortisol levels were negatively correlated with performance on the Wisconsin Card Sorting Task, a measure of executive function (Egeland et al., 2005).

In sum, pharmacological, physical, psychological, and non-laboratory stressors can interfere with normal, or optimal, cognitive function in animal models and humans. In particular, the evidence suggests that chronic or excessive GC levels impair cognition in tasks requiring executive resources, such as working memory.

3.4 Estradiol can Facilitate and Impair Memory Processes

Just as seen with GCs, E2 can exert both negative and positive effects on cognitive function. This dichotomy is highlighted in reports of E2 treatment on cognition in post-menopausal women, with some reporting negative effects (Espeland et al., 2004; Mulnard et al., 2000; Rapp et al., 2003; Resnick et al., 2006; Shumaker et al., 2003) and others reporting benefits (Baker et al., 2012; Duff and Hampson, 2000; Maki et al., 2001; Miller et al., 2002; Smith et al., 2001; Wolf and Kirschbaum, 2002; Wolf et al., 1999), which receive support from basic

science reports of estrogenic fortification of the central nervous system (Brinton et al., 2000; Chen et al., 2006; Hosoda et al., 2001; Pike, 1999). To reconcile these mixed reports, Brinton (2005) proposed a healthy cell bias of estrogen action, stating that E2 would be advantageous to a population of healthy cells, but detrimental to a population of declining or already injured cells. To test this hypothesis, Brinton and colleagues compared treatment and prevention models *in vitro*. In treatment models, E2 is applied to neurons either at the time of, or after, application of the deleterious agent (e.g. beta-amyloid; A β), whereas prevention models pretreat cells with E2 *before* exposure to the agent. Consistent with the hypothesis, neurons *pretreated* with E2 prior to A β protein exposure showed significantly less cell damage and death compared with cultures exposed to A β alone. Whereas neurons simultaneously treated with E2 and A β protein did not receive any estrogenic protection, nor did neurons exposed to the toxic protein for 1 to 2 days prior to E2 exposure. However, introduction of E2 to the cultures after 5 days of A β exposure resulted in greater cell death than the A β -alone cultures (Chen et al., 2006). This pattern suggests that the reports of negative estrogenic effects may be related to the sample population selected; particularly in studies with women many years past menopause (Espeland et al., 2004; Rapp et al., 2003; Resnick et al., 2006; Shumaker et al., 2003), and those using women with documented neural decline (Mulnard et al., 2000)³.

3.5 Estradiol Facilitates Memory Processes in Animals

In contrast to these negative reports of E2 on cognition, E2 often facilitates learning and memory in animals. Ovariectomized (OVX) female animals made fewer working memory errors during a radial arm maze if treated with E2 after OVX compared with OVX females not treated with E2 (Bimonte and Denenberg, 1999; Daniel et al., 1997; Fader et al., 1999). Likewise, females receiving only a sham surgery (ovaries left intact and therefore still producing E2) exhibited a greater number of consecutive correct choices on a radial arm maze, compared with OVX-alone females (Wilson et al., 1999; See Table 2 for effects on working memory).

Benefits of E2 also have been observed in other tasks. Compared with OVX-alone females, ovariectomized animals treated with estradiol benzoate displayed better retention for the location of a hidden platform in a delayed match-to-place task across 10-, 30-, and 100-minute delay intervals (Sandstrom and Williams, 2001). Likewise, E2 treated animals also required less time to reach learning criteria of a delayed-match-to-position task than OVX-alone animals (Gibbs, 1999). The same pattern was observed in an object recognition task. This version of the object recognition task exposed animals to a total of 4 novel objects. Animals were first allowed to explore a pair of 2 novel objects, after some delay they were exposed to another 2 objects (one object from pair #1 and the third novel object); after a second delay period the animals were exposed to 2 additional objects (one being the second object from pair #1 and the fourth novel object). The animal is said to recognize an object if it spends less time exploring that object compared with time spent exploring the novel

³Negative effects of E2 also have been reported in younger adult women, where E2 levels are negatively associated with memory for emotional items. Attempting to explain this negative association, Sakaki and Mather (2012) proposed that the reported decreased neural activation of the amygdala during *high* E2 phases of the menstrual cycle, relative to low E2 phases (Goldstein et al., 2010; Goldstein et al., 2005), is responsible for the negative relationship between E2 levels and emotional memory.

object. Ovariectomy followed by 21 days of E2 treatment enhanced retention for the first-tested object after a 3-hour delay and of the second-tested object after a 6-hour delay, compared with non-E2 treated mice (Vaucher et al., 2002; See Table 3 for effects on other cognitive processes).

3.6 Estradiol Facilitates Working Memory and Executive Function in Peri- and Post-Menopausal Women

Akin to the animal studies comparing performance between OVX and OVX + E2 females, the largest effects of E2 are observed in post-menopausal women taking some form of E2 replacement as they no longer produce E2 at the same levels as younger women.

Few studies have examined the cognitive effects of naturally declining E2 levels during menopause in the absence of hormone replacement. In one such study, women within five years of menopause and beyond five years of menopause did not differ on measures of sustained attention, category generation, or episodic memory. However, higher E2 levels in women within five years of menopause were associated with better performance in measures of mental flexibility and planning (i.e., executive function) compared with the steady low E2 levels in the women beyond five years of menopause (Elsabagh et al., 2007), an effect maintained after controlling for age and IQ of women within and beyond 5 years of menopause. Endogenous levels of E2 in postmenopausal women not taking hormone supplements also were positively correlated with performance on the Stroop task, another measure of executive function (Wolf and Kirschbaum, 2002; See Table 3 for effects on executive functions).

When women received 2 weeks of transdermal E2 treatment, those experiencing higher endogenous E2 levels in response to the dosage performed better on the delayed recall portion of a verbal memory test than women experiencing lower dose-induced endogenous E2 levels (Wolf et al., 1999). Women randomly assigned to transdermal hormone replacement experienced improved performance on the Stroop task, spatial working memory (Baker et al., 2012; Krug et al., 2006), digit ordering (Krug et al., 2006), and delayed recall (Baker et al., 2012). Similarly, women already using hormone replacement also made fewer errors on non-spatial working memory (sequential memory of an unfamiliar story, digit ordering and digit span backward) and spatial working memory tasks (Duff and Hampson, 2000), performed better on measures of verbal fluency and working memory (Miller et al., 2002), and displayed better performance on a non-verbal memory task (Smith et al., 2001) compared with women not taking any hormone supplements (See Table 2 for effects on working memory).

However, there also are reports of no effect (Maki et al., 2001; Pefanco et al., 2007; Smith et al., 2001; Yaffe et al., 2006), or negative effects of various E2 supplementation regimens (Espeland et al., 2004; Mulnard et al., 2000; Rapp et al., 2003; Resnick et al., 2006; Shumaker et al., 2003). The tasks in these studies assessed similar cognitive domains as the executive tasks discussed above which benefited from E2 treatment, as well as tasks not involving executive function, such as attention and short-term episodic memory (e.g., Newhouse et al., 2010). The most widely reported negative findings come from the Women's Health Initiative (WHI) studies. At the time of inception, the Women's Health

Initiative Memory Study (WHIMS) and Women's Health Initiative Study of Cognitive Aging (WHISCA) were the largest randomized studies testing claims that post-menopausal hormone treatment maintained cognitive function otherwise observed to decline after menopause. However, the WHIMS and WHISCA studies found that conjugated equine estrogens negatively affected performance on various cognitive measures, such as the California Verbal Learning Test (a measure of verbal retention and recall; Resnick et al., 2006) and the Modified Mini Mental State Exam (a measure of global cognitive function; Espeland et al., 2004; Rapp et al., 2003), as well as increasing the risk of developing dementia (Shumaker et al., 2003). The Alzheimer's Disease Cooperative Study, a randomized, double-blind, placebo-controlled, clinical trial, found similar negative effects of E2 treatment, reporting a faster progression of Alzheimer's in older women with preexisting mild to moderate Alzheimer's disease who were on E2 supplementation (Mulnard et al., 2000).

One key difference between the results of the Alzheimer's Disease Cooperative Study and WHI studies and the positive effects reported above is that, on average, the women reported to experience positive effects began treatment closer to the age of menopause than the women in the Alzheimer's Disease Cooperative Study and WHI studies. For example, E2-associated enhancements in working memory were observed in women who began hormone treatment during peri-menopause or soon after menopause (Duff and Hampson, 2000); those showing benefits in semantic fluency, attention, and working memory were on average 63 years old but had been using E2 supplements for an average of 12 years (Miller et al., 2002), and those displaying better performance on a non-verbal memory task initiated E2 replacement within two years of menopause (Smith et al., 2001).

The above-described findings of E2-induced cognitive impairment were also surprising based on basic science reports of E2-related protection of the central nervous system (Brinton et al., 2000; Chen et al., 2006; Hosoda et al., 2001; Pike, 1999). Together with the proposal for a critical window for E2 treatment initiation (for review see, Maki, 2013; see also, Resnick and Henderson, 2002), Brinton's (2005) healthy-cell bias of E2 action (described above) can account for the disparity between basic science experiments and human studies. Findings that introduction of E2 after 5 days of A β exposure exacerbates A β -induced damage (Chen et al., 2006) suggest that E2 treatment would exacerbate existing damage in the neural tissue of participants in both studies, particularly in those women with preexisting Alzheimer's disease, presumably leading to declines in cognitive capacity.

4. Glucocorticoids and Estradiol Differentially Affect Neuronal Viability and Action

Consistent with the pattern discussed for cognition, exposure to GCs, either via direct administration or stress exposure, typically results in negative effects such as cell damage or death (Behl et al., 1997; Gerlach and McEwen, 1972; MacPherson et al., 2005; McEwen et al., 1968; Stein-Behrens et al., 1992; Tombaugh et al., 1992; Tynan et al., 2010; Woolley et al., 1990b) whereas E2 exposure results in positive effects such as cell growth and protection (Brinton et al., 2000; Chen et al., 2006; Gerstner et al., 2007; Hao et al., 2007;

Saravia et al., 2007; See Table 3)⁴. The following sections review these effects of GC on neuronal viability and action, and then contrast those with the effects of E2.

4.1 Glucocorticoids Damage Neurons in the Hippocampus and Prefrontal Cortex of Animals

The hippocampus and prefrontal cortex are the neural regions most sensitive to the deleterious effects of GCs and stress, as might be expected based on the effects of GCs on cognition and these regions' involvement in short-term memory (Alonso et al., 2002; Cabeza et al., 2002), long-term memory (Alonso et al., 2002), and working memory (Cabeza et al., 2002; Curtis and D'Esposito, 2003; for review see, Laroche et al., 2000).

The hippocampus contains a high concentration of GC receptors (Gerlach and McEwen, 1972; McEwen et al., 1968) and is thus highly sensitive to the negative effects of GCs. Exposing the hippocampus to GCs exacerbates damage induced by other neuronal insults, such as A β and glutamate (Behl, 1998; for reviews see, Sapolsky, 1990; Sapolsky, 1999, 2000). For example, exposure of embryonic hippocampal neuron cultures to A β or glutamate experienced greater rates of cell death if pretreated with dexamethasone (a synthetic GC) or corticosterone (the primary GC in rodents) for 24 hours than cultures not pretreated with GC (Behl et al., 1997). Fetal hippocampal cultures preexposed to corticosterone followed by exposure to hypoxic or hypoglycemic environments (Tombaugh et al., 1992) and adult males preexposed to corticosterone followed by kainic acid exposure (Stein-Behrens et al., 1992) also experienced exacerbated damage (See Table 4 for effects on neuronal survival in concert with other insults).

In addition to exacerbating damage by other insults, GC exposure alone, induced via chronic or severe acute stress exposure, has been shown to upregulate pro-inflammatory cytokines throughout the brain and damage hippocampal neurons. Although normal circulating GC levels have inherently protective anti-inflammatory effects (for reviews see, Chrousos, 2009; Elenkov and Chrousos, 2002; see also, Nadeau and Rivest, 2003), male rats exposed to either severe acute stress (inescapable tail- or foot-shock) or chronic stress (restraint) exhibited increased expression of pro-inflammatory immune-response cytokines such as interleukin-1H (Blandino Jr et al., 2006), interleukin-1 β (O'Connor et al., 2003) and activated microglia (Tynan et al., 2010), throughout various brain regions. Increases in microglial markers in response to this type of chronic GC exposure indicate increased inflammation to stressors as part of an immune response (Frank et al., 2007).

Chronic GC and stress exposure also affect neuronal morphology and function in the brain. The hippocampal CA3 region appears particularly susceptible. Male rats experience decreased dendrite length and branch points after twenty-one days of corticosterone injections (Woolley et al., 1990b) and apoptosis and cell damage after dexamethasone administration (Haynes et al., 2003). Male and OVX female rats also experience decreased apical dendrite length and branch points in the region after three weeks of daily restraint

⁴Although chronic or excessive GC exposure is detrimental, it must be noted that GCs are necessary and even beneficial for neuronal proliferation and function when present in basal physiological levels or administered in low doses (Anacker et al., 2012; for review see, de Kloet et al., 1999; see also, Nadeau and Rivest, 2003).

stress (Galea et al., 1997; McLaughlin et al., 2010; Watanabe et al., 1992), and male vervet monkeys exposed to chronic social stress exhibited fewer CA3 hippocampal pyramidal cells plus atrophy of dendritic branches of the surviving cells compared with non-stressed cohorts (Uno et al., 1989; See Table 5 for effects on brain tissue).

The same effects have been observed in other regions such as the striatum (Haynes et al., 2003) and layers II and III of the prefrontal cortex (Brown et al., 2005; Cook and Wellman, 2004; Wellman, 2001), where repeated acute and chronic restraint stress decreased apical dendrite branch number and length in male rats. Male rats also displayed decreased volume of layers I and II of medial prefrontal cortex after chronic unpredictable stress, presumably a result of dendritic retraction (Cerqueira et al., 2007). This effect on prefrontal cortical tissue is at least partially a result of GC action, as infusion of the GC receptor agonist RU 28362 into the region interfered with T-maze performance similarly to systemic injection of corticosterone in male rodents (Roozendaal, 2004), but infusion of the GC receptor antagonist RU 38486 into the region blocked stress-induced cognitive interference during a delayed win-shift task in males (Butts et al., 2011). That the medial prefrontal cortex displays the same degree of restructuring after repeated acute stress and not just chronic stress, suggests that this region is highly sensitive to stressful events (for review see, Holmes and Wellman, 2009) and perhaps even more sensitive than the hippocampus. Greater sensitivity of the prefrontal cortex may explain why cognition related to executive function appears to be more sensitive to the effects of stress or stress hormone exposure than other types of cognition, and why acute stressors in the laboratory can successfully interfere with these cognitive processes in humans.

4.2 Glucocorticoids and Stress Affect Brain Activity in Humans

Just as GC and stress can alter neural processes in animals, neuroimaging work in humans also indicates that pharmacological induction of stress and exposure to acute stressors results in marked alterations of brain activity in the aforementioned regions. Typically, men and women activate the dorsolateral and ventrolateral prefrontal cortex during working memory tasks (Cabeza et al., 2002; Owen et al., 2005; Porcelli et al., 2008; Qin et al., 2009). The hippocampus, on the other hand, is significantly co-activated with the default mode network (Greicius et al., 2004), a network of regions active during internally focused tasks that deactivates during tasks requiring external focus, such as working memory tasks (Esposito et al., 2006; Qin et al., 2009). In line with its relation to the default mode network, men experiencing deactivation of the hippocampus during a working memory task exhibited enhanced working memory performance (Cousijn et al., 2012). However, administration of hydrocortisone results in attenuated activation in ventrolateral prefrontal cortex and attenuated deactivation in the hippocampus during performance of a high-cognitive-load n-back task in men and women (Symonds et al., 2012). Similarly, women who viewed aversive video clips prior to performing an n-back task experienced attenuated activation of the dorsolateral prefrontal cortex and attenuated deactivation of the hippocampus during performance of the working memory task (Qin et al., 2009; See Table 6 for effects on brain activation).

These studies show that GC exposure alters brain activity in prefrontal cortex and hippocampal regions during working memory tasks, indicating that GC impairs executive control mechanisms critical for maintaining and accessing information in memory via altering brain activation patterns observed in non-stressful situations. Taken further, the results in animals suggests that prolonged exposure to stress may result in more than just altered brain activation, but to more long-lasting consequences such as neural damage.

4.3 Estradiol Protects Neurons in the Hippocampus and Prefrontal Cortex of Animals

Although E2 can increase tissue vulnerability to a population of already damaged cells, in the absence of preexisting toxicity and damage E2 promotes neurogenesis and is neuroprotective. For instance, fetal primary hippocampal neurons exposed to conjugated-equine estrogens (Brinton et al., 2000) or E2 (Chen et al., 2006) showed increased number, and length, of neurites, as well as increased secondary branching and branch length when compared with cultures free of any estrogenic compounds (See Table 4 for effects on neuronal survival in concert with other insults).

E2 also promotes neurogenesis in mature animals (See Table 5 for effects on neuronal morphology). E2 treatment increased cell proliferation and decreased cell death in the dentate gyrus region of the hippocampus of middle-aged male mice (Saravia et al., 2007), essentially slowing down the rate of age-related declines in cell proliferation of this hippocampal region. Estrogenic promotion of cell proliferation in the hippocampus could aid in the maintenance of hippocampal-dependent learning and memory processes. For instance, in adult animals mature granule cells appear to only respond to specific stimuli or stimuli ranges, while new granule cells have a lower threshold for response to a wider range of stimuli (Marin-Burgin et al., 2012) possibly allowing them to better form new connections as learning occurs.

In addition to promoting neurogenesis and maintaining neural integrity, E2 also is neuroprotective. *In vitro* studies find that pretreatment of hippocampal fetal neurons with either E2 (Chen et al., 2006) or conjugated-equine estrogens (Brinton et al., 2000) blocked the neurodegeneration caused by A β exposure, decreased the degree of apoptosis, and attenuated the A β -induced decrease in ATP levels. Similar estrogenic protection has been reported against hydrogen peroxide challenge and glutamate excitotoxicity (Brinton et al., 2000). Estradiol also protects oligodendrocytes, the glial cells responsible for myelinating axons within the central nervous system, from hyperoxic insults in rat pups, suggesting E2 can act as a line of defense against insults threatening proper myelination during development (Gerstner et al., 2007).

In vivo studies report similar patterns of neuroprotection by E2 in hippocampus and entorhinal cortex, as well as neurogenesis in prefrontal cortex. Estradiol pretreatment blocked the marked dose-dependent neuron loss in the entorhinal cortex and hippocampus induced by kainic acid in OVX-alone females (Hoffman et al., 2003). In males, intracerebroventricular injection of E2 ten minutes prior to quinolinic acid injection protected the CA1 region of the hippocampus (Kuroki et al., 2001). Estradiol pretreatment also protected male mice from the stereotypical motor deficits accompanying methylmercury exposure and reduced the amount of lipid peroxidation in the cerebellum

(Malagutti et al., 2009), suggested to be a result of E2's protection of the antioxidant glutathione (Malagutti et al., 2009). Estradiol may also protect the brain from age-related declines; ovariectomy followed by cyclic E2 replacement increased dendrite spine density in layer III of dorsolateral prefrontal cortex in young-adult female rhesus monkeys and restored density in aged female monkeys, compared with female rhesus monkeys not treated with E2 (Hao et al., 2007).

These findings indicate that E2 protects the brain via at least 3 mechanisms: 1) regulation of excitatory ion influx, 2) antioxidant systems, and 3) maintaining proper energy metabolism despite facing trauma. Furthermore, E2 appears to make use of various sub-mechanisms in modulating these protective effects. Molecular mechanisms of E2 protection include 1) upregulation of extra-cellular signal-regulated kinase (ERK) phosphorylation in the face of quinolinic acid exposure, as blockade of E2-induced upregulation of ERK phosphorylation made E2 ineffective at inhibiting quinolinic acid damage (Kuroki et al., 2001), 2) E2 reduction of oxidative stress and maintenance of respiratory capacity of mitochondria faced with various mitochondrial-specific toxins (Yao et al., 2011), and 3) blockade of glutamate-induced and hydrogen peroxide-induced cell loss by increasing nitric oxide synthesis (Andozia et al., 2010; Mannella et al., 2009).

Other ways E2 limits excitotoxicity include 1) enhancing the transport of extra-cellular lactate into neurons in the face of glutamate challenge (Mendelowitsch et al., 2001), 2) blockade of NMDA-induced rises of calcium influx via direct (versus genomic) inhibition of NMDA receptors (Weaver Jr et al., 1997), 3) reductions in hydrogen peroxide- and glutamate-induced cell death (Singer et al., 1998; Zhao et al., 2004) and cerebral ischemia-induced cell death (Dubal et al., 1999) via upregulation of the anti-apoptotic protein bcl-2, 4) protection against A β exposure associated with upregulation of another anti-apoptotic protein from the bcl family, bcl-x_L (Pike, 1999), 5) blockade of glutamate-induced and A β -induced cell damage by increasing the tolerance of mitochondria to calcium and lengthening the time of mitochondrial tolerance through upregulation of bcl-2 (Nilsen and Brinton, 2003; Nilsen et al., 2006), and 6) blockade of glutamate-induced cell death via inhibition of the excitotoxic levels of calcium influx associated with exposure to high levels of glutamate (Andozia et al., 2010; Mannella et al., 2009; Nilsen et al., 2002).

4.4 Estradiol Affects Brain Structure and Activity in Humans

Women exhibit natural fluctuations of circulating E2 levels during the monthly menstrual cycle, and dramatic decreases in E2 levels during menopause. Notably, E2 is produced not only in the reproductive tract but also in various brain regions, including the hippocampus (Mukai et al., 2006)⁵. This begs the question of whether and how brain levels of E2 relate to levels circulating elsewhere in the body. Rodents either left intact or gonadectomized followed by E2 replacement or vehicle showed treatment-dependent levels of E2 in the hippocampus (Barker and Galea, 2009), suggesting that local production of E2 in the hippocampus is influenced by circulating levels of the hormone. In addition, there is intriguing initial evidence that hippocampal volume fluctuates along with E2 levels during

⁵One source of local E2 production in the brain is aromatization of testosterone to estradiol (Simpson, 2003; Simpson, 2002).

the menstrual cycle (Protopopescu et al., 2008). In this study, women showed greater gray matter volume in the right anterior hippocampus (overlapping with anterior parahippocampal gyrus) during the high E2, low progesterone, late follicular phase (10-12 day after the onset of menses) than during in the high progesterone, low E2, late luteal phase (1-5 days before the onset of menses; See Table 5 for effects on neuronal morphology).

Given that hippocampal dendrite spine density (Gould et al., 1990; Woolley et al., 1990a; Woolley and McEwen, 1993) and synapse density (Woolley and McEwen, 1992) are modulated by E2 levels, as a function of NMDA receptor activation (Woolley and McEwen, 1994), E2 should be able to modulate the ability of the hippocampus to exhibit long-term potentiation during these different E2 phases. However, this may be disrupted in post-menopausal women, who no longer produce significant levels of E2. Evidence that E2 level within the hippocampus is influenced by circulating E2 levels in the body (Barker and Galea, 2009) suggests that post-menopausal women may produce less E2 in the hippocampus as a result of decreased circulating levels of E2⁶. Coupled with evidence for E2-related hippocampal neuroprotection, loss of E2 after menopause may make the hippocampus more vulnerable to degeneration. Thus, application of E2 close to the time of menopause may help maintain E2 production in the hippocampus, thereby prolonging the integrity of the structure and aiding in maintenance of cognitive function.

Neuroimaging studies in human females support the basic science findings of estrogenic neuroprotection (for review see, Smith and Zubieta, 2001). In a structural magnetic resonance imaging study, post-menopausal women using unopposed E2 for 20+ years experienced less ventricular enlargement and white matter loss than women who did not use E2 (Ha et al., 2007), possibly partially resulting from the estrogenic protection of oligodendrocytes and subsequent promotion of myelination observed in developmental animal models (Gerstner et al., 2007).

Functional magnetic resonance imaging (fMRI) studies have also reported greater activity in regions associated with the specific tasks being tested (See Table 6 for effects on brain activation). Women who began combination E2/progestin treatment during peri-menopause performed better on a verbal recognition task, which was associated with greater activation in the hippocampus compared with nonusers (Maki et al., 2011). Ten years of uninterrupted combination hormone treatment, initiated within two years of menopause, also was associated with greater frontal cortex and hippocampal activation during a visual working memory task performed during an fMRI scan, which positively correlated with performance on the task, compared with never-users (Berent-Spillson et al., 2010).

Other post-menopausal women following the same treatment guidelines as in Berent-Spillson et al. (2010) exhibited higher cholinergic activity in the hippocampus and posterior cingulate at rest, compared with never-users (Smith et al., 2011). Maintenance of cholinergic activity has important implications for cognitive preservation since the neurotransmitter is

⁶Prior to menopause, both circulating levels and local production via aromatization should influence local levels of E2 in women's brains. The loss of one source of E2 after menopause might then result in lower levels of the hormone in the brain as only one source of E2 remains.

intimately involved in attention, learning and memory (for reviews see, Klinkenberg et al., 2011; Micheau and Marighetto, 2011).

Although reintroduction of E2 after onset of Alzheimer's disease can exacerbate symptoms, early initiation of hormone treatment might protect carriers of the ApoE-e4 allele, an allele associated with greater risk for Alzheimer's, from developing the disease. E2 use in post-menopausal women, both with and without the ApoE-e4 allele, prevented declines in glucose metabolism in the posterior cingulate (Rasgon et al., 2005), an area associated with early declines in metabolic function for those with preclinical Alzheimer's disease (Minoshima et al., 1997), relative to carriers and non-carriers of the e4 allele not on an E2 regimen.

5. Interactions between Estradiol and Glucocorticoids in Neurons and the Stress Response

Thus far we have reviewed findings from cell-, animal-, and human-based research in order to understand how E2 and GCs affect neuronal integrity and function, as well as cognition. In this section we will discuss how E2 may modulate the effect of GC in the brain and stress response. However, it should be noted that the physiological response to stress differs between female rodents and women. For instance, female rodents display larger GC responses to stressors than males (for review see, Handa et al., 1994; see also, McCormick et al., 2002; Seale et al., 2004a; Seale et al., 2004b; Suzuki et al., 2001), attributed to a combination of excitatory actions of estradiol on the stress response system in females (Burgess and Handa, 1992; Suzuki et al., 2001; Weiser and Handa, 2009) and inhibitory actions of testosterone on the stress response system in males (Bingaman et al., 1994; Suzuki et al., 2001), while women tend to display smaller GC responses than men (Dixon et al., 2004; Kirschbaum et al., 1999; Kirschbaum et al., 1992; Kudielka et al., 2004; Kudielka et al., 1998; Kudielka and Kirschbaum, 2005). The contradicting GC response profiles of rodents and humans limits the translational generalization of stress response profiles and suggests the way in which the two systems interact to affect cognition may differ across species. Thus, discussion of E2 influences on the stress response, and implications for peri- and post-menopausal women, will be limited to studies with humans. However, we will begin with a review of cell and animal research examining how E2 and GCs interact to change the effect of one another on neurons.

5.1 Estradiol Offers Protection Against Glucocorticoid Damage

In isolation, E2 exerts protective effects in the same regions where GCs are damaging; furthermore, E2 can directly protect neural tissue from the damaging effects of GC exposure in male and female rodents (See Table 7). Males pretreated with E2 prior to dexamethasone administration exhibited attenuated apoptosis and cell damage in the striatum and hippocampus than those not receiving pretreatment (Haynes et al., 2003). Males also experienced hippocampal protection and possible protection of learning and memory processes from stress such that hippocampal slices from males exposed to a stressor prior to termination displayed greater long-term potentiation when bathed in a medium containing E2 than slices bathed in a control solution of artificial cerebrospinal fluid (Foy et al., 2008).

Similar protective trends were seen in intact females compared with intact males exposed to chronic restraint stress. Despite exhibiting higher corticosterone responses to the stressor and less success at habituation, stressed females experienced less neuronal damage in the CA3 region of the hippocampus than males. Stressed females experienced loss of basal branch points, but not branch length, while stressed males experienced a significant loss of both apical branch points and length, compared with control animals (Galea et al., 1997). In OVX females, chronic restraint stress decreased cell number (Takuma et al., 2007) and apical dendrite complexity (McLaughlin et al., 2010) in the CA3 region of the hippocampus and layers II and III of the medial prefrontal cortex (Garrett and Wellman, 2009), effects prevented by E2 replacement after OVX (Garrett and Wellman, 2009; McLaughlin et al., 2010; Takuma et al., 2007).

One mechanism of action for the estrogenic neuroprotection against GC exposure may be the ability of E2 to regulate GC receptor concentration. Typically, chronic stress or stress hormone administration results in downregulation of GC receptors (Ferrini et al., 1995; Herman et al., 1989; Sapolsky et al., 1984; for review see, Seckl and Olsson, 1995). The resulting reduction of receptors leads to an inability to accurately detect GC levels, hindering the ability of the HPA axis to shut down the stress response (Sapolsky et al., 1986). E2 administration, however, upregulates GC receptors both in the absence of stress (Ferrini and De Nicola, 1991), and in the face of chronic corticosterone administration (Ferrini et al., 1995). Estrogenic upregulation of GC receptors should allow the brain to better detect lower levels of GC, making the HPA axis more responsive, thereby leading to more rapid discontinuation of the stress response, and thus to neural protection by way of reducing exposure time to GC.

5.2 Estradiol and Stress Response Interactions in Humans

In humans, sex differences provide one line of evidence for estrogenic blunting of HPA axis activation, as women display dampened HPA responses to stressors. Naturally cycling young-adult women (i.e., not using hormone contraception) show lower adrenocorticotrophic hormone (ACTH) (Faraday et al., 2005; Kirschbaum et al., 1999; Kudielka et al., 1998) and biologically active free cortisol (Davis and Emory, 1995; Kirschbaum et al., 1999; Kirschbaum et al., 1992; Kudielka et al., 2004; Kudielka et al., 1998) responses compared with young-adult men. Likewise, women tested during the high E2 phase of the menstrual cycle showed lower stress-related activation of the anterior cingulate gyrus, orbitofrontal cortex, medial and ventromedial prefrontal cortex, amygdala, and hippocampus than men (Goldstein et al., 2010), with similar activational differences in women tested during the low E2 versus higher E2 phases of the menstrual cycle (Goldstein et al., 2005). Furthermore, women using hormone contraception show additionally blunted cortisol responses compared with naturally cycling women, possibly as a result of the ethinyl estradiol in these forms of contraception (Kirschbaum et al., 1999; Nielsen et al., 2014; Nielsen et al., 2013). Evidence in young-adult men also suggests that E2 dampens sympathetic responses to a stressor, with just one day of E2 treatment resulting in blunted systolic blood pressure, pulse rate, epinephrine, and norepinephrine responses to a mental stressor (Del Rio et al., 1994).

Similar dampening of the stress response has been observed in post-menopausal women on E2 treatment. Transdermal or oral E2 interventions spanning 1 day (Del Rio et al., 1998), 3 weeks (Ceresini et al., 2000), 1 month (Puder et al., 2001), 6 weeks (Lindheim et al., 1992), and 8 weeks (Komesaroff et al., 1999) reduced the epinephrine (Ceresini et al., 2000; Del Rio et al., 1998), norepinephrine (Ceresini et al., 2000), diastolic blood pressure (Ceresini et al., 2000; Komesaroff et al., 1999; Lindheim et al., 1992), systolic blood pressure (Komesaroff et al., 1999; Lindheim et al., 1992), ACTH and cortisol (Puder et al., 2001) responses to various stressors, such as mental stressors (Ceresini et al., 2000; Del Rio et al., 1998; Lindheim et al., 1992), the cold pressor task (Lindheim et al., 1992), and an endotoxin challenge (Puder et al., 2001).

Together, the findings indicate that E2 blunts the magnitude of the HPA axis response to stressors in women. Taking into account the negative effects of prolonged or heightened stress hormone exposure, the loss of E2 during and after menopause may put post-menopausal women at greater risk of experiencing stress-induced alterations in cognitive function, as a result of dramatically reduced E2 levels. However, the use of hormone supplements during and after the menopause transition should alleviate the potential increased negative effects of stress exposure on cognitive function.

6. Estradiol May Counteract Negative Effects of Glucocorticoids in Middle-Aged and Older Females

Aging is associated with a number of bodily changes. Among these changes are alterations in the function of the stress response system, particularly the HPA axis. A common paradigm for testing HPA axis sensitivity to stressors in aged subjects is to administer real or synthetic GCs and measure how quickly the HPA axis responds to and adjusts subsequent hormone release. In one model, animals are given injections of dexamethasone (DEX), which activates the negative feedback system of the HPA axis, and reduces subsequent release of the animal's endogenous corticosterone response to a stressor. In young-adult animals, this procedure results in a significant reduction of corticosterone release in response to a stress, however, aged male rodents fail to show this DEX-induced corticosterone suppression to a stressor, (Ferrini et al., 1999; Riegle and Hess, 1972). This age-related failure to inhibit HPA axis reactivity also is observed in female rhesus monkeys subjected to corticotropin releasing hormone (CRH), ACTH, and DEX tests (Goncharova and Lapin, 2002) and in a CRH test performed in male and female dogs (Reul et al., 1991). Additionally, aged male rats also displayed prolonged corticosterone secretion after termination of an acute stressor compared with young animals (Ferrini et al., 1999; Sapolsky et al., 1983), with similar results observed in older male and female rodents exposed to chronic stress (Riegle, 1973; Sapolsky et al., 1983).

One possible explanation for the heightened and prolonged stress response in aged animals may be age-related downregulation of GC receptors in the brain, particularly in the hippocampus (Ferrini et al., 1999; Mizoguchi et al., 2009; Sapolsky et al., 1986). Loss of GC receptors in the aged hippocampus, a structure intimately involved in termination of the stress response, leads to a failure to accurately detect circulating GC levels and thus a failure to downregulate the HPA axis during and after stress exposure. The resulting excess GC

exposure leads to hippocampal damage and subsequent loss of more GC receptors, leading to a cycle of more GC exposure, damage, and receptor loss (Sapolsky et al., 1986).

Consistent with this pattern, some reports indicate older adult humans also exhibit higher basal cortisol, and potentiated and prolonged stress responses as measured by various biomarkers, such as cortisol and salivary alpha amylase (a biomarker for norepinephrine; Almela et al., 2011; for salivary alpha amylase as a marker for norepinephrine response see, Chatterton et al., 1996; see also, Dodt et al., 1991; Wilkinson et al., 2001). While human studies on brain levels of GC receptor concentration are few and far between, there is preliminary work showing older adults do experience GC receptor downregulation in both the brain and bloodstream. For instance, older adults had reduced levels of GC receptor mRNA in plasma compared with younger adults (Grasso et al., 1997), and showed lower GC receptor concentration in the dorsolateral prefrontal cortex than the expression displayed in the brains of adolescents and adults in a postmortem study (Perlman et al., 2007). Reductions of GC mRNA in plasma and the brain may explain failures of older adults to downregulate the HPA axis after a brief stressful event.

Women, however, may experience protection from this cycle of decline in GC receptor concentration, HPA hyperactivity, and ensuing neural damage, by extending E2 presence and action beyond menopause. The loss of E2 after menopause likely leads to potentiated age-related changes in HPA axis function due to the loss of estrogenic protection. However, prolonging E2 exposure beyond menopause will likely maintain proper HPA axis function. For instance, recall that E2 administration upregulated GC receptors in the hippocampus of older male rats, compared with older animals not treated with E2 (Ferrini et al., 1999). This type of estrogenic upregulation of GC receptors in the hippocampus could help prevent or reverse the age-related hyperactivity of the HPA axis as more receptors would be available to detect circulating GC levels and spare the hippocampus, and other brain regions, such as the prefrontal cortex, from further damage.

Importantly for human females, E2 has been shown to counter each of the age-related dysfunctions (e.g., hippocampal damage, receptor downregulation, and HPA axis hyperactivity) described above. Yet, despite the protective effects of E2 treatment after menopause, and the attenuating effects of E2 on the stress response, Newhouse et al. (2010) reported worse performance on various measures of attention and for memory of hard word pairs in a verbal paired associates task in women assigned to three months of E2 treatment, compared with women who had been assigned to placebo, after completion of the TSST. However, similar to the WHI study, some caveats apply. Women assigned to E2 were an average age of 65.18 years old, with an age range of 52-83, and years since menopause averaged 14.57 years, with a range of 1-31 years. Thus, while the age and time-since-menopause ranges surely captured some women within the first years since their last menses, it also captured women well beyond the climacteric transition. Similar to the WHI studies, the inclusion of women outside the critical window (for review see, Maki, 2013; see also, Resnick and Henderson, 2002) could have activated effects of the healthy cell bias (Brinton, 2005) and resulted in less protection from stress and greater impairments in attention and short-term episodic memory.

Insofar as E2 protects the hippocampus and prefrontal cortex, and consequently the negative feedback system of the HPA axis, extension of E2 exposure past menopause may be a possible mechanism for the relatively enhanced cognitive function observed in post-menopausal women who initiated hormone treatment close to the time of menopause.

7. Conclusion

Stress is encountered by all people on a daily basis. With regard to women, there is clear evidence that the systems governing stress and sex hormones intimately interact and even exert opposing effects on various bodily systems. Given the countervailing effects of the two hormones, the decline in proper GC regulation with aging, and the ability of E2 to directly mitigate potential negative effects of GC exposure, there is the potential for increased and/or accelerated neuronal and cognitive decline post-menopause due to increased HPA reactivity to stress exposure and lack of E2 protection.

This line of research is relevant to half of the global population, as all women should eventually experience menopause, and while the exact parameters of if and when E2 replacement is suitable vary from woman to woman, it is nonetheless important to understand all potential benefits and risks of E2 replacement, such as whether E2 can limit age-related changes in the stress response system. There are also many questions that remain to be answered. One such question, discussed here, is when estradiol replacement should begin relative to menopause in order to enhance potential benefits and limit potential dangers. Others needing to be addressed and discussed elsewhere in relation to other domains, include whether different forms of estrogens follow different time and dose curves before affecting the domain of interest, or whether some forms of estrogens and estradiol lead to negative versus positive effects (Barha et al., 2010; for review see, Barha and Galea, 2010; see also, Barha and Galea, 2013; Barha et al., 2009; McClure et al., 2013). These points illustrate how the current state of estradiol-treatment research is incomplete and in need of more intensive research. Working toward uncovering how the interaction between GC and E2 changes after menopause, and the implications of these changes elsewhere in the peri- and post-menopausal woman's body, is necessary for providing the most complete understanding of how estradiol treatment may affect women's health.

Acknowledgements

This work was supported by grant R01AG038043 from the National Institute on Aging. We would like to thank the reviewers for their comprehensive notes and suggestions, as well as Shawn E. Nielsen, Ph.D. for her comments on the final draft of this review.

Abbreviations

GC	Glucocorticoids
(HPA) Axis	Hypothalamic-Pituitary-Adrenal
(HPG) Axis	Hypothalamic-Pituitary-Gonadal
E2	17 β -Estradiol

TSST	Trier Social Stress Test
CPT	Cold Pressor Task
Aβ	Beta-Amyloid
OVX	Ovariectomized
WHI	Women's Health Initiative
WHIMS	Women's Health Initiative Memory Study
WHISCA	Women's Health Initiative Study of Cognitive Aging
fMRI	Functional Magnetic Resonance Imaging
ACTH	Adrenocorticotrophic Hormone
DEX	Dexamethasone
CRH	Corticotropin Releasing Hormone

REFERENCES

- Al'Absi M, Hugdahl K, Lovallo WR. Adrenocortical stress responses and altered working memory performance. *Psychophysiology*. 2002; 39:95–99. [PubMed: 12206301]
- Alexander JK, Hillier A, Smith RM, Tivarus ME, Beversdorf DQ. Beta-adrenergic modulation of cognitive flexibility during stress. *Journal of Cognitive Neuroscience*. 2007; 19:468–478. [PubMed: 17335395]
- Almela M, Hidalgo V, Villada C, van der Meij L, Espín L, Gómez-Amor J, Salvador A. Salivary alpha-amylase response to acute psychosocial stress: The impact of age. *Biological Psychology*. 2011; 87:421–429. [PubMed: 21664412]
- Alonso M, Vianna MRM, Depino AM, Mello e Souza T, Pereira P, Szapiro G, Viola H, Pitossi F, Izquierdo I, Medina JH. Bdnf-triggered events in the rat hippocampus are required for both short- and long-term memory formation. *Hippocampus*. 2002; 12:551–560. [PubMed: 12201640]
- Anacker C, Cattaneo A, Luoni A, Musaelyan K, Zunszain PA, Milanese E, Rybka J, Berry A, Cirulli F, Thuret S, Price J, Riva MA, Gennarelli M, Pariante CM. Glucocorticoid-related molecular signaling pathways regulating hippocampal neurogenesis. *Neuropsychopharmacology*. 2012; 38:872–883. [PubMed: 23303060]
- Anderson KJ. Impulsivity, caffeine, and task difficulty: A within-subjects test of the yerkes-dodson law. *Personality and Individual Differences*. 1994; 16:813–829.
- Andozia MB, Vieira CS, Franceschini SA, Tolloi MRT, de Sá MFS, Ferriani RA. Ethinylestradiol and estradiol have different effects on oxidative stress and nitric oxide synthesis in human endothelial cell cultures. *Fertility and sterility*. 2010; 94:1578–1582. [PubMed: 19909948]
- Arbel I, Kadar T, Silberman M, Levy A. The effects of long-term corticosterone administration on hippocampal morphology and cognitive performance of middle-aged rats. *Brain Research*. 1994; 657
- Baker LD, Asthana S, Cholerton BA, Wilkinson CW, Plymate SR, Green PS, Merriam GR, Fishel MA, Watson GS, Cherrier MM, Kletke ML, Mehta PD, Craft S. Cognitive response to estradiol in postmenopausal women is modified by high cortisol. *Neurobiology of Aging*. 2012; 33:829.e829–829.e820. [PubMed: 21855173]
- Barha CK, Dalton GL, Galea LA. Low doses of 17[alpha]-estradiol and 17[beta]-estradiol facilitate, whereas higher doses of estrone and 17[alpha]- and 17[beta]-estradiol impair, contextual fear conditioning in adult female rats. *Neuropsychopharmacology*. 2010; 35:547–559. [PubMed: 19847162]

- Barha CK, Galea LA. Influence of different estrogens on neuroplasticity and cognition in the hippocampus. *Biochimica et Biophysica Acta (BBA) - General Subjects*. 2010; 1800:1056–1067. [PubMed: 20100545]
- Barha CK, Galea LA. The hormone therapy, premarin, impairs hippocampus-dependent spatial learning and memory and reduces activation of new granule neurons in response to memory in female rats. *Neurobiol Aging*. 2013; 34:986–1004. [PubMed: 22938820]
- Barha CK, Lieblich SE, Galea LA. Different forms of oestrogen rapidly upregulate cell proliferation in the dentate gyrus of adult female rats. *Journal of Neuroendocrinology*. 2009; 21:155–166. [PubMed: 19076272]
- Barker JM, Galea LAM. Sex and regional differences in estradiol content in the prefrontal cortex, amygdala and hippocampus of adult male and female rats. *General and Comparative Endocrinology*. 2009; 164:77–84. [PubMed: 19457436]
- Beck KD, Luine VN. Sex differences in behavioral and neurochemical profiles after chronic stress: Role of housing conditions. *Physiology & behavior*. 2002; 75:661–673. [PubMed: 12020731]
- Beck KD, Luine VN. Evidence for sex-specific shifting of neural processes underlying learning and memory following stress. *Physiology & Behavior*. 2010; 99:204–211. [PubMed: 19376144]
- Becker JB, Monteggia LM, Perrot-Sinal TS, Romeo RD, Taylor JR, Yehuda R, Bale TL. Stress and disease: Is being female a predisposing factor? *The Journal of Neuroscience*. 2007; 27:11851–11855. [PubMed: 17978023]
- Behl C. Effects of glucocorticoids on oxidative stress-induced hippocampal cell death: Implications for the pathogenesis of alzheimer's disease. *Experimental Gerontology*. 1998; 33:689–696. [PubMed: 9951616]
- Behl C, Lezoualch F, Trapp T, Widmann M, Skutella T, Holsboer F. Glucocorticoids enhance oxidative stress-induced cell death in hippocampal neurons in vitro. *Endocrinology*. 1997; 138:101–106. [PubMed: 8977391]
- Berent-Spillon A, Persad CC, Love T, Tkaczyk A, Wang H, Reame NK, Frey KA, Zubieta J-K, Smith YR. Early menopausal hormone use influences brain regions used for visual working memory. *Menopause*. 2010; 1
- Bimonte HA, Denenberg V, H. Estradiol facilitates performance as working memory load increases. *Psychoneuroendocrinology*. 1999; 24:161–173. [PubMed: 10101725]
- Bingaman E, Magnuson D, Gray T, Handa R. Androgen inhibits the increases in hypothalamic corticotropin-releasing hormone (crh) and crh-Immunoreactivity following gonadectomy. *Neuroendocrinology*. 1994; 59:228–234. [PubMed: 8159272]
- Blandino P Jr, Barnum CJ, Deak T. The involvement of norepinephrine and microglia in hypothalamic and splenic il-1 β responses to stress. *Journal of neuroimmunology*. 2006; 173:87–95. [PubMed: 16386803]
- Bowman RE. Chronic stress effects on memory: Sex differences in performance and monoaminergic activity. *Hormones and Behavior*. 2003; 43:48–59. [PubMed: 12614634]
- Bowman RE, Ferguson D, Luine VN. Effects of chronic restraint stress and estradiol on open field activity, spatial memory, and monoaminergic neurotransmitters in ovariectomized rats. *Neuroscience*. 2002; 113:401–410. [PubMed: 12127097]
- Bowman RE, Micik R, Gautreaux C, Fernandez L, Luine VN. Sex-dependent changes in anxiety, memory, and monoamines following one week of stress. *Physiology & Behavior*. 2009; 97:21–29. [PubMed: 19419681]
- Breen KM, Karsch FJ. New insights regarding glucocorticoids, stress and gonadotropin suppression. *Frontiers in Neuroendocrinology*. 2006; 27:233–245. [PubMed: 16712908]
- Brien T. Human corticosteroid binding globulin. *Clinical Endocrinology*. 1981; 14:193–212. [PubMed: 7021007]
- Brinton RD. Investigative models for determining hormone therapy-induced outcomes in brain: Evidence in support of a healthy cell bias of estrogen action. *Annals of the New York Academy of Sciences*. 2005; 1052:57–74. [PubMed: 16024751]
- Brinton RD, Chen S, Montoya M, Hsieh D, Minaya J, Kim J, Chu H-P. The women's health initiative estrogen replacement therapy is neurotrophic and neuroprotective. *Neurobiology of Aging*. 2000; 21:475–496. [PubMed: 10858597]

- Brown SM, Henning S, Wellman CL. Mild, short-term stress alters dendritic morphology in rat medial prefrontal cortex. *Cerebral Cortex*. 2005; 15:1714–1722. [PubMed: 15703248]
- Buchanan TW, Lovallo WR. Enhanced memory for emotional material following stress-level cortisol treatment in humans. *Psychoneuroendocrinology*. 2001; 26:307–317. [PubMed: 11166493]
- Bullinger M, Naber D, Pickar D, Cohen RM, Kalin NH, Pert A, Bunney WE Jr. Endocrine effects of the cold pressor test: Relationships to subjective pain appraisal and coping. *Psychiatry Research*. 1984; 12:227–233. [PubMed: 6093173]
- Burgess LH, Handa RJ. Chronic estrogen-induced alterations in adrenocorticotropin and corticosterone secretion, and glucocorticoid receptor-mediated functions in female rats. *Endocrinology*. 1992; 131:1261–1269. [PubMed: 1324155]
- Burke C, Roulet F. Increased exposure of tissues to cortisol in late pregnancy. *British medical journal*. 1970; 1:657. [PubMed: 5443967]
- Butts KA, Weinberg J, Young AH, Phillips AG. Glucocorticoid receptors in the prefrontal cortex regulate stress-evoked dopamine efflux and aspects of executive function. *Proceedings of the National Academy of Sciences*. 2011; 108:18459–18464.
- Cabeza R, Dolcos F, Graham R, Nyberg L. Similarities and differences in the neural correlates of episodic memory retrieval and working memory. *NeuroImage*. 2002; 16:317–330. [PubMed: 12030819]
- Ceresini G, Freddi M, Marganti S, Rebecchi I, Modena AB, Rinaldi M, Manca C, Amaducci A, Del Rio G, Valenti G. The effects of transdermal estradiol on the response to mental stress in postmenopausal women: A randomized trial. *American Journal of Medicine*. 2000; 109:463–468. [PubMed: 11042235]
- Cerqueira JJ, Mailliet F, Almeida OF, Jay TM, Sousa N. The prefrontal cortex as a key target of the maladaptive response to stress. *The Journal of Neuroscience*. 2007; 27:2781–2787. [PubMed: 17360899]
- Chatterton RT, Vogelsong KM, Lu Y.c. Ellman AB, Hudgens GA. Salivary α -amylase as a measure of endogenous adrenergic activity. *Clinical Physiology*. 1996; 16:433–448. [PubMed: 8842578]
- Chen S, Nilsen J, Brinton RD. Dose and temporal pattern of estrogen exposure determines neuroprotective outcome in hippocampal neurons: Therapeutic implications. *Endocrinology*. 2006; 147:5303–5313. [PubMed: 16916950]
- Chrousos GP. Stress and disorders of the stress system. *Nature Reviews Endocrinology*. 2009; 5:374–381.
- Coburn-Litvak PS, Pothakos K, Tata DA, McCloskey DP, Anderson BJ. Chronic administration of corticosterone impairs spatial reference memory before spatial working memory in rats. *Neurobiology of Learning and Memory*. 2003; 80:11–23. [PubMed: 12737930]
- Cook SC, Wellman CL. Chronic stress alters dendritic morphology in rat medial prefrontal cortex. *Journal of Neurobiology*. 2004; 60:236–248. [PubMed: 15266654]
- Cousijn H, Rijpkema M, Qin S, van Wingen GA, Fernández G. Phasic deactivation of the medial temporal lobe enables working memory processing under stress. *NeuroImage*. 2012; 59:1161–1167. [PubMed: 21983180]
- Curtis CE, D'Esposito M. Persistent activity in the prefrontal cortex during working memory. *Trends in Cognitive Sciences*. 2003; 7:415–423. [PubMed: 12963473]
- Daniel JM, Fader AJ, Spencer AL, Dohanich GP. Estrogen enhances performance of female rats during acquisition of a radial arm maze. *Hormones and Behavior*. 1997; 32:217–225. [PubMed: 9454673]
- Davis M, Emory E. Sex differences in neonatal stress reactivity. *Child Development*. 1995; 66:14–27. [PubMed: 7497822]
- de Kloet ER, Oitzl MS, Joëls M. Stress and cognition: Are corticosteroids good or bad guys? *Trends in Neurosciences*. 1999; 22:422–426. [PubMed: 10481183]
- Del Rio G, Velardo A, Menozzi R, Zizzo G, Tavernari V, Venneri MG, Marrama P, Petraglia F. Acute estradiol and progesterone administration reduced cardiovascular and catecholamine responses to mental stress in menopausal women. *Neuroendocrinology*. 1998; 67:269–274. [PubMed: 9588696]

- Del Rio G, Velardo A, Zizzo G, Avogaro A, Cipolli C, Della Casa L, Marrama P, Macdonald IA. Effect of estradiol on the sympathoadrenal response to mental stress in normal men. *Journal of Clinical Endocrinology & Metabolism*. 1994; 79:836–840. [PubMed: 8077370]
- Delmas PD, Bjarnason NH, Mitlak BH, Ravoux A, Shah AS, Huster WJ, Draper M, Christiansen C. Effects of raloxifene on bone mineral density, serum cholesterol, concentration, and uterine endometrium in postmenopausal women. *The new england journal of medicine*. 1997; 337:1641–1647. [PubMed: 9385122]
- Diamond DM, Bennett MC, Fleshner M, Rose GM. Inverted-u relationship between the level of peripheral corticosterone and the magnitude of hippocampal primed burst potentiation. *Hippocampus*. 1992; 2:421–430. [PubMed: 1308198]
- Diamond DM, Fleshner M, Ingersoll N, Rose GM. Psychological stress impairs spatial working memory: Relevance to electrophysiological studies of hippocampal function. *Behavioral Neuroscience*. 1996; 110:661–672. [PubMed: 8864259]
- Dixon KE, Thorn BE, Ward LC. An evaluation of sex differences in psychological and physiological responses to experimentally-induced pain: A path analytic description. *Pain*. 2004; 112:188–196. [PubMed: 15494200]
- Dodson JD. The relation of strength of stimulus to rapidity of habit-formation in the kitten. *Journal of Animal Behavior*. 1915; 5:330–336.
- Dotz C, Dittman J, Hruba J, Spath-Schwalbe E, Born J, Schuttler R, Fehm HL. Different regulation of adrenocorticotropin and cortisol in young, mentally healthy elderly, and patients with senile dementia of alzheimer's type. *Journal of Clinical Endocrinology & Metabolism*. 1991; 72:272–276. [PubMed: 1846870]
- Dubal DB, Shughrue PJ, Wilson ME, Merchenthaler I, Wise PM. Estradiol modulates bcl-2 in cerebral ischemia: A potential role for estrogen receptors. *The Journal of neuroscience*. 1999; 19:6385–6393. [PubMed: 10414967]
- Duff SJ, Hampson E. A beneficial effect of estrogen on working memory in postmenopausal women taking hormone replacement therapy. *Hormones and Behavior*. 2000; 38:262–276. [PubMed: 11104644]
- Duncko R, Johnson L, Merikangas K, Grillon C. Working memory performance after acute exposure to the cold pressor stress in healthy volunteers. *Neurobiology of Learning and Memory*. 2009; 91:377–381. [PubMed: 19340949]
- Dunn GA, Morgan CP, Bale TL. Sex-specificity in transgenerational epigenetic programming. *Hormones and Behavior*. 2011; 59:290–295. [PubMed: 20483359]
- Edelson JT, Robertson GL. The effect of the cold pressor test on vasopressin secretion in man. *Psychoneuroendocrinology*. 1986; 11:307–316. [PubMed: 3786636]
- Egeland J, Lund A, Landrø NI, Rund BR, Sundet K, Asbjørnsen A, Mjellem N, Roness A, Stordal KI. Cortisol level predicts executive and memory function in depression, symptom level predicts psychomotor speed. *Acta Psychiatrica Scandinavica*. 2005; 112:434–441. [PubMed: 16279872]
- Elenkov IJ, Chrousos GP. Stress hormones, proinflammatory and antiinflammatory cytokines, and autoimmunity. *Annals of the New York Academy of Sciences*. 2002; 966:290–303. [PubMed: 12114286]
- Elsabagh S, Hartley D, File S. Cognitive function in late versus early postmenopausal stage. *Maturitas*. 2007; 56:84–93. [PubMed: 16857328]
- Elzinga BM, Roelofs K. Cortisol-induced impairments of working memory require acute sympathetic activation. *Behavioral Neuroscience*. 2005; 119:98–103. [PubMed: 15727516]
- Espeland MA, Rapp SR, Shumaker SA, Brunner R, Manson JE, Sherwin BB, Hsia J, Margolis KL, Hogan PE, Wallace R, Dailey M, Freeman R, Hays J. Conjugated equine estrogens and global cognitive function in postmenopausal women. *Journal of the American Medical Association*. 2004; 291:2959–2968. [PubMed: 15213207]
- Espósito F, Bertolino A, Scarabino T, Latorre V, Blasi G, Papolizio T, Tedeschi G, Cirillo S, Goebel R, Di Salle F. Independent component model of the default-mode brain function: Assessing the impact of active thinking. *Brain Research Bulletin*. 2006; 70:263–269. [PubMed: 17027761]

- Fader AJ, Johnson PE, Dohanich GP. Estrogen improves working but not reference memory and prevents amnesic effects of scopolamine on a radial-arm maze. *Pharmacology Biochemistry and Behavior*. 1999; 62:711–717.
- Faraday M, Blakeman K, Grunberg N. Strain and sex alter effects of stress and nicotine on feeding, body weight, and hpa axis hormones. *Pharmacology Biochemistry and Behavior*. 2005; 80:577–589.
- Felson DT, Zhang Y, Hannan MT, Kiel DP, Wilson PWF, Anderson JJ. The effect of postmenopause estrogen therapy on bone density in elderly women. *The new england journal of medicine*. 1993; 329:1141–1146. [PubMed: 8377776]
- Ferrini M, De Nicola AF. Estrogens up-regulate type i and type ii glucocorticoid receptors in brain regions from ovariectomized rats. *Life Sciences*. 1991; 48:2593–2601. [PubMed: 2046482]
- Ferrini M, Lima A, De Nicola AF. Estradiol abolishes autologous down regulation of glucocorticoid receptors in brain. *Life Sciences*. 1995; 57:2403–2412. [PubMed: 8847961]
- Ferrini M, Piroli G, Frontera M, Falbo A, Lima A, De Nicola AF. Estrogens normalize the hypothalamic-pituitary-adrenal axis response to stress and increase glucocorticoid receptor immunoreactivity in hippocampus of aging male rats. *Neuroendocrinology*. 1999; 69:129–137. [PubMed: 9986926]
- Filipovic D, Zlatkovic J, Gass P, Inta D. The differential effects of acute vs. Chronic stress and their combination on hippocampal parvalbumin and inducible heat shock protein 70 expression. *Neuroscience*. 2013; 236:47–54. [PubMed: 23352936]
- Foy MR, Baudry M, Foy JG, Thompson RF. 17 β -estradiol modifies stress-induced and age-related changes in hippocampal synaptic plasticity. *Behavioral Neuroscience*. 2008; 122:301–309. [PubMed: 18410170]
- Frank MG, Baratta MV, Sprunger DB, Watkins LR, Maier SF. Microglia serve as a neuroimmune substrate for stress-induced potentiation of cns pro-inflammatory cytokine responses. *Brain, Behavior, and Immunity*. 2007; 21:47–59.
- Galea LA, McEwen BS, Tanapat P, Deak T, Spencer RL, Dhabhar FS. Sex differences in dendritic atrophy of ca3 pyramidal neurons in response to chronic restraint stress. *Neuroscience*. 1997; 81:689–697. [PubMed: 9316021]
- Garrett JE, Wellman CL. Chronic stress effects on dendritic morphology in medial prefrontal cortex: Sex differences and estrogen dependence. *Neuroscience*. 2009; 162:195–207. [PubMed: 19401219]
- Gerlach JL, McEwen BS. Rat brain binds adrenal steroid hormone: Radioautography of hippocampus with corticosterone. *Science*. 1972; 175:1133–1136. [PubMed: 5060047]
- Gerstner B, Sifringer M, Dzietko M, Schüller A, Lee J, Simons S, Obladen M, Volpe JJ, Rosenberg PA, Felderhoff-Mueser U. Estradiol attenuates hyperoxia-induced cell death in the developing white matter. *Annals of Neurology*. 2007; 61:562–573. [PubMed: 17427919]
- Gibbs RB. Estrogen replacement enhances acquisition of a spatial memory task and reduces deficits associated with hippocampal muscarinic receptor inhibition. *Hormones and Behavior*. 1999; 36:222–233. [PubMed: 10603286]
- Goel N, Bale TL. Examining the intersection of sex and stress in modelling neuropsychiatric disorders. *Journal of Neuroendocrinology*. 2009; 21:415–420. [PubMed: 19187468]
- Goldstein JM, Jerram M, Abbs B, Whitfield-Gabrieli S, Makris N. Sex differences in stress response circuitry activation dependent on female hormonal cycle. *Journal of Neuroscience*. 2010; 30:431–438. [PubMed: 20071507]
- Goldstein JM, Jerram M, Poldrack R, Ahern T, Kennedy DN, Seidman LJ, Makris N. Hormonal cycle modulates arousal circuitry in women using functional magnetic resonance imaging. *Journal of Neuroscience*. 2005; 25:9309–9316. [PubMed: 16207891]
- Goncharova ND, Lapin BA. Effects of aging on hypothalamic–pituitary–adrenal system function in non-human primates. *Mechanisms of ageing and development*. 2002; 123:1191–1201. [PubMed: 12044968]
- Gore AC, Attardi B, DeFranco DB. Glucocorticoid repression of the reproductive axis: Effects on *gnrh* and gonadotropin subunit mRNA levels. *Molecular and Cellular Endocrinology*. 2006; 256:40–48. [PubMed: 16839661]

- Gould E, Woolley CS, Frankfurt M, McEwen BS. Gonadal steroids regulate dendritic spine density in hippocampal pyramidal cells in adulthood. *The Journal of Neuroscience*. 1990; 10:1286–1291. [PubMed: 2329377]
- Grasso G, Lodi L, Lupo C, Muscettola M. Glucocorticoid receptors in human peripheral blood mononuclear cells in relation to age and to sport activity. *Life Sciences*. 1997; 61:301–308. [PubMed: 9217290]
- Green JS, Stanforth PR, Rankinen T, Leon AS, Rao DC, Skinner JS, Bouchard C, Wilmore JH. The effects of exercise training on abdominal visceral fat, body composition, and indicators of the metabolic syndrome in postmenopausal women with and without estrogen replacement therapy: The heritage family study. *Metabolism*. 2004; 53:1192–1196. [PubMed: 15334383]
- Greicius MD, Srivastava G, Reiss AL, Menon V, Raichle ME. Default-mode network activity distinguishes alzheimer's disease from healthy aging. *Proceedings of the National Academy of Sciences*. 2004; 101:4637–4642.
- Ha DM, Xu J, Janowsky JS. Preliminary evidence that long-term estrogen use reduces white matter loss in aging. *Neurobiology of Aging*. 2007; 28:1936–1940. [PubMed: 17030475]
- Handa RJ, Burgess LH, Kerr JE, O'Keefe JA. Gonadal steroid hormone receptors and sex differences in the hypothalamo-pituitary-adrenal axis. *Hormones and Behavior*. 1994; 28:464–476. [PubMed: 7729815]
- Hao J. Estrogen alters spine number and morphology in prefrontal cortex of aged female rhesus monkeys. *Journal of Neuroscience*. 2006; 26:2571–2578. [PubMed: 16510735]
- Hao J, Rapp PR, Janssen WGM, Lou W, Lasley BL, Hof PR, Morrison JH. Interactive effects of age and estrogen on cognition and pyramidal neurons in monkey prefrontal cortex. *Proceedings of the National Academy of Sciences*. 2007; 104:11465–11470.
- Haynes LE, Lendon CL, Barber DJ, Mitchell IJ. 17 β -oestradiol attenuates dexamethasone-induced lethal and sublethal neuronal damage in the striatum and hippocampus. *Neuroscience*. 2003; 120:799–806. [PubMed: 12895519]
- Herbison AE. Multimodal influence of estrogen upon gonadotropin-releasing hormone neurons. *Endocrine reviews*. 1998; 19:302–330. [PubMed: 9626556]
- Herman JP, Patel PD, Akil H, Watson SJ. Localization and regulation of glucocorticoid and mineralocorticoid receptor mRNAs in the hippocampal formation of the rat. *Molecular Endocrinology*. 1989; 3:1886–1894. [PubMed: 2558306]
- Hillner KM, Neumann ID, Couillard-Despres S, Aigner L, Slattery DA. Sex-dependent regulation of hippocampal neurogenesis under basal and chronic stress conditions in rats. *Hippocampus*. 2013; 23:476–487. [PubMed: 23504963]
- Hoffman G, Moore N, Fiksum G, Murphy AZ. Ovarian steroid modulation of seizure severity and hippocampal cell death after kainic acid treatment. *Experimental Neurology*. 2003; 182:124–134. [PubMed: 12821382]
- Holmes A, Wellman CL. Stress-induced prefrontal reorganization and executive dysfunction in rodents. *Neuroscience & Biobehavioral Reviews*. 2009; 33:773–783. [PubMed: 19111570]
- Hosoda T, Nakajima H, Honjo H. Estrogen protects neuronal cells from amyloid beta-induced apoptotic cell death. *NeuroReport*. 2001; 12:1965–1970. [PubMed: 11435931]
- Hsueh AJW, Erickson GF. Glucocorticoid inhibition of fsh-induced estrogen production in cultured rat granulosa cells. *Steroids*. 1978; 32:639–648. [PubMed: 734698]
- Jia D, O'Brien CA, Stewart SA, Manolagas SC, Weinstein RS. Glucocorticoids act directly on osteoclasts to increase their life span and reduce bone density. *Endocrinology*. 2006; 147:5592–5599. [PubMed: 16935844]
- Jones, EE. The female reproductive system. In: Boron, WF.; Boulpaep, EL., editors. *Medical Physiology*. 2 ed.. Elsevier Health Sciences; 2009.
- Kirschbaum C, Kudielka B, Gaab J, Schommer NC, Hellhammer DH. Impact of gender, menstrual cycle phase, and oral contraceptives on the activity of the hypothalamus-pituitary-adrenal axis. *Psychosomatic Medicine*. 1999; 61:154–162. [PubMed: 10204967]
- Kirschbaum C, Wust S, Hellhammer DH. Consistent sex differences in cortisol responses to psychological stress. *Psychosomatic Medicine*. 1992; 54:648–657. [PubMed: 1454958]

- Kleen JK, Sitomer MT, Killeen PR, Conrad CD. Chronic stress impairs spatial memory and motivation for reward without disrupting motor ability and motivation to explore. *Behavioral Neuroscience*. 2006; 120:842–851. [PubMed: 16893290]
- Klinkenberg I, Sambeth A, Blokland A. Acetylcholine and attention. *Behavioural Brain Research*. 2011; 221:430–442. [PubMed: 21108972]
- Komesaroff PA, Elser MD, Sudhir K. Estrogen supplementation attenuates glucocorticoid and catecholamine responses to mental stress in perimenopausal women. *Journal of Clinical Endocrinology & Metabolism*. 1999; 84:606–610. [PubMed: 10022424]
- Krotkiewski M, Bjorntorp P, Sjostrom L, Smith U. Impact of obesity on metabolism in men and women: Importance of regional adipose tissue distribution. *Journal of Clinical Investigation*. 1983; 83:1150–1162. [PubMed: 6350364]
- Krug R, Born J, Rasch B. A 3-day estrogen treatment improves prefrontal cortex-dependent cognitive function in postmenopausal women. *Psychoneuroendocrinology*. 2006; 31:965–975. [PubMed: 16831520]
- Kudielka B, Buske-Kirschbaum A, Hellhammer DH, Kirschbaum C. Hpa axis responses to laboratory psychosocial stress in healthy elderly adults, younger adults, and children: Impact of age and gender. *Psychoneuroendocrinology*. 2004; 29:83–98. [PubMed: 14575731]
- Kudielka B, Hellhammer J, Hellhammer DH, Wolf O, Pirke K, Varadi E, Pilz J, Kirschbaum C. Sex differences in endocrine and psychological responses to psychosocial stress in healthy elderly subjects and the impact of a 2-week dehydroepiandrosterone treatment. *Journal of Clinical Endocrinology & Metabolism*. 1998; 83:1756–1761. [PubMed: 9589688]
- Kudielka B, Kirschbaum C. Sex differences in hpa axis responses to stress: A review. *Biological Psychology*. 2005; 69:113–132. [PubMed: 15740829]
- Kuhlmann S, Piel M, Wolf OT. Impaired memory retrieval after psychosocial stress in healthy young men. *The Journal of Neuroscience*. 2005; 25:2977–2982. [PubMed: 15772357]
- Kuroki Y, Fukushima K, Kanda Y, Mizuno K, Watanabe Y. Neuroprotection by estrogen via extracellular signal-regulated kinase against quinolinic acid-induced cell death in the rat hippocampus. *European Journal of Neuroscience*. 2001; 13:472–476. [PubMed: 11168553]
- Laroche S, Davis S, Jay TM. Plasticity at hippocampal to prefrontal cortex synapses: Dual roles in working memory and consolidation. *Hippocampus*. 2000; 10:438–446. [PubMed: 10985283]
- Lee Y-A, Goto Y. Chronic stress effects on working memory: Association with prefrontal cortical tyrosine hydroxylase. *Behavioural brain research*. 2015
- Lighthall NR, Mather M, Gorlick MA. Acute stress increases sex differences in risk seeking in the balloon analogue risk task. *PLoS ONE*. 2009; 4
- Lighthall NR, Sakaki M, Vasunilashorn S, Nga L, Somayajula S, Chen EY, Samii N, Mather M. Gender differences in reward-related decision processing under stress. *Social Cognitive and Affective Neuroscience*. 2011
- Lindheim SR, Legro RS, Bernstein L, Stanczyk FZ, Vijod MA, Presser SC, Lobo RA. Behavioral stress responses in premenopausal and postmenopausal women and the effects of estrogen. *American Journal of Obstetrics and Gynecology*. 1992; 167:1831–1836. [PubMed: 1471706]
- Liston C, McEwen B, Casey B. Psychosocial stress reversibly disrupts prefrontal processing and attentional control. *Proceedings of the National Academy of Sciences*. 2009; 106:912–917.
- Luethi M, Meier B, Sandi C. Stress effects on working memory, explicit memory, and implicit memory for neutral and emotional stimuli in healthy men. *Frontiers in Behavioral Neuroscience*. 2009; 2:1–9.
- Luine VN. Sex differences in chronic stress effects on memory in rats. *Stress: The International Journal on the Biology of Stress*. 2002; 5:205–216.
- Lupien SJ, Gillin CJ, Hauger RL. Working memory is more sensitive than declarative memory to the acute effects of corticosteroids: A dose-response study in humans. *Behavioral Neuroscience*. 1999; 113:420–430. [PubMed: 10443770]
- Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature reviews. Neuroscience*. 2009; 10:434–445. [PubMed: 19401723]

- MacPherson A, Dinkel K, Sapolsky R. Glucocorticoids worsen excitotoxin-induced expression of pro-inflammatory cytokines in hippocampal cultures. *Experimental Neurology*. 2005; 194:376–383. [PubMed: 16022865]
- Magarinos AM, Li CJ, Gal Toth J, Bath KG, Jing D, Lee FS, McEwen BS. Effect of brain-derived neurotrophic factor haploinsufficiency on stress-induced remodeling of hippocampal neurons. *Hippocampus*. 2011; 21:253–264. [PubMed: 20095008]
- Maki PM. Critical window hypothesis of hormone therapy and cognition: A scientific update on clinical studies. *Menopause*. 2013; 20:695–709. [PubMed: 23715379]
- Maki PM, Dennerstein L, Clark M, Guthrie J, LaMontagne P, Fornelli D, Little D, Henderson VW, Resnick SM. Perimenopausal use of hormone therapy is associated with enhanced memory and hippocampal function later in life. *Brain Research*. 2011; 1379:232–243. [PubMed: 21078303]
- Maki PM, Zonderman AB, Resnick SM. Enhanced verbal memory in nondemented elderly woman receiving hormone-replacement therapy. *American Journal of Psychiatry*. 2001; 158:227–233. [PubMed: 11156805]
- Malagutti KS, da Silva AP, Braga HC, Mitozo PA, Soares dos Santos AR, Dafre AL, de Bem AF, Farina M. 17 β -estradiol decreases methylmercury-induced neurotoxicity in male mice. *Environmental Toxicology and Pharmacology*. 2009; 27:293–297. [PubMed: 21783955]
- Mannella P, Sanchez AM, Giretti MS, Genazzani AR, Simoncini T. Oestrogen and progestins differently prevent glutamate toxicity in cortical neurons depending on prior hormonal exposure via the induction of neural nitric oxide synthase. *Steroids*. 2009; 74:650–656. [PubMed: 19463685]
- Marin-Burgin A, Mongiat LA, Pardi MB, Schinder AF. Unique processing during a period of high excitation/inhibition balance in adult-born neurons. *Science*. 2012; 335:1238–1242. [PubMed: 22282476]
- Mateo JM. Inverted-u shape relationship between cortisol and learning in ground squirrels. *Neurobiology of Learning and Memory*. 2008; 89:582–590. [PubMed: 18164635]
- Mather M, Lighthall NR, Nga L, Gorlick MA. Sex differences in how stress affects brain activity during face viewing. *NeuroReport*. 2010; 21:933–937. [PubMed: 20808182]
- McClure RE, Barha CK, Galea LA. 17 β -estradiol, but not estrone, increases the survival and activation of new neurons in the hippocampus in response to spatial memory in adult female rats. *Hormones and behavior*. 2013; 63:144–157. [PubMed: 23063473]
- McCormick CM, Linkroum W, Sallinen BJ, Miller NW. Peripheral and central sex steroids have differential effects on the hpa axis of male and female rats. *Stress: The International Journal on the Biology of Stress*. 2002; 5:235–247.
- McEwen BS, Weiss JM, Schwartz LS. Selective retention of corticosterone by limbic structures in the rat brain. *Nature*. 1968; 220:911–912. [PubMed: 4301849]
- McGuinness OP, Fugiwara T, Murrell S, Bracy D, Neal D, O'Connor D, Cherrington AD. Impact of chronic stress hormone infusion on hepatic carbohydrate metabolism in the conscious dog. *Endocrinology & metabolism*. 1993; 28:E314–E322.
- McLaughlin KJ, Wilson JO, Harman J, Wright RL, Wiczorek L, Gomez J, Korol DL, Conrad CD. Chronic 17 β -estradiol or cholesterol prevents stress-induced hippocampal ca3 dendritic retraction in ovariectomized female rats: Possible correspondence between ca1 spine properties and spatial acquisition. *Hippocampus*. 2010:NA–NA.
- McMorris T, Sproule J, Turner A, Hale BJ. Acute, intermediate intensity exercise, and speed and accuracy in working memory tasks: A meta-analytical comparison of effects. *Physiology & behavior*. 2011; 102:421–428. [PubMed: 21163278]
- Mendelovitch A, Ritz M-F, Ros J, Langemann H, Gratzl O. 17 β -estradiol reduces cortical lesion size in the glutamate excitotoxicity model by enhancing extracellular lactate: A new neuroprotective pathway. *Brain Research*. 2001; 901:230–236. [PubMed: 11368971]
- Micheau J, Marighetto A. Acetylcholine and memory: A long, complex and chaotic but still living relationship. *Behavioural Brain Research*. 2011; 221:424–429. [PubMed: 21130809]
- Miller KJ, Conney JC, Rasgon NL, Fairbanks LA, Small GW. Mood symptoms and cognitive performance in women estrogen users and nonusers and men. *Journal of the American Geriatrics Society*. 2002; 50:1826–1830. [PubMed: 12410901]

- Minoshima S, Giordani B, Berent S, Frey KA, Foster NL, Kuhl DE. Metabolic reduction in the posterior cingulate cortex in very early alzheimer's disease. *Annals of Neurology*. 1997; 42:85–94. [PubMed: 9225689]
- Mizoguchi K, Ikeda R, Shoji H, Tanaka Y, Maruyama W, Tabira T. Aging attenuates glucocorticoid negative feedback in rat brain. *Neuroscience*. 2009; 159:259–270. [PubMed: 19141312]
- Mueller BR, Bale TL. Early prenatal stress impact on coping strategies and learning performance is sex dependent. *Physiology & Behavior*. 2007; 91:55–65. [PubMed: 17367828]
- Mueller BR, Bale TL. Sex-specific programming of offspring emotionality after stress early in pregnancy. *The Journal of Neuroscience*. 2008; 28:9055–9065. [PubMed: 18768700]
- Mulnard RA, Cotman CW, Kawas C, van Dyck CH, Sano M, Doody R, Koss E, Pfeiffer E, Jin S, Gamst A, Grundman M, Thomas R, Thal LJ. Estrogen replacement therapy for treatment of mild to moderate alzheimer disease: A randomized controlled trial. *Journal of the American Medical Association*. 2000; 283:1007–1015. [PubMed: 10697060]
- Musatov S, Chen W, Pfaff DW, Mobbs CV, Yang XJ, Clegg DJ, Kaplitt MG, Ogawa S. Silencing of estrogen receptor in the ventromedial nucleus of hypothalamus leads to metabolic syndrome. *Proceedings of the National Academy of Sciences*. 2007; 104:2501–2506.
- Nadeau S, Rivest S. Glucocorticoids play a fundamental role in protecting the brain during innate immune response. *The Journal of Neuroscience*. 2003; 23:5536–5544. [PubMed: 12843254]
- Newhouse PA, Dumas J, Wilkins H, Coderre E, Sites CK, Naylor M, Benkelfat C, Young SN. Estrogen treatment impairs cognitive performance after psychosocial stress and monoamine depletion in postmenopausal women. *Menopause*. 2010; 17:860–873. [PubMed: 20616673]
- Nielsen SE, Ahmed I, Cahill L. Postlearning stress differentially affects memory for emotional gist and detail in naturally cycling women and women on hormonal contraceptives. 2014
- Nielsen SE, Segal SK, Worden IV, Yim IS, Cahill L. Hormonal contraception use alters stress responses and emotional memory. *Biological Psychology*. 2013; 92:257–266. [PubMed: 23131613]
- Nilsen J, Brinton RD. Mechanism of estrogen-mediated neuroprotection: Regulation of mitochondrial calcium and bcl-2 expression. *Proceedings of the National Academy of Sciences*. 2003; 100:2842–2847.
- Nilsen J, Chen S, Brinton RD. Dual action of estrogen on glutamate-induced calcium signaling: Mechanisms requiring interaction between estrogen receptors and src/mitogen activated protein kinase pathway. *Brain Research*. 2002; 930:216–234. [PubMed: 11879813]
- Nilsen J, Chen S, Irwin RW, Iwamoto S, Brinton RD. Estrogen protects neuronal cells from amyloid beta-induced apoptosis via regulation of mitochondrial proteins and function. *BMC neuroscience*. 2006; 7:74. [PubMed: 17083736]
- O'Brien CA, Jia D, Plotkin LI, Bellido T, Powers CC, Stewart SA, Manolagas SC, Weinstein RS. Glucocorticoids act directly on osteoblasts and osteocytes to induce their apoptosis and reduce bone formation and strength. *Endocrinology*. 2004; 145:1835–1841. [PubMed: 14691012]
- O'Connor KA, Johnson JD, Hansen MK, Wieseler Frank JL, Maksimova E, Watkins LR, Maier SF. Peripheral and central proinflammatory cytokine response to a severe acute stressor. *Brain Research*. 2003; 991:123–132. [PubMed: 14575884]
- Oei NYL, Everaerd WTAM, Elzinga BM, van Well S, Bermond B. Psychosocial stress impairs working memory at high loads: An association with cortisol levels and memory retrieval. *Stress: The International Journal on the Biology of Stress*. 2006; 9:133–141.
- Öhman L, Nordin S, Bergdahl J, Birgander LS, Neely AS. Cognitive function in outpatients with perceived chronic stress. *Scandinavian journal of work, environment & health*. 2007:223–232.
- Owen AM, McMillan KM, Laird AR, Bullmore E. N-back working memory paradigm: A meta-analysis of normative functional neuroimaging studies. *Human Brain Mapping*. 2005; 25:46–59. [PubMed: 15846822]
- Park CR, Campbell AM, Diamond DM. Chronic psychosocial stress impairs learning and memory and increases sensitivity to yohimbine in adult rats. *Biological Psychiatry*. 2001; 50
- Pasquali R, Vicennati V, Cacciari M, Pagotto U. The hypothalamic-pituitary-adrenal axis activity in obesity and the metabolic syndrome. *Annals of the New York Academy of Sciences*. 2006; 1083:111–128. [PubMed: 17148736]

- Payne JD, Jackson ED, Hoscheidt S, Ryan L, Jacobs WJ, Nadel L. Stress administered prior to encoding impairs neutral but enhances emotional long-term episodic memories. *Learning & Memory*. 2007; 14:861–868. [PubMed: 18086830]
- Pefanco MA, Kenny AM, Kaplan RF, Kuchel G, Walsh S, Kleppinger A, Prestwood K. The effect of 3-year treatment with 0.25 mg/day of micronized 17 β -estradiol on cognitive function in older postmenopausal women. *Journal of the American Geriatrics Society*. 2007; 55:426–431. [PubMed: 17341247]
- Perlman WR, Webster MJ, Herman MM, Kleinman JE, Weickert CS. Age-related differences in glucocorticoid receptor mRNA levels in the human brain. *Neurobiology of Aging*. 2007; 28:447–458. [PubMed: 16540204]
- Pike CJ. Estrogen modulates neuronal bcl-xl expression and b-amyloid-induced apoptosis: Relevance to alzheimer's disease. *Journal of Neurochemistry*. 1999; 72:1552–1563. [PubMed: 10098861]
- Porcelli AJ, Cruz D, Wenberg K, Patterson MD, Biswal BB, Rypma B. The effects of acute stress on human prefrontal working memory systems. *Physiology & Behavior*. 2008; 95:282–289. [PubMed: 18692209]
- Protopopescu X, Butler T, Pan H, Root J, Altemus M, Polanecsky M, McEwen BS, Silbersweig D, Stern E. Hippocampal structural changes across the menstrual cycle. *Hippocampus*. 2008; 18:985–988. [PubMed: 18767068]
- Puder JJ, Freda PU, Goland RS, Wardlaw SL. Estrogen modulates the hypothalamic-pituitary-adrenal and inflammatory cytokine responses to endotoxin in women. *Journal of Clinical Endocrinology & Metabolism*. 2001; 86:2403–2408. [PubMed: 11397831]
- Purves, D.; Augustine, GJ.; Fitzpatrick, D.; Hall, WC.; LaMantia, A-S.; McNamara, JO.; Williams, SM. *Neuroscience*. 3rd ed.. Sinauer; Sunderland, MA, U.S.A.: 2004.
- Qin S, Hermans EJ, van Marle HJF, Luo J, Fernández G. Acute psychological stress reduces working memory-related activity in the dorsolateral prefrontal cortex. *Biological Psychiatry*. 2009; 66:25–32. [PubMed: 19403118]
- Rannevik G, Carlström K, Jeppsson S, Bjerre B, Svanberg L. A prospective long-term study in women from pre-menopause to post-menopause: Changing profiles of gonadotrophins, oestrogens, and androgens. *Maturitas*. 1986; 8:297–307. [PubMed: 2952867]
- Rapp SR, Espeland MA, Shumaker SA, Henderson VW, Brunner RL, Manson JE, Gass MLS, Stefanick ML, Lane DS, Hays J, Johnson KC, Coker LH, Dailey M, Bowen D. Effect of estrogen plus progestin on global cognitive function in postmenopausal women the women's health initiative memory study: A randomized controlled trial. *Journal of the American Medical Association*. 2003; 289:2663–2672. [PubMed: 12771113]
- Rasgon NL, Silverman D, Siddarth P, Miller K, Ercoli LM, Elman S, Lavretsky H, Huang SC, Phelps ME, Small GW. Estrogen use and brain metabolic change in postmenopausal women. *Neurobiology of Aging*. 2005; 26:229–235. [PubMed: 15582750]
- Rebuffe-Scrive M, Walsh UA, McEwen BS, Rodin J. Effect of chronic stress and exogenous glucocorticoids on regional fat distribution and metabolism. *Physiology & Behavior*. 1992; 52:583–590. [PubMed: 1409924]
- Resnick SM, Henderson VW. Hormone therapy and risk of alzheimer disease: A critical time. *Journal of the American Medical Association*. 2002; 288:2170–2172. [PubMed: 12413378]
- Resnick SM, Maki PM, Rapp SR, Espeland MA, Brunner R, Coker LH, Granek IA, Hogan P, Ockene JK, Shumaker SA. Effects of combination estrogen plus progestin hormone treatment on cognition and affect. *Journal of Clinical Endocrinology & Metabolism*. 2006; 91:1802–1810. [PubMed: 16522699]
- Reul JMHM, Rothuizen J, de Kloet ER. Age-related changes in the dog hypothalamic-pituitary-adrenocortical system: Neuroendocrine activity and corticosteroid receptors. *Journal of Steroid biochemistry and molecular biology*. 1991; 40:63–69. [PubMed: 1659883]
- Riegle GD. Chronic stress effects on adrenocortical responsiveness in young and aged rats. *Neuroendocrinology*. 1973; 11:1–10. [PubMed: 4349439]
- Riegle GD, Hess GD. Chronic and acute dexamethasone suppression of stress activation of the adrenal cortex in young and aged rats. *Neuroendocrinology*. 1972; 9:175–187. [PubMed: 5059845]

- Rivier C, Rivest S. Effect of stress on the activity of the hypothalamic-pituitary-gonadal axis: Peripheral and central mechanisms. *Biology of reproduction*. 1991; 45:523–532. [PubMed: 1661182]
- Roozendaal B. The basolateral amygdala interacts with the medial prefrontal cortex in regulating glucocorticoid effects on working memory impairment. *Journal of Neuroscience*. 2004; 24:1385–1392. [PubMed: 14960610]
- Rosmond R. Role of stress in the pathogenesis of the metabolic syndrome. *Psychoneuroendocrinology*. 2005; 30:1–10. [PubMed: 15358437]
- Sakaki M, Mather M. How emotion and reward processing change across the menstrual cycle. *social and personality psychology compass*. 2012; 6:1–17. [PubMed: 22737180]
- Salehi B, Cordero MI, Sandi C. Learning under stress: The inverted-u-shape function revisited. *Learning & Memory*. 2010; 17:522–530. [PubMed: 20884754]
- Sandstrom NJ, Williams CL. Memory retention is modulated by acute estradiol and progesterone replacement. *Behavioral Neuroscience*. 2001; 115:384–393. [PubMed: 11345963]
- Sapolsky RM. Chapter 2 glucocorticoids, hippocampal damage and the glutamatergic synapse. 1990; 86:13–23.
- Sapolsky RM. Glucocorticoids, stress, and their adverse neurological effects: Relevance to aging. *Experimental Gerontology*. 1999; 34:721–732. [PubMed: 10579633]
- Sapolsky RM. Glucocorticoids and hippocampal atrophy in neuropsychiatric disorders. *Archives of general psychiatry*. 2000; 57:925–935. [PubMed: 11015810]
- Sapolsky RM, Krey LC, McEwen BS. The adrenocortical stress-response in the aged male rat: Impairment of recovery from stress. *Experimental Gerontology*. 1983; 18:55–64. [PubMed: 6683660]
- Sapolsky RM, Krey LC, McEwen BS. Stress down-regulates corticosterone receptors in a site-specific manner in the brain. *Endocrinology*. 1984; 114:287–292. [PubMed: 6690273]
- Sapolsky RM, Krey LC, McEwen BS. The neuroendocrinology of stress and aging: The glucocorticoid cascade hypothesis. *Endocrine Reviews*. 1986; 7:284–301. [PubMed: 3527687]
- Saravia F, Beauquis J, Pietranera L, De Nicola AF. Neuroprotective effects of estradiol in hippocampal neurons and glia of middle age mice. *Psychoneuroendocrinology*. 2007; 32:480–492. [PubMed: 17459595]
- Scholz U, La Marca R, Nater UM, Aberle I, Ehlert U, Hornung R, Martin M, Kliegel M. Go no-go performance under psychosocial stress: Beneficial effects of implementation intentions. *Neurobiology of Learning and Memory*. 2009; 91:89–92. [PubMed: 18817886]
- Schoofs D, Preuß D, Wolf OT. Psychosocial stress induces working memory impairments in an n-back paradigm. *Psychoneuroendocrinology*. 2008; 33:643–653. [PubMed: 18359168]
- Schoofs D, Wolf OT, Smeets T. Cold pressor stress impairs performance on working memory tasks requiring executive functions in healthy young men. *Behavioral Neuroscience*. 2009; 123:1066–1075. [PubMed: 19824773]
- Seale J, Wood S, Atkinson H, Bate E, Lightman S, Ingram C, Jessop D, Harbuz M. Gonadectomy reverses the sexually diergic patterns of circadian and stress-induced hypothalamic-pituitary-adrenal axis activity in male and female rats. *Journal of neuroendocrinology*. 2004a; 16:516–524. [PubMed: 15189326]
- Seale J, Wood S, Atkinson H, Harbuz M, Lightman S. Gonadal steroid replacement reverses gonadectomy-induced changes in the corticosterone pulse profile and stress-induced hypothalamic-pituitary-adrenal axis activity of male and female rats. *Journal of neuroendocrinology*. 2004b; 16:989–998. [PubMed: 15667454]
- Seckl JR, Olsson T. Glucocorticoid hypersecretion and the age-impaired hippocampus: Cause or effect? *Journal of endocrinology*. 1995; 145:201–211. [PubMed: 7616153]
- Selye H. The effect of adaptation to various damaging agents on the female sex organs in the rat. *Endocrinology*. 1939; 25:615–624.
- Shaywitz SE, Naftolin F, Zelterman D, Marchione KE, Holahan JM, Palter SF, Shaywitz BA. Better oral reading and short-term memory in midlife, postmenopausal women taking estrogen. *Menopause*. 2003; 10:420–426. [PubMed: 14501603]

- Shors TJ. Learning during stressful times. *Learning & Memory*. 2004; 11:137–144. [PubMed: 15054128]
- Shumaker SA, Legault C, Rapp SR, Thal L, Wallace RB, Ockene JK, Hendrix SL, Jones BN, Assaf AR, Jackson RD, Kotchen J, Wassertheil-Smoller S, Wactawski-Wende J. Estrogen plus progestin and the incidence of dementia and mild cognitive impairment in postmenopausal women. The women's health initiative memory study: A randomized controlled trial. *Journal of the American Medical Association*. 2003; 289:2651–2662. [PubMed: 12771112]
- Simpson E. Sources of estrogen and their importance. *The Journal of steroid biochemistry and molecular biology*. 2003; 86:225–230. [PubMed: 14623515]
- Simpson ER. Aromatization of androgens in women: Current concepts and findings. *Fertility and sterility*. 2002; 77:6–10.
- Singer CA, Rogers KL, Dorsa DM. Modulation of bcl-2 expression: A potential component of estrogen protection in nt2 neurons. *Neuroreport*. 1998; 9:2565–2568. [PubMed: 9721933]
- Smeets T, Jelicic M, Merckelbach H. The effect of acute stress on memory depends on word valence. *International Journal of Psychophysiology*. 2006; 62:30–37. [PubMed: 16388863]
- Smith YR, Bowen L, Love TM, Berent-Spillion A, Frey KA, Persad CC, Reame NK, Koeppe RA, Zubieta JK. Early initiation of hormone therapy in menopausal women is associated with increased hippocampal and posterior cingulate cholinergic activity. *Journal of Clinical Endocrinology & Metabolism*. 2011; 96:E1761–E1770. [PubMed: 21865354]
- Smith YR, Giordani B, Lajiness-O'Neill R, Zubieta J-K. Long-term estrogen replacement is associated with improved nonverbal memory and attentional measures in postmenopausal women. *Menopause*. 2001; 76:1101–1107.
- Smith YR, Zubieta J-K. Neuroimaging of aging and estrogen effects on central nervous system physiology. *Fertility and Sterility*. 2001; 76:651–659. [PubMed: 11591393]
- Sowers M, Crutchfield M, Bandekar R, Randolph JF, Shapiro B, Schork MA, Jannausch M. Bone mineral density and its change in pre- and perimenopausal white women: The michigan bone health study. *Journal of Bone and Mineral Research*. 1998; 13:1134–1140. [PubMed: 9661077]
- Starcke K, Wolf OT, Markowitsch HJ, Brand M. Anticipatory stress influences decision making under explicit risk conditions. *Behavioral neuroscience*. 2008; 122:1352. [PubMed: 19045954]
- Stein-Behrens BA, Elliott EM, Miller CA, Schilling JW, Newcombe R, Sapolsky RM. Glucocorticoids exacerbate kainic acid-induced extracellular accumulation of excitatory amino acids in the rat hippocampus. *Journal of Neurochemistry*. 1992; 58:1730–1735. [PubMed: 1560229]
- Suzuki S, Lund TD, Price RH, Handa RJ. Sex differences in the hypothalamo-pituitary-adrenal axis: Novel roles for androgen and estrogen receptors. *Recent research developments in endocrinology*. 2001:69–86.
- Symonds CS, McKie S, Elliott R, William Deakin JF, Anderson IM. Detection of the acute effects of hydrocortisone in the hippocampus using pharmacological fmri. *European Neuropsychopharmacology*. 2012
- Takuma K, Matsuo A, Himeno Y, Hoshina Y, Ohno Y, Funatsu Y, Arai S, Kamei H, Mizoguchi H, Nagai T. 17 β -estradiol attenuates hippocampal neuronal loss and cognitive dysfunction induced by chronic restraint stress in ovariectomized rats. *Neuroscience*. 2007; 146:60–68. [PubMed: 17320294]
- Tata DA, Anderson BJ. The effects of chronic glucocorticoid exposure on dendritic length, synapse numbers and glial volume in animal models: Implications for hippocampal volume reductions in depression. *Physiol Behav*. 2010; 99:186–193. [PubMed: 19786041]
- Tetsuka M. Actions of glucocorticoid and their regulatory mechanisms in the ovary. *Animal Science Journal*. 2007; 78:112–120.
- Tilbrook A, Turner A, Clarke I. Effects of stress on reproduction in non-rodent mammals: The role of glucocorticoids and sex differences. *Reviews of reproduction*. 2000; 5:105–113. [PubMed: 10864855]
- Tombaugh GC, Yang SH, Swanson RA, Sapolsky RM. Glucocorticoids exacerbate hypoxic and hypoglycemic hippocampal injury in vitro: Biochemical correlates and a role for astrocytes. *Journal of Neurochemistry*. 1992; 59:137–146. [PubMed: 1613495]

- Tynan RJ, Beynon SB, Hinwood M, Johnson SJ, Nilsson M, Woods JJ, Walker FR. Chronic stress-induced disruption of the astrocyte network is driven by structural atrophy and not loss of astrocytes. *Acta Neuropathol.* 2013; 126:75–91. [PubMed: 23512378]
- Tynan RJ, Naicker S, Hinwood M, Nalivaiko E, Buller KM, Pow DV, Day TA, Walker FR. Chronic stress alters the density and morphology of microglia in a subset of stress-responsive brain regions. *Brain, Behavior, and Immunity.* 2010; 24:1058–1068.
- Uno H, Tarara R, Else JG, Suleman MA, Sapolsky RM. Hippocampal damage associated with prolonged and fatal stress in primates. *The Journal of Neuroscience.* 1989; 9:1705–1711. [PubMed: 2723746]
- Vaucher E, Reymond I, Najaffe R, Kar S, Quirion R, Miller MM, Franklin KBJ. Estrogen effects on object memory and cholinergic receptors in young and old female mice. *Neurobiology of Aging.* 2002; 23:87–95. [PubMed: 11755023]
- Velázquez-Zamora DA, Garcia-Segura LM, González-Burgos I. Effects of selective estrogen receptor modulators on allocentric working memory performance and on dendritic spines in medial prefrontal cortex pyramidal neurons of ovariectomized rats. *Hormones and Behavior.* 2012
- Watanabe Y, Gould E, McEwen BS. Stress induces atrophy of apical dendrites of hippocampal ca3 pyramidal neurons. *Brain Research.* 1992; 588:341–343. [PubMed: 1393587]
- Weaver CE Jr, Park-Chung M, Gibbs TT, Farb DH. 17 β -estradiol protects against nmda-induced excitotoxicity by direct inhibition of nmda receptors. *Brain Research.* 1997; 761:338–341. [PubMed: 9252035]
- Weiser MJ, Handa RJ. Estrogen impairs glucocorticoid dependent negative feedback on the hypothalamic–pituitary–adrenal axis via estrogen receptor alpha within the hypothalamus. *Neuroscience.* 2009; 159:883–895. [PubMed: 19166915]
- Wellman CL. Dendritic reorganization in pyramidal neurons in medial prefrontal cortex after chronic corticosterone administration. *Journal of Neurobiology.* 2001; 49:245–253. [PubMed: 11745662]
- Wilkinson CW, Petrie EC, Murray SR, Colasurdo EA, Raskind MA, Peskind ER. Human glucocorticoid feedback inhibition is reduced in older individuals: Evening study. *Journal of Clinical Endocrinology & Metabolism.* 2001; 86:545–550. [PubMed: 11158007]
- Wilson IA, Puolivali J, Heikkinen T, Riekkinen P Jr. Estrogen and nmda receptor antagonism: Effects upon reference and working memory. *European Journal of Pharmacology.* 1999; 381:93–99. [PubMed: 10554875]
- Wolf OT, Kirschbaum C. Endogenous estradiol and testosterone levels are associated with cognitive performance in older women and men. *Hormones and Behavior.* 2002; 41:259–266. [PubMed: 11971659]
- Wolf OT, Kudielka B, Hellhammer DH, Torber S, McEwen BS, Kirschbaum C. Two weeks of transdermal estradiol treatment in postmenopausal elderly women and its effect on memory and mood: Verbal memory changes are associated with the treatment induced estradiol levels. *Psychoneuroendocrinology.* 1999; 24:727–741. [PubMed: 10451908]
- Wolf OT, Minnebusch D, Daum I. Stress impairs acquisition of delay eyeblink conditioning in men and women. *Neurobiology of Learning and Memory.* 2009; 91:431–436. [PubMed: 19049887]
- Woolley CS, Gould E, Frankfurt M, McEwen BS. Naturally occurring fluctuation in dendritic spine density on adult hippocampal pyramidal neurons. *The Journal of Neuroscience.* 1990a; 10:4035–4039. [PubMed: 2269895]
- Woolley CS, Gould E, McEwen BS. Exposure to excess glucocorticoids alters dendritic morphology of adult hippocampal pyramidal neurons. *Brain Research.* 1990b; 531:225–231. [PubMed: 1705153]
- Woolley CS, McEwen BS. Estradiol mediates fluctuation in hippocampal synapse density during the estrous cycle in the adult rat. *The Journal of Neuroscience.* 1992; 12:2549–2554. [PubMed: 1613547]
- Woolley CS, McEwen BS. Roles of estradiol and progesterone in regulation of hippocampal dendritic spine density during the estrous cycle in the rat. *The Journal of Comparative Neurology.* 1993; 336:293–306. [PubMed: 8245220]

- Woolley CS, McEwen BS. Estradiol regulates hippocampal dendritic spine density via an *n*-methyl-d-aspartate receptor-dependent mechanism. *The Journal of Neuroscience*. 1994; 14:7680–7687. [PubMed: 7996203]
- Yaffe K, Vittinghoff E, Ensrud KE, Johnson KC, Diem S, Hanes V, Grady D. Effects of ultra-low-dose transdermal estradiol on cognition and health-related quality of life. *Archives of Neurology*. 2006; 63:945–950. [PubMed: 16831962]
- Yao J, Chen S, Cadenas E, Brinton RD. Estrogen protection against mitochondrial toxin-induced cell death in hippocampal neurons: Antagonism by progesterone. *Brain research*. 2011; 1379:2–10. [PubMed: 21134358]
- Yerkes RM, Dodson JD. The relation of strength of stimulus to rapidity of habit-formation. *Journal of Comparative Neurology and Psychology*. 1908; 18:459–482.
- Young AH, Sahakian BJ, Robbins TW, Cowen PJ. The effects of chronic administration of hydrocortisone on cognitive function in normal male volunteers. *Psychopharmacology*. 1999; 145:260–266. [PubMed: 10494574]
- Zhao L, Wu T.-w. Brinton RD. Estrogen receptor subtypes alpha and beta contribute to neuroprotection and increased bcl-2 expression in primary hippocampal neurons. *Brain Research*. 2004; 1010:22–34. [PubMed: 15126114]

Highlights

- Exposure to glucocorticoids can be detrimental to cognition and neural integrity
- Estradiol exerts protective effects on cognition and neural integrity
- Estradiol also can mitigate the magnitude of the stress response
- Protective effects of estradiol may mitigate stress-induced glucocorticoid damage
- This may be an additional benefit of post-menopausal estradiol treatment

TABLE 1

Effects of Glucocorticoids and Estradiol in the Body

Metabolic Syndrome and Related Indices in Animal Models			
Species	Index	Effect of, or relationship with, Glucocorticoids	Effect of, or relationship with, Estradiol
Human	Metabolic Syndrome	↑ (Pasquali et al., 2006; Rosmond, 2005)	/
	Abdominal visceral fat	↑ (Green et al., 2004)	/
	Triglyceride, Glucose, and Insulin	/	↓ (Krotkiewski et al., 1983)
	Bone Mineral Density	/	↑ (Delmas et al., 1997; Felson et al., 1993; Sowers et al., 1998)
Rat & Mouse	Metabolic Syndrome	↑ (Musatov et al., 2007)	/
	Intraabdominal fat	↑ (Rebuffe-Scrive et al., 1992)	/
	Insulin	↑ (Rebuffe-Scrive et al., 1992)	/
	Osteoblasts	↓ (O'Brien et al., 2004)	/
	Osteocytes	↓ (O'Brien et al., 2004)	/
	Osteoclasts	↑ (Jia et al., 2006)	/
Dog	Glucose and Insulin	/	↓ (McGuinness et al., 1993)

"/" indicates unknown, or not investigated in cited articles

TABLE 2

Countervailing effects of glucocorticoids and estradiol on working memory

Working Memory			
Species	Task	Effect of Stress or Glucocorticoid On Performance	Effect of Estradiol On Performance
Human	Digit Ordering	/	↑ (Females) (Duff and Hampson, 2000; Krug et al., 2006)
	Digit Span	↓ (Males & Females) (Elzinga and Roelofs, 2005; Schoofs et al., 2009)	↑ (Females) (Miller et al., 2002)
	N-Back	↓ (Males) (Schoofs et al., 2008)	/
	Reading Span	↓ (Males) (Luethi et al., 2009)	/
	Sternberg Item Recognition	↓ (Males) (Duncko et al., 2009; Lupien et al., 1999; Oei et al., 2006)	/
	Visuospatial Sketchpad	↓ (Males) (Young et al., 1999)	/
	Trail Making Test B	↓ (Males & Females) (Ohman et al., 2007)	/
Rat	Delayed Match-to-Place & Delayed Match-to-Position	/	↑ (Females) (Gibbs, 1999; Sandstrom and Williams, 2001)
	Radial Arm Maze & Delayed Win-Shift Task	↓ (Males) (Arbel et al., 1994; Butts et al., 2011; Diamond et al., 1996)	↑ (Females) (Bimonte and Denenberg, 1999; Daniel et al., 1997; Fader et al., 1999; Wilson et al., 1999)
	Radial Arm Water Maze & Morris Water Maze	↓ (Males) (Cerqueira et al., 2007; Park et al., 2001)	/
	Y-Maze & T-maze (delayed alternation)	↓ (Males) (Coburn-Litvak et al., 2003; Kleen et al., 2006; Lee and Goto, 2015; Roozendaal, 2004)	/

"/" indicates unknown, or not investigated in cited articles

TABLE 3

Countervailing effects of glucocorticoids and estradiol on executive function and other cognitive processes

Executive Processes			
Species	Task	Effect of Stress or Glucocorticoid On Performance	Effect of Estradiol On Performance
Human	Cognitive Flexibility, Mental Flexibility, & Planning	↓ (Males & Females) (Alexander et al., 2007)	↑ (Females) (Elsabagh et al., 2007)
	Go No-Go	↓ (Males) (Scholz et al., 2009)	/
	Stroop Task	/	↑ (Females) (Baker et al., 2012; Krug et al., 2006; Wolf and Kirschbaum, 2002)
	Verbal Fluency	/	↑ (Females) (Miller et al., 2002)
	Mental Arithmetic	↓ (Males) (Al'Absi et al., 2002)	/
	Attention Shift	↓ (Males & Females) (Liston et al., 2009)	/
	Decision Making	↓ (Sex not stated) (Starcke et al., 2008)	/
	Wisconsin Card Sorting Task	↓ (Males & Females) (Egeland et al., 2005)	
Rat	Reversal Learning	↓ (Males) (Cerqueira et al., 2007)	/
Other Cognitive Processes			
Human	Delayed Recall	/	↑ (Females) (Baker et al., 2012)
	Paired Associates	↓ (Males) (Young et al., 1999)	/
	California Verbal Learning Test	/	↓ (Females) (Resnick et al., 2006)
	Modified Mini Mental State Exam	/	↓ (Females) (Espeland et al., 2004; Rapp et al., 2003)
Rat	Object Recognition	/	↑ (Females) (Vaucher et al., 2002)
	Reference Memory	↓ (Males) (Cerqueira et al., 2007)	/

"/" indicates unknown, or not investigated in cited articles

TABLE 4

Effects of Glucocorticoids and Estradiol on Neuronal Viability

Effects of Glucocorticoids and Estradiol on Neuronal Survival After Exposure to Insults				
Species	Neural Region or Neural Tissue	Insult	Viability after glucocorticoid with other insults	Viability after estradiol with other insults
Mice	Cerebellum	Methylmercury	/	↑ (Males) (Malagutti et al., 2009)
Rat	Fetal Hippocampal Cultures	A β	↓ (Behl et al., 1997)	↑ (Brinton et al., 2000; Chen et al., 2006)
		Hypoxia	↓ (Tombaugh et al., 1992)	/
		Hypoglycemia	↓ (Tombaugh et al., 1992)	/
		Hydrogen Peroxide	/	↑ (Brinton et al., 2000)
		Glutamate Excitotoxicity	/	↑ (Brinton et al., 2000)
	Hippocampus	Kainic Acid	↓ (Males) (Stein-Behrens et al., 1992)	↑ (Females) (Hoffman et al., 2003)
		Quinolinic Acid	/	↑ (Males) (Kuroki et al., 2001)
Rhesus Monkey	Prefrontal Cortex	Aging	/	↑ (Females) (Hao et al., 2007)

"/" indicates unknown, or not investigated in cited articles

TABLE 5

Effects of Glucocorticoids and Estradiol on Neuronal Morphology

Effects of Glucocorticoids and Estradiol on Neuronal Morphology				
Species	Neural Region or Neural Tissue	Outcome of Interest	Effect of Glucocorticoid or Stress on Neural Tissue	Effect of Estradiol on Neural Tissue
Rat	Various regions	Inflammatory Markers	↑ (Males) (Blandino Jr et al., 2006; MacPherson et al., 2005; O'Connor et al., 2003; Tynan et al., 2010)	/
	Hippocampus or Fetal Hippocampal Cultures	Dendrite Length	↓ (Males & Females) (Galea et al., 1997; McLaughlin et al., 2010; Watanabe et al., 1992; Woolley et al., 1990b)	↑ (Fetal Cultures) (Brinton et al., 2000; Chen et al., 2006)
		Dendrite Branch Points	↓ (Males & Females) (Galea et al., 1997; McLaughlin et al., 2010; Watanabe et al., 1992; Woolley et al., 1990b)	↑ (Fetal Cultures) (Brinton et al., 2000; Chen et al., 2006)
		Dendrite Spine Density	/	↑ (Females) (Gould et al., 1990; Woolley et al., 1990a; Woolley and McEwen, 1993)
		Synaptic Density	/	↑ (Females) (Woolley and McEwen, 1992)
		Neuron Number	↓ (Males) (Haynes et al., 2003)	↑ (Males & Fetal Cultures) (Brinton et al., 2000; Chen et al., 2006; Saravia et al., 2007)
	Striatum	Neuron Number	↓ (Males) (Haynes et al., 2003)	/
	Prefrontal Cortex	Dendrite Length	↓ (Males) (Brown et al., 2005; Cook and Wellman, 2004; Wellman, 2001)	/
		Dendrite Branch Points	↓ (Males) (Brown et al., 2005; Cook and Wellman, 2004; Wellman, 2001)	/
Vervet Monkey	Hippocampus	Neuron Number	↓ (Males) (Uno et al., 1989)	/
		Dendrite Branch Points	↓ (Males) (Uno et al., 1989)	/
Human	Hippocampus	Area Volume	/	↑ (Females) (Protopopescu et al., 2008)

"/" indicates unknown, or not investigated in cited articles

TABLE 6

Effects of Glucocorticoids and Estradiol on Brain Activation

Effects of Glucocorticoids and Estradiol on Brain Activation				
Species	Neural Region or Neural Tissue	Outcome of Interest	Effects of Glucocorticoids on Brain Activation	Effects of Estradiol on Brain Activation
Human	Prefrontal Cortex	Increased BOLD activation during working memory task	↓ (Males & Females) (Qin et al., 2009; Symonds et al., 2012)	/
	Hippocampus	Decreased BOLD activation during working memory task	↓ (Females) (Qin et al., 2009; Symonds et al., 2012)	/
		BOLD activation during a verbal recognition task	/	↑ (Females) (Maki et al., 2011)
		Cholinergic activity	/	↑ (Females) (Smith et al., 2011)

“/” indicates unknown, or not investigated in cited articles

TABLE 7

Estrogenic Modulation of Glucocorticoid Effects

Modulation of the Glucocorticoid or Stress effects on Neural Tissue by Estradiol			
Species	Neural Region	Effect of Glucocorticoid Exposure	Modulated Effect by Estradiol
Rat	Hippocampus	↓ (Males) (Haynes et al., 2003)	Blocked (Haynes et al., 2003)
		↓LTP (Males) (Foy et al., 2008)	Attenuated (Foy et al., 2008)
		↓ Dendritic Complexity (Males & OVX Females) (Galea et al., 1997; McLaughlin et al., 2010)	Attenuated or Blocked (Intact Females & OVX+E2 replacement) (Galea et al., 1997; McLaughlin et al., 2010)
		↓ Cell Number (OVX Females) (Takuma et al., 2007)	Blocked (Takuma et al., 2007)
	Striatum	↓ (Males) (Haynes et al., 2003)	Blocked (Haynes et al., 2003)
	Prefrontal Cortex	↓ (OVX Females) (Garrett and Wellman, 2009)	Blocked (Garrett and Wellman, 2009)

“/” indicates unknown, or not investigated in cited articles