System-Level Planning, Coordination, and Communication

Care of the Critically Ill and Injured During Pandemics and Disasters: CHEST Consensus Statement

Jeffrey R. Dichter, MD; Robert K. Kanter, MD; David Dries, MD; Valerie Luyckx, MD; Matthew L. Lim, MD; John Wilgis, MD; Michael R. Anderson, MD, MBA; Babak Sarani, MD; Nathaniel Hupert, MD; Ryan Mutter, MD; Asha V. Devereaux, MD, MPH, FCCP; Michael D. Christian, MD, FRCPC, FCCP; and Niranjana Kissoon, MBBS, FRCPC; on behalf of the Task Force for Mass Critical Care

BACKGROUND: System-level planning involves uniting hospitals and health systems, local/regional government agencies, emergency medical services, and other health-care entities involved in coordinating and enabling care in a major disaster. We reviewed the literature and sought expert opinions concerning system-level planning and engagement for mass critical care due to disasters or pandemics and offer suggestions for system-planning, coordination, communication, and response. The suggestions in this chapter are important for all of those involved in a pandemic or disaster with multiple critically ill or injured patients, including front-line clinicians, hospital administrators, and public health or government officials.

METHODS: The American College of Chest Physicians (CHEST) consensus statement development process was followed in developing suggestions. Task Force members met in person to develop nine key questions believed to be most relevant for system-planning, coordination, and communication. A systematic literature review was then performed for relevant articles and documents, reports, and other publications reported since 1993. No studies of sufficient quality were identified upon which to make evidence-based recommendations. Therefore, the panel developed expert opinion-based suggestions using a modified Delphi process.

RESULTS: Suggestions were developed and grouped according to the following thematic elements: (1) national government support of health-care coalitions/regional health authorities (HC/RHAs), (2) teamwork within HC/RHAs, (3) system-level communication, (4) system-level surge capacity and capability, (5) pediatric patients and special populations, (6) HC/RHAs and networks, (7) models of advanced regional care systems, and (8) the use of simulation for preparedness and planning.

CONCLUSIONS: System-level planning is essential to provide care for large numbers of critically ill patients because of disaster or pandemic. It also entails a departure from the routine, independent system and involves all levels from health-care institutions to regional health authorities. National government support is critical, as are robust communication systems and advanced planning supported by realistic exercises.

CHEST 2014; 146(4_Suppl):e87S-e102S

ABBREVIATIONS: HC/RHA = health-care coalition/regional health authority

Summary of Suggestions

National Government Support of Health-care Coalitions/Regional Health Authorities—Policy

1a. We suggest political leadership at national levels should support health-care preparedness through financial assistance, support of market driven incentives, and preparedness requirements to health-care coalitions/regional health authorities (HC/RHAs).

1b. We suggest national governments should support the development of responsive and nimble disaster/pandemic research processes that can both organize and assess information from prior disasters/pandemics, acquire real-time data in an ongoing one to provide situational awareness, and which can also learn from and support international disaster relief efforts.

1c. We suggest national, state/province/regional, and city/district governments should:

- **Working with health-care experts and leadership, develop formal legal disaster/pandemic activation mechanisms to initiate, implement, and support disaster/pandemic plans and standards of care for HC/RHAs and healthcare professionals; and legally initiate step down termination procedures and processes as conditions and criteria warrant in the recovery phase**
- **Work with health-care experts and leadership in the greater health-care community to develop and refine specific “trigger” criteria for formal legal activation and step down termination procedures and pro-**

cesses of disaster/pandemic plans and standards of care.

1d. We suggest local governments and government agencies should be formal partners in their local health-care coalition(s), and be actively engaged with their ongoing preparedness and response activities.

Teamwork Within HC/RHAs—Foundational Principles

2. We suggest health-care coalition partners should work together, with the following objectives:

2a. HC/RHA clinical and administrative leaders from all partners meet together on a routine, scheduled basis. Clinician leaders must include critical care medicine experts.

2b. HC/RHA clinical and administrative leaders from all partners work together at least yearly with primary focus on developing and updating joint disaster/pandemic preparedness plans based on likely events (Hazard Vulnerability Analyses).

2c. HC/RHA clinical and administrative leaders from all partners jointly practice activation and implementation of disaster/pandemic plans and standards of care through exercises.

2d. HC/RHA partners activate their communication and collaboration mechanisms for virtually all actual or potential surge events, or unusual or large scale planned or unplanned events requiring cooperation, to ensure optimal responses and enable experience working together.

Revision accepted May 1, 2014; originally published Online First August 21, 2014.

AFFILIATIONS: From Allina Health (Dr Dichter), Minneapolis, MN; Aurora Healthcare (Dr Dichter), Milwaukee, WI; SUNY Upstate Medical University (Dr Kanter), Syracuse, NY; Columbia University (Dr Kanter), New York, NY; HealthPartners Medical Group (Dr Dries), University of Minnesota, Minneapolis, MN; International Society of Nephrology (Dr Luyckx), University of Alberta, Edmonton, AB, Canada; Office of Global Affairs (Dr Lim), Department of Health and Human Services, Washington, DC; Florida Hospital Association (Dr Wilgis), Orlando, FL; Case Western Reserve University (Dr Anderson), Cleveland, OH; George Washington University (Dr Sarani), Washington, DC; Weill Medical College (Dr Hupert) Cornell University, New York, NY; New York-Presbyterian Hospital (Dr Hupert), New York, NY; Agency for Healthcare Quality Research (Dr Mutter), Washington, DC; Sharp Hospital (Dr Devereaux), Coronado, CA; Royal Canadian Medical Service (Dr Christian), Canadian Armed Forces and Mount Sinai Hospital, Toronto, ON, Canada; and BC Children's Hospital and Sunnyville Health Centre (Dr Kisson), University of British Columbia, Vancouver, BC, Canada.

FUNDING/SUPPORT: This publication was supported by the Cooperative Agreement Number 1U90TP00591-01 from the Centers of Disease

Control and Prevention, and through a research sub award agreement through the Department of Health and Human Services [Grant 1 - HPPEP070013-01-00] from the Office of Preparedness of Emergency Operations. In addition, this publication was supported by a grant from the University of California–Davis.

COI grids reflecting the conflicts of interest that were current as of the date of the conference and voting are posted in the online supplementary materials.

DISCLAIMER: American College of Chest Physicians guidelines and consensus statements are intended for general information only, are not medical advice, and do not replace professional care and physician advice, which always should be sought for any medical condition. The complete disclaimer for this consensus statement can be accessed at <http://dx.doi.org/10.1378/chest.1464S1>.

CORRESPONDENCE TO: Jeffrey R. Dichter, MD, Allina Health, 550 Osborne Rd NE, Fridley, MN 55432; e-mail: jeffrey.dichter@allina.com

© 2014 AMERICAN COLLEGE OF CHEST PHYSICIANS. Reproduction of this article is prohibited without written permission from the American College of Chest Physicians. See online for more details.

DOI: 10.1378/chest.14-0738

2e. HC/RHAs identify clinical experts to oversee and address the needs of specific populations, especially pediatrics, and also specialty populations such as trauma, burns, oncologic, etc.

2f. HC/RHA clinical and administrative leadership should be defined by position, not specific personnel, consistent with Incident Command System (ICS) nomenclature or equivalent, and designed with appropriate redundancy.

System-Level Communication—Foundational Principles

3a. We suggest HC/RHAs should have secure online and/or published directories for all partners' clinical and administrative leadership, with emergency contact information (phone numbers, e-mail addresses, pagers, cell phone texting preferences, other means) and current call schedules.

3b. We suggest HC/RHA's should have defined communication vehicles which may include (but are not limited to): dedicated secure health-care coalition web sites; conference call lines and teleconferencing technologies (eg, Skype, others); hospital phones (land lines and cell phones); pagers, hand-held walkie-talkies, ham radios, or other similar means of communication; telemedicine technologies, such as E-ICU, integrated into their disaster plans.

3c. We suggest HC/RHA partners should attempt to routinely use those agreed upon communication vehicles when working together.

3d. We suggest all agreed upon communication vehicles should be tested on a scheduled basis, with objective criteria to validate the test.

3e. We suggest the choice of communication vehicles and testing may be based on likely disaster/pandemic events (Hazard Vulnerability Analyses), and/or other appropriate considerations.

3f. We suggest developing defined disaster/pandemic plans for monitoring and leveraging popular social media (eg, Twitter, Facebook, others) during all actual or potential surge events, or unusual or large scale planned or unplanned events requiring cooperation, as both a means for gathering and transmitting information, as appropriate.

3g. We suggest HC/RHAs should have defined communication tools designated for each level of

organizational leadership, which should be consistent with ICS structure or equivalent.

System-Level Surge Capacity and Capability

4a. We suggest HC/RHA surge objectives should be consistent with individual hospital surge goals and include the capability to surge to:

- Up to 200% above routine maximal capacity based on the nature and severity of the disaster (contingency to crisis)
- Up to the limit of the total number of ventilators available to coalition partners.
- Up to projected patient loads in a slow onset, slow evolving disaster.

4b. We suggest HC/RHAs should be able to monitor and track their defined surge capacity supplies and equipment, ideally "real time" and electronically, with the intent of being able to use all HC/RHA assets. These supplies and equipment may include identified caches of important medications or equipment, and bed availability among partners.

4c. We suggest HC/RHAs should have the ability to track the number of available ICU capable personnel ("force multipliers") and other designated specialist "resources" (eg, pediatric and special populations) through their partner hospitals. Partners with telemedicine capability (such as tele-ICU's) should have plans for how to use this resource to optimize the use of pediatric and specialty expertise across hospitals served by the telemedicine resource.

4d. We suggest HC/RHAs should have defined policies and procedures for emergency privileging for all health-care professionals designated as coalition resources.

4e. We suggest fair and adequate reimbursement for expenditures and loss of revenue related to delivery of acute critical care services during a disaster or pandemic must be ensured. This should include the guarantee of payments from governmental sources, as well as by insurance companies and other payers of health-care services.

Pediatric Patients and Specialty Populations

5a. We suggest HC/RHAs have identified, and be familiar with, the following pediatric disaster/pandemic designated resources including, but not limited to:

- Pediatric consultative specialists available by dedicated phone line support and/or dedicated video or telemedicine consultation.
- Designated pediatric surge personnel (eg, pediatric hospitalists, others) available to non-pediatric hospitals and health systems to support surge in contingency or crisis level events, with a defined plan for how to activate this resource when needed.
- Identified pediatric capable transport resources for allocation and matching of pediatric patients to available HC/RHA pediatric resources.
- Knowledge of available key supplies, medications, and other pediatric assets; location of these assets with a defined process for how they may be accessed urgently; and ability to monitor when asset reserves fall below a defined critical threshold.
- Pediatric educational resources. If web-based, they should be found on HC/RHA websites, or with links to appropriate resources. If published, resources should be readily available to all partners.

5b. We suggest HC/RHAs should have plans to provide care for specialty populations routinely found in their catchment area or region in parallel as described for pediatrics. Resources should include consultative services, potential surge personnel, transport resources, specialty supplies/medications, and educational resources. These populations include but are not limited to trauma, nephrology, burns, oncologic patients.

5c. Health-care coalitions, health systems, and hospitals identify patients with high-level chronic disease care needs, such as a home ventilator, home oxygen, chronic dialysis, and work to ensure their needs are met at home to help prevent these patients from having to seek assistance at hospitals.

HC/RHAs and Networks

6a. We suggest during a disaster requiring transfer of patients, whether from emergency medicine departments or inpatient areas, transferring partners may have initial choice of where patients are referred based on traditional referral patterns. However, HC/RHA leadership must oversee this process, and be able to intercede as both a resource and with the authority to redirect transfers based on anticipated or actual events. Defined health-care coalition coordination processes and transfer resources should be planned and identified ahead of time.

6b. Health-care coalitions should designate neighboring health-care coalitions as potential partners

during a contingency or crisis event, and have readily available leadership contact information, and knowledge of these potential partners' size and capabilities.

Models of Advanced Regional Care Systems

7. Advanced Regional Care Systems instituted within large hospitals, and across hospitals, health systems and HC/RHAs, will have the greatest chance for success if they are established with the following goals:

- Clear and transparent objectives for what those programs are to accomplish, and the programs are well integrated and accepted across their hospital and health system partners.
- Have administrative and financial resources sufficient to support the objectives desired
- Are evaluated based on objective outcome measures and best-practice process indicators, and strive for consistent data definitions and goals, which facilitate outcome comparison with other systems.
- Are driven by an impassioned performance improvement culture.
- Have effective communication systems and processes across their hospitals, health systems, and health-care coalitions/regional health authority partners, between potential pre- and post-hospital partners, and with patients and families.
- Develop clear expectations and supportive clinical and educational resources for patients and their families, especially those patients with chronic medical illnesses.

The Use of Simulation for Preparedness and Planning

8. We suggest hospitals, health systems, and HC/RHAs promote the use of computer modeling to gain insight into their operational capabilities and limitations, in the following ways:

- Support the creation of computer models utilizing industry templates in collaboration with their own administrative, clinical, and technical resource experts from participating system partners. Models should include government and military resources when applicable, and include provision of maintenance of chronically ill patient populations.
- Collaborate with modelers in the design, implementation, and testing of these models; and with the interpretation and application of these results.
- Support the data requirements for such system models, and develop repositories for operationally

relevant data that can be used in future modeling efforts.

- **Leverage their relationships with national, regional, and local governments and public health agencies and emergency medical service providers to obtain necessary data on the transportation and patient logistic components of such models as required.**

Introduction

System-level planning is focused on health-care delivery that unites all local health-care organizations, including hospitals, health systems, local/regional government agencies, emergency medical services, and virtually all other health-care entities that may be called on to deliver care in a major disaster or pandemic. In the United States, this is identified as a health-care coalition (sometimes referred to as a tier 2 or Multi-agency Coordination System)¹⁻³ and internationally referenced in this publication as a regional health authority (RHA).

Although community-based health-care coalition/RHAs (HC/RHAs) may not have specific operational authority and thus not be considered a response organization, in collaboration with emergency management and other government partners, an HC/RHA can cooperatively support medical surge planning and response through information sharing and resource coordination.^{1,2,4,5} Specific hospital accreditation requirements may also

define a hospital's role in a community-based emergency medical system.⁶

HC/RHA partners may not routinely work together and may even be competitors but usually will cooperate during a disaster/pandemic. The power of their effectiveness relies on teamwork. Thus, the task force chose to address system-level planning with the intent to develop suggestions to facilitate teamwork among HC/RHAs in conjunction with their government partners to highlight the necessary infrastructure to provide effective mass critical care.

Government support, teamwork, and system-level communication are basic infrastructure for HC/RHAs, and because of their importance, elements of these three principles are briefly reviewed early in this document. Effective teamwork is the foundation for increasing HC/RHA success in all endeavors. Finally, we offer suggestions developed from examples of currently effective advanced regional care systems and simulation models such that these may offer insight to base future system-level design.

The suggestions in this chapter are important for all of those involved in a disaster or pandemic with multiple critically ill patients, including front-line clinicians, hospital administrators, and public health or government officials. Although it is important for all providers to be familiar with the system requirements of disaster and pandemic response, Table 1 provides an overview of the suggestions most of interest to each of the above groups.

Materials and Methods

The Systems Planning panel followed the American College of Chest Physicians (CHEST) Guidelines Oversight Committee's methodology to develop suggestions, based on a consensus development process (see "Methodology" article by Ornelas et al⁷ in this consensus statement). The Systems Planning panel developed nine key questions (see e-Appendix 1 for key questions list and corresponding search terms and

results). A systematic literature review was then performed for relevant articles and documents, reports, and other publications reported since 1993 to 2012; English-language papers were included, and non-English-language papers were excluded. No studies of sufficient quality were identified upon which to make evidence-based recommendations. Therefore, the panel developed expert-opinion-based suggestions utilizing a modified Delphi process.

Results

National Government Support of HC/RHAs—Policy

1a. We suggest political leadership at national levels should support health-care preparedness through financial assistance, support of market driven incentives, and preparedness requirements to HC/RHAs.

1b. We suggest national governments should support the development of responsive and nimble disaster/pandemic research processes that can both organize and assess information from prior disasters/pandemics, acquire real-time data in an ongoing

one to provide situational awareness, and which can also learn from and support international disaster relief efforts.

1c. We suggest national, state/province/regional, and city/district governments should:

- **Working with health-care experts and leadership, develop formal legal disaster/pandemic activation mechanisms to initiate, implement, and support disaster/pandemic plans and standards of care for health-care coalitions/regional health authorities and health-care professionals; and legally initiate**

TABLE 1] List of Suggestions, With Recommended Category of Health-care Professionals for Which Each Set of Suggestions Is Intended

Suggestion No.	Primary Target Audience		
	Clinicians	Hospital Administrators	Public Health and Government
1			✓
2	✓	✓	
3	✓	✓	
4	✓	✓	
5	✓	✓	
6	✓	✓	
7	✓	✓	✓
8	✓	✓	✓

step down termination procedures and processes as conditions and criteria warrant in the recovery phase

- **Work with health-care experts and leadership in the greater health-care community to develop and refine specific “trigger” criteria for formal legal activation and step down termination procedures and processes of disaster/pandemic plans and standards of care.**

1d. We suggest local governments and government agencies should be formal partners in their local health-care coalition(s), and be actively engaged with their ongoing preparedness and response activities.

Since 2002, health-care coalitions in the United States have developed because of financial support and leadership from the federal government.^{4,8} Success in disaster preparedness relies on this level of leadership and coordination. National governments should financially support HC/RHAs with market-led incentives when appropriate and guide establishment of national standards for preparedness and response (Fig 1).

Active government participation in health-care coalitions in the United States ensures integration with local public health authorities, emergency medical services, and health-care providers such as nursing and mental health facilities, which have previously been omitted from the emergency preparedness community.^{4,8} Government-health care partnerships are an important component of both preparedness and successful response capabilities.⁹ These partnerships are important because HC/RHAs are vulnerable to disruptions of normal patient care during virtually any disaster, leading to potential mismatch of need and resources.^{5,10} Although dedicated resources for developing triggers

GOVERNMENT

Figure 1 – National governments and suggestions for support of disaster preparedness (top three boxes) and suggestions for continuing to work with local governments and health-care coalition/regional health authorities to facilitate preparedness planning (bottom two boxes).

for implementing and terminating crisis standards of care are available, the work of developing them is a responsibility of local health-care and government leadership.^{11,12} Government leadership, in collaboration with health-care experts, is needed to define specific criteria for activation of emergency response plans, including when “crisis standards of care” are invoked and when they are terminated, and provide guidance for providers and health-care systems¹⁰ (see “Legal Preparedness” article by Courtney et al¹³ in this consensus statement).

There is little scientific evidence to guide disaster preparation and practice. During the 2009 influenza A(H1N1) pandemic, insufficient research infrastructure to initiate clinical studies meant vital questions remained unanswered. Similar gaps have been noted in other disasters, indicating the importance of public health emergency response research.^{14–16} We suggest national governments support the creation of a clinical research infrastructure, including predeveloped protocols, data collection instruments, ethical oversight, and funding to conduct necessary research during acute public health emergencies.¹⁶

Teamwork Within HC/RHAs—Foundational Principles

2. We suggest health-care coalition partners should work together, with the following objectives:

2a. HC/RHA clinical and administrative leaders from all partners meet together on a routine, scheduled basis. Clinician leaders must include critical care medicine experts.

2b. HC/RHA clinical and administrative leaders from all partners work together at least yearly with primary focus on developing and updating joint disaster/pandemic preparedness plans based on likely events (Hazard Vulnerability Analyses).

2c. HC/RHA clinical and administrative leaders from all partners jointly practice activation and implementation of disaster/pandemic plans and standards of care through exercises.

2d. HC/RHA partners activate their communication and collaboration mechanisms for virtually all actual or potential surge events, or unusual or large scale planned or unplanned events requiring cooperation, to ensure optimal responses and enable experience working together.

2e. HC/RHAs identify clinical experts to oversee and address the needs of specific populations, especially pediatrics, and also specialty populations such as trauma, burns, oncologic, etc.

2f. HC/RHA clinical and administrative leadership should be defined by position, not specific personnel, consistent with Incident Command System (ICS) nomenclature or equivalent, and designed with appropriate redundancy.

Effective teamwork among all partners is critical for optimal performance of the HC/RHAs in preparedness and response activities.^{1,4,5,8} As evidenced during Hurricane Sandy,¹⁶ the foundation of any team is personal relationships and the development of trust through common work toward mutual goals.¹⁷ Although smaller in scale, trauma networks and telemedicine systems are associated with improved outcomes through regional coordination, with consensus recommendations supporting similar systems for acute coronary syndromes and stroke.¹⁷⁻²⁰ Administrative and clinical leaders meeting, working, and performing disaster exercises together on a routine, scheduled basis serve to create effective teams through professional and social interactions. Meeting annually is believed to be the minimum required to develop effective relationships.

Intensivist expertise in planning and response is also important in pandemics and disasters.^{21,22} Therefore, a critical care physician's expertise should be part of a HC/RHA's leadership planning and operational

structure to aid in patient triage and transfer; help prevent undertriage and overtriage; help coordinate resources and information; serve forward medical response teams (through the National Disaster Medical System or equivalent organizations); and provide a level of portable critical care that can assist local providers and coalitions with inpatient triage, treatment, and transport^{5,21} (see "Surge Capacity Principles" article by Hick et al,²³ "Surge Capacity Logistics" article by Einav et al,²⁴ and "Evacuation of the ICU" article by King et al²⁵ in this consensus statement).

System-Level Communication—Foundational Principles

3a. We suggest HC/RHAs should have secure online and/or published directories for all partners' clinical and administrative leadership, with emergency contact information (phone numbers, e-mail addresses, pagers, cell phone texting preferences, other means) and current call schedules.

3b. We suggest HC/RHA's should have defined communication vehicles which may include (but are not limited to): dedicated secure health-care coalition web sites; conference call lines and teleconferencing technologies (eg, Skype, others); hospital phones (land lines and cell phones); pagers, hand-held walkie-talkies, ham radios, or other similar means of communication; telemedicine technologies, such as E-ICU, integrated into their disaster plans.

3c. We suggest HC/RHA partners should attempt to routinely use those agreed upon communication vehicles when working together.

3d. We suggest all agreed upon communication vehicles should be tested on a scheduled basis, with objective criteria to validate the test.

3e. We suggest the choice of communication vehicles and testing may be based on likely disaster/pandemic events (Hazard Vulnerability Analyses), and/or other appropriate considerations.

3f. We suggest developing defined disaster/pandemic plans for monitoring and leveraging popular social media (eg, Twitter, Facebook, others) during all actual or potential surge events, or unusual or large scale planned or unplanned events requiring cooperation, as both a means for gathering and transmitting information, as appropriate.

3g. We suggest HC/RHAs should have defined communication tools designated for each level of

organizational leadership, which should be consistent with ICS structure or equivalent.

HC/RHAs are the most important source of accurate, real-time data, with hospitals being the primary interface with patients, the community, and government agencies.^{1,4} They are the information clearinghouses on locations and levels of supplies, medications, beds, personnel, and virtually all resources necessary to coordinate an effective mass critical care disaster response and interface with the media and public (Fig 2).^{1,4,26,27}

Limited information describes how HC/RHA communication systems might function,^{26,27} but we suggest on the use of uniform shared communication systems of varying sophistication. Regardless of which communication modalities are selected, redundancy is important, and back-up systems should always be tested and readily available. Emergency contact information should be easily and quickly accessible, with multiple ways of finding personnel. Web-based or network-centric computer information systems are powerful options, given the volume of information encountered in any disaster and the urgent need to access and assimilate it by partners and government at all levels.^{3,28,29} Additionally, telemedicine technology may be used to improve situational awareness, assist in field triage, and augment local medical surge capacity by providing specialty expertise immediately to the scene of an event.³⁰

Partners should test technologies and processes to verify that they function as planned. We suggest this be done yearly at minimum but strongly encourage partners to use them at every opportunity, from planned large-scale

events and moderate surge events to disasters. Although research on emergency network communication is proposed, no defined standards exist.³¹ Individually developed metrics should focus on the choice of communication vehicles and their accessibility within appropriate time frames under disaster-like circumstances.

Social media, such as Facebook and Twitter, are a powerful way to disseminate information at a stunningly rapid pace during disasters.³² This technology is an important new communication tool, and we suggest HC/RHAs consult with social media experts or software companies in developing strategies to use them.

System-Level Surge Capacity and Capability

4a. We suggest HC/RHA surge objectives should be consistent with individual hospital surge goals and include the capability to surge to:

- Up to 200% above routine maximal capacity based on the nature and severity of the disaster (contingency to crisis)
- Up to the limit of the total number of ventilators available to coalition partners.
- Up to projected patient loads in a slow onset, slow evolving disaster.

4b. We suggest HC/RHAs should be able to monitor and track their defined surge capacity supplies and equipment, ideally “real time” and electronically, with the intent of being able to use all HC/RHA assets. These supplies and equipment may include identified caches of important medications or equipment, and bed availability among partners.

4c. We suggest HC/RHAs should have the ability to track the number of available ICU capable personnel (“force multipliers”) and other designated specialist “resources” (eg, pediatric and special populations) through their partner hospitals. Partners with telemedicine capability (such as tele-ICU’s) should have plans for how to use this resource to optimize the use of pediatric and specialty expertise across hospitals served by the telemedicine resource.

4d. We suggest HC/RHAs should have defined policies and procedures for emergency privileging for all health-care professionals designated as coalition resources.

4e. We suggest fair and adequate reimbursement for expenditures and loss of revenue related to delivery of acute critical care services during a disaster or

Figure 2 – Summary of suggestions for health-care coalition/regional health authority leadership regarding communication.

pandemic must be ensured. This should include the guarantee of payments from governmental sources, as well as by insurance companies and other payers of health-care services.

In responding to any disaster or pandemic, the task force strongly believes HC/RHAs will be most prepared if they have quantitative knowledge of where resources are available to facilitate access, mobilization, and effective use (Fig 3). These resources include beds, equipment, supplies, and personnel. Suggested system-level surge goals are the same as for individual hospitals and range from conventional (20% above capacity) to contingency (up to 100% above capacity), to crisis surge levels (up to 200% above capacity) (see “Surge Capacity Principles” and “Surge Capacity Logistics” articles by Hick et al²³ and Einav et al,²⁴ respectively, in this consensus statement).³³ Most hospitals and regions have limited reserve bed capacity under routine circumstances, with only 25% maximum number of open beds, which equates to 500 beds per million population.³⁴⁻³⁶ Pediatric beds are scarcer, or may be geographically or otherwise unavailable, with altered standards of care likely needed to accommodate even a moderate disaster surge.^{10,36-38}

Given these considerations, the task force strongly believed it important for HC/RHA leadership to have knowledge of all partner hospitals’ actual and potential bed availability, to be considered as surge capacity. This engagement should occur at the earliest opportunity, especially if there are, or are likely to be, a high number of resource-intense pediatric or specialty care casualties (eg, burns, and so forth) to provide extensive resources quickly, redirect transfers, and provide for the greatest possible breadth of care.

Figure 3 – Summary of suggestions for health-care coalition/regional health authority leadership regarding the number and location of surge resources, and important surge priorities.

In planning transfer of critically ill patients, ED-housed patients may be at less risk than those already admitted, but at increasing risk the longer they reside in an ED.^{39,40} Patients with high acuity, such as those who are mechanically ventilated, may benefit from transfers to tertiary care centers, but specific ICU type (eg, medical ICU, cardiac care unit) is less likely to impact care.⁴¹⁻⁴³ Coordination of patient transfer to facilities with preidentified centers of excellence in critical care is an opportunity to improve patient outcomes further.^{44,45} Critically ill patients not (immediately) triaged to critical care, typically based on various exclusion criteria (see “Triage” article by Christian et al⁴⁶ in this consensus statement⁴⁶), are at increased risk of death.⁴⁷

Surge capacity is limited by the need for one ventilator per mass critical care patient. A recent inventory determined that US hospitals own a supply of 62,188 full-feature ventilators and 98,738 less-advanced ventilators (total, 160,926)⁴⁸ to serve 93,955 ICU beds.⁴⁹ Thus, insufficient ventilators are available to even double usual ICU capability in a national public health emergency without access to regional or national stockpiles. Despite the limitation in ventilator supply and the high occupancy of existing ICU beds, only 29% of average US hospital ICU beds are occupied with a patient on a ventilator.⁵⁰ Other systems, such as county (state-run) hospitals in the United States, Canada, and Europe, routinely run at near 100% capacity with ventilated patients. Thus, the ability of existing critical care services to accommodate substantial mass critical care surges varies widely and can only be established by knowing ventilator capacity and their location.

We suggest mass critical care local operational planning include an inventory of the number of ventilators, including full-feature ventilators, specialized ventilators for newborns, and less-advanced ventilators with limited versatility that nonetheless can be lifesaving in patients with respiratory failure. Suggestions regarding the types and number of medications, critical care supplies, and equipment are found elsewhere in this supplement (see “Surge Capacity Logistics” article by Einav et al²⁴ in this consensus statement).⁵¹ Beds, ventilators, medications, supplies, and equipment should be tracked electronically by HC/RHAs, with knowledge of both their absolute numbers and whereabouts immediately available to system leaders.

Potential critical care surge personnel (force multipliers) should be identified with current contact information (see “Surge Capacity Logistics” article by Einav et al²⁴ in this consensus statement). Given the foreseeable shortage

of intensivists, hospitalists may be the likely group to augment intensive care staffing, although any hospital-based provider may be recruited.^{52,53} Training and just-in-time education for these health-care professionals must be planned so as to provide critical care in an emergency.

Some precedent and support for credentialing health-care professional volunteers in a major disaster exist (See “Legal Preparedness” article by Courtney et al¹³ in this consensus statement).^{54,55} To the degree possible under local law, a mechanism for emergency credentialing of health-care professional volunteers should exist, which may even include HC/RHA partners simply recognizing credentialing processes of other health-care organizations to benefit from each other’s professional pools.

Pediatric Patients and Specialty Populations

5a. We suggest HC/RHA have identified, and be familiar with, the following pediatric disaster/pandemic designated resources including, but not limited to:

- **Pediatric consultative specialists available by dedicated phone line support and/or dedicated video or telemedicine consultation.**
- **Designated pediatric surge personnel (eg, pediatric hospitalists, others) available to non-pediatric hospitals and health systems to support surge in contingency or crisis level events, with a defined plan for how to activate this resource when needed.**
- **Identified pediatric capable transport resources for allocation and matching of pediatric patients to available HC/RHA pediatric resources.**
- **Knowledge of available key supplies, medications, and other pediatric assets; location of these assets with a defined process for how they may be accessed urgently; and ability to monitor when asset reserves fall below a defined critical threshold.**
- **Pediatric educational resources. If web-based, they should be found on HC/RHA websites, or with links to appropriate resources. If published, resources should be readily available to all partners.**

5b. We suggest HC/RHAs should have plans to provide care for specialty populations routinely found in their catchment area or region in parallel as described for pediatrics. Resources should include consultative services, potential surge personnel, transport resources, specialty supplies/medications, and educational resources. These populations include but are not limited to trauma, nephrology, burns, oncologic patients.

5c. Health-care coalitions, health systems, and hospitals identify patients with high-level chronic disease care needs, such as a home ventilator, home oxygen, chronic dialysis, and work to ensure their needs are met at home to help prevent these patients from having to seek assistance at hospitals.

Pediatric hospitals provide superior care for severe pediatric trauma,^{56,57} as well as for critically ill children⁵⁸⁻⁶⁰ and neonates,^{61,62} with better outcomes than nonpediatric facilities. Most children (> 90%) are cared for in nonpediatric EDs, and one-half of these departments see < 10 children each day.⁶³ Fewer than 20% of hospital beds are intended for the care of children, and these are clustered at a minority of hospitals.³⁶ Pediatric critical care beds account for < 10% of all ICU beds in the United States and are clustered at an even smaller number of hospitals.^{49,64} As a result, professional organizations have long recommended that pediatric emergency and critical care resources be regionalized so that resuscitation and stabilization are available close to every community; common low-risk conditions are treated at community hospitals, and high-risk complex conditions are transferred to pediatric hospitals.⁶⁵

Children account for 25% of the population in the United States. It is plausible that in public health emergencies, pediatric surges will exceed age-specific resources even if pediatric patients are proportional to their numbers in the general population. The Pediatric Emergency Mass Critical Care Task Force⁶⁶ recently endorsed CHEST’s mass critical care framework,⁶⁷ and we continue to recommend that pediatric-specific disaster planning is essential. However, to accommodate surges of children, pediatric disaster/pandemic services must be well integrated into regional systems of care.⁶⁶⁻⁶⁸ In a disaster or pandemic, pediatric care is usual care, not specialty care. All HC/RHAs must be prepared to provide care for large numbers of pediatric patients across their region, and operational plans must account for unique local circumstances (Fig 4).¹⁰ Pediatric consultants should be available to all regional partners by phone or telemedicine for remote consultation. Potential benefits of telemedicine communications include improved situational awareness, assistance in field triage, and timely expert consultation.³⁰ Telemedicine communications may be vulnerable to damage in some disasters.⁶⁹

Nonpediatric hospitals must survey and identify staff with pediatric capability, organize contact information, and develop job assignments for them. All members of the hospital team who will provide care to children will

Figure 4 – Summary of suggestions for health-care coalition/regional health authority leadership regarding pediatric resources (which may also be used for special populations).

benefit from drills and education as well as from just-in-time education tailored to the circumstances of the emergency. Internet resources may be invaluable to support just-in-time education, but electronic access may be unreliable during emergencies, so consideration should be given for printed educational resources.

Interhospital transport resources may also be limited in pandemics and disasters. Decision makers must evaluate the merits of sending critical care staff, supplies, and equipment from uninvolved remote sites to a localized disaster vs transporting the patients from nonpediatric hospitals to a pediatric center. Priorities must be determined for those who should be transferred first (one-on-one transport staffing for the most severely ill patients may exhaust limited critical care resources). In a disaster or pandemic, optimizing population outcomes may require transport strategies distinct from those used in ordinary circumstances, adapting to event-specific local needs and resources (see “Evacuation of the ICU” article by King et al²⁵ in this consensus statement).

Supplies and medications unique to pediatric populations may rapidly be depleted in a large disaster, and the regions’ hospitals should have knowledge and access to these resources at other sites. The levels of supplies should ideally be monitored, as described in the surge article (see “Surge Capacity Logistics” article by Einav et al²⁴ in this consensus statement).

New pediatric-specific evidence may illustrate approaches that also pertain to integrating care of other vulnerable populations into regional systems. HC/RHAs should define their local vulnerable populations, optimally with plans to meet their needs at home rather than in a hospital (see “Special Populations” article by

Dries et al⁷⁰ in this consensus statement). The task force suggests that planning for these vulnerable populations should also mirror the approach described above for pediatric patients.

HC/RHAs and Networks

6a. We suggest during a disaster requiring transfer of patients, whether from emergency medicine departments or inpatient areas, transferring partners may have initial choice of where patients are referred based on traditional referral patterns. However, HC/RHA leadership must oversee this process, and be able to intercede as both a resource and with the authority to redirect transfers based on anticipated or actual events. Defined health-care coalition coordination processes and transfer resources should be planned and identified ahead of time.

6b. Health-care coalitions should designate neighboring health-care coalitions as potential partners during a contingency or crisis event, and have readily available leadership contact information, and knowledge of these potential partners’ size and capabilities.

Hospitals commonly transfer and receive patients along an informal yet structured network based on availability of services, with more-resourced hospitals tending to receive more referrals.^{44,45} Approximately 4.5% of Medicare patients admitted to an ICU ultimately undergo an interhospital critical care transfer.⁴⁵ If a major referral hospital was incapacitated, accepting transfers would halt, and transfer of its own patients would likely be devastating to usual referral networks.⁴⁵ During a major disaster or pandemic, HC/RHA leadership needs to access knowledge of all regional ICU and hospital beds quickly, as discussed previously.

The movement of patients, whether to evacuate a hospital or to decompress a facility to increase acute care capacity, will be enhanced by a regionalized, coordinated, and accountable emergency care system.^{19,71} Neighboring HC/RHA partners are also crucial allies for accepting potential patient transfers; hence, the task force has suggested that all HC/RHAs be familiar with their neighboring coalitions’ leadership, capacity, and organizational capabilities.

Models of Advanced Regional Care Systems

7. Advanced Regional Care Systems instituted within large hospitals, and across hospitals, health systems and HC/RHAs, will have the greatest chance for

success if they are established with the following goals:

- **Clear and transparent objectives for what those programs are to accomplish, and the programs are well integrated and accepted across their hospital and health system partners.**
- **Have administrative and financial resources sufficient to support the objectives desired**
- **Are evaluated based on objective outcome measures and best-practice process indicators, and strive for consistent data definitions and goals, which facilitate outcome comparison with other systems.**
- **Are driven by an impassioned performance improvement culture.**
- **Have effective communication systems and processes across their hospitals, health systems, and HC/RHA partners, between potential pre- and post- hospital partners, and with patients and families.**
- **Develop clear expectations and supportive clinical and educational resources for patients and their families, especially those patients with chronic medical illnesses.**

Advanced regional care systems may be defined as large clinical programs that function across hospitals, health systems, and/or regions and are characterized by established program objectives and performance expectations (Fig 5, e-Table 1). Successful examples include trauma systems, recognized as the standard for best care to injured patients⁷²; telemedicine ICU programs, now numbering > 300 programs in the United States and providing high-quality ICU care for up to 8% of nongovernment ICU beds⁷³; and the Renal Disaster Relief Task Force, with a proven history of

providing urgent dialysis to disaster victims (usually after earthquakes) in less-developed countries^{74,75} (see e-Appendix 1 for further discussion of these examples).

Successful foundations of these programs may be viewed as a five-level pyramid (Fig 5), with the foundation being strong leadership in establishing and planning the vision of the program and securing the necessary financial and political resources.^{17,20,74-77} Successive levels include the culture that drives the clinical systems of care; the clinical data and outcomes used for quality assessment and continuous improvement; and, finally, technology, which is driven primarily by program objectives.^{30,76,78} Quality of patient care is at the top of the pyramid, with mortality data currently the best objective care measure (e-Table 1), with functional outcomes as a high priority, and with patient and family experience being increasingly important.^{72,75,79-84} (Please see e-Appendix 1 for further discussion of advanced regional care systems in relation to trauma networks, telemedicine ICU programs, and the Renal Disaster Relief Task Force.)

The Use of Simulation for Preparedness and Planning

8. We suggest hospitals, health systems, and HC/RHAs promote the use of computer modeling to gain insight into their operational capabilities and limitations, in the following ways:

- **Support the creation of computer models utilizing industry templates in collaboration with their own administrative, clinical, and technical resource experts from participating system partners. Models should include government and military resources when applicable, and include provision of maintenance of chronically ill patient populations.**
- **Collaborate with modelers in the design, implementation, and testing of these models; and with the interpretation and application of these results.**
- **Support the data requirements for such system models, and develop repositories for operationally relevant data that can be used in future modeling efforts.**
- **Leverage their relationships with national, regional, and local governments and public health agencies and emergency medical service providers to obtain necessary data on the transportation and patient logistic components of such models as required.**

Computer-based modeling is a widely used and powerful tool that helps complex organizations understand their own processes, resources, and capacity in the face of variability in demand and/or supply of services and

Figure 5 – Pyramid illustrating levels of priorities for successful advanced regional care systems.

materiel.⁸⁵ Mathematical and simulation models have been used to optimize well-defined processes, such as the allocation and movement of prehospital transportation resources,⁸⁶⁻⁸⁸ forecast bed demand and critical resources given normal and exceptional use patterns,^{89,90} and represent health-care resources from the individual facility⁹¹ to a national scale.⁹² Health-system models may use a variety of methods, ranging from queuing theory,⁹³ linear programming,⁸⁶ and spreadsheet modeling⁹⁴ to a variety of computational techniques including Monte Carlo,⁹¹ discrete-event,⁸⁹ and agent-based simulation.⁹⁵

Computer modeling provides benefits for health-system planning and integration. First, efforts to create in silico models of current or planned systems require preliminary interactions among key participants to define capabilities and requirements.⁹⁶ Quantifying current use and resource requirements may help in preparation for surge requirements during disasters/pandemics and provide the rationale for future collaboration and resource-sharing arrangements. Once calibrated, models may be used to explore an evidence-based, system-wide approach to health-system planning and response at both the prehospital and hospital-based levels without causing disruption to current activities.^{87,88}

While “off-the-shelf” modeling platforms are available, the following two-step approach for developing simulations is suggested: (1) Health-care leaders should first specify the operational questions to be answered. For example, Levi and colleagues⁹² (see later) created their model to answer the question of whether bed number was a good marker for operative capability. (2) Once the scope and purpose are clear, health-care leaders may consider reaching out to regional engineering schools, which typically have experience in developing policy-relevant and technically sophisticated simulation with models of complex real-world processes.

An example illustrating both the potential benefits and limits of using modeling for planning mass casualty care is the pioneering study of Levi et al,⁹² in which they developed a novel simulation model of Israeli trauma management to assess how well hospitals’ stated bed capacity correlates with projected operative capability under high-volume scenarios. This exercise required extensive data collection, including historical review of injury number and type (eg, orthopedic, neurologic) and also extensive and detailed information about actual hospital size, bed number, staffing (anesthesiologists and surgeons), and use across the country. Two strategies for load reduction at major trauma centers were tested:

alternating day admissions and triage of less severe cases to outlying hospitals. The unexpected finding was that uniform policies did not always lead to improvement in operative capability and that the gains that did occur were not always proportional to hospital size during the emergency scenario. The health-system-level insights gained through this modeling exercise would have been difficult to ascertain otherwise except in the setting of an actual national disaster. Additionally, access to high-quality, realistic hospital-level data helped create this policy-relevant model.

Conclusions

The Task Force provides an enhanced framework for HC/RHA leaders to optimize teamwork and effective planning in pandemics and disasters. We suggest focusing on including joint communication tools and processes; identification and sharing surge resources, including location, quantity, and how to access them quickly; planning for pediatric populations as usual (not special) care; and understanding internal transfer networks to optimize potential transfers both within their region and to neighboring HC/RHAs. Task force members strongly believe that critical care specialists should be represented in HC/RHA leadership and be actively involved in all disaster/pandemic planning for large numbers of potential critically ill patients. The Task Force also promotes modeling of advanced regional care systems and simulation to provide insight into the development of future processes.

Acknowledgments

Author contributions: J. R. D. had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. J. R. D., R. K. K., D. D., V. L., M. L. L., J. W., M. A., B. S., N. H., R. M., A. V. D., M. D. C., and N. K. contributed to the development of PICO questions; J. R. D., R. K. K., D. D., V. L., M. L. L., J. W., M. A., B. S., N. H., R. M., and N. K. conducted the literature review; J. R. D., R. K. K., D. D., V. L., M. L. L., J. W., M. A., B. S., N. H., R. M., A. V. D., M. D. C., and N. K. contributed to development of expert opinion suggestion; J. R. D., R. K. K., D. D., V. L., M. L. L., J. W., M. A., B. S., N. H., R. M., A. V. D., M. D. C., and N. K. contributed to the conception and design, or acquisition of data, or analysis and interpretation of data from the Delphi process; J. R. D., R. K. K., D. D., V. L., M. L. L., J. W., M. A., B. S., N. H., and R. M. developed and drafted the manuscript; and J. R. D., A. V. D., M. D. C., and N. K. revised the manuscript critically for important intellectual content.

Financial/nonfinancial disclosures: The authors have reported to *CHEST* that no potential conflicts of interest exist with any companies/organizations whose products or services may be discussed in this article.

Endorsements: This consensus statement is endorsed by the American Association of Critical-Care Nurses, American Association for Respiratory Care, American College of Surgeons Committee on Trauma, International Society of Nephrology, Society for Academic Emergency Medicine, Society of Critical Care Medicine, Society of Hospital Medicine, World Federation of Pediatric Intensive and Critical Care Societies, World Federation of Societies of Intensive and Critical Care Medicine.

Role of sponsors: The American College of Chest Physicians was solely responsible for the development of these guidelines. The remaining supporters played no role in the development process. External supporting organizations cannot recommend panelists or topics, nor are they allowed prepublication access to the manuscripts and recommendations. Further details on the Conflict of Interest Policy are available online at <http://chestnet.org>.

Other contributions: We thank Marge Weller, Unity Hospital Administrative Assistant, Minneapolis, MN for her assistance in developing the figures used in the article. The views expressed in this consensus statement are those of the author (M. L. L.) and do not reflect the official policy or position of the Department of the Navy, Department of Defense, or the US Government. The opinions of the author (R. M.) do not represent the policy of either the Substance Abuse and Mental Health Services Administration (SAMHSA) or the US Department of Health and Human Services (DHHS) and no official endorsement by SAMHSA or DHHS is intended or should be inferred. Information disclosed by Mr Wilgis and the Florida Hospital Association related to hospital and health system emergency management and emergency preparedness, prevention, protection, response, or recovery is rendered as advisory only and does not show preference to any company, product, government agency, or representative. Any reference to specific commercial products, process, service, manufacturer, or company does not constitute its endorsement or recommendation by Mr Wilgis or the FHA. The opinions expressed within this manuscript are solely those of the author (M. D. C.) and do not represent the official position or policy of the Royal Canadian Medical Service, Canadian Armed Forces, or the Department of National Defence.

Additional information: The e-Appendix and e-Table can be found in the Supplemental Materials section of the online article.

Collaborators: Executive Committee: Michael D. Christian, MD, FRCPC, FCCP; Asha V. Devereaux, MD, MPH, FCCP, co-chair; Jeffrey R. Dichter, MD, co-chair; Niranjan Kissoon, MBBS, FRCPC; Lewis Robinson, MD, PhD; Panelists: Dennis Amundson, DO, FCCP; Michael R. Anderson, MD; Robert Balk, MD, FCCP; Wanda D. Barfield, MD, MPH; Martha Bartz, MSN, RN, CCRN; Josh Benditt, MD; William Beninati, MD; Kenneth A. Berkowitz, MD, FCCP; Lee Daugherty Biddison, MD, MPH; Dana Branner, MD; Richard D Branson, MSc, RRT; Frederick M. Burkle Jr, MD, MPH, DTM; Bruce A. Cairns, MD; Brendan G. Carr, MD; Brooke Courtney, JD, MPH; Lisa D. DeDecker, RN, MS; COL Marla J. De Jong, PhD, RN [USAF]; Guillermo Dominguez-Cherit, MD; David Dries, MD; Sharon Einav, MD; Brian L. Erstad, PharmD; Mill Etienne, MD; Daniel B. Fagbuyi, MD; Ray Fang, MD; Henry Feldman, MD; Hernando Garzon, MD; James Geiling, MD, MPH, FCCP; Charles D. Gomersall, MBBS; Colin K. Grissom, MD, FCCP; Dan Hanfling, MD; John L. Hick, MD; James G. Hodge Jr, JD, LL.M.; Nathaniel Hupert, MD; David Inghar, MD, FCCP; Robert K. Kanter, MD; Mary A. King, MD, MPH, FCCP; Robert N. Kuhnley, RRT; James Lawler, MD; Sharon Leung, MD; Deborah A. Levy, PhD, MPH; Matthew L. Lim, MD; Alicia Livinski, MA, MPH; Valerie Luyckx, MD; David Marcozzi, MD; Justine Medina, RN, MS; David A. Miramontes, MD; Ryan Mutter, PhD; Alexander S. Niven, MD, FCCP; Matthew S. Penn, JD, MLIS; Paul E. Pepe, MD, MPH; Tia Powell, MD; David Prezant, MD, FCCP; Mary Jane Reed, MD, FCCP; Preston Rich, MD; Dario Rodriguez, Jr, MSc, RRT; Beth E. Roxland, JD, MBioethics; Babak Sarani, MD; Umair A. Shah, MD, MPH; Peter Skippen, MBBS; Charles L. Sprung, MD; Italo Subbarao, DO, MBA; Daniel Talmor, MD; Eric S. Toner, MD; Pritish K. Tosh, MD; Jeffrey S. Upperman, MD; Timothy M. Uyeki, MD, MPH, MPP; Leonard J. Weireter Jr, MD; T. Eoin West, MD, MPH, FCCP; John Wilgis, RRT, MBA; ACCP Staff: Joe Ornelas, MS; Deborah McBride; David Reid; Content Experts: Amado Baez, MD; Marie Baldisseri, MD; James S. Blumenstock, MA; Art Cooper, MD; Tim Ellender, MD; Clare Helminiak, MD, MPH; Edgar Jimenez, MD; Steve Krug, MD; Joe Lamana, MD; Henry Masur, MD; L. Rudo Mathivha, MBChB; Michael T. Osterholm, PhD, MPH; H. Neal Reynolds, MD; Christian Sandrock, MD, FCCP; Armand Sprecher, MD, MPH; Andrew Tillyard, MD; Douglas White, MD; Robert Wise, MD; Kevin Yeskey, MD.

References

- Barbera JA, Macintyre AG. Medical surge capacity and capability: a management system for integrating medical and health resources during large-scale emergencies. US Department of Health and Human Services. Washington, DC: The CNA Corporation Institute for Public Research; 2007.
- Hanfling D. Role of regional healthcare coalitions in managing and coordinating disaster response. Paper presented at: 2013 Workshop on Nationwide Response to an Improvised Nuclear Device Attack; January 23-24, 2013; Washington DC.
- McMaster R, Baber C. Multi-agency operations: cooperation during flooding. *Appl Ergon*. 2012;43(1):38-47.
- Courtney B, Toner E, Waldhorn R, et al. Healthcare coalitions: the new foundation for national healthcare preparedness and response for catastrophic health emergencies. *Biosecur Bioterror*. 2009;7(2):153-163.
- Farmer JC, Carlton PK Jr. Providing critical care during a disaster: the interface between disaster response agencies and hospitals. *Crit Care Med*. 2006;34(3)(suppl):S56-S59.
- Sauer LM, McCarthy ML, Knebel A, Brewster P. Major influences on hospital emergency management and disaster preparedness. *Disaster Med Public Health Prep*. 2009;3(suppl 2):S68-S73.
- Ornelas J, Dichter JR, Devereaux AV, Kissoon N, Livinski A, Christian MD; on behalf of the Task Force for Mass Critical Care. Methodology: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. *Chest*. 2014;146(4_suppl):35S-41S.
- Rambhia KJ, Waldhorn RE, Selck F, Mehta AK, Franco C, Toner ES. A survey of hospitals to determine the prevalence and characteristics of healthcare coalitions for emergency preparedness and response. *Biosecur Bioterror*. 2012;10(3):304-313.
- Maldin B, Lam C, Franco C, et al. Regional approaches to hospital preparedness. *Biosecur Bioterror*. 2007;5(1):43-53.
- King MA, Koelemay K, Zimmerman J, Rubinson L. Geographical maldistribution of pediatric medical resources in Seattle-King county. *Prehosp Disaster Med*. 2010;25(4):326-332.
- Hanfling D, Hick JL, Stroud C, eds; Committee on Crisis Standards of Care. *Crisis Standards of Care: A Toolkit for Indicators and Triggers*. Washington, DC: The National Academies Press; 2013.
- Stroud C, Altevogt BM, Nadig L, Hougan M. Institute of Medicine (US) Forum on Medical and Public Health Preparedness for Catastrophic Events. *Crisis Standards of Care: Summary of a Workshop Series*. Washington, DC: The National Academies Press; 2010.
- Courtney B, Hodge JG Jr, Toner ES, et al; on behalf of the Task Force for Mass Critical Care. Legal preparedness: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. *Chest*. 2014;146(4_suppl):e134S-e144S.
- Ortiz JR, Rudd KE, Clark DV, Jacob ST, West TE. Clinical research during a public health emergency: a systematic review of severe pandemic influenza management. *Crit Care Med*. 2013;41(5):1345-1352.
- Morton M, Levy JL. Challenges in disaster data collection during recent disasters. *Prehosp Disaster Med*. 2011;26(3):196-201.
- Lurie N, Manolio T, Patterson AP, Collins F, Frieden T. Research as a part of public health emergency response. *N Engl J Med*. 2013;368(13):1251-1255.
- Edlich RF, Wish JR, Britt LD, Long WB. An organized approach to trauma care: legacy of R Adams Cowley. *J Long Term Eff Med Implants*. 2004;14(6):481-511.
- Lansink KW, Leenen LP. Do designated trauma systems improve outcome? *Curr Opin Crit Care*. 2007;13(6):686-690.
- Carr BG, Matthew Edwards J, Martinez R. Regionalized care for time-critical conditions: lessons learned from existing networks. *Acad Emerg Med*. 2010;17(12):1354-1358.
- Lilly CM, Cody S, Zhao H, et al; University of Massachusetts Memorial Critical Care Operations Group. Hospital mortality, length of stay, and preventable complications among critically ill patients before and after tele-ICU reengineering of critical care processes. *JAMA*. 2011;305(21):2175-2183.
- Shirley PJ, Mandersloot G. Clinical review: the role of the intensive care physician in mass casualty incidents: planning, organisation, and leadership. *Crit Care*. 2008;12(3):214.
- Weick KE, Sutcliffe KM. *Managing the Unexpected*. San Francisco, CA: John Wiley and Sons, Inc; 2007.
- Hick JL, Einav S, Hanfling D, et al; on behalf of the Task Force for Mass Critical Care. Surge capacity principles: care of the critically

- ill and injured during pandemics and disasters: CHEST consensus statement. *Chest*. 2014;146(4_suppl):e1S-e16S.
24. Einav S, Hick JL, Hanfling D, et al; on behalf of the Task Force for Mass Critical Care. Surge capacity logistics: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. *Chest*. 2014;146(4_suppl):e17S-e43S.
25. King MA, Niven AS, Beninati W, et al; on behalf of the Task Force for Mass Critical Care. Evacuation of the ICU: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. *Chest*. 2014;146(4_suppl):e44S-e60S.
26. Booth CM, Stewart TE. Communication in the Toronto critical care community: important lessons learned during SARS. *Crit Care*. 2003;7(6):405-406.
27. Hawryluck L, Lapinsky SE, Stewart TE. Clinical review: SARS - lessons in disaster management. *Crit Care*. 2005;9(4):384-389.
28. von Lubitz DK, Beakley JE, Patricelli F. 'All hazards approach' to disaster management: the role of information and knowledge management, Boyd's OODA Loop, and network-centricity. *Disasters*. 2008;32(4):561-585.
29. Stanovich ML. "Network-centric" emergency response: the challenges for a new command and control paradigm. Institute for Security, Technology, and Society website. <http://www.ists.dartmouth.edu/library/191.pdf>. Accessed September 28, 2013.
30. Xiong W, Bair A, Sandrock C, Wang S, Siddiqui J, Hupert N. Implementing telemedicine in medical emergency response: concept of operation for a regional telemedicine hub. *J Med Syst*. 2012;36(3):1651-1660.
31. Glickman SW, Kit Delgado M, Hirshon JM, et al. Defining and measuring successful emergency care networks: a research agenda. *Acad Emerg Med*. 2010;17(12):1297-1305.
32. Bagrow JP, Wang D, Barabási AL. Collective response of human populations to large-scale emergencies. *PLoS ONE*. 2011;6(3):e17680.
33. Hick JL, Barbera JA, Kelen GD. Refining surge capacity: conventional, contingency, and crisis capacity. *Disaster Med Public Health Prep*. 2009;3(suppl 2):S59-S67.
34. Kaji AH, Lewis RJ. Hospital disaster preparedness in Los Angeles County. *Acad Emerg Med*. 2006;13(11):1198-1203.
35. Dayton C, Ibrahim J, Augenbraun M, et al. Integrated plan to augment surge capacity. *Prehosp Disaster Med*. 2008;23(2):113-119.
36. Kanter RK, Moran JR. Hospital emergency surge capacity: an empiric New York statewide study. *Ann Emerg Med*. 2007;50(3):314-319.
37. Kanter RK, Moran JR. Pediatric hospital and intensive care unit capacity in regional disasters: expanding capacity by altering standards of care. *Pediatrics*. 2007;119(1):94-100.
38. Kanter RK. Strategies to improve pediatric disaster surge response: potential mortality reduction and tradeoffs. *Crit Care Med*. 2007;35(12):2837-2842.
39. Chalfin DB, Trzeciak S, Likourezos A, Baumann BM, Dellinger RP; DELAY-ED study group. Impact of delayed transfer of critically ill patients from the emergency department to the intensive care unit. *Crit Care Med*. 2007;35(6):1477-1483.
40. Gerber DR, Schorr C, Ahmed I, Dellinger RP, Parrillo J. Location of patients before transfer to a tertiary care intensive care unit: impact on outcome. *J Crit Care*. 2009;24(1):108-113.
41. Kahn JM, Linde-Zwirble WT, Wunsch H, et al. Potential value of regionalized intensive care for mechanically ventilated medical patients. *Am J Respir Crit Care Med*. 2008;177(3):285-291.
42. Nguyen YL, Kahn JM, Angus DC. Reorganizing adult critical care delivery: the role of regionalization, telemedicine, and community outreach. *Am J Respir Crit Care Med*. 2010;181(11):1164-1169.
43. Sidlow R, Aggarwal V. "The MICU is full": one hospital's experience with an overflow triage policy. *Jt Comm J Qual Patient Saf*. 2011;37(10):456-460.
44. Iwashyna TJ, Christie JD, Kahn JM, Asch DA. Uncharted paths: hospital networks in critical care. *Chest*. 2009;135(3):827-833.
45. Iwashyna TJ, Christie JD, Moody J, Kahn JM, Asch DA. The structure of critical care transfer networks. *Med Care*. 2009;47(7):787-793.
46. Christian MD, Sprung CL, King MA, et al; on behalf of the Task Force for Mass Critical Care. Triage: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. *Chest*. 2014;146(4_suppl):e61S-e74S.
47. Sinuff T, Kahn moui K, Cook DJ, Luce JM, Levy MM; Values Ethics and Rationing in Critical Care Task Force. Rationing critical care beds: a systematic review. *Crit Care Med*. 2004;32(7):1588-1597.
48. Rubinson L, Vaughn F, Nelson S, et al. Mechanical ventilators in US acute care hospitals. *Disaster Med Public Health Prep*. 2010;4(3):199-206.
49. Halpern NA, Pastores SM. Critical care medicine in the United States 2000-2005: an analysis of bed numbers, occupancy rates, payer mix, and costs. *Crit Care Med*. 2010;38(1):65-71.
50. Wunsch H, Wagner J, Herlim M, Chong DH, Kramer AA, Halpern SD. ICU occupancy and mechanical ventilator use in the United States. *Crit Care Med*. 2013;41(12):2712-2719.
51. Rubinson L, Hick JL, Curtis JR, et al; Task Force for Mass Critical Care. Definitive care for the critically ill during a disaster: medical resources for surge capacity: from a Task Force for Mass Critical Care summit meeting, January 26-27, 2007, Chicago, IL. *Chest*. 2008;133(suppl 5):32S-50S.
52. Siegal EM, Dressler DD, Dichter JR, Gorman MJ, Lipsett PA. Training a hospitalist workforce to address the intensivist shortage in American hospitals: a position paper from the Society of Hospital Medicine and the Society of Critical Care Medicine. *Crit Care Med*. 2012;40(6):1952-1956.
53. Wise KR, Akopov VA, Williams BR Jr., Ido MS, Leeper KV Jr., Dressler DD. Hospitalists and intensivists in the medical ICU: a prospective observational study comparing mortality and length of stay between two staffing models. *J Hosp Med*. 2012;7(3):183-189.
54. Minnesota Statutes regarding out of state license holders; powers, duties. Office of the Revisor of Statutes website. <https://www.revisor.mn.gov/statutes/?id=12.42>. Accessed October 12, 2013.
55. Hospital staffing and surge capacity during a disaster event. *Research Brief*. May, 2007.
56. Osler TM, Vane DW, Tepas JJ, Rogers FB, Shackford SR, Badger GJ. Do pediatric trauma centers have better survival rates than adult trauma centers? An examination of the National Pediatric Trauma Registry. *J Trauma*. 2001;50(1):96-101.
57. Densmore JC, Lim HJ, Oldham KT, Guice KS. Outcomes and delivery of care in pediatric injury. *J Pediatr Surg*. 2006;41(1):92-98.
58. Pollack MM, Alexander SR, Clarke N, Ruttimann UE, Tesselaar HM, Bachulis AC. Improved outcomes from tertiary center pediatric intensive care: a statewide comparison of tertiary and nontertiary care facilities. *Crit Care Med*. 1991;19(2):150-159.
59. Tilford JM, Simpson PM, Green JW, Lensing S, Fiser DH. Volume-outcome relationships in pediatric intensive care units. *Pediatrics*. 2000;106(2 pt 1):289-294.
60. Ruttimann UE, Patel KM, Pollack MM. Relevance of diagnostic diversity and patient volumes for quality and length of stay in pediatric intensive care units. *Pediatr Crit Care Med*. 2000;1(2):133-139.
61. Phibbs CS, Baker LC, Caughey AB, Daniels B, Schmitt SK, Phibbs RH. Level and volume of neonatal intensive care and mortality in very-low-birth-weight infants. *N Engl J Med*. 2007;356(21):2165-2175.
62. Goodman DC, Fisher ES, Little GA, Stukel TA, Chang CH, Schoendorf KS. The relation between the availability of neonatal intensive care and neonatal mortality. *N Engl J Med*. 2002;346(20):1538-1544.
63. Gausche-Hill M, Schmitz C, Lewis RJ. Pediatric preparedness of US emergency departments: a 2003 survey. *Pediatrics*. 2007;120(6):1229-1237.
64. Randolph GD, Murray M, Swanson JA, Margolis PA. Behind schedule: improving access to care for children one practice at a time. *Pediatrics*. 2004;113(3 pt 1):e230-e237.
65. Council of the Society of Critical Care Medicine. Consensus report for regionalization of services for critically ill or injured children. *Crit Care Med*. 2000;28(1):236-239.

66. Kissoon N; Task Force for Pediatric Emergency Mass Critical Care. Deliberations and recommendations of the Pediatric Emergency Mass Critical Care Task Force: executive summary. *Pediatr Crit Care Med*. 2011;12(suppl 6):S103-108.
67. Devereaux A, Christian MD, Dichter JR, Geiling JA, Robinson L; Task Force for Mass Critical Care. Summary of suggestions from the Task Force for Mass Critical Care summit, January 26-27, 2007. *Chest*. 2008;133(5_suppl):1S-7S.
68. National Commission on Children and Disasters. *2010 Report to the President and Congress*. Rockville, MD: Agency for Healthcare Research and Quality; October 2010.
69. Kim J, Kim D, Jung S, et al. Implementation and performance evaluation of mobile ad hoc network for Emergency Telemedicine System in disaster areas. *Conf Proc IEEE Eng Med Biol Soc*. 2009; 2009:1663-1666.
70. Dries D, Reed MJ, Kissoon N, et al; on behalf of the Task Force for Mass Critical Care. Special populations: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. *Chest*. 2014;146(4_suppl):e75S-e86S.
71. Durairaj L, Will JG, Torner JC, Doebbeling BN. Prognostic factors for mortality following interhospital transfers to the medical intensive care unit of a tertiary referral center. *Crit Care Med*. 2003;31(7): 1981-1986.
72. MacKenzie EJ, Rivara FP, Jurkovich GJ, et al. A national evaluation of the effect of trauma-center care on mortality. *N Engl J Med*. 2006; 354(4):366-378.
73. Lilly CM, Fisher KA, Ries M, et al. A national ICU telemedicine survey: validation and results. *Chest*. 2012;142(1):40-47.
74. Vanholder R, Sever MS, Lameire N. The role of the Renal Disaster Relief Task Force. *Nat Clin Pract Nephrol*. 2008;4(7):347.
75. Vanholder R, Borniche D, Claus S, et al. When the earth trembles in the Americas: the experience of Haiti and Chile 2010. *Nephron Clin Pract*. 2011;117(3):c184-c197.
76. Lilly CM, Thomas EJ. Tele-ICU: experience to date. *J Intensive Care Med*. 2010;25(1):16-22.
77. ACS Committee on Trauma. *Resources for Optimal Care of the Injured Patient 2006*. Chicago, IL: American College of Surgeons; 2006.
78. Reynolds HN, Bander J, McCarthy M. Different systems and formats for tele-ICU coverage: designing a tele-ICU system to optimize functionality and investment. *Crit Care Nurs Q*. 2012;35(4):364-377.
79. Young LB, Chan PS, Lu X, Nallamothu BK, Sasson C, Cram PM. Impact of telemedicine intensive care unit coverage on patient outcomes: a systematic review and meta-analysis. *Arch Intern Med*. 2011;171(6):498-506.
80. Wilcox ME, Adhikari NK. The effect of telemedicine in critically ill patients: systematic review and meta-analysis. *Crit Care*. 2012;16(4): R127.
81. Zawada ET Jr, Herr P, Larson D, Fromm R, Kapaska D, Erickson D. Impact of an intensive care unit telemedicine program on a rural health care system. *Postgrad Med*. 2009;121(3):160-170.
82. Clay Mann N, Mullins RJ, Hedges JR, Rowland D, Arthur M, Zechin AD. Mortality among seriously injured patients treated in remote rural trauma centers before and after implementation of a statewide trauma system. *Med Care*. 2001;39(7):643-653.
83. Shafi S, Rayan N, Barnes S, Fleming N, Gentilello LM, Ballard D. Moving from "optimal resources" to "optimal care" at trauma centers. *J Trauma Acute Care Surg*. 2012;72(4):870-877.
84. Gabbe BJ, Simpson PM, Sutherland AM, Wolfe R, Lyons RA, Cameron PA. Evaluating time points for measuring recovery after major trauma in adults. *Ann Surg*. 2013;257(1):166-172.
85. Lattimer V, Brailsford S, Turnbull J, et al. Reviewing emergency care systems I: insights from system dynamics modelling. *Emerg Med J*. 2004;21(6):685-691.
86. Branas CC, MacKenzie EJ, ReVelle CS. A trauma resource allocation model for ambulances and hospitals. *Health Serv Res*. 2000;35(2):489-507.
87. Amram O, Schuurman N, Hedley N, Hameed SM. A web-based model to support patient-to-hospital allocation in mass casualty incidents. *J Trauma Acute Care Surg*. 2012;72(5):1323-1328.
88. Yi W, Özdamar L. A dynamic logistics coordination model for evacuation and support in disaster response activities. *Eur J Oper Res*. 2007;179(3):1177-1193.
89. Zhu Z, Hen BH, Teow KL. Estimating ICU bed capacity using discrete event simulation. *Int J Health Care Qual Assur*. 2012;25(2): 134-144.
90. Barthel ER, Pierce JR, Goodhue CJ, Ford HR, Grikscheit TC, Upperman JS. Availability of a pediatric trauma center in a disaster surge decreases triage time of the pediatric surge population: a population kinetics model. *Theor Biol Med Model*. 2011; 8(1):38.
91. Hirshberg A, Frykberg ER, Mattox KL, Stein M. Triage and trauma workload in mass casualty: a computer model. *J Trauma*. 2010;69(5): 1074-1081.
92. Levi L, Bregman D, Geva H, Revah M. Does number of beds reflect the surgical capability of hospitals in wartime and disaster? The use of a simulation technique at a national level. *Prehosp Disaster Med*. 1997;12(4):300-304.
93. Fawcett W, Oliveira CS. Casualty treatment after earthquake disasters: development of a regional simulation model. *Disasters*. 2000;24(3): 271-287.
94. Hupert N, Wattson D, Cuomo J, Hollingsworth E, Neukermans K, Xiong W. Predicting hospital surge after a large-scale anthrax attack: a model-based analysis of CDC's cities readiness initiative prophylaxis recommendations. *Med Decis Making*. 2009;29(4):424-437.
95. Smith SW, Portelli I, Narzisi G, et al. A novel approach to multihazard modeling and simulation. *Disaster Med Public Health Prep*. 2009;3(2): 75-87.
96. Powell T, Hanfling D, Gostin LO. Emergency preparedness and public health: the lessons of Hurricane Sandy. *JAMA*. 2012;308(24): 2569-2570.