Published in final edited form as:

FEBS J. 2015 July ; 282(14): 2646–2660. doi:10.1111/febs.13304.

Non-cognate DNA damage prevents formation of active conformation of Y-family DNA polymerases DinB and pol kappa

Philip Nevin, Xueguang Lu, Ke Zhang, John R. Engen, and Penny J. Beuning*

Department of Chemistry and Chemical Biology, Northeastern University, 360 Huntington Avenue, Boston, MA 02115, USA

Abstract

Y-family DNA polymerases are specialized to copy damaged DNA and are associated with increased mutagenesis due to their low fidelity. It is believed that the mechanism of nucleotide selection by Y-family DNA polymerases involves conformational changes preceding nucleotidyl transfer but there is limited experimental evidence for such structural changes. In particular, nucleotide-induced conformational changes in bacterial or eukaryotic Y-family DNA polymerases have to date not been extensively characterized. Using hydrogen/deuterium exchange mass spectrometry, we demonstrate here that the *Escherichia coli* Y-family DNA polymerase DinB and its human ortholog DNA polymerase κ undergo a conserved nucleotide-induced conformational change in the presence of undamaged DNA and the correct incoming nucleotide. Notably, this holds true for damaged DNA containing *N*²-furfuryl-deoxyguanosine that is efficiently copied by these two polymerases, but not for damaged DNA containing the major groove modification *O*⁶-methyl-deoxyguanosine that is a poor substrate. Our observations suggest that DinB and pol κ utilize a common mechanism for nucleotide selection involving a conserved pre-chemical conformational transition promoted by the correct nucleotide and only preferred DNA substrates.

Abstract

Correspondence. Penny J. Beuning, Department of Chemistry and Chemical Biology, Northeastern University; 360 Huntington Ave, 102 Hurtig Hall, Boston, MA 02115, USA, Fax: +1 617 373 8795, Tel: +1 617 373 2865, p.beuning@neu.edu.

Conflicts of interest

J.R.E. is a paid consultant of the Waters Corporation.

AUTHOR CONTRIBUTIONS

P.N., J.R.E., and P.J.B. conceived of and planned the experiments, analyzed the data, and wrote the paper. P.N. performed all the experiments. X.L. and K.Z. synthesized DNA substrates. All authors edited and approved the manuscript.

SUPPLEMENTARY INFORMATION

Fig. S1. Peptic peptides (gray bars) for which deuterium uptake was followed in HDX-MS experiments for DinB and pol κ

Fig. S2. Peptide-level HDX of free pol κ mapped on the apo-structure

Fig. S3. Differences in deuterium uptake between different states mapped on crystal structures

Fig. S4. Representative mass spectra of deuterated peptides

Table S1. Complete HDX data set with raw values

Keywords

Nucleotide selection; substrate specificity; conformational change; DNA replication; hydrogen exchange mass spectrometry

INTRODUCTION

Faithful DNA replication relies on the ability of DNA polymerases (pols) to select the correct nucleotide from a pool of potential substrates. The striking specificity of most pols far surpasses that expected solely from the thermodynamics of formation of any single base pair [1]. Indeed, structural and kinetic data indicate that many pols utilize conformational changes as kinetic checkpoints to select Watson-Crick base pairs [2, 3]. However, for damage bypass pols, such as those of the Y family, the mechanism of nucleotide selection has been less clear.

Y-family pols confer cellular resistance to DNA damage by carrying out translesion synthesis, in which they copy damage in the DNA template that normally blocks high-fidelity pols. This damage-tolerance mechanism is often associated with increased mutagenesis due to the relatively high error rate of Y-family pols [4]. The Y-family pol *Escherichia coli* DinB and its human ortholog DNA polymerase κ (pol κ) are specialized for bypass of certain bulky minor groove adducts at the N^2 position of guanine, such as N^2 -furfurylguanine (N^2 fG) [5] or those caused by polycyclic aromatic hydrocarbons [6–8], and are inhibited by certain major groove adducts [9, 10], although there are some differences in their specificities for different types of DNA damage [10, 11]. DinB has been shown to be required for generating mutations in *ampD* that confer resistance to β -lactam antibiotics [12], whereas pol κ has been shown to be involved in the tolerance of DNA interstrand cross-links induced by the chemotherapeutic agent mitomycin C [13].

The specialized functions of Y-family pols are related to their unique structural features. Like replicative DNA polymerases, members of the Y family adopt a fold resembling a right hand [14], but with a less structurally-constrained active site. In addition to the fingers

(nucleotide binding), palm (catalysis), and thumb (DNA binding) domains [14], Y-family pols possess a little finger (LF) domain (also known as polymerase-associated domain, PAD), which makes contacts with the major groove of DNA, albeit not at the nascent base pair, and may determine DNA damage specificity [4, 15–17]. In addition, the Y-family fingers domain is smaller than that of high-fidelity pols resulting in a wider active site that can accommodate lesions in the DNA template. DinB and the catalytic core of pol κ (sequence ID 32%) are structurally very similar (RMSD ~ 1 Å) with the exception of the N-clasp of pol κ , an N-terminal extension that wraps around the thumb domain and DNA and is important for polymerase activity [18, 19]. Domains beyond the catalytic core of pol κ are important for protein-protein interactions and cellular localization [20].

Comparison of crystal structures of A, B, X, and RT family pols in various stages of the catalytic cycle have revealed a dNTP-induced open-to-closed transition of the fingers domain preceding the chemistry step [3], which for many pols is believed to be a fidelity checkpoint. Whereas relatively large conformational changes of Y-family DNA polymerases are observed upon binding DNA [18, 21–24], crystal structures of binary pol-DNA and ternary pol-DNA-dNTP complexes of Y-family pols are highly similar, regardless of the complementarity of the incoming dNTP and the templating base [19, 23, 25–28], which has led to the suggestion that Y-family pols are in a closed conformation prior to dNTP binding [27, 29]. Yet, other kinetic studies have suggested a non-covalent step preceding chemistry for yeast pol η [30], human pol η [31], human pol κ [32], and the archaeal DinB orthologs *S. solfataricus* Dpo4 [33, 34] and Dbh [35]. Stopped-flow fluorescence studies of pre- and post-catalytic Dpo4 have indicated subtle movements of all four domains relative to DNA [36–38] and hydrogen/deuterium exchange (HDX) experiments of Dpo4 indicated dNTP-dependent changes in the dynamics of the fingers and thumb domains [39]. Only rather subtle dNTP-induced conformational changes in Y-family DNA pols have been observed. Indeed, the difference in RMSD in the crystal structures of DinB ternary complexes with correct or incorrect incoming nucleotide is 0.35–0.43 Å [19], which is similar (0.4 Å) to ternary complex structures of DinB bound to undamaged DNA or DNA containing *N*²-furfuryl-dG [40]. Dpo4 undergoes an apparent movement of domains upon binding DNA, but very little or no change in FRET efficiency is observed upon formation of a ternary complex [41]. Moreover, Dpo4 is relatively non-specific in its lesion bypass activity, whereas we and others have shown that DinB and pol κ are specific for minor groove adducts and discriminate against certain major groove adducts [9, 10, 42]. In light of the limited understanding of the role of conformational changes in the mechanism of nucleotide selection and the DNA damage specificity of bacterial and eukaryotic Y-family pols, it is of interest to characterize conformational changes in DinB and pol κ .

In this study, we probed the conformations of DinB and the catalytic core of pol κ in complex with undamaged and damaged DNA and various nucleotide substrates using HDX coupled with mass spectrometry (MS), as we recently carried out to probe sugar selectivity by these pols [43]. HDX-MS probes the accessibility and H-bonding of protein backbone amide hydrogens in solution, and is sensitive to structural changes that alter the environment of the backbone amide hydrogens [44–46]. Disruption of H-bonds, e.g. due to fast backbone fluctuations or slower structural rearrangements, may increase the probability of exchange

for the hydrogen involved in those bonds, resulting in unique HDX patterns for different protein conformations [47, 48]. HDX-MS can offer insight into the conformational dynamics of proteins that are complementary to those gained from crystallography [49], which can be biased towards the induced fit hypothesis of protein-ligand interactions. We have compared the extent of deuterium incorporation at the peptide level for pol alone, binary pol-DNA complexes, and ternary pol-DNA-dNTP complexes to test the hypothesis that the conformation of DinB and pol κ depend on the identity of the bound substrate. Our observations suggest that DinB and pol κ select the correct nucleotide via a common mechanism involving structural changes of regions both near to and far from the active site. In particular, these pols adopt a specific conformation only with the correct incoming nucleotide and with preferred DNA templates, that is, undamaged DNA or DNA containing N^2 fG as the template base; such changes are not observed with pols bound to damaged DNA that is a poor substrate, in this case, DNA with the major groove modification O^6 -methyl-dG (O^6 meG) as the template base, even in the presence of the correct incoming nucleotide. These observations provide new insights into the dynamics of Y-family pols and contribute to the understanding of pol nucleotide selection and DNA damage specificity.

RESULTS

To characterize the local dynamics of DinB and the catalytic core of pol κ , standard HDX-MS experiments were carried out [43, 50]. We exposed polymerase alone (E), binary pol-DNA complexes (ED), or ternary pol-DNA-dNTP complexes (EDN) to deuterated buffer, quenched the HDX reaction after periods of time ranging from 10 s to 4 h, digested the proteins into peptides under slow-exchange (quench) conditions, and determined the mass increase of each peptide by MS. The measured mass increase is due only to the deuteration of backbone amides because sidechain deuterons are washed away during the analysis due to fast HDX rates [45]. Although some deuterium (here between 25–35% based on deuterated peptide standards) is lost from peptides during analysis even in quench conditions, by comparing the deuteration level of all peptides under identical experimental conditions (a so-called relative deuterium exchange measurement), the loss of label was statistically the same for different forms (i.e., E, ED, EDN, etc.) of the proteins [50]. Measured differences in relative deuterium levels can therefore be attributed to conformational changes under the labeling conditions for each state. After analyzing the entire data set of undeuterated and deuterated peptides in all states, the sequence coverage for DinB and pol κ was 83% and 86%, respectively (Fig. S1).

Localized deuterium uptake of DinB and pol κ in the absence of substrates

To visualize the local deuterium uptake of the free pols in solution, we mapped the peptide-level HDX data for free DinB and free pol κ onto their crystal structures (Fig. 1), using the respective ternary complex crystal structures [18, 19] as there are no structural data for DinB in the absence of DNA and incoming nucleotide. Table S1 contains the relative deuterium levels for all peptides. Figure S2 shows the HDX data for free pol κ mapped on the structure of the apo-enzyme [22], which lacks most of the N-clasp. The deuterium levels for most regions of free DinB and free pol κ were relatively low after 10 s of exchange, indicating protection of backbone amides from HDX, as we previously reported [43]. The deuterium

levels increased over time with different rates in different parts of the proteins. For example, regions in the fingers domain reached relatively high deuterium levels after 10 min whereas most regions in the palm domain remained protected during the same period of time. The linker connecting the LF and thumb domains in DinB was almost fully deuterated after 10 s, consistent with the absence of secondary structure (we were not able to identify peptides in the corresponding linker of pol κ). The slowest exchanging regions of DinB were located in the LF and palm domains. We observed a similar trend in pol κ , with the palm domain exhibiting the slowest deuterium incorporation. In fact, DinB residues 122–133 and the corresponding pol κ residues 293–307, which are part of an α helix located on the surface of the palm domain, were the slowest exchanging region with almost no backbone amides exchanged after 4 h in D₂O. Similarly, previous HDX experiments by others with Dpo4 revealed the palm domain to be the most resistant to exchange in the free polymerase [39].

The N-clasp of pol κ (residues 19–67) as well as the large palm domain loop insertion (residues 222–283), which are not present in DinB, were fully deuterated after only 10 s of HDX (Figs. 1 and 2). This result indicates a high degree of exposure and dynamics which is consistent with the fact that the only pol κ construct crystallized in the absence of DNA is a truncation lacking residues 1–67 [22] and that the palm domain loop insertion is unresolved in pol κ crystal structures [18, 22]. In comparison to DinB, the thumb domain of pol κ exhibited slower HDX, consistent with the additional, and presumably stabilizing, contacts made on the outside of the pol κ thumb domain by the α helix that links the N-clasp to the palm (Fig. 2, peptides IV and V). Residues in this α helix were relatively protected (Fig. 2, IV and V), displaying deuterium uptake that is to be expected for structured regions of a protein. These observations are consistent with the idea that the N-clasp of pol κ is disordered in the absence of DNA, as hypothesized previously [18], and that structure in this unique N-terminal extension (pol κ residues 1–100) begins at residue 68 in the absence of DNA [22].

DNA binding protects DinB and pol κ from deuterium exchange

We next probed how DNA (a 2',3'-dideoxy-terminated 13-mer primer annealed to a 18-mer template) affects the deuterium uptake of DinB and pol κ by comparing the free pols (E) to a binary pol-DNA complex (ED) of each polymerase. We used a similar strategy to test how other substrates affect deuterium uptake as described in subsequent sections. For each comparison, HDX data were measured for two states (for example, E versus ED or ED versus EDN, Fig. 3a) and then the deuterium levels for each peptide at each deuterium exposure time in one state (e.g., ED) were subtracted from the deuterium levels for that same peptide and exposure time in another state (e.g., E), as shown at the top of each panel in Figure 3. Differences in deuterium level were color coded according to the scale shown. The differences were mapped onto the crystal structures (Fig. 3c,d,f,g, and Fig. S3) to visualize the three-dimensional locations of the changes in deuterium uptake in panels b and e of Figure 3. The entire data set for all comparisons with raw values is provided in Table S1.

The first comparison involved free pols (E) and their binary complexes (ED). Formation of a binary complex resulted in protection of several backbone amides in all domains of DinB

and pol κ (Fig. 3), and increased thermodynamic stability for both proteins (Fig. 4). Regions exhibiting lower HDX due to DNA binding were located in the large DNA binding cleft in the polymerases but also distal to DNA (e.g. DinB residues 16–24, 77–88, 132–147, and 262–270; pol κ residues 68–84, 115–133, 165–180, and 442–457; Fig. 3). Although differences in relative deuterium levels between the free pol and the binary complex followed similar trends overall for DinB and pol κ , the HDX of the palm and thumb domains of pol κ was less affected by DNA binding than those domains of DinB (compare Fig. 3b, left with 3e, left, and Fig. 3c and 3f). This observation is consistent with the relatively lower deuterium uptake of the palm and thumb domains of free pol κ (Fig. 1) and the relatively lower increase in T_m due to DNA binding (Fig. 4). Both DinB and pol κ LF domains showed strong protection from exchange upon DNA binding; consistent with this observation, the crystal structures of pol κ show a dramatic conformational change in the orientation of the LF (PAD) domain upon DNA binding [18, 22]. Deuterium incorporation in the N-terminal residues 19–31 of the pol κ N-clasp was unaffected by formation of the binary pol κ -DNA complex (Fig. 2, I and Fig. 3e). In contrast, N-clasp residues 54–67 and 68–84 exhibited significant protection in the binary complex and residues 32–41 showed modest protection in the presence of DNA (Fig. 2, III, IV and Fig. 3e), consistent with idea that the N-clasp is ordered only in the presence of DNA. Overall, our observations indicate that formation of the binary pol-DNA complex changes the conformation of DinB and pol κ from a relatively dynamic and/or open structure to a more rigid and/or less solvent accessible structure.

Ternary complex with correct dNTP is characterized by unique HDX pattern

Next we measured the deuterium incorporation of ternary pol-DNA-dNTP complexes (EDN) in the presence of different dNTPs to test how the conformation may change depending on the identity of the incoming dNTP. Formation of a ternary DinB complex in the presence of dGTP, which represents an incorrect dNTP, resulted in similar relative deuterium levels to those measured for the binary DinB complex (Fig. 3b, middle). Deuterium differences of 0.6–1.0 Da were observed for a few DinB peptides at only some time points (Fig. 3) and these changes were quite modest relative to those observed comparing free to binary pol-DNA complex. For all peptides with changes in deuterium incorporation, there was slightly more deuterium incorporation in the DinB ternary complex than in the binary complex. We made similar observations using dTTP as the incorrect dNTP, although the subtle changes we observed were protection rather than exposure, consistent with previous observations that pyrimidines are preferred by DinB as the incoming nucleotide [5] (Table S1). These observations indicate that the backbone dynamics and/or H-bonding network of DinB ternary complexes with incorrect dNTPs is similar to that of DinB binary complexes with no dNTP. In contrast to the DinB ternary complex with incorrect incoming nucleotide dGTP, the corresponding pol κ complex with incorrect dNTP showed protection from HDX relative to the binary complex, as lower deuterium levels were found for a few pol κ peptides throughout the protein (Fig. 3e, middle). A number of peptides representing four regions of pol κ showed reduced deuteration greater than 1 Da at a few time points in the incorrect pol κ ternary complex relative to the binary complex.

Thermal shift assays indicated an increased thermodynamic stability of DinB and pol κ upon formation of a correct ternary complex in relation to incorrect ternary complexes (Fig. 4). To further probe these changes, we measured the deuterium levels of ternary complexes with dCTP, which is the correct incoming dNTP based on the DNA sequence used. There was protection from HDX in a number of regions of both DinB (Fig. 3b, right and 3d) and pol κ (Fig. 3e, right and 3g) due to formation of a correct ternary complex. For both DinB (Fig. 3d) and pol κ (Fig. 3g), protection occurred in the fingers domain, which is located near dNTP binding site. In addition, certain regions of the thumb, palm, and LF domains were protected in both DinB and pol κ . Specifically, DinB palm residues 148–162 and the corresponding pol κ residues 320–332 were protected in the correct ternary complex in relation to the binary or incorrect ternary complex (Fig. 3b, right and Figure 3e, right). DinB thumb residues 173–192 and the corresponding pol κ residues 345–364 were also protected in the presence of correct dNTP. This region is located distal to the active site and does not make contacts with the dNTP, suggesting that the protection in this region represents an allosteric change in structure (Fig. 3d and 3g). For both DinB and pol κ , protection was also observed in a region of the LF domain making contacts with DNA 25–30 Å away from the active site (residues 283–307 in DinB; residues 419–431 and 442–477 in pol κ), further highlighting the allosteric nature of the structural changes associated with correct ternary complex formation. The HDX of the N-terminal residues of the N-clasp in pol κ (residues 19–49), which may make contacts with the active site and determine damage specificity [51], was not affected by binding to nucleotide substrates (Figs 2 and 3). However, N-clasp residues 54–67 were protected more in the presence of correct dNTPs than in the presence of incorrect dNTPs, suggesting a role for the N-clasp in specificity for the incoming nucleotide.

Only preferred template lesions and correct dNTPs promote “active” state protected from exchange

DinB and pol κ efficiently and accurately bypass the minor groove lesion *N*²-furfuryl-dG [5, 40] but are inhibited by major groove lesions [9, 10, 42]. Using primer extension assays, we confirmed that both DinB and pol κ are inhibited by *O*⁶meG as the template base [52] (Fig. 5a). In order to probe possible conformational changes in DinB and pol κ when binding to damaged DNA templates, we repeated HDX-MS experiments with DNA templates containing either *N*²fG or *O*⁶meG as the template base. The relative deuterium levels of binary DinB complexes with DNA templates containing either *O*⁶meG or *N*²fG were similar to those of the binary complex with undamaged DNA (Fig. 5b, top). For pol κ , the presence of either the *O*⁶meG or *N*²fG lesions resulted in lower HDX for a few regions in the N-clasp, thumb, and LF (Fig. 5b, bottom), although the HDX for the majority of identified peptides was unaffected. Binding of the incorrect nucleotide dGTP to the binary complexes with the *N*²fG template (Fig. 5b, middle panels) resulted in only subtle changes in the HDX, as observed also with undamaged DNA (Fig. 3). On the other hand, for both DinB and pol κ , the correct ternary complex with the *N*²fG template and dCTP exhibited deuterium incorporation similar to the ternary complex in the presence of undamaged DNA and the correct nucleotide dCTP (Fig. 5b, middle panels).

Both dTTP and dCTP can form stable base pairs with *O*⁶meG [53] and pol κ inserts dCTP and dTTP opposite *O*⁶meG [42, 52]. The ternary complex of DinB or pol κ in the presence

of the O^6 meG template and the correct nucleotide dTTP or dCTP exhibited similar deuterium uptake as the incorrect ternary complexes (Fig. 5b, right panels). In agreement with this observation both DinB and pol κ were inhibited by O^6 meG relative to N^2 fG or undamaged DNA in primer extension assays (Fig. 5a). Thus, only the correct nucleotide in the presence of DNA templates that are preferred substrates, including damaged DNA, induces specific HDX changes in DinB and pol κ . This suggests that the changes in deuterium incorporation we observed in the presence of the correct dNTP and preferred DNA, either undamaged DNA or DNA containing a lesion that is efficiently bypassed by DinB and pol κ , represent a structural change in the polymerases towards an “active” state. A lesion that cannot be bypassed, O^6 meG, even with incoming dNTPs that can form a base pair, did not induce any HDX changes.

DISCUSSION

The substrate specificity of pols governs the fidelity of genome duplication and is believed to be achieved by a mechanism involving substrate-induced or -selected conformational changes of the polymerase prior to the chemistry step [3], although the rate-limiting step in the reaction cycle is under debate [54–56]. At first glance, the relatively high error frequencies of Y-family pols may appear consistent with the lack of distinct conformations upon forming the correct base pair in the available crystal structures [18, 19, 23, 25–28] since distinct conformations would require well-defined end states separated by a free energy barrier that could act as a kinetic checkpoint for substrate specificity. However, the nucleotide selectivity of DinB and pol κ surpasses that expected solely from the energetics of base pairing by several orders of magnitude [57, 58] leading to the hypothesis that their intrinsic molecular features govern nucleotide selection.

Structural information for DinB is limited to the ternary complex [19], but for pol κ [18, 22] and Dpo4 [23] comparison of apo and ternary crystal structures suggest a large movement of the LF domain upon DNA binding. The prominent protection that we observed in the LF of DinB and pol κ upon DNA binding is consistent with a transition of the LF towards DNA. However, our observations indicate that the LF is intrinsically protected from HDX in the absence of DNA, consistent with a globular and/or stable structure. Thus, the LF is probably only mobile in the sense that it can undergo global transitions because it is attached to the pol via a flexible linker. Our observation that the HDX in the palm domain was relatively low (Fig. 1) and relatively unaffected by DNA binding (Fig. 3) points to a model where the palm domain acts as a stable core with the more mobile fingers, thumb and LF domains binding the DNA duplex. Thus, the human right hand analogy translates both to the tertiary structures and dynamics of DinB and pol κ . The analysis of Dpo4 similarly showed more modest changes in HDX of the palm domain than in other domains in the presence of DNA [39]. In contrast to our observation that no regions of DinB or pol κ showed increased HDX in the presence of unmodified DNA, the linker between the polymerase and the little finger domains of Dpo4 exhibited increased HDX in the presence of DNA [39]. Thus, while we observed conserved changes in HDX in DinB and pol κ with a range of substrates, there are some modest differences between these two polymerases and Dpo4.

Crystal structures of Y-family pols in ternary complexes likely represent structural snapshots of well-defined low-energy states in the reaction pathway. We hypothesized that there exist additional conformations of DinB and pol κ in solution. Indeed, our HDX-MS data indicate that for DinB and pol κ , ternary complexes in the presence of preferred template bases and the correct dNTP adopt a unique “active” state that is characterized by reduced HDX in several regions of the proteins, including regions distal to the dNTP binding site (Fig. 3). Notably, this holds true for damaged DNA that is efficiently copied by these two polymerases, but not for a DNA template that is a poor substrate (Fig. 5). The formation of this “active” state involves structural changes in all domains of DinB and pol κ rather than in any specific domain.

The question remains as to what the cause of reduced HDX is in the “active” state. A reduction in the rate of HDX due to correct dNTP binding could be caused by a reduction in the intrinsic breathing motions of the polymerase structure in proximity to the dNTP so that those regions visit exposed, HDX-competent states less frequently. However, our observations that the dNTP-dependent HDX changes in DinB and pol κ are specific and occur both proximal and distal to the active site suggest that formation of the “active” state also involves a conformational change resulting in a different H-bond pattern causing slower HDX. This would be analogous to the dNTP-induced conformational change observed for A-, B-, RT-, and X-family pols [3]. The significant reduction of HDX in the fingers domain due to correct dNTP binding (Figs. 3 and 5) is consistent with a closing transition although only atomic-level structural studies can determine the exact nature of this structural change. It is also possible that the “active” conformation of DinB and pol κ exists in the absence of substrates when the pols are in their free and dynamic states, as has been shown for pol I [59]. In accordance with the conformational selection hypothesis [60, 61] the “active” conformation would then be promoted or selected by the correct dNTP and preferred DNA substrates.

Two prominent regions involved in the correct dNTP-associated conformation of DinB and pol κ are located in the palm (DinB residues 148–162; pol κ residues 320–332) and thumb (DinB residues 173–192; pol κ residues 345–364). The palm contains a conserved lysine (DinB K157; pol κ K328) that may make contacts with the terminal phosphate of the dNTP [18, 19] and has been shown to be important for activity of DinB [62]. Because these two regions are connected via numerous sidechain interactions, it is possible that correct dNTP binding induces or selects a structural change in this region of DinB and pol κ to align important active site residues and the dNTP prior to catalysis. The dNTP-induced HDX changes in this region are strikingly similar between DinB and pol κ and a previous study indicated that the corresponding palm and thumb regions in the archaeal DinB ortholog Dpo4 exhibit dNTP-dependent changes in HDX [39]. Thus, dNTP-dependent changes in HDX in these regions of the palm and thumb appear to be general for Y-family pols of the DinB subfamily. On the other hand, two α -helices in the thumb domain of Dpo4 were sensitive to the identity of dNTP, resulting in greater exposure to HDX in the presence of the correct dNTP as well as the incorrect dATP [39]; only subtle exposure in this region of the thumb domain was observed for DinB, mainly with the incorrect dNTP (Figs. 3 and 5). Furthermore, it has been shown that Dpo4 can exist in a binary preinsertion complex, in which the terminal nucleotide of the primer occupies the dNTP binding site [63] and it is

possible that DinB and pol κ may transition between the binary complex and a similar preinsertion complex. The conformational change upon correct dNTP binding would then involve polymerase translocation or rotation of the DNA to accommodate the nascent base pair, which could explain the allosteric changes in the LF domain, located 25–30 Å from the active site. A similar translocation event has been proposed to occur during ternary complex formation of yeast pol η [21].

We propose that the “active” conformation that we identify here by HDX-MS represents a state similar to the ternary complex structures of DinB and pol κ observed by crystallography [18, 19], and that DinB and pol κ do not exist in stable closed conformations in the absence of DNA and correct nucleotide substrates. Figure 6 shows a proposed reaction pathway for DinB and pol κ with the favored protein conformation at each step indicated. First, the free pol binds DNA and forms a relatively closed and stable binary complex. Upon ternary complex formation, there is a conserved conformational change from an “inactive” ternary complex to an “active” ternary complex that is favored only for correct dNTP incorporation opposite preferred DNA lesions or undamaged DNA. In the case of incorrect dNTP incorporation or non-preferred DNA damage, a non-productive ternary complex or dNTP dissociation is favored, resulting in a slower reaction. This model is similar to that proposed for many high-fidelity pols [64]. It is possible that the different conformations inferred from our HDX studies may be present as minority populations in any of the states. Thus, the “active” conformation may or may not be an intermediate in the reaction pathway for incorrect dNTPs or non-preferred DNA.

Here we demonstrate that the structural dynamics of DinB and pol κ change upon substrate binding. This highlights that Y-family pols exist in an ensemble of conformations that depends on the identity of the bound substrates. The unique HDX profile observed only in the presence of non-blocking template bases and the correct dNTP suggests the existence of a functionally relevant conformational change related to nucleotide selection. These observations indicate that DinB and pol κ utilize a common mechanism for nucleotide selection involving a conserved structural change preceding the chemistry step.

MATERIALS AND METHODS

Proteins and DNA

E. coli DinB and the catalytic core of human pol κ (linker residues GPGS followed by pol κ residues 19–526, hereafter referred to as pol κ , a gift from Prof. Fred Guengerich, Vanderbilt University) were purified as described previously [65, 66]. Protein purity and intact mass was verified using an LCT-premier mass spectrometer with an electrospray ionization source (Waters, Milford, MA, USA). Unmodified DNA was from Eurofins Operon (Huntsville, AL, USA); DNA containing modified bases was made with reagents from Glen Research (Sterling, VA, USA). The sequence of the 13-mer primer was 5'-GCTTGCCGGACGC-3'. HDX and thermal shift experiments that included DNA were performed with a 2',3'-dideoxy-terminated primer to avoid nucleotide incorporation, as described previously [18, 39]. All DNA templates had the sequence 5'-CCTGGXCGTCCGGCAACG-3' where X was G or a modified base. The template containing a single *O*⁶-methylguanine was prepared by solid-phase synthesis using *O*⁶meG

phosphoramidite (Glen Research) according to manufacturer instructions. The template containing a single *N*²-furfurylguanine was prepared by the convertible nucleoside approach [67] after synthesis of an oligonucleotide with 2-fluoro-*O*⁶-(*p*-nitrophenylethyl)- 2'-deoxyinosine phosphoramidite (Glen Research) at the desired position. Briefly, fully protected resin-bound oligonucleotide was treated with furfurylamine (Alfa Aesar, Ward Hill, MA, USA) at 22 °C for 48 h followed by deprotection using first 1,8-diazabicyclo[5.4.0]undec-7-ene (Fluka, St. Louis, MO, USA) and then ammonium hydroxide (Fisher, Pittsburgh, PA, USA). All oligonucleotides were purified by denaturing polyacrylamide gel electrophoresis using the crush and soak method [68].

HDX-MS

Proteins (25 μM) were allowed to equilibrate in the presence or absence of primer/template DNA (46 μM) and dNTPs (5 mM) in 20 μL of binding buffer (25 mM Hepes, pH 7.5, 20 mM NaCl, 5 mM MgSO₄, and 5 mM dithiothreitol) for at least 30 min at 22 °C. Based on reported dissociation constants [9, 18], the percent DNA-bound polymerase was >98%. HDX reactions were initiated by diluting 3 μL of the mixture 17-fold with D₂O buffer (25 mM Hepes, pH 7.5, 20 mM NaCl, 5 mM MgSO₄) at 22 °C and quenched after defined periods of time (10 s, 1 min, 10 min, 1 h, and 4 h) by addition of an equal volume of ice-cold quench buffer (0.1 M sodium phosphate, 4 M GuHCl, 0.05 M TCEP, pH 2.0). The final pH of the quenched reactions was 2.7.

The quenched HDX reactions were immediately injected into a nanoACQUITY system with HDX technology for UPLC separation (Waters) [69] and passed through a 2.1 mm × 50 mm stainless-steel column packed with immobilized pepsin [70] with a flow rate of 100 μL/min in 0.1% formic acid at 15 °C. Peptides were trapped and desalted on an ACQUITY UPLC BEH C18 1.7-μm 2.1 mm × 5 mm VanGuard Pre-column (Waters) kept at 0 °C. After 4 min, the trap was placed in line with an ACQUITY UPLC BEH C18 1.7-μm 1.0 mm × 100 mm column (Waters) and peptides were separated at 0 °C using a 7 min linear gradient (8 – 40% acetonitrile, 0.1% formic acid, pH 2.5) at a flow rate of 40 μL/min and directed into a Synapt G1 mass spectrometer with an electrospray ionization source (Waters). Mass spectra were acquired in positive ion mode over an *m/z* range of 50 – 1,900 under optimized deuterium recovery conditions. Due to poor response in positive ionization electrospray, DNA were not monitored during HDX-MS. Continuous lock-mass correction was performed using Glu-fibrinogen B peptide standard (Sigma, St. Louis, MO, USA).

Peptide identification and data analysis

The peptides produced in triplicate undeuterated control samples were identified by a combination of accurate mass and collision-induced dissociation in data-independent acquisition mode (MS^E) [71] aided by PLGS software (Waters). Only peptides identified in at least two replicates were included in the analysis. Mass spectra were extracted and analyzed using DynamX 2.0 (Waters). Representative mass spectra are shown in Figure S4. Relative deuterium levels were calculated by subtracting the centroid of the isotope distribution of the undeuterated peptide from that of deuterated peptides. As all measurements were comparisons between different biological or bound states, and were performed under nearly identical experimental conditions, the deuterium levels were not

corrected for back-exchange; therefore, all results are reported as relative, as described further in Ref. [50]. The fractional relative deuterium uptake was calculated by dividing the relative deuterium uptake of each peptide by its number of backbone amides after subtracting one for the N-terminus and one for each proline residue [45]. To assess the statistical significance of measured differences, we determined the uncertainty in the relative deuterium uptake with triplicate independent HDX-MS experiments for DinB, DinB-DNA, DinB-DNA-dCTP, free pol κ , pol κ -DNA-dGTP, and pol κ -DNA-dCTP on separate days and with multiple different preparations of protein, DNA, and labeling buffer. The mean intensity-weighted standard deviation of the relative deuterium uptake was 0.15 Da and was calculated for each peptide and time point using multiple charge states in each replicate. A 98% confidence interval for the mean relative deuterium uptake of ± 0.6 Da was calculated as described [72] and used as a threshold for significance, i.e. changes larger than 0.6 Da were considered significant at 98% confidence.

Thermal shift assays

Samples containing 3–10 μ M protein in the presence or absence of primer/template DNA (2-fold molar excess) and dNTP (1 mM) in 15 μ L assay buffer (25 mM Hepes, pH 7.5, 20 mM NaCl, 5 mM MgSO₄, and 3 mM dithiothreitol) were incubated for 30 min at room temperature before addition of 0.5 μ L of Sypro Orange (Invitrogen, Grand Island, NY, USA) to a final concentration of 10 $\times$. Samples were analyzed using a CFX 96 Real-Time system (Bio-Rad, Hercules, CA, USA) as described [73].

Primer extension assays

Primer extension assays were carried out as described previously [9] using ³²P-labeled 13-mer primer annealed to 18-mer template. Reactions contained a final concentration of 10 nM DinB or pol κ , 100 nM primer/template DNA, 1 mM dNTPs, 7.5 mM MgSO₄, 30 mM Hepes (pH 7.5), 20 mM NaCl, 2 mM β -mercaptoethanol, 1% (w/v) bovine serum albumin, and 4% glycerol. Reaction products were separated by denaturing 16% polyacrylamide gel electrophoresis and analyzed by phosphorimaging.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We give special thanks to Prof. Fred Guengerich (Vanderbilt University) for providing the pol κ expression plasmid and to Prof. Robert Eoff (University of Arkansas) for helpful advice. This work was supported by generous financial support from the National Science Foundation [CAREER Award, MCB-0845033 to P.J.B.]; the National Institutes of Health [R01-GM101135 to J.R.E.]; a research collaboration with the Waters Corporation, the American Cancer Society [Research Scholar Grant RSG-12-161-01-DMC to P.J.B.]; Research Corporation for Science Advancement [Cottrell Scholar Award to P.J.B.]; and the Northeastern University Office of the Provost.

Abbreviations

pol	DNA polymerase
LF	little finger

N²fG	N ² -furfurylguanine
O⁶meG	O ⁶ -methylguanine
HDX	hydrogen/deuterium exchange

REFERENCES

1. McCulloch SD, Kunkel TA. The fidelity of DNA synthesis by eukaryotic replicative and translesion synthesis polymerases. *Cell Res.* 2008; 18:148–161. [PubMed: 18166979]
2. Echols H, Goodman MF. Fidelity mechanisms in DNA replication. *Annu Rev Biochem.* 1991; 60:477–511. [PubMed: 1883202]
3. Joyce CM, Benkovic SJ. DNA polymerase fidelity: kinetics, structure, and checkpoints. *Biochemistry.* 2004; 43:14317–14324. [PubMed: 15533035]
4. Goodman MF, Woodgate R. Translesion DNA polymerases. *Cold Spring Harb Perspect Biol.* 2013; 5:a010363. [PubMed: 23838442]
5. Jarosz DF, Godoy VG, DeLaney JC, Essigmann JM, Walker GC. A single amino acid governs enhanced activity of DinB DNA polymerases on damaged templates. *Nature.* 2006; 439:225–228. [PubMed: 16407906]
6. Shen X, Sayer JM, Kroth H, Ponten I, O'Donnell M, Woodgate R, Jerina DM, Goodman MF. Efficiency and accuracy of SOS-induced DNA polymerases replicating benzo[a]pyrene-7,8-diol 9,10-epoxide A and G adducts. *J Biol Chem.* 2002; 277:5265–5274. [PubMed: 11734560]
7. Avkin S, Goldsmith M, Velasco-Miguel S, Geacintov N, Friedberg EC, Livneh Z. Quantitative analysis of translesion DNA synthesis across a benzo[a]pyrene-guanine adduct in mammalian cells: the role of DNA polymerase kappa. *J Biol Chem.* 2004; 279:53298–53305. [PubMed: 15475561]
8. Yuan B, Cao H, Jiang Y, Hong H, Wang Y. Efficient and accurate bypass of N2-(1-carboxyethyl)-2'-deoxyguanosine by DinB DNA polymerase in vitro and in vivo. *Proc Natl Acad Sci U S A.* 2008; 105:8679–8684. [PubMed: 18562283]
9. Walsh JM, Bouamaied I, Brown T, Wilhelmsson LM, Beuning PJ. Discrimination against the cytosine analog tC by *Escherichia coli* DNA polymerase IV DinB. *J Mol Biol.* 2011; 409:89–100. [PubMed: 21477595]
10. Walsh JM, Ippoliti PJ, Ronayne EA, Rozners E, Beuning PJ. Discrimination against major groove adducts by Y-family polymerases of the DinB subfamily. *DNA repair.* 2013; 12:713–722. [PubMed: 23791649]
11. Suzuki N, Okashi E, Hayashi K, Ohmori H, Grollman AP, Shibutani S. Translesional Synthesis Past Acetylaminofluorene-Derived DNA Adducts Catalyzed by Human DNA Polymerase κ and *Escherichia coli* DNA Polymerase IV. *Biochemistry.* 2001; 40:15176–15183. [PubMed: 11735400]
12. Petrosino JF, Galhardo RS, Morales LD, Rosenberg SM. Stress-induced beta-lactam antibiotic resistance mutation and sequences of stationary-phase mutations in the *Escherichia coli* chromosome. *J Bacteriol.* 2009; 191:5881–5889. [PubMed: 19648247]
13. Minko IG, Harbut MB, Kozekov ID, Kozekova A, Jakobs PM, Olson SB, Moses RE, Harris TM, Rizzo CJ, Lloyd RS. Role for DNA polymerase kappa in the processing of N2-N2-guanine interstrand cross-links. *J Biol Chem.* 2008; 283:17075–17082. [PubMed: 18434313]
14. Steitz TA. DNA polymerases: structural diversity and common mechanisms. *J Biol Chem.* 1999; 274:17395–17398. [PubMed: 10364165]
15. Pata JD. Structural diversity of the Y-family DNA polymerases. *Biochim Biophys Acta.* 2010; 1804:1124–1135. [PubMed: 20123134]
16. Wilson RC, Jackson MA, Pata JD. Y-family polymerase conformation is a major determinant of fidelity and translesion specificity. *Structure.* 2013; 21:20–31. [PubMed: 23245850]
17. Boudsocq F, Kokoska RJ, Plosky BS, Vaisman A, Ling H, Kunkel TA, Yang W, Woodgate R. Investigating the role of the little finger domain of Y-family DNA polymerases in low fidelity synthesis and translesion replication. *J Biol Chem.* 2004; 279:32932–32940. [PubMed: 15155753]

18. Lone S, Townson SA, Uljon SN, Johnson RE, Brahma A, Nair DT, Prakash S, Prakash L, Aggarwal AK. Human DNA polymerase kappa encircles DNA: implications for mismatch extension and lesion bypass. *Mol Cell*. 2007; 25:601–614. [PubMed: 17317631]
19. Sharma A, Kottur J, Narayanan N, Nair DT. A strategically located serine residue is critical for the mutator activity of DNA polymerase IV from *Escherichia coli*. *Nucleic Acids Res*. 2013; 41:5104–5114. [PubMed: 23525461]
20. Ohmori H, Ohashi E, Ogi T. Mammalian Pol kappa: regulation of its expression and lesion substrates. *Adv Protein Chem*. 2004; 69:265–278. [PubMed: 15588846]
21. Alt A, Lammens K, Chiocchini C, Lammens A, Pieck JC, Kuch D, Hopfner KP, Carell T. Bypass of DNA lesions generated during anticancer treatment with cisplatin by DNA polymerase eta. *Science*. 2007; 318:967–970. [PubMed: 17991862]
22. Uljon SN, Johnson RE, Edwards TA, Prakash S, Prakash L, Aggarwal AK. Crystal Structure of the Catalytic Core of Human DNA Polymerase Kappa. *Structure*. 2004; 12:1395–1404. [PubMed: 15296733]
23. Wong JH, Fiala KA, Suo Z, Ling H. Snapshots of a Y-family DNA polymerase in replication: substrate-induced conformational transitions and implications for fidelity of Dpo4. *J Mol Biol*. 2008; 379:317–330. [PubMed: 18448122]
24. Silverstein TD, Johnson RE, Jain R, Prakash L, Prakash S, Aggarwal AK. Structural basis for the suppression of skin cancers by DNA polymerase eta. *Nature*. 2010; 465:1039–1043. [PubMed: 20577207]
25. Ling H, Boudsocq F, Woodgate R, Yang W. Crystal Structure of a Y-Family DNA Polymerase in Action: A Mechanism for Error-Prone and Lesion-Bypass Replication. *Cell*. 2001; 107:91–102. [PubMed: 11595188]
26. Zhang H, Eoff RL, Kozekov ID, Rizzo CJ, Egli M, Guengerich FP. Structure-function relationships in miscoding by *Sulfolobus solfataricus* DNA polymerase Dpo4: guanine N2,N2-dimethyl substitution produces inactive and miscoding polymerase complexes. *J Biol Chem*. 2009; 284:17687–17699. [PubMed: 19542237]
27. Ummat A, Silverstein TD, Jain R, Buku A, Johnson RE, Prakash L, Prakash S, Aggarwal AK. Human DNA polymerase eta is pre-aligned for dNTP binding and catalysis. *J Mol Biol*. 2012; 415:627–634. [PubMed: 22154937]
28. Biertumpfel C, Zhao Y, Kondo Y, Ramon-Maiques S, Gregory M, Lee JY, Masutani C, Lehmann AR, Hanaoka F, Yang W. Structure and mechanism of human DNA polymerase eta. *Nature*. 2010; 465:1044–1048. [PubMed: 20577208]
29. Zhou BL, Pata JD, Steitz TA. Crystal structure of a DinB lesion bypass DNA polymerase catalytic fragment reveals a classic polymerase catalytic domain. *Mol Cell*. 2001; 8:427–437. [PubMed: 11545744]
30. Washington MT, Prakash L, Prakash S. Yeast DNA polymerase eta utilizes an induced-fit mechanism of nucleotide incorporation. *Cell*. 2001; 107:917–927. [PubMed: 11779467]
31. Washington MT, Johnson RE, Prakash L, Prakash S. The mechanism of nucleotide incorporation by human DNA polymerase eta differs from that of the yeast enzyme. *Mol Cell Biol*. 2003; 23:8316–8322. [PubMed: 14585988]
32. Zhao L, Pence MG, Eoff RL, Yuan S, Fercu CA, Guengerich FP. Elucidation of kinetic mechanisms of human translesion DNA polymerase kappa using tryptophan mutants. *FEBS J*. 2014; 281:4394–4410. [PubMed: 25065501]
33. Fiala KA, Suo Z. Mechanism of DNA polymerization catalyzed by *Sulfolobus solfataricus* P2 DNA polymerase IV. *Biochemistry*. 2004; 43:2116–2125. [PubMed: 14967051]
34. Zhang H, Guengerich FP. Effect of N2-guanyl modifications on early steps in catalysis of polymerization by *Sulfolobus solfataricus* P2 DNA polymerase Dpo4 T239W. *J Mol Biol*. 2010; 395:1007–1018. [PubMed: 19969000]
35. DeLucia AM, Grindley ND, Joyce CM. Conformational changes during normal and error-prone incorporation of nucleotides by a Y-family DNA polymerase detected by 2-aminopurine fluorescence. *Biochemistry*. 2007; 46:10790–10803. [PubMed: 17725324]
36. Xu C, Maxwell BA, Brown JA, Zhang L, Suo Z. Global conformational dynamics of a Y-family DNA polymerase during catalysis. *PLoS Biol*. 2009; 7:e1000225. [PubMed: 19859523]

37. Beckman JW, Wang Q, Guengerich FP. Kinetic analysis of correct nucleotide insertion by a Y-family DNA polymerase reveals conformational changes both prior to and following phosphodiester bond formation as detected by tryptophan fluorescence. *J Biol Chem.* 2008; 283:36711–36723. [PubMed: 18984592]
38. Maxwell BA, Xu C, Suo Z. DNA lesion alters global conformational dynamics of Y-family DNA polymerase during catalysis. *J Biol Chem.* 2012; 287:13040–13047. [PubMed: 22362779]
39. Eoff RL, Sanchez-Ponce R, Guengerich FP. Conformational changes during nucleotide selection by *Sulfolobus solfataricus* DNA polymerase Dpo4. *J Biol Chem.* 2009; 284:21090–21099. [PubMed: 19515847]
40. Kottur J, Sharma A, Gore KR, Narayanan N, Samanta B, Pradeepkumar PI, Nair DT. Unique Structural Features in DNA Polymerase IV Enable Efficient Bypass of the N(2) Adduct Induced by the Nitrofurazone Antibiotic. *Structure.* 2015; 23:56–67. [PubMed: 25497730]
41. Maxwell BA, Xu C, Suo Z. Conformational Dynamics of a Y-Family DNA Polymerase during Substrate Binding and Catalysis As Revealed by Interdomain Forster Resonance Energy Transfer. *Biochemistry.* 2014; 53:1768–1778. [PubMed: 24568554]
42. Choi JY, Chowdhury G, Zang H, Angel KC, Vu CC, Peterson LA, Guengerich FP. Translesion synthesis across O6-alkylguanine DNA adducts by recombinant human DNA polymerases. *J Biol Chem.* 2006; 281:38244–38256. [PubMed: 17050527]
43. Nevin P, Engen JR, Beuning PJ. Steric gate residues of Y-family DNA polymerases DinB and pol kappa are crucial for dNTP-induced conformational change. *DNA repair.* 2015
44. Katta V, Chait BT. Conformational changes in proteins probed by hydrogen-exchange electrospray-ionization mass spectrometry. *Rapid Commun Mass Spectrom.* 1991; 5:214–217. [PubMed: 1666528]
45. Zhang Z, Smith DL. Determination of amide hydrogen exchange by mass spectrometry: a new tool for protein structure elucidation. *Protein Sci.* 1993; 2:522–531. [PubMed: 8390883]
46. Konermann L, Vahidi S, Sowole MA. Mass spectrometry methods for studying structure and dynamics of biological macromolecules. *Anal Chem.* 2014; 86:213–232. [PubMed: 24304427]
47. Englander SW, Downer NW, Teitelbaum H. Hydrogen exchange. *Annu Rev Biochem.* 1972; 41:903–924. [PubMed: 4563445]
48. Percy AJ, Rey M, Burns KM, Schriemer DC. Probing protein interactions with hydrogen/deuterium exchange and mass spectrometry—a review. *Anal Chim Acta.* 2012; 721:7–21. [PubMed: 22405295]
49. Badireddy S, Yunfeng G, Ritchie M, Akamine P, Wu J, Kim CW, Taylor SS, Qingsong L, Swaminathan K, Anand GS. Cyclic AMP analog blocks kinase activation by stabilizing inactive conformation: conformational selection highlights a new concept in allosteric inhibitor design. *Mol Cell Proteomics.* 2011; 10:M110 004390. [PubMed: 21081668]
50. Wales TE, Engen JR. Hydrogen exchange mass spectrometry for the analysis of protein dynamics. *Mass Spectrom Rev.* 2006; 25:158–170. [PubMed: 16208684]
51. Jia L, Geacintov NE, Broyde S. The N-clasp of human DNA polymerase kappa promotes blockage or error-free bypass of adenine- or guanine-benzo[a]pyrenyl lesions. *Nucleic Acids Res.* 2008; 36:6571–6584. [PubMed: 18931375]
52. Haracska L, Prakash L, Prakash S. Role of human DNA polymerase kappa as an extender in translesion synthesis. *Proc Natl Acad Sci U S A.* 2002; 99:16000–16005. [PubMed: 12444249]
53. Warren JJ, Forsberg LJ, Beese LS. The structural basis for the mutagenicity of O(6)-methyl-guanine lesions. *Proc Natl Acad Sci U S A.* 2006; 103:19701–19706. [PubMed: 17179038]
54. Schlick T, Arora K, Beard WA, Wilson SH. Perspective: pre-chemistry conformational changes in DNA polymerase mechanisms. *Theor Chem Acc.* 2012; 131:1287–1295. [PubMed: 23459563]
55. Prasad BR, Kamerlin SCL, Florián J, Warshel A. Prechemistry barriers and checkpoints do not contribute to fidelity and catalysis as long as they are not rate limiting. *Theor Chem Acc.* 2012; 131:1288–1302.
56. Maxwell BA, Suo Z. Recent insight into the kinetic mechanisms and conformational dynamics of Y-Family DNA polymerases. *Biochemistry.* 2014; 53:2804–2814. [PubMed: 24716482]

57. Kobayashi S, Valentine MR, Pham P, O'Donnell M, Goodman MF. Fidelity of *Escherichia coli* DNA polymerase IV. Preferential generation of small deletion mutations by dNTP-stabilized misalignment. *J Biol Chem*. 2002; 277:34198–34207. [PubMed: 12097328]
58. Ohashi E, Bebenek K, Matsuda T, Feaver WJ, Gerlach VL, Friedberg EC, Ohmori H, Kunkel TA. Fidelity and processivity of DNA synthesis by DNA polymerase kappa, the product of the human DINB1 gene. *J Biol Chem*. 2000; 275:39678–39684. [PubMed: 11006276]
59. Rothwell PJ, Mitaksov V, Waksman G. Motions of the fingers subdomain of klenTaq1 are fast and not rate limiting: implications for the molecular basis of fidelity in DNA polymerases. *Mol Cell*. 2005; 19:345–355. [PubMed: 16061181]
60. Boehr DD, Nussinov R, Wright PE. The role of dynamic conformational ensembles in biomolecular recognition. *Nat Chem Biol*. 2009; 5:789–796. [PubMed: 19841628]
61. Nussinov R, Tsai CJ. Unraveling structural mechanisms of allosteric drug action. *Trends Pharmacol Sci*. 2014; 35:256–264. [PubMed: 24742712]
62. Walsh JM, Parasuram R, Rajput PR, Rozners E, Ondrechen MJ, Beuning PJ. Effects of non-catalytic, distal amino acid residues on activity of *E. coli* DinB (DNA polymerase IV). *Environ Mol Mutagen*. 2012; 53:766–776. [PubMed: 23034734]
63. Rechkoblit O, Malinina L, Cheng Y, Kuryavii V, Broyde S, Geacintov NE, Patel DJ. Stepwise translocation of Dpo4 polymerase during error-free bypass of an oxoG lesion. *PLoS Biol*. 2006; 4:e11. [PubMed: 16379496]
64. Tsai YC, Johnson KA. A new paradigm for DNA polymerase specificity. *Biochemistry*. 2006; 45:9675–9687. [PubMed: 16893169]
65. Beuning PJ, Simon SM, Godoy VG, Jarosz DF, Walker GC. Characterization of *Escherichia coli* translesion synthesis polymerases and their accessory factors. *Methods Enzymol*. 2006; 408:318–340. [PubMed: 16793378]
66. Irimia A, Eoff RL, Guengerich FP, Egli M. Structural and functional elucidation of the mechanism promoting error-prone synthesis by human DNA polymerase kappa opposite the 7,8-dihydro-8-oxo-2'-deoxyguanosine adduct. *J Biol Chem*. 2009; 284:22467–22480. [PubMed: 19542228]
67. Erlanson DA, Chen L, Verdine GL. DNA Methylation through a Locally Unpaired Intermediate. *J Am Chem Soc*. 1993; 115:12583–12584.
68. Sambrook, J.; Fritsch, EF.; Maniatis, T. *Molecular Cloning--A laboratory manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 1989.
69. Wales TE, Fadgen KE, Gerhardt GC, Engen JR. High-speed and high-resolution UPLC separation at zero degrees Celsius. *Anal Chem*. 2008; 80:6815–6820. [PubMed: 18672890]
70. Wang L, Pan H, Smith DL. Hydrogen exchange-mass spectrometry: optimization of digestion conditions. *Mol Cell Proteomics*. 2002; 1:132–138. [PubMed: 12096131]
71. Geromanos SJ, Vissers JP, Silva JC, Dorschel CA, Li GZ, Gorenstein MV, Bateman RH, Langridge JI. The detection, correlation, and comparison of peptide precursor and product ions from data independent LC-MS with data dependant LC-MS/MS. *Proteomics*. 2009; 9:1683–1695. [PubMed: 19294628]
72. Houde D, Berkowitz SA, Engen JR. The utility of hydrogen/deuterium exchange mass spectrometry in biopharmaceutical comparability studies. *J Pharm Sci*. 2011; 100:2071–2086. [PubMed: 21491437]
73. Fang J, Nevin P, Kairys V, Venclovas C, Engen JR, Beuning PJ. Conformational analysis of processivity clamps in solution demonstrates that tertiary structure does not correlate with protein dynamics. *Structure*. 2014; 22:572–581. [PubMed: 24613485]

Fig. 1.

Local deuterium uptake for DinB and pol κ in the absence of substrates. Peptide-level HDX data for DinB (*top*) and pol κ (*bottom*) are mapped together with atomic-level structural information to visualize the intrinsic local HDX of the free polymerases in solution. For each time point, the percent deuterium uptake for each peptide is displayed as a color gradient on the respective ternary crystal structures (DinB, PDB: 4IRC; pol κ , PDB: 2OH2). No correction for deuterium back-exchange was performed (see Materials and Methods). The results are the average of three replicate experiments. DNA and nucleotides are not shown. The part of the pol κ palm loop insertion that is unresolved in the crystal structure (residues 225–281) is shown as a dashed line and colored according to its relative percent deuterium uptake. Portions of this figure were previously published in Nevin, et al. [43] and are reprinted here with permission of Elsevier.

Fig. 2.

Substrate-dependent HDX in pol κ N-clasp domain. (a) Deuterium uptake for pol κ N-clasp peptides as a function of time for free pol κ (black diamonds), pol κ -DNA (gray triangles, dashed line), pol κ -DNA-dGTP_{incorrect} (blue squares), and pol κ -DNA-dCTP_{correct} (green circles). Deuterium levels are not corrected for back-exchange and the theoretical maximum number of deuterated amides in each peptide equals its number of residues after subtracting one for the N-terminus and one for each proline residue and is set as the maximum on the Y axis. Error bars represent variation of triplicate experiments. (b) Peptides highlighted on the crystal structure of pol κ (PDB: 2OH2) according to the colors indicated in (a).

Fig. 3.

DinB and pol κ are protected from exchange in the presence of undamaged DNA and the correct dNTP. (a) Minimal model for nucleotide incorporation. The free pol is denoted E, the binary pol-DNA complex ED, and the ternary pol-DNA-dNTP complex EDN. The states in the box were probed by HDX-MS. (b) Differences in deuterium uptake between the indicated DinB states. The templating base was G and the incorrect and correct dNTPs were dGTP and dCTP, respectively. For each peptide and time point, the relative deuterium level of the second state was subtracted from the first and displayed as a color gradient. The data are the average at least two replicate experiments. Differences >0.6 Da are significant at 98% confidence. (c and d) Regions of DinB with deuterium uptake differences between the indicated states of at least 1 Da at any time point displayed as a color gradient on ternary complex crystal structure (PDB: 4IRC). In these images, brown indicates regions for which no data were obtained. (e) Differences in deuterium uptake between the indicated pol κ states calculated and displayed as in (b). (f and g) Regions of pol κ with deuterium uptake

differences between the indicated states of at least 1 Da at any time point displayed as a color gradient on ternary complex crystal structure (PDB: 2OH2).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 4.

Thermodynamic stability of DinB and pol κ in different states. Melting transition midpoints (T_m) were determined from thermal denaturation transitions of DinB and pol κ in complex with the indicated substrates. The templating base was G. The data are the average of three experiments and error bars represent $\pm$ SD.

Fig. 5.

Only preferred template lesions and correct dNTPs promote “active” conformation. (a) Primer extension opposite G, N^2 fG, and O^6 meG by DinB (top) and pol κ (bottom). Reactions were quenched after 0, 5, 30, and 60 min and analyzed by 16% denaturing polyacrylamide gel electrophoresis and subsequent phosphorimaging (b) Average differential deuterium uptake between indicated states of DinB (top) and pol κ (bottom) at each indicated time point (0.17, 1, 10, 60, 240 min) for peptides (same as in Figure 3) displayed as a color gradient. Differences >0.6 Da are significant at 98% confidence. For

each peptide and time point, the relative deuterium level of the second state was subtracted from the first. The templating base and the incoming dNTP are indicated.

Fig. 6.

Proposed reaction pathway for DinB and pol κ . The free pol binds DNA which results in a relatively closed and stable binary complex. In the case of correct dNTP incorporation opposite preferred DNA bases, there is a conserved conformational change towards a relatively closed and “active” ternary complex resulting in fast nucleotide incorporation. In the case of incorrect dNTP incorporation or non-preferred DNA, a non-productive ternary complex is favored, resulting in a slower reaction.