Published in final edited form as:

Cancer Immunol Immunother. 2015 August ; 64(8): 977–987. doi:10.1007/s00262-015-1706-4.

Extended evaluation of a Phase 1/2 trial on dosing, safety, immunogenicity, and overall survival after immunizations with an advanced generation Ad5 [E1-, E2b-]-CEA(6D) vaccine in late stage colorectal cancer

Joseph P. Balint¹, Elizabeth S. Gabitzsch^{1,*}, Adrian Rice¹, Yvette Latchman¹, Younong Xu¹, Gerald L. Messerschmidt¹, Arvind Chaudhry², Michael A. Morse³, and Frank R. Jones¹

¹Etubics Corporation, Seattle, WA

²Medical Oncology Associates, Spokane, WA

³Department of Medicine, Duke University Medical Center, Durham, NC

Abstract

A phase 1/2 clinical trial evaluating dosing, safety, immunogenicity, and overall survival on metastatic colorectal cancer (mCRC) patients after immunotherapy with an advanced generation Ad5 [E1-, E2b-]-CEA(6D) vaccine was performed. We report our extended observations on long-term overall survival and further immune analyses on a subset of treated patients including assessment of cytolytic T cell responses, T-regulatory (Treg) to T-effector (Teff) cell ratios, flow cytometry on peripheral blood mononuclear cells (PBMC), and determination of HLA-A2 status. An overall survival of 20% (median survival of 11 months) was observed during long-term follow-up and no long-term adverse effects were reported. Cytolytic T cell responses increased after immunizations and cell-mediated immune (CMI) responses were induced whether or not patients were HLA-A2 positive or Ad5 immune. PBMC samples from a small subset of patients were available for follow-up immune analyses. It was observed that the levels of carcinoembryonic antigen (CEA) specific CMI activity decreased from their peak values during follow-up in 5 patients analyzed. Preliminary results revealed that activated CD4+ and CD8+ T cells were detected in a post immunization sample exhibiting high CMI activity. Treg to Teff cell ratios were assessed and samples from 3 of 5 patients exhibited a decrease in Treg to Teff cell ratio during the treatment protocol. Based upon the favorable safety and immunogenicity data obtained, we plan to perform an extensive immunologic and survival analysis on mCRC patients to be enrolled in a randomized/controlled clinical trial that investigates Ad5 [E1-, E2b-]-CEA(6D) as a single agent with booster immunizations.

*Correspondence: Elizabeth S. Gabitzsch., Etubics Corporation, 410 West Harrison Street, Suite 100, Seattle, WA, 98119, USA, Tel.: 206-838-5110, Fax: 206-838-2978, beth@etubics.com.

Conflict of Interest Statement

Joseph P. Balint is a shareholder and employee of Etubics and has stock options in the Company. Elizabeth S. Gabitzsch is a shareholder and employee of Etubics and has stock options in the Company. Adrian Rice is an employee of Etubics and has stock options in the Company. Yvette Latchman is an employee of Etubics and has stock options in the Company. Younong Xu has stock options in the Company. Gerry L. Messerschmidt is an employee of Etubics and has stock options in the Company. Frank R Jones is a shareholder and employee of Etubics and has stock options in the Company. Arvind Chaudhry and Michael Morse have no conflicts of interest related to this work.

Keywords

Ad5 [E1-, E2b-]-CEA(6D); Colorectal cancer; Immunotherapy; Phase1/2 Trial

Introduction

As the knowledge on identification and function of tumor associated antigens (TAA) increases, so has the development of immune-based therapies that target TAA to treat various cancers. Development of gene-based vaccines to treat cancer and/or prevent tumor recurrence has become an area of significant basic research and clinical evaluation. In contrast to therapeutic monoclonal antibodies that “passively immunize” to treat certain cancers, some anti-cancer vaccines are designed to “actively” stimulate cell-mediated immunity (CMI) and antibody responses following as a response to immunotherapy. Among the anti-cancer vaccines being investigated and developed, those utilizing recombinant specific viral based delivery platforms are being exploited because of their unique ability to also generate significant CMI responses, essential in achieving antitumor immune killing [1–3]. Early generation recombinant non-replicating adenovirus serotype-5 (Ad5) based vector platforms with deletions in the early 1 (E1) gene and early 3 (E3) gene regions (Ad5 [E1-]) emerged as prior leading candidates for such immunotherapy [3–6]. Moreover, Ad5-based vaccines are an important pharmaceutical consideration because they can be produced in large quantities and the viral genome is non-integrating remaining episomal thus eliminating the possibility of permanent gene insertion in the host [6].

Unfortunately, one of the major challenges facing previous generation Ad5 [E1-]-based vectors is the presence of pre-existing immunity to Ad5 that mitigates their immunizing capability. The preponderance of humans exhibit neutralizing antibody against Ad5, the most widely used subtype for human vaccines, with two-thirds of humans studied having humoral and lympho-proliferative responses against Ad5 [7,8]. This immunity inhibits immunization and especially re-immunization (boost) with Ad5-based vectors and precludes immunization of a vaccinee against a second disease antigen as well. Furthermore, previous Ad5 [E1-]-based vectors are not effective in cases where repeated homologous immunizations to maintain and/or increase anti-cancer specific T-cell (CD4+ and CD8+) and humoral immunity are required for continued killing of cancer cells. To avoid the Ad immunization barrier, we have constructed an improved and advanced generation Ad5-based vector platform. The Ad5 [E1-, E2b-] vector platform is novel having additional deletions in the early gene 2b (E2b) region by removing the DNA polymerase (pol) and the pre terminal protein (pTP) genes and is propagated in the E.C7 human cell line [9–12]. In various animal studies, we have reported that cancer and infectious disease targeting vaccines based on the Ad5 [E1-, E2b-] vector platform can be used in multiple homologous immunization regimens designed to induce and increase CMI responses despite the presence of pre-existing and/or vector induced immunity [13–22].

We have previously reported on the development and clinical use in advanced stage mCRC patients with the Ad5 [E1-, E2b-]-based cancer vaccine designed to target carcinoembryonic antigen (CEA) expressing tumors [13,14,23]. In adult humans, CEA expression is normally

low and mainly restricted to the mucosa of the colon [24,25]; however, CEA is over expressed in all adenocarcinomas of the colon and rectum [26] and CEA has been classified as a priority cancer antigen biomarker [27]. Several immunotherapy platforms targeting CEA are being developed and there is mounting clinical evidence indicating that targeting CEA can safely be overcome despite pre-existing tolerance [28]. The vaccine we developed utilizes a modification in the 9 amino acid sequence of the CAP1 region of CEA that is incorporated into our recombinant vector platform and referred to as Ad5 [E1-, E2b-]-CEA(6D) [23]. CAP1(6D) is a peptide analog of CAP1 and its sequence includes a heteroclitic (non-anchor position) mutation, resulting in an amino acid change from asparagine to aspartic acid, to enhance recognition by the T-cell receptor without any change in binding to HLA-A2. Compared with the non-mutated CAP1 epitope, CAP1(6D) has been shown to enhance sensitization of cytotoxic T lymphocytes (CTL) by 100 to 1,000 times [29,30]. In pre-clinical studies by us and others, multiple homologous immunizations with Ad5 [E1-, E2b-]-CEA(6D) induced CEA-specific CMI responses with antitumor activity despite the presence of existing Ad5 immunity in mice [13,14]. We have proceeded to perform a dose escalation/safety phase 1/2 trial and reported results on metastatic colorectal cancer (mCRC) patients treated with the Ad5 [E1-, E2b-]-CEA(6D) [23]. Escalating doses of vaccine up to 5×10^{11} viral particles (VP) of Ad5 [E1-, E2b-]-CEA(6D) were injected subcutaneously (SQ) 3 times and were shown to have minimal toxicity. Employing ELISpot assays on PBMC samples to detect interferon-gamma (IFN- γ) secreting cells, CEA specific CMI responses were induced. Importantly, these CMI responses were observed despite the presence of pre-existing Ad5 immunity in a majority of patients and did not appear to affect the induction of CEA specific CMI responses. At the time of publication, overall survival of mCRC patients was 48% at 12 months. We now present further results on long-term overall survival in treated mCRC patients and report our extended findings on additional immune analyses of PBMC samples from treated patients.

Materials and Methods

Patients and Clinical Protocol

The clinical trial was approved by all local institutional review boards where the study was conducted under an FDA approved investigational new drug application (IND14325) and registered at ClinicalTrials.gov (NCT01147965). Informed consent was obtained from all individual participants included in the study. Details of the trial were previously reported [23]. Briefly, 32 patients with mCRC cancer, median age 57.5 (range 38–77) who had failed a median of three prior anti-tumor regimens (range 2–5), had a median Karnofsky performance status of 90% (range 70–100%), and had a range of 1–4 sites of metastatic disease, were enrolled. Eligibility requirements included metastatic cancer expressing CEA and adequate hematologic, renal, and hepatic function. Clinical activity was assessed according to Response Evaluation Criteria in Solid Tumors (RECIST 1.0 criteria) [31] using computed tomography (CT) or magnetic resonance imaging (MRI) scans obtained at baseline and after completion of treatments. Toxicity was assessed according to the National Cancer Institute Common Terminology Criteria for Adverse Events (CTCAE) version 4.0 [32].

A total of 5 cohorts were evaluated for dose-limiting toxicity (DLT). The Ad5 [E1-, E2b-]-CEA(6D) doses were delivered subcutaneously (SQ) to patients as follows: cohort 1: dose of 1×10^9 VP SQ in the same thigh every 3 weeks for 3 treatments; cohort 2: dose of 1×10^{10} VP SQ every 3 weeks for 3 treatments; cohort 3: dose of 1×10^{11} VP SQ every 3 weeks for 3 treatments, cohort 4, a phase 2 expansion: dose of 1×10^{11} VP SQ every 3 weeks for 3 treatments. After completing the phase 2 expansion cohort 4, an additional cohort (cohort 5) received a dose of 5×10^{11} VP SQ every 3 weeks for 3 treatments to determine safety of the highest dose. Serum and PBMC were collected from patients just prior to the immunizations at weeks 0, 3, 6, and three weeks (week 9) following the last treatment. Blood was also collected from some patients up to 12 months following treatment. The serum was frozen at -60°C and PBMC were frozen in liquid nitrogen until immune assays were performed.

Immunotherapeutic Vaccine

Clinical grade Ad5 [E1-, E2b-]-CEA(6D) (product referred to as ETBX-011) was constructed as previously described [14,23] and manufactured using the E.C7 human cell line (9–11) under GMP at Sigma Aldrich Fine Chemicals (SAFC), Carlsbad, California. The vaccine was stored at -20°C and thawed just prior to use.

ELISpot assays for IFN- γ and granzyme B secreting cells

The ELISpot assay for IFN- γ -secreting lymphocytes was adapted from our previous studies and performed as described [14,23]. An ELISpot assay for granzyme B-secreting lymphocytes to detect cytolytic T cell activity [33,34] was also performed. Briefly, isolated PBMC (2×10^5 cells/well) from individual patient samples were incubated 36–40 hours with a CEA peptide pool (15mers with 11aa overlap covering full length CEA with the CAP1-6D modification; $0.1 \mu\text{g}/\text{well}$) to stimulate IFN- γ or granzyme B-secreting cells. Specificity of the responses was demonstrated by the lack of reactivity with the irrelevant antigens β -galactosidase and HIV-gag. Cells stimulated with concanavalin A (Con A) at a concentration of $0.25 \mu\text{g}/\text{well}$ served as positive controls. Colored spot-forming cells (SFC) were counted using an Immunospot ELISpot plate reader (Cellular Technology, Shaker Heights, OH), and responses were considered to be positive if 50 SFC were detected/ 10^6 cells after subtraction of the negative control number and SFC were twofold higher than those in the negative control wells.

Analyses of PBMC by immunofluorescence (flow) cytometry

To assess CD4+ and CD8+ T cell responses, PMBC samples from individual patients were assayed for IFN- γ and tumor necrosis factor alpha (TNF- α) expression using a BD Accuri™ C6 Flow Cytometer and intracellular cytokine staining methods previously described [35,36]. Briefly, 10^6 PBMC cells/well were incubated 6 hours with $2.0 \mu\text{g}/\text{ml}$ CEA peptide pool, $2.0 \mu\text{g}/\text{ml}$ SIV nef negative control peptide pool, or media alone. A protein transport inhibitor (GolgiStop) was added for the final 4 hours of the stimulation. After stimulation, cells were stained for CD4 and CD8, fixed, permeabilized, stained for IFN- γ and TNF- α , and then analyzed by flow cytometry.

Regulatory T (Treg) cells and effector T (Teff) cells were analyzed by staining PBMC for CD4, CD25 and Foxp3 with the Human Regulatory T Cell Staining Kit #2 (eBiosciences).

The ratio of regulatory T cells to effector T cells (Treg/Teff) was defined as the ratio of CD4+CD25+Foxp3+ cells to CD4+CD25-Foxp3- cells.

Flow cytometry was also employed to assess HLA-A2 expression on patient PBMC samples. Staining with fluorochrome-conjugated anti-HLA-A2 antibody was performed as recommended by the manufacturer (BD Biosciences).

Determination of Ad5 neutralizing antibody (NAb) titers in sera

Endpoint Ad5 NAb titers were determined as previously described [14,23]. Briefly, dilutions of heat-inactivated test sera in 100 μ L of DMEM containing 10% fetal calf serum were mixed with 4×10^7 VP of Ad5 [E1-]-null and incubated for 60 min at room temperature. The samples were added to microwells containing HEK293 cells cultured in DMEM containing 10 % heat-inactivated fetal calf serum at 2×10^3 cells/well for 24 hours at 37°C in 5% CO₂. The mixture was incubated for an additional 72 hours at 37°C in 5% CO₂. An MTS tetrazolium bioreduction assay (Promega Corp. Madison, WI) [37] was used to measure cell killing and endpoint Ad5 NAb titers.

Statistics

Statistical analyses were performed employing PRISM Software (GraphPad). The long-term survival curve was made employing Kaplan–Meier plots (PRISM Software, GraphPad).

Results

Summary of dose administration, adverse events, and initial survival

The majority of patients (28/32, 87.5%) were able to receive all three immunizations. Four patients with rapid disease progression did not receive all 3 injections (1 patient received only 1 dose and 3 patients received 2 doses). There were no dose-limiting toxicities and no serious adverse events (SAE) that resulted in treatment discontinuation at any dose level of the vaccine and there were no dropouts due to treatment. The most common adverse effect (AE) was a self-limited, injection site reaction. Other reactions occurred with less than a 10% incidence and included fever, flu-like symptoms, anorexia, chills, nausea, and headache. These symptoms were self-limiting and did not require intervention other than symptomatic control measures such as acetaminophen. Routine hematology and chemistry studies showed no significant biologic changes during the immunization period and these above results have been previously reported [23]. During our longer-term follow-up, no treatment related adverse effects were reported.

Results on extended follow-up overall survival data and post treatment therapies

Survival data and post treatment therapies that patients received are presented in Table I. Kaplan-Meier overall survival plots are presented in Figure 1. Twenty-nine month overall survival of the intent-to-treat population (32 patients) was 20% (Figure 1a) with a median survival time of 11 months from informed consent/first injection. For the subset of 28 patients that received all 3 immunizations, the 29-month survival was 23% (Figure 1b) with a median survival time of 13 months. For the 22 patients optimally dosed with the two highest doses of vaccine (1 and 5×10^{11}) and receiving all 3 immunizations (see Table I), the

28-month overall survival was 19% (Figure 1c). Median overall survival was 13 months in the optimally treated patients. Since there was no active control group in the study, comparisons for significance in survival time cannot be made. There were 3 stable disease events observed immediately after completion of treatment and over the course of long-term follow-up, we observed that a total of 6 disease stabilizations were recorded.

Post treatment therapies were also recorded during follow-up (see Table I). For 7 patients follow-up therapy information was not available, for 8 patients there were no further therapies, 4 patients received other unknown chemotherapy, 2 patients received radiation treatment, 1 patient received radiation treatment and entered another clinical trial, 2 patients entered another clinical trial, 1 patient received treatment with Avastin, 1 patient received Avastin and Xeliri, 1 patient received Avastin and Xeloda, 1 patient received Avastin, Xeloda, and FOLFOX, 1 patient received Xeloda, 1 patient received FOLFOX, 1 patient received treatment with Nexavar, and 1 patient received Erbitux.

Immune analyses

We previously reported on ELISpot analyses of PBMC samples for CMI activity as assessed by numbers of IFN- γ secreting cells after stimulation with CEA peptide pools (23). We observed a dose response effect with the highest magnitude CEA-specific CMI responses occurring in patients who received the highest dose of Ad5 [E1-, E2b-]-CEA(6D). Of the doses received, 0/3 (0%) patients in cohort 1, 1/4 (25 %) patients in cohort 2, 10/19 (53%) cohort 3/phase II, and 4/6 (67%) patients in cohort 5 exhibited positive CEA directed CMI responses. As previously reported [23], the level of pre-existing Ad5 neutralizing immunity did not appear to affect the induced CEA directed CMI responses.

For this study, we performed additional flow cytometry analyses on patient PBMC. A sample from only one patient (patient 013, week 3) that exhibited high CMI activity (1000 IFN- γ secreting SFC) during treatment was observed to be positive for polyfunctional T cells. Although the week 9 sample from patient 507 also exhibited high CMI activity, we were unable to detect polyfunctional T cells.

The flow cytometry ICS profile of the PBMC from patient 013 after exposure to CEA peptides showed the presence of:

- CD8+/IFN- γ + cells (3.5%),
- CD8+/IFN- γ + /TNF- α + cells (1.0%),
- CD4+/IFN- γ + cells (0.4%), and
- CD4+/IFN- γ + /TNF- α + cells (0.2%).

Specificity of the response was demonstrated by the lack of PBMC reactivity with SIV-nef peptides.

In further follow-up, we obtained PBMC samples from a subset of 5 patients that could be tested post-treatment for extended CMI responses. It was observed that the levels of CEA specific CMI activity decreased from their peak values during follow-up in the 5 patients analyzed (Figure 2). We were able to perform further analyses on a large number of PBMC

samples pre (week 0) and post (week 6–9) treatment and assess CTL activity employing ELISpot assays for granzyme B secretion. Twenty-seven evaluable patient PBMC samples were tested. As shown in Figure 3, significantly ($P < 0.05$, *Wilcoxon test*) elevated levels of granzyme B secretion activity were observed in post immunization PBMC samples as compared to their respective baseline pre-immunization PBMC samples.

Analysis of peripheral blood T regulatory (Treg) to T effector (Teff) cell ratios pre and post-treatment (week 9) in 15 patients was examined employing flow cytometry. No change in Treg to Teff cell ratio occurred over the course of the 9-week treatment evaluation period. The mean Treg to Teff cell ratio at baseline was 0.062 ± 0.02 (SD) and the mean Treg to Teff cell ratio at week 9 was 0.07 ± 0.03 . Of the 5 patient PBMC samples available for longer-term assessment, one patient exhibited an increase in Treg to Teff cell ratio, one patient exhibited an increase in Treg to Teff cell ratio that decreased to near baseline level at last follow-up, and samples from 3 patients exhibited a decrease in Treg to Teff cell ratio during the treatment protocol with one of these patients showing a return to near baseline ratio after 52 weeks (Figure 4).

Patient PBMC samples were tested for HLA-A2 positivity by flow cytometry and 61% of the samples tested were HLA-A2 positive. When the highest CMI response level (as measured by IFN- γ secreting cells) achieved per patient during treatment was assessed in association with the presence of HLA-A2, there was no significant difference observed between HLA-A2+ and HLA-A2- patients (HLA-A2+ = 264.6 ± 119.0 IFN- γ secreting cells versus HLA-A2- = 165.6 ± 108.1 IFN- γ secreting cells; $p = 0.5809$).

Ad5 NAb titers were assessed in follow-up serum samples from 6 patients. Patients 005, 024, 501, and 507 had pre-existing Ad5 endpoint titers of 200, 200, 100, and 50, respectively. Patients 031 and 500 did not exhibit Ad5 NAb activity at baseline. As shown in Figure 5, Ad5 NAb titers increased at week 9 and then decreased thereafter. In patient 507, Ad5 NAb titers decreased to below baseline titer but were within 1 titer of the baseline value. In the remaining 5 patients (005, 024, 031, 500, and 501), the Ad5 NAb titers decreased but remained above baseline levels at last follow-up.

Discussion

We previously reported our initial study findings on the dosing, safety, immunogenicity, and clinical findings of a phase 1/2 clinical trial evaluating the use of an Ad5 [E1-, E2b-]-CEA(6D) in mCRC patients [23]. Patient demographics were similar to those reported in other clinical trials with mCRC patients [38–40]. Study findings indicated that the Ad5 [E1-, E2b-]-CEA(6D) immunotherapeutic product as a single agent was well tolerated and could easily be administered. It was reported that CEA specific CMI responses were induced and increased over the course of three immunizations. In addition, a dose response effect was observed. Pre-existing Ad5 immunity was present in 61% of patients and the presence of Ad5 immunity did not appear to affect the CEA immunogenicity. Although no objective anti-tumor responses occurred by RECIST, there were 3 stable disease events observed immediately after completion of treatment. Over the course of long-term follow-up, we noted that a total of 6 disease stabilizations were recorded. Previously we reported a 48%

survival at 12 months follow-up [23] and in our extended follow-up for overall survival, we noted that on an intent-to-treat analysis the 29-month survival was 20% with a favorable median overall survival of 11 months. However, one should note that these overall survival data are confounded by the fact that the patients received other anti-cancer treatment(s) after they went off protocol. Also, it is of interest to note that the FDA approved cancer vaccine Sipuleucel-T did not show a significant anti-tumor impact but did demonstrate overall survival benefit [41–43]. Whether or not we observe similar results can only be demonstrated by testing the Ad5 [E1-, E2b-]-CEA(6D) immunotherapeutic as a single agent to determine its true clinical benefit in a larger randomized/controlled phase 2b study wherein patients are followed long-term and which we plan to perform.

In our further analyses of patient PBMC samples, ELISpot assays were employed to determine CEA directed CTL activity pre and post immunization. ELISpot granzyme B assays were used since this assay provides a measure of functional CTL activity [33,34]. We observed a significant increase in CTL activity after immunizations with Ad5 [E1-, E2b-]-CEA(6D) indicating that CEA specific CTL activity was induced. Also, we observed granzyme B secreting activity in baseline PBMC samples and we speculate that this reflects endogenous CTL activity induced by the presence of CEA expressing tumors. Thus, in addition to the induction of CEA directed CMI responses as previously reported [23], we also observed increases in CEA directed CTL activity in treated patients.

Our flow cytometry data indicated that a high level of CMI activity, as measured by ELISpot IFN- γ secreting cells, must be achieved before we could detect the presence of activated T cells by ICS. In two patients, PBMC samples were sufficiently evaluable and we were able to detect activated CD4+ and CD8+ T cells in a post immunization sample from one patient. Although the sample size is very small, this data may provide evidence that the Ad5 [E1-, E2b-]-CEA(6D) can induce robust CMI responses and we plan to study this on a larger patient sample size in a phase 2b trial. Moreover, if one were to treat mCRC patients earlier in the course of their disease, more robust CD4+ and CD8+ populations might be achieved.

Because of the importance of Tregs in influencing anti-tumor CMI responses [44], Treg to Teff ratios were determined in patient PBMC samples. Of the 5 patient samples tested, three patients exhibited a decrease in the Treg to Teff cell ratio during the immunization period. Since we also observed that levels of CMI activity decreased during follow-up in the 5 patients analyzed, it may well be that we need to administer additional booster immunizations in order to maintain elevated levels of CEA directed CMI responses that generate activated T cells with beneficial changes in Treg to Teff cell ratios. Based upon the observation that CMI responses waned by 26 weeks (23 weeks after the last immunization), we plan to administer “booster” immunizations at 4-month intervals in our next phase 2b study to determine if we can maintain induced CMI responses. Immune tests including ELISpot assays for IFN- γ and granzyme B secreting cells and flow cytometry tests for activated CD4+, CD8+, Treg, and Teff cells will be studied as surrogate parameters of immune function in planned future trials. We expect that a larger randomized mCRC patient trial will give us much needed additional information and provide a greater understanding of the immune mechanisms(s) generated by immunizations with Ad5 [E1-, E2b-]-CEA(6D).

Since the modified immunogenic HLA-A2 restricted CAP1(6D) epitope of CEA was incorporated into our vaccine, we investigated the association between induced CMI responses and the presence or absence of HLA-A2 in PBMC samples tested. Although the average CMI responses were relatively higher in HLA-A2+ PBMC samples, the difference was not significant when compared to values obtained in HLA-A2- samples. Thus, in this small study CMI responses did not appear to be HLA-A2+ restricted. Again a larger trial may elucidate this difference.

In our analyses of Ad5 NAb titers on follow-up serum samples from 6 patients, we observed that the Ad5 NAb titers decreased from their peak values after immunizations ended. This is similar to our findings with CMI responses in which the responses declined after treatment was stopped. We previously reported on the analyses of Ad5 NAb titers in non-human primates (NHP) studies investigating Ad5 [E1-, E2b-]-HIV/SIV vaccines [18,21]. In these prior studies employing NHP with pre-existing Ad5 immunity, we observed that multiple homologous immunizations in with Ad5 [E1-, E2b-]-HIV/SIV vaccines resulted in an increase in Ad5 NAb titers during vaccinations; however, after immunizations ended, Ad5 NAb titers decreased from their peak values. Thus, the present observations are in agreement with our earlier findings using this platform-based vaccine on NHP.

In conclusion, the clinical data obtained using the Ad5 [E1-, E2b-]-CEA(6D) demonstrate that this product can be easily and safely administered to mCRC patients to induce CEA directed CMI responses. Although no objective antitumor responses were observed, there appeared to be favorable survival with an observed median overall survival of 11 months. A small number of patient PBMC samples were available for follow-up immune analyses and although the sample size was small, the results obtained indicate that the generated CEA directed CMI responses induced might require additional booster immunizations to maintain the induced immune response. This will be an important factor to consider especially in the context of immunotherapy vaccines. We plan to conduct a randomized/controlled phase 2b clinical trial to evaluate immunogenicity and include booster immunizations to determine the effect of Ad5 [E1-, E2b-]-CEA (6D) as a single agent on overall survival in mCRC patients. Also, it maybe important to investigate the use of this immunotherapeutic agent in an adjuvant setting enrolling recently resected early stage colorectal cancer patients to assess the time to progression. The results of such clinical trials will allow us to evaluate the true clinical benefit of using Ad5 [E1-, E2b-]-CEA(6D) as an immunotherapeutic.

Acknowledgements

Younong Xu was an employee of Etubics during performance of the study and is now an employee of Johnson & Johnson. The authors wish to thank Susan Thorburn for her assistance in the patient data collection and Ms. Carol Jones for her excellent assistance with grants management.

This study was funded by National Cancer Institute Small Business Innovative Research Grant 1R43CA134063, National Cancer Institute Small Business Innovative Research Grant 2R44CA134063, National Cancer Institute Small Business Innovative Research Contract HHSN261200900059C, and National Cancer Institute Small Business Innovative Research Contract HHSN261201100097C.

Abbreviations

Ad5	Adenovirus serotype-5.
Ad5 [E1-]	Adenovirus serotype-5 (Ad5) based vector platforms with deletions in the early 1 (E1) gene and early 3 (E3) gene regions.
Ad5 [E1-, E2b-]	Ad5 [E1-] with additional deletions in the early 2 (E2) gene region.
Ad5 [E1-, E2b-]-CEA(6D)	Ad5 [E1-, E2b-] containing the CAP1(6D) gene.
AE	Adverse effects.
CAP1	Nine amino acid sequence of CEA.
CAP1(6D)	Peptide analog sequence of CAP1.
CEA	Carcinoembryonic antigen.
CMI	Cell mediated immune.
Con A	Concanavalin A.
CT	Computed tomography.
CTCAE	Common Terminology Criteria for Adverse Events.
CTL	Cytotoxic T lymphocytes.
E1	Early 1 gene region.
E2b	Early 2b gene region.
E3	Early 3 gene region.
IFN-γ	Interferon gamma.
mCRC	Metastatic colorectal cancer.
MRI	Magnetic resonance imaging.
NAb	Neutralizing antibody.
NHP	Non-human primates.
PBMC	Peripheral blood mononuclear cells.
pol	Polymerase gene.
pTP	Pre-terminal protein gene.
RECIST	Response Evaluation Criteria in Solid Tumors.
SAE	Serious adverse effects.
SAFC	Sigma Aldrich Fine Chemicals.
SFC	Spot forming cells.

SQ	Subcutaneously.
TAA	Tumor associated antigens.
Teff	T effector lymphocytes.
Treg	T regulatory lymphocytes.
TNF-α	Tumor necrosis factor alpha.
VP	Viral particles.

References Cited

1. Vergati M, Intrivici C, Huen N-Y, Schlom J, Tsang KY. Strategies for Cancer Vaccine Development. *J Biomed and Biotechnol*. 2010
2. Palena C, Schlom J. Vaccines against Human Carcinomas: Strategies to Improve Antitumor Immune Responses. *J Biomed and Biotechnol*. 2010
3. Schlom J. Therapeutic Cancer Vaccines: Current Status and Moving Forward. *J Natl Cancer Inst*. 2012; 104:599–513. [PubMed: 22395641]
4. Tatsis N, Ertl HCJ. Adenoviruses as vaccine vectors. *Mol Ther*. 2004; 10:616–629. [PubMed: 15451446]
5. Bangari DS, Mittal SK. Development of nonhuman adenoviruses as vaccine vectors. *Vaccine*. 2006; 24:849–862. [PubMed: 16297508]
6. Campos SK, Barry MA. Current advances and future challenges in adenoviral vector biology and targeting. *Curr Gene Ther*. 2007; 7:189–204. [PubMed: 17584037]
7. Chirmule N, Probert K, Magosin S, Qian Y, Qian R, Wilson J. Immune responses to adenovirus and adeno-associated virus in humans. *Gene Ther*. 1999; 6:1574–1583. [PubMed: 10490767]
8. Barouch DH, Kikic SV, Weverling GJ, et al. International seroepidemiology of adenovirus serotypes 5, 26, 35, and 48 in pediatric and adult populations. *Vaccine*. 2011; 29:5203–5209. [PubMed: 21619905]
9. Amalfitano A, Chamberlain JS. Isolation And Characterization Of Packaging Cell Lines That Coexpress The Adenovirus E1, DNA Polymerase, And Preterminal Proteins: Implications For Gene Therapy. *Gene Ther*. 1997; 4:258–263. [PubMed: 9135740]
10. Amalfitano A, Hauser MA, Hu H, Serra D, Begy CR, Chamberlain JS. Production And Characterization Of Improved Adenovirus Vectors With The E1, E2b, And E3 Genes Deleted. *J Virol*. 1998; 72:926–933. [PubMed: 9444984]
11. Hartigan-O'Connor D, Amalfitano A, Chamberlain JS. Improved Production of Gutted Adenovirus in Cells Expressing Adenovirus Preterminal Protein and DNA Polymerase. *J Virol*. 1999; 73:7835–7841. [PubMed: 10438876]
12. Seregin SS, Amalfitano A. Overcoming pre-existing Adenovirus immunity by genetic engineering of Adenovirus based vectors. *Expert Opin Biol Ther*. 2009; 9:1–11. [PubMed: 19063688]
13. Osada T, Yang XY, Hartman ZC, et al. Optimization of vaccine responses with an E1, E2b and E3-deleted Ad5 vector circumvents pre-existing anti-vector immunity. *Cancer Gene Ther*. 2009; 16:673–682. [PubMed: 19229288]
14. Gabitzsch ES, Xu Y, Balint JP Jr, Hartman ZC, Lyerly HK, Jones FR. Anti-tumor immunity despite immunity to adenovirus using a novel adenoviral vector Ad5 [E1-, E2b-]-CEA. *Cancer Immunol Immunother*. 2010; 59:1131–1135. [PubMed: 20361185]
15. Gabitzsch ES, Xu Y, Yoshida LH, Balint J, Gayle RB, Amalfitano A, Jones FR. A preliminary and comparative evaluation of a novel Ad5 [E1-, E2b-] recombinant based vaccine used to induce cell mediated immune responses. *Immunol Lett*. 2009; 122:44–51. [PubMed: 19073216]
16. Gabitzsch ES, Yu X, Yoshida LH, Balint J, Amalfitano A, Jones FR. Novel adenovirus type 5 vaccine platform induces cellular immunity against HIV-Gag, Pol, Nef despite the presence of Ad5 immunity. *Vaccine*. 2009; 27:6394–6398. [PubMed: 19559110]

17. Gabitzsch ES, Xu Y, Balcitis S, Balint JP Jr, Jones FR. An Ad5 [E1-, E2b-]-HER2/neu vector induces immune responses and inhibits HER2/neu expressing tumor progression in Ad5 immune mice. *Cancer Gene Ther.* 2010; 18:326–335. [PubMed: 21233857]
18. Gabitzsch ES, Xu Y, Balint JP Jr, Balcitis S, Sanders-Beer B, Jones FR. Induction and Comparison of SIV immunity in Ad5 Naïve and Ad5 Immune Non-human Primates using an Ad5 [E1-, E2b-] based vaccine. *Vaccine.* 2011; 29:8101–8107. [PubMed: 21864618]
19. Jones FR, Gabitzsch ES, Xu Y, et al. Prevention of influenza virus shedding and protection from lethal H1N1 challenge using a consensus H1N1 HA and NA adenovirus vector vaccine. *Vaccine.* 2009; 29:7020–7026. 2011. [PubMed: 21821082]
20. Gabitzsch ES, Jones FR. New Recombinant Ad5 Vector Overcomes Ad5 Immunity Allowing for Multiple Safe, Homologous Immunizations. *J Clin Cell Immunol.* 2011; S4:001
21. Gabitzsch ES, Balint JP Jr, Xu Y, et al. Control of SIV infection and subsequent induction of pandemic H1N1 immunity in rhesus macaques using an Ad5 [E1-, E2b-] vector platform. *Vaccine.* 2012; 30:7265–7270. [PubMed: 23041546]
22. Wieking BG, Vermeer DW, Spanos WC, et al. A non-oncogenic HPV 16 E6/E7 vaccine enhances treatment of HPV expressing tumors. *Cancer Gene Ther.* 2012; 19:667–674. [PubMed: 22918471]
23. Morse MM, Chaudhry A, Gabitzsch ES, et al. Novel Adenoviral Vector Induces T Cell Responses Despite Anti-Adenoviral Neutralizing Antibodies in Colorectal Cancer Patients. *Cancer Immunol Immunother.* 2013; 62:1293–1201. [PubMed: 23624851]
24. Thompson JA, Grunert F, Zimmermann W. Carcinoembryonic antigen gene functions and expression in normal and malignant tissues. *Semin Cancer Biol.* 1991; 9:67–81.
25. Berinstein NL. Carcinoembryonic antigen as a target for therapeutic anticancer vaccines: a review. *J Clin Oncol.* 2002; 20:2197–2107. [PubMed: 11956282]
26. Cheever MA, Allison JP, Ferris AS, et al. The Prioritization of Cancer Antigens: A National Cancer Institute Pilot Project for the Acceleration of Translational Research. *Clin Cancer Res.* 2009; 15:5323–5337. [PubMed: 19723653]
27. Gameiro SR, Jammeh ML, Hodge JW. Cancer vaccines targeting family: molecular biology and clinical perspectives. *J Clin Lab Anal.* 2013; 5:344–366.
28. Hammarstrom S. The carcinoembryonic antigen (CEA) family: structures, suggested carcinoembryonic antigen: state-of-the-art and future promise. *Expert Rev Vaccines.* 1999; 12:617–629.
29. Zaremba S, Barzaga E, Zhu M, Soares N, Tsang KY, Schlom J. Identification of an enhancer agonist cytotoxic T lymphocyte peptide from human carcinoembryonic antigen. *Cancer Res.* 1997; 57:4570–4577. [PubMed: 9377571]
30. Tangri S, Ishioka GY, Huang X, et al. Structural features of peptide analogs of human histocompatibility leukocyte antigen class I epitopes that are more potent and immunogenic than wild type peptide. *J Exp Med.* 2001; 194:833–846. [PubMed: 11560998]
31. Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumors: revised RECIST guideline (version 1.1). *Eur J Cancer.* 2009; 45:228–247. [PubMed: 19097774]
32. CTEP Cancer Therapy Evaluation Program. [Accessed 10 Feb, 2012] CTCAE and CTC Website. 2010. http://ctep.cancer.gov/protocolDevelopment/electronic_applications/ctc.htm
33. Rininsland FH, Helms T, Asaad RJ, Boehm BO, Tary-Lehmann M. Granzyme B ELISPOT assay for ex vivo measurements of T cell immunity. *J Immunol Methods.* 2000; 240:143–155. [PubMed: 10854609]
34. Shafer-Weaver K, Sayers T, Strobl S, et al. The Granzyme B ELISPOT assay: an alternative to the ⁵¹Cr-release assay for monitoring cell-mediated cytotoxicity. *J Transl Med.* 2003; 1:14–23. [PubMed: 14697097]
35. Hobeika AC, Clay TM, Mosca PJ, Lyerly HK, Morse MA. Quantitating therapeutically relevant T-cell responses to cancer vaccines. *Crit Rev Immunol.* 2001; 21:287–297. [PubMed: 11642610]
36. Hobeika AC, Morse MA, Osada T, et al. Enumerating antigen-specific T-cell responses in peripheral blood: a comparison of peptide MHC Tetramer, ELISpot, and intracellular cytokine analysis. *J Immunother.* 2005; 28:63–72. [PubMed: 15614046]
37. Cory AH, Owen TC, Barltrop JA, Cory JG. Use of an aqueous soluble tetrazolium/formazan assay for cell growth assays in culture. *Cancer Comm.* 1991; 3:207–212.

38. Jonker DJ, O'Callaghan CJ, Karapetis CS, et al. Cetuximab for the treatment of colorectal cancer. *N Engl J Med*. 2007; 357:2040–2048. [PubMed: 18003960]
39. Karapetis CS, Khambata-Ford S, Jonker DJ, et al. K-ras mutations and benefit from Cetuximab in advanced colorectal cancer. *N Engl J Med*. 2008; 359:1757–1765. [PubMed: 18946061]
40. Van Cutsem E, Peeters M, Siena S, et al. Open-label Phase III trial of panitumumab plus best supportive care compared with best supportive care alone in patients with chemotherapy-refractory metastatic colorectal cancer. *J Clin Oncol*. 2007; 25:1658–1664. [PubMed: 17470858]
41. Small EJ, Schellhammer PF, Higano CS, et al. Placebo-controlled phase III trial of immunologic therapy with sipuleucel-T (APC8015) in patients with metastatic, asymptomatic hormone refractory. *J Clin Oncol*. 2006; 24:3089–3094. [PubMed: 16809734]
42. Higano CS, Schellhammer PF, Small EJ, et al. Integrated data from 2 randomized, double-blind, placebo-controlled, phase 3 trials of active cellular immunotherapy with sipuleucel-T in advanced prostate cancer. *Cancer*. 2009; 115:3670–3679. [PubMed: 19536890]
43. Kantoff PW, Higano CS, Shore ND, et al. Sipuleucel-T immunotherapy for castration-resistant prostate cancer. *N Engl J Med*. 2010; 363:411–422. [PubMed: 20818862]
44. Beyer M, Schultze JL. Regulatory T cells in cancer. *Blood*. 2006; 108:804–811. [PubMed: 16861339]

Fig. 1.

Kaplan-Meier survival plots on long-term overall survival of treated mCRC patients. Panel **a** represents all treated patients. Panel **b** represents patients that received all 3 treatments. Panel **c** represents patients immunized 3 times with the 2 highest doses of vaccine. There were 23 events during the study.

Fig. 2.

Profiles of CEA directed CMI responses in 5 mCRC patients during and after the course of immunotherapy. Note the decrease in CMI responses from their peak values after immunizations ended. Values are Mean $\pm$ SEM.

Fig. 3. CTL responses (ELISpot granzyme B secretion) were assessed Pre (week 0) and Post (week 6–9) treatment. Note the increase in CTL activity after immunizations of mCRC patients ($P < 0.05$ Wilcoxon test). Values are Mean $\pm$ SEM.

Fig. 4.

Treg/Teff ratios in 5 mCRC patients during and after immunotherapy. Ratios increased in 1 patient (031) and increased with a return to baseline ratio in another patient (501). Ratios decreased with a return to baseline ratio at 52 weeks in 1 patient (005). Ratios were decreased in 2 of the other 5 patients (024 and 507).

Fig. 5.
Ad5 NAb titers over time in 6 mCRC patients immunized with Ad5 [E1-, E2b-]-CEA(6D).
Titers increased in all patients by week 9 (3 weeks after the last immunization) and then decreased thereafter.

Table 1

Overall Survival and Post-Treatment Therapies

Patient ID #	Cohort	Number of Immunizations	Tumor Type	Survival Time (Months) ^a	Post-Treatment Therapies ^b
002	1	3	colon	2.75 (--)	None
003	1	3	rectal	8.75 (--)	R
004	1	3	colon	11.25 (--)	A, F, Xld
005	2	3	colo-rectal	17.8 (--)	OCT, R
007	2	1	colon	1.25 (--)	None
008	2	3	colo-rectal	29 (+)	None
010	2	3	colon	12 (--)	OCT
011	3	3	rectal	13.2 (--)	R
012	3	3	colon	28 (+)	OUC
013	3	3	rectal	3.5 (--)	N
015	3	3	colo-rectal	10 (--)	E
016	3	3	colon	5.75 (--)	OUC
017	3	3	rectal	2.75 (--)	None
500	3	3	colon	26.6 (--)	Not Reported
501	4	3	rectal	25.5 (+)	OUC
502	4	2	colon	2.75 (--)	Not Reported
504	4	3	colon	7.25 (--)	Not Reported
506	4	3	colon	2.5 (--)	Not Reported
507	4	3	rectal	24 (+)	Not Reported
019	4	3	colon	17.5 (--)	A, Xli
020	4	3	colon	24.5 (+)	A
021	4	3	colon	25.25 (+)	A, Xld
023	4	3	colon	4.25 (--)	None
024	4	3	colon	23 (+)	None
025	4	3	colon	7.25 (--)	None
026	4	2	colon	4.2 (--)	None

Patient ID #	Cohort	Number of Immunizations	Tumor Type	Survival Time (Months) ^a	Post-Treatment Therapies ^b
030	5	3	colo-rectal	15.7 (--)	OCT
031	5	3	rectal	15.5 (+)	OUC
032	5	3	rectal	5.6 (--)	F
033	5	3	rectal	4.9 (--)	Not Reported
034	5	3	colon	13.75 (+)	Xld
035	5	2	colon	2.0 (--)	Not Reported

^a (+) Alive at last follow-up; (--) Not alive.

^b Post-Treatment Therapy Abbreviations: A (Avastin), E (Eribitux), N (Nexavar), F (FOLFOX), N (Nexavar), OCT (other clinical trial), OUC (other unknown chemotherapy), R (radiation), Xli (Xeliri), Xld (Xeloda).