Published in final edited form as:

J Neuroendocrinol. 2011 March ; 23(3): 244–253. doi:10.1111/j.1365-2826.2010.02099.x.

Neuronal nitric oxide synthase as a substrate for the evolution of pseudosexual behaviour in parthenogenetic whiptail lizard

Lauren A. O'Connell^{1,2}, Bryan J. Matthews², and David Crews^{1,2,*}

¹ Institute for Cell and Molecular Biology, University of Texas at Austin, Austin, TX 78705, USA

² Section of Integrative Biology, University of Texas at Austin, Austin, TX 78705, USA

Abstract

The evolution of neuroendocrine mechanisms governing sex-typical behaviour is poorly understood. An outstanding animal model is the whiptail lizard (*Cnemidophorus*) as both the ancestral and descendent species still exist. The ancestral little striped whiptail, *C. inornatus*, consists of males and females, which exhibit sex-specific mating behaviours. The descendent desert grassland whiptail, *C. uniparens*, consists only of females that alternately exhibit both female-like and male-like pseudosexual behaviour. Castrated male *C. inornatus* will mount a conspecific in response to exogenous androgen, but some are also sensitive to progesterone. This polymorphism in progesterone sensitivity in the ancestral species may have been involved in evolution of progesterone-mediated male-typical behaviour in the descendant unisexual lizards. We tested if progesterone activates the typically androgenic signalling pathway by investigating hormonal regulation of neuronal nitric oxide synthase (nNOS) using *in situ* hybridization and NADPH diaphorase histochemistry, a stain for nNOS protein. NADPH diaphorase is widely distributed throughout the brain of both species, but only in the periventricular nucleus of the preoptic area (pvPOA) are there differences between mounting and non-mounting individuals. The number of cells expressing nNOS mRNA and NADPH diaphorase is higher in the pvPOA of individuals that mount in response to progesterone or androgen. Further, the nNOS promoter has both androgen and progesterone response elements, and NADPH diaphorase colocalizes with the progesterone receptor in the pvPOA. These data suggest that a polymorphism in progesterone sensitivity in the sexual ancestor reflects a differential regulation of nNOS and may account for the male-typical behaviour in unisexual whiptail lizards.

Keywords

progesterone receptor; neuronal nitric oxide synthase; preoptic area; sexual behaviour; parthenogenesis

* All correspondence and requests for reprints should be addressed to: David Crews Section of Integrative Biology University of Texas at Austin, Austin, TX 78712 Phone: 512-471-1113 crews@mail.utexas.edu.

Disclosure Statement: The authors have nothing to disclose.

INTRODUCTION

The evolution of neural mechanisms which govern social decision-making are poorly understood. However, insight into the proximate mechanisms underlying behavioural outputs to social stimuli is of primary importance for understanding the evolution and selection of behavioural traits (1). Whiptail lizards (genus *Cnemidophorus*) are an excellent model system for the study of the evolution of neuroendocrine mechanisms mediating sexually dimorphic behaviour (2), as both ancestral and descendant species still exist. Here, we explore the neuroendocrine mechanisms governing reproductive behaviours of whiptail lizards in an effort to delineate the molecular substrates for the evolution of pseudocopulatory behaviour in a parthenogenetic species.

The all-female whiptail lizard, *Cnemidophorus uniparens*, consists of clonal descendants (3), and displays both male-like mounting behaviour and female-like receptivity during discrete phases of the ovarian cycle (4, 5). This male-typical behaviour is indistinguishable from the mounting behaviour seen in males of their direct ancestor, *C. inornatus* (4). This behavioural adaptation serves to stimulate and synchronize reproductive activity in conspecific females (6). The display of male-typical behaviour in these females is associated with the postovulatory phase of the ovarian cycle when progesterone levels are high (5,7). During the preovulatory phase when estrogen levels are high, *C. uniparens* are receptive to mounting from conspecifics, similar to the ancestral *C. inornatus* females. Androgens are not detectable in the circulation at any time of the ovarian cycle in females of either species (8). Unlike *C. uniparens*, the ancestral *C. inornatus* females do not display mounting behaviour during the postovulatory phase when progesterone levels are high.

The regulation of male-typical mounting behaviour by progesterone in *C. uniparens* has an evolutionary precursor in *C. inornatus* males, where progesterone, acting either alone or synergistically with androgens, stimulates mounting behaviour (9). This effect of progesterone is mimicked by R5020, a non-metabolizable progesterone receptor (PR) agonist, and abolished by the anti-progestin RU486, suggesting that progestins mediate this behaviour at the level of PR rather than via progestin metabolites such as androgens (10). In both *C. inornatus* and *C. uniparens*, PR is expressed in two regions of the preoptic area: the medial preoptic area (mPOA) and the periventricular nucleus (pvPOA) (11). Implantation of progesterone directly into the POA elicits the full repertoire of mounting and intromission in *C. inornatus* males that had previously been established as sensitive to progesterone (12). However, the molecular actions of the ligand-activated PR that promote male-typical behaviour are not known.

To understand how progesterone may facilitate male-typical mounting behaviour in both the ancestral and descendent whiptail species, we first searched for a candidate molecule that facilitates male-typical behaviours in other vertebrates and is regulated by both progesterone and androgens in different physiological contexts. One model suggests that androgenic upregulation of neuronal nitric oxide synthase (nNOS) may be a key cellular event that allows the expression of male-typical mounting behaviour in rodents (13). It has been established that expression of male-typical copulatory behaviour in male *C. inornatus* is associated with an androgen-driven increase in nNOS mRNA and protein levels in the POA,

and that the mounting behaviour is suppressed when nNOS is inhibited in *C. uniparens* (14,15). In addition to being regulated by androgens, nNOS is also regulated by progesterone in the context of rat female-typical sexual behaviour (16).

In this study, we investigate the role of nNOS in facilitating male-typical copulatory behaviour, and its regulation by progesterone in the ancestral and descendant whiptail lizard species. To better understand where nNOS may be acting to facilitate male-like copulatory behaviour, we first mapped its distribution in the whiptail brain using NADPH diaphorase histochemistry, a marker for nNOS protein activity (17). We then analyzed nNOS mRNA and protein levels in hormonally manipulated *C. inornatus* males and *C. uniparens* by *in situ* hybridization and quantified NADPH diaphorase-positive cells in the pvPOA. We then determined if progesterone could potentially regulate nNOS expression by analyzing the nNOS promoter for progesterone receptor response elements and colocalized nNOS and PR in the pvPOA. Finally, to investigate species differences in mating behaviour, we compared the number of NADPH diaphorase-positive cells in the pvPOA of *C. inornatus* females and *C. uniparens* across the ovarian cycle. Our findings indicate that nNOS is upregulated in individuals that mount in response to progesterone on a level comparable to testosterone-induced levels and that PR may directly regulate nNOS transcription. These data suggest that progesterone-mediated mechanisms of facilitating male-typical copulatory behaviour are similar to testosterone induction of mounting behaviour and further suggest that nNOS may have been recruited in the evolution of pseudocopulatory behaviour in *C. uniparens*.

MATERIALS AND METHODS

Animals

Adult *C. inornatus* were captured in the vicinity of Sanderson, TX in May 2008 and transported to the University of Texas at Austin campus. Adult *C. uniparens* were captured in the vicinity of Portal, Arizona in June 2008, housed for no more than three weeks at the Southwestern Research Station, and then transported to the University of Texas at Austin. Lizards are housed in environmentally controlled chambers in terraria with *ad libitum* water and food in the form of crickets (18,19).

Behavioural Tests and Surgeries

C. inornatus—After one to two weeks to acclimatize to the laboratory environment, sexually active males were identified as previously described (14,19). These sexually active males were castrated and returned to their original tanks as previously described (20). Seven weeks after castration, they were given three ten-minute tests with receptive females to verify the absence of courtship behaviour, and then implanted subcutaneously capsules 12mm Silastic capsules (1.47mm diameter, 1.96 mm outer diameter) packed with crystalline testosterone, progesterone or with empty implants (controls) as previously described (18,19) to result in physiologically relevant levels of testosterone and progesterone in whiptail lizards and to induce male-like copulatory behaviour. All surgical procedures were carried out under hypothermic anesthesia. Eighteen days later, males were tested once a day for five consecutive days with receptive females in the home cage of the focal male for reinstatement of copulatory behaviour. After testing they were immediately anaesthetized by

hypothermia, and sacrificed by decapitation. Males were sacrificed within five min of the final behavioural observation to avoid the effect of recent behaviour on nNOS induction. This has been shown to be an appropriate time-window as *egr-1* expression, an immediate early gene marker for neuronal activity, has a 10 to 30 min delay from the last behavioural experience (21). Males whose sexual behaviour was reinstated with systemic progesterone were classified as progesterone-sensitive; those in which sexual behaviour was not reinstated were classified as progesterone-insensitive. Reinstatement of sexual behaviour was considered mounting and taking a stereotyped doughnut-posture in three out of five tests. Males classified as progesterone-insensitive did not mount in any of the five tests. Roughly 40% of tested males were progesterone-sensitive. In all cases, castrated males implanted with blank capsules did not display mounting behaviour to a receptive female. One male implanted with testosterone did not mount in response to a receptive female and was excluded from further analysis. All other castrated males implanted with testosterone mounted a receptive female. At the time of sacrifice, all males were inspected to confirm complete castration.

Female *C. inornatus* and *C. uniparens*: Preovulatory (PreOv) and postovulatory (PostOv) ovarian states were determined by abdominal palpation. Female-like receptivity was tested as previously described (22). Ovarian morphology was noted after sacrificing the experimental animal. All PreOv animals were characterised by the presence of developing follicles and receptive behaviour. PostOv animals had corpora lutea and were not receptive.

In hormonally manipulated groups, adult *C. uniparens* were ovariectomized and implanted with an empty 12mm Silastic capsule (OVX+Blank) or with a capsule packed with testosterone (OVX+T). Animals were behaviourally tested eighteen days after surgery. In all cases, OVX+Blank animals failed to show either male-like mounting behaviour or female-like receptive behaviour. Some OVX+T animals failed to display mounting behaviour to a receptive individual ($n = 3$) and were excluded from further analysis.

We adopted the neuroanatomical nomenclature of Young and Crews (11). All procedures were approved by the University of Texas IACUC.

nNOS Promoter Cloning and Analysis

Genomic DNA from whiptail livers was isolated by phenol/chloroform extraction. The whiptail nNOS promoters were cloned using GenomeWalker (Clontech, Palo Alto, CA, USA) following the manufacturer's instructions. The outer and nested gene specific primer used for nNOS promoter cloning were 5'-TGGAAGATTCAAGCCTTTAGCCTCAGTAAG and 5'-AACCATTGGACACACACTTTGGGCGG, respectively. The promoter sequence was submitted to Genbank (FJ517552). Putative steroid hormone receptor binding sites were identified using MatInspector (Genomatix).

***In situ* Hybridization**

Brains were removed, embedded in OCT and stored at -80°C until sectioning. Brains were then sectioned on a cryostat at $20\text{ }\mu\text{m}$ and thaw-mounted onto Super-Frost Plus slides (Erie Scientific Co., Portsmouth, NH) in three series. Sections were fixed as previously described

(14) and stored at -80°C . Riboprobes were reverse-transcribed in the presence of Fluorescein-labelled UTP (Roche, Indianapolis, IN) using a T7/SP6 Maxiscript in vitro transcription kit (Ambion, Austin, TX) to produce antisense or sense Fluorescein-labelled riboprobes. The template used to make the nNOS probe was the same as in Sanderson et al. (14), and was 641bp in length (Genbank accession DQ141603). Slides were then warmed to room temperature, air dried, and pre-equilibrated in hybridization buffer (50% formamide, 5x SSC, 5x Denhardt's solution, 125 mg/mL Baker's yeast tRNA, 250 mg/mL denatured herring sperm DNA) for 2 h at 60°C . Sections were then incubated in riboprobe overnight at the same temperature. Experimental slides were exposed to anti-sense Fluorescein-labelled probe, whereas control slides were incubated with sense Fluorescein-labelled probe. After RNase A treatment at 37°C for 15 min, sections were washed in a decreasing series of SSC and equilibrated in 150 mM NaCl/100 mM Tris (pH 7.5) at room temperature before incubation in 1:1000 anti-Fluorescein-alkaline phosphatase Fab fragments (Roche) in 0.5% Tween 20/PBS for 2 h at room temperature. Sections were then washed in 100 mM Tris (pH 7.5). Chromogenic product was formed using BM Purple (Roche) at room temperature until desired darkness was achieved and was terminated simultaneously for all slides. For the nNOS *in situ* hybridization, the sections were then dehydrated, delipidated, and coverslipped under Permount (Fisher).

NADPH Diaphorase Histochemistry

Brains were removed, fixed in 4% paraformaldehyde at 4°C overnight, cryoprotected in 30% sucrose overnight at 4°C , and then embedded in OCT and stored at -80°C until sectioning. Brains were then sliced on a cryostat at $20\text{ }\mu\text{m}$ and thaw-mounted onto Super-Frost Plus slides (Erie Scientific Co., Portsmouth, NH) in three series. Slides were taken from the -80°C , air-dried briefly, and processed for NADPH diaphorase histochemistry as previously described (14). Briefly, slides were fixed for 20 min in chilled 4% paraformaldehyde in 0.1 M phosphate buffer, pH 7.4 (PB). After two washes in PB, slides were incubated for 2 hours at 37°C in PB containing 0.3% triton-X, 0.1 mg/mL nitro blue tetrazolium, and 0.25 mg/mL β -NADPH. Samples intended for co-localization were immediately processed for PR immunohistochemistry. All other sections were then rinsed, dehydrated and cover-slipped with Permount. For control sections, all procedures were the same except that β -NADPH was omitted.

Dual immunohistochemistry for PR

After processing for NADPH diaphorase histochemistry, sections were then rinsed in PBS, and incubated in 3% hydrogen peroxide in PBS for 20 min. After washing in PBS, antigen retrieval was performed by incubating in boiling citrate buffer (10mM Citric Acid, 0.05% Tween 20, pH 6.0). After two min, the boiling citrate buffer was replaced two times and incubated for five min each, followed by a PBS wash. After blocking for 1 hour in blocking solution (5% normal goat serum and 0.3% TritonX-100 in PBS), sections were incubated in primary antibody (PR 1:500, abcam 2767, monoclonal antibody raised against chicken PR) in PBS with 2% normal goat serum and 0.3% Triton-X at room temperature overnight. Sections were then rinsed, incubated for 2 hours in a biotinylated goat anti-mouse secondary antibody (1:200, Vector Laboratories), rinsed again and, after treatment with the ABC peroxidase staining kit (Vector Laboratories) according to the manufacturer's instructions,

immunoreactivity was visualized using 3,3'-diaminobenzidine (DAB) substrate (Vector Labs). Sections were then dehydrated and cover-slipped with Permount (Fisher Scientific, Itasca, IL). For control sections, all procedures were the same except that primary antibody was omitted.

Western Blot characterization of the PR antibody—To determine whether this antibody would bind specifically to the lizard antigens, we extracted protein from whole brain using a Mammalian Cell Lysis kit (Sigma) according to the manufacturer's instructions. Whole brain protein extract was run on an SDS-PAGE gel in replicate, in which one half of the gel was used for downstream Western blotting and the other half was exposed to Coomassie stain to verify protein presence. Whole brain extract on the gel was transferred onto a nitrocellulose membrane overnight. The membrane was then blocked in 5% dry milk in wash buffer (0.5% TritonX-100, 0.1% Tween-20 in 1X Tris-buffered saline [TBS]) and incubated in primary antibody (1:2000 PR in 1X TBS and 2% NaN₃) for one hour. After incubation, the membrane was washed five times for three min each in wash buffer, and then incubated in goat-anti-mouse HRP-conjugated antibody (Southern Biotech) in blocking solution for 30 min. After washing five times for three min each with wash buffer, the membrane was exposed to HRP substrate (Immobilon Western, Millipore) and exposed to film for five minutes. Using the PR antibody, one band was visualized at the predicted size of 130 kD, putatively representing PR.

Densitometric Analysis of nNOS Expression

Densitometric analysis of expression levels was measured as in Zhao et al. (23) and Zhang et al. (24). Briefly, images of nNOS *in situ* hybridization slides were captured using a Nikon Eclipse 80i microscope with a Nikon 12-bit 2-megapixel monochrome camera (DS-2MBWc) and analyzed using Nikon NIS-Elements BR 2.30 program. An investigator blind to group status measured the mean optical density of a defined region using grid sizes of 100 micron by 100 micron obtained from 2-3 individual sections from each animal. The background optical density was measured simultaneously to account for nonspecific hybridization using a grid size of 100 micron by 100 micron. The background was defined on each section as an adjacent area of tissue that did not contain a specific hybridization signal. The specific hybridization signal was obtained by subtracting the background optical density from the optical density in the specifically labelled nuclei. The mean optical density of the target POA subnuclei was averaged for each section per individual. The values from each treatment group were averaged to obtain the final mean $\pm$ SEM. Non-radioactive and radioactive *in situ* hybridization has been shown to give similar results in previous studies (25).

Cell Counting

Cells labelled by NADPH diaphorase histochemistry were counted using the Fractionator routine of the Stereo Investigator software package (Microbrightfield, Williston, VT, USA). A region of interest was defined under low power, and then, under higher magnification, positive cells were counted that fell within 50 μ m-square counting frames. The computer placed counting frames systematically every 80 μ m (i.e. with 30 μ m between each counting frame) within the region, after a randomly chosen start-site. Cell cytoplasm containing

nNOS protein was clearly marked by dark purple staining after diaphorase histochemistry, and was counted using a 20X objective. Slides were coded and processed by a blinded observer.

Statistical Analysis

All statistical analysis was conducted using JMP for Mac with significance set at $p < 0.05$. For *in situ* hybridization and cell counting data, Kruskal-Wallis was conducted using hormonal state as the independent variable and mRNA level or cell number as dependent variables. Kruskal-Wallis was chosen because most of our data sets were not normally distributed and such non-parametric tests are more conservative for small sample sizes. Mann-Whitney U tests were applied post-hoc.

RESULTS

General distribution of nNOS

As distribution of nNOS within the whiptail brain is unknown, we used NADPH diaphorase histochemistry to visualize nNOS cells and fibers (26). To our knowledge there are no commercially available antibodies that recognize the reptilian nNOS protein. Here we present the distribution of NADPH diaphorase histochemistry (Figure 1) throughout the brain of the whiptail lizard. For this study we used *C. uniparens* individuals displaying either female-like or male-like sexual behaviours and *C. inornatus* males. The only qualitative (total presence or absence) difference between the groups was in the pvPOA.

In the rostral telencephalon, there are NADPH diaphorase positive cell bodies dorsal to cortex medialis (CXM) and cortex dorsalis (CXD), as well as in the dorsal ventricular ridge (DVR) (Figure 1A). More caudally, the number of NADPH diaphorase positive cells increase in the DVR. There are also some cell bodies and fibers positive for NADPH diaphorase in the nucleus accumbens (ACC) and the ventral striatum (STR; Figure 1B). Caudal to the striatum, these cell bodies and fibers appear to wrap around the lateral forebrain bundle (LFB; Figure 1C). There are a few cells positive for NADPH diaphorase staining in the external amygdaloid nucleus (AME), but no fibers. Interestingly, the pvPOA contains many cells and fibers positive for NADPH diaphorase (Figure 1C1), but staining is completely absent from the mPOA. The caudal portion of the pvPOA only has cells positive for NADPH diaphorase histochemistry in the dorsal part surrounding the third ventricle (Figure 1D). Near this portion of the caudal POA, NADPH diaphorase-positive cells and fibers are no longer alongside the LFB, although there are cells in both the central amygdaloid nucleus (AMC) and AME and fibers only in the AME. There are also a few cells and fibers positive for NADPH diaphorase staining in the spheris lateralis nucleus (NSL), although no reactivity was seen here in the more rostral regions of the NSL.

In the diencephalon, there are few cells and fibers positive for NADPH diaphorase histochemistry in the lateral hypothalamic area (LHA) and dorsolateral anterior nucleus (DL) (Figure 1E). More caudally there are cells and fibers positive for NADPH histochemistry in the lentiformis thalami pars plicata (LTP; Figure 1F). In the hypothalamic regions, the ventromedial nucleus of the hypothalamus (VMN, Figure 1F1) only has fibers

while the dorsal hypothalamus (DH) only has cells positive for NADPH diaphorase histochemistry.

Hormonal modulation of nNOS and facilitation of male-typical behaviour

To determine the relationship between nNOS and progesterone-facilitated mounting behaviour, we quantified nNOS mRNA levels in individuals with different hormonal conditions. *In situ* hybridization for nNOS in the pvPOA was performed in both *C. inornatus* males and *C. uniparens* animals that displayed mounting behaviour in response to progesterone or testosterone compared to non-mounting individuals. Progesterone-sensitive (n=8) and testosterone-treated (n=3) male *C. inornatus* have comparable nNOS mRNA levels in the pvPOA that are significantly greater than progesterone-insensitive (n=10) and castrated + blank-implanted (n=10) *C. inornatus* males, none of which mount (Kruskal-Wallis test: Chi square value=21.624, df=3, $p<0.0001$, post-hoc Mann-Whitney U test, $p<0.004$) (Figure 2A). Similarly, intact PostOv *C. uniparens* (n=14) have nNOS expression levels in the pvPOA comparable to ovariectomized individuals implanted with testosterone (OVX+T, n=10) (Figure 2B). Both PostOv and OVX+T groups have significantly higher levels of nNOS in the pvPOA than PreOv (n=12) and OVX+Blank (n=10) groups (Kruskal-Wallis test: Chi square value=25.931, df=3, $p<0.0001$, post hoc Mann-Whitney U test, $p<0.004$).

Colocalization of nNOS with PR

We next examined the number of cells positive for nNOS protein in the pvPOA by NADPH diaphorase histochemistry in PreOv (n=13) and PostOv (n=13) *C. uniparens*. nNOS-positive cell number within the pvPOA is significantly higher in PostOv *C. uniparens* compared to PreOv individuals (Wilcoxon Rank Sum: Chi Square value= 8.153, df=1, $p<0.0001$, Figure 3C), which supports the results found with mRNA. We also colocalized cells in the pvPOA with nNOS cells by immunohistochemistry for PR. We found that almost 90% of nNOS-positive cells colocalized with PR in the pvPOA of *C. uniparens* (Figure 3B). Taken together, this suggests that nNOS plays a role in progesterone-mediated facilitation of mounting behaviour in a similar mechanism to testosterone-facilitation.

Hormonal regulation of the nNOS promoter

To determine if PR can regulate transcription of nNOS, we analyzed the promoter regions of both the rat and the whiptail lizard nNOS gene, and found several putative steroid hormone receptor binding sites (Figure 4), which are relatively conserved between rats and whiptails. In addition to many putative androgen and oestrogen receptor response elements (ARE and ERE), two putative PREs are located upstream of the whiptail translation start site, while the rat nNOS promoter has a single putative PRE. To our knowledge, this is the first analysis of the nNOS promoter region in any species, and supports our hypothesis that transcription of nNOS may be directly controlled by steroid hormone receptors in whiptail lizards.

Species differences in nNOS across the ovarian cycle

We next investigated the between-species differences of female whiptails in nNOS regulation in different ovarian stages, as females of both the ancestral and descendent

species have high levels of progesterone during the post-ovulatory phase of the ovarian cycle, but only the parthenogens respond to progesterone with male-like mounting behaviour. To better understand the molecular basis for this species difference, nNOS-positive cells were quantified within the pvPOA of naturally cycling *C. uniparens* and *C. inornatus* females. PostOv *C. uniparens* (n=18) have significantly more NADPH diaphorase-positive cells in the pvPOA than PreOv *C. uniparens* (n=12) and both PreOv (n=9) and PostOv (n=9) *C. inornatus* (Kruskal Wallis test: Chi square value=24.222, df=3, $p<0.0001$; post-hoc Mann Whitney U test, $p<0.002$) (Figure 5). There was no significant difference between PostOv and PreOv *C. inornatus* females (post-hoc Mann Whitney U test, $p=0.136$). These numbers in PostOv *C. uniparens* are comparable to those found in testosterone-implanted *C. inornatus* males (14), suggesting that nNOS is upregulated in PostOv *C. uniparens* and facilitates male-like mounting behaviour.

DISCUSSION

By investigating species differences in the regulation of a gene crucial for the display of male-typical mounting behaviour, we have shown that nNOS is upregulated in individuals displaying mounting behaviour by both progestins and androgens. Although nNOS cell bodies and fibers are widely distributed throughout the whiptail brain, we found qualitative differences (absence/presence) between different reproductive states only in the pvPOA. We discuss here the role of the pvPOA in whiptail lizards, modulation of nNOS by steroid hormones, and the evolutionary implications of these results.

nNOS distribution and role of the pvPOA

NADPH diaphorase and cells immunoreactive to nNOS are comparable both in brain and peripheral tissues (17). The distribution of NADPH diaphorase-positive cells and fibers were similar to that reported (27) for the gecko, with the exception that NADPH diaphorase cells were not observed in the whiptail cortex. The only region where we observed qualitative differences (complete absence or presence) in the number of cells positive for NADPH diaphorase between sex, species, and ovarian state was the pvPOA, which became the brain region of focus for this study. However, it should be noted that quantitative differences in nNOS was not examined in other brain regions that could potentially be regulating reproductive behaviour. The presence of NADPH diaphorase cells in the POA seems to be highly conserved across vertebrates, including mammals (28), birds (29), frogs (30), fish (31). We hypothesize that the pvPOA in whiptails is more similar to the mammalian mPOA due to its central role in mediating male-typical copulatory behaviour (32,33) and the neurochemical profile of steroid hormone receptors and nNOS expression (11,28, present study).

Interestingly, whiptail lizards, gecko (27), and turtle (34) show NADPH diaphorase fibers but not cell bodies within the ventromedial hypothalamus. This is in contrast to results seen in birds (29), mammals (35), and frogs (30), where cell-bodies positive for NADPH diaphorase are clearly visible. This observation is interesting, as the ventromedial hypothalamus is crucial for the display of female-typical receptivity in mammals and reptiles (36,37) and nNOS plays a role in facilitating female receptivity in rats (16). It will

be interesting to see if nNOS plays a role in female receptivity in whiptail lizards as well as male-typical copulatory behaviour.

Integration of behavioural and neuroendocrine mechanisms

The role of nitric oxide in male sexual behaviour has been extensively studied in mammals. nNOS expression in the rat POA is reduced by castration (38) and increased by either testosterone or oestradiol (39). Hull and colleagues (40) have proposed an elegant model of the hormonal gating of male sexual behaviour in which testosterone (after aromatization to oestradiol) facilitates an upregulation of nNOS. Sensory stimuli from the amygdala trigger glutamate release that stimulates NMDA receptors, and after an influx of calcium, calmodulin activates nNOS. Synthesis of nitric oxide then increases the level of dopamine in the POA, which increases the probability of a copulatory response. These same androgen-mediated mechanisms seem to play out in male *C. inornatus* as well (14) and on a time course that parallels reinstatement of copulatory behaviour after castration. Although work in rodent models with nNOS facilitating male sexual behaviour has mainly been limited to androgens and oestrogens, it would be interesting to investigate whether progesterone also plays a role in the nNOS pathway in rodents, as progesterone also facilitates mounting behaviour in male mammals (41).

The regulation of nNOS by oestrogens and androgens has been well studied in rodents in the context of sexual behaviour. However, the regulation of nNOS by either androgens or oestrogens seems to be sex and steroid specific, as oestradiol and progesterone will increase nNOS in the POA in female rats but not male rats (42). In female rats, both oestradiol and progesterone upregulate nNOS protein in the POA and hypothalamus both *in vivo* and *in vitro* hypothalamic cells (43). In male rats, androgens are required for the development of the nNOS system (44), while oestradiol seems to be necessary for the role of nNOS is facilitating sexual behaviour in adults (39).

Based on the insights gained from mammals, we can speculate on the synergistic actions of progesterone and nitric oxide in the whiptail lizards. It seems that progesterone via the progesterone receptor has taken on a new role in the parthenogens to facilitate male-typical mounting behaviour, and that this behaviour requires nitric oxide (15,45). Presumably, PR upregulates nNOS, which through a cascade of events influences dopamine levels in the POA, and leads to an increased probability of mounting behaviour. This is important, as the copulatory response to a receptive female is influenced by the dopamine availability in the POA of male rats (13). However, at this point we cannot exclude possible local steroid synthesis that may be regulating nNOS in the whiptail pvPOA. We have shown that the nNOS promoter has putative steroid hormone response elements sensitive to oestradiol and androgens as well as progestins, although these results will be confirmed with promoter assays in future studies. The results presented here strongly support the interaction between PR and nNOS in promoting male-typical sexual behaviour in the all-female whiptail species, *C. uniparens*, although to firmly establish a functional relationship pharmacological studies need to be conducted.

In addition to providing insights into the role of nNOS in sexual behaviour, nNOS knockout mice have also revealed the role of nNOS in regulating aggression, possibly via the

serotonin system, as these mice are highly aggressive and will repeatedly mount unreceptive females (46,47). Serotonin metabolism is drastically decreased in nNOS knockout mice, and these animals also have deficits in serotonin receptor expression throughout several brain regions (48). PostOv *C. uniparens* individuals have lower serotonin levels in the POA compared to PreOv individuals and infusions of serotonin directly into the POA increased mount latency in testosterone-implanted individuals (22). Given the interaction of serotonin and nitric oxide, it would be interesting to look at the dynamics of these two molecules in the context of regulating sexual behaviour in whiptails.

Molecular substrates of evolution

Through the comparative analysis of the nNOS promoter region, we have found more putative progesterone response elements (PREs) in the nNOS promoter region of whiptails compared to rats. Rodriguez-Trelles and colleagues (49) have suggested that promoter sequences (more so than coding regions) are a key substrate for selection and the evolution of between-species differences in phenotype. Given that a transcription factor such as PR interacts with numerous promoters, it seems more likely that evolutionary novel regulation of genes required for induction of male-typical mounting behaviours by PR is due to changes in promoter sequences (gain or loss of PREs) rather than the transcription factor specificity for its response element (50), which the data presented here support.

The work presented here have opened a new avenue of possible research into mechanisms that govern species differences in gene expression. *C. inornatus* females and *C. uniparens* both have a progesterone surge during the postovulatory phase of the ovarian cycle (5,7), and PR is present in the pvPOA of both species during the postovulatory phase (present study). However, only in *C. uniparens* does progesterone via PR elicit the male-typical mounting behaviour. Additionally, only *C. uniparens* has high levels of nNOS present in the pvPOA during the postovulatory phase. This observation begs the question how does PR upregulate nNOS in *C. uniparens* but not the *C. inornatus* females? This question can also be asked of P-sensitive and P-insensitive *C. inornatus* males. We hypothesize that differences in regulation of the promoter sequences and the epigenome could be regulating the polymorphism to progesterone sensitivity and will be thoroughly investigated in the future.

Conclusion

Evolutionary change results from exploitation of variation leading to the emergence of new molecules and, as in this case, coopting endocrine signals for new functions (8). These data support the hypothesis that a progesterone-sensitive *C. inornatus* individual played a major role in the hybridization events that gave rise to the all-female *C. uniparens* species and, in the absence of androgens, the evolution of a novel neuroendocrine mechanism controlling pseudosexual mounting in response to high levels of progesterone. Future work will investigate the role of nNOS in progesterone-facilitation of male-typical mounting behaviour using pharmacological manipulations of both nNOS and PR, as well as female receptivity in both the ancestral and descendant species.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We thank Nicholas Sanderson, Vicky Huang, Yuiko Matsumoto, and Ryan Wong for technical contributions; Christina Shoemaker and Nicholas Sanderson for discussions; Brian Dias and Vicky Huang for help collecting lizards; Molly Cummings and Hans Hofmann for generosity in using their equipment; Juan Dominguez and Lin Huffman for comments on earlier versions of this manuscript; and Fernando Nottebohm and Andrea Gore for editorial suggestions. Research was supported by MH41770 to D.C.

REFERENCES

1. Tinbergen N. On Aims and Methods in Ethology. *Zeitschrift für Tierpsychologie*. 1963; 20:410–433.
2. Crews D. Evolution of neuroendocrine mechanisms that regulate sexual behaviour. *Trends Endocrin Metab*. 2005; 16:354–361.
3. Cantant, R.; Collins, JT. Reptiles and amphibians: Eastern and Central North America. Houghton Mifflin Co; New York: 2003.
4. Crews D, Fitzgerald KY. “Sexual” behavior in parthenogenetic lizards (*Cnemidophorus*). *Proc Natl Acad Sci USA*. 1980; 77:499–502. [PubMed: 16592761]
5. Moore MC, Whittier JM, Billy AJ, Crews D. Male-like behavior in an all-female lizard: Relationship to ovarian cycle. *Animal Behav*. 1985; 33:284–289.
6. Crews D, Grassman M, Lindzey J. Behavioral facilitation of reproduction in sexual and unisexual whiptail lizards. *Proc Natl Acad Sci USA*. 1986; 83:499–502.
7. Moore MC, Whittier JM, Crews D. Sex steroid hormones during the ovarian cycle of an all-female, parthenogenetic lizard and their correlation with pseudosexual behaviour. *Gen Comp Endocr*. 1985; 60:144–153. [PubMed: 4065527]
8. Crews D, Moore MC. Evolution of mechanisms controlling mating behaviour. *Science*. 1986; 231:121–125. [PubMed: 3941893]
9. Lindzey J, Crews D. Interactions between progesterone and androgens in the stimulation of sex behaviors in male little striped whiptail lizards, *Cnemidophorus inornatus*. *Gen Comp Endocr*. 1992; 86:52–58. [PubMed: 1505730]
10. Lindzey J, Crews D. Effects of progestins on sexual behavior in castrated lizards (*Cnemidophorus inornatus*). *J Endocr*. 1988; 119:265–271. [PubMed: 3199058]
11. Young LJ, Lopreato GF, Horan K, Crews D. Cloning and *in situ* hybridization analysis of estrogen receptor, progesterone receptor, and androgen receptor expression in the brain of whiptail lizards (*Cnemidophorus uniparens* and *C. inornatus*). *J Comp Neurol*. 1994; 347:288–300. [PubMed: 7814669]
12. Crews D, Godwin J, Hartman V, Grammar M, Prediger EA, Shephard R. Intrahypothalamic implantation of progesterone in castrated male whiptail lizards (*Cnemidophorus inornatus*) elicits courtship and copulatory behavior and affects androgen receptor and progesterone receptor-mRNA expression in the brain. *J Neurosci*. 1996; 16:7347–7352. [PubMed: 8929441]
13. Hull EM, Dominguez JM. Getting his act together: roles of glutamate, nitric oxide, and dopamine in the medial preoptic area. *Brain Res*. 2006; 1126:66–75. [PubMed: 16963001]
14. Sanderson NS, Le B, Zhou Z, Crews D. Preoptic neuronal nitric oxide synthase induction by testosterone is consistent with a role in gating male-typical copulatory behaviour. *Eur J Neurosci*. 2008; 27:183–901. [PubMed: 18184320]
15. Sanderson NS, Weissler E, Crews D. The nitric oxide synthase inhibitor L-NAME suppresses androgen-induced male-like pseudocopulatory behavior in whiptail lizards. *Brain Res*. 2005; 1052:236–239. [PubMed: 16023092]

16. Mani SK, Allen JM, Rettori V, McCann SM, O'Malley BW, Clark JH. Nitric oxide mediates sexual behavior in female rats. *Proc Natl Acad Sci USA*. 1994; 91:6468–6472. [PubMed: 7517551]
17. Dawson TM, Bredt DS, Fotuhi M, Hwang PM, Snyder SH. Nitric oxide synthase and neuronal NADPH diaphorase are identical in brain and peripheral tissues. *Proc Natl Acad Sci U S A*. 1991; 88:7797–7801. [PubMed: 1715581]
18. Lindzey J, Crews D. Hormonal control of courtship and copulatory behavior in male *Cnemidophorus inornatus*, a direct sexual ancestor of a unisexual, parthenogenetic lizard. *Gen Comp Endocr*. 1986; 64:411–418. [PubMed: 3803894]
19. Sakata JT, Woolley SC, Gupta A, Crews D. Differential effects of testosterone and progesterone on the activation and retention of courtship behavior in sexual and parthenogenetic *Cnemidophorus* whiptail lizards. *Horm Behav*. 2003; 43:523–530. [PubMed: 12799168]
20. Wade J, Crews D. The relationship between reproductive state and “sexually” dimorphic brain areas in sexually reproducing and parthenogenetic whiptail lizards. *J Comp Neurol*. 1991; 309:507–514. [PubMed: 1918445]
21. Burmeister SS, Fernald RD. Evolutionary conservation of the *egr-1* immediate-early gene response in a teleost. *J Comp Neurol*. 2005; 481:220–232. [PubMed: 15562507]
22. Dias BG, Crews D. Regulation of pseudosexual behavior in the parthenogenetic whiptail lizard, *Cnemidophorus uniparens*. *Endocrinology*. 2008; 149:4622–4631. [PubMed: 18483155]
23. Zhao C, Fujinaga R, Yanai A, Kokubu K, Takeshita Y, Watanabe Y, Shinoda K. Sex steroidal regulation of aromatase mRNA expression in adult male rat brain: a quantitative non-radioactive *in situ* hybridization study. *Cell Tissue Res*. 2008; 332:381–391. [PubMed: 18401595]
24. Zhang L, Ravipati A, Joseph J, Roth GS. Aging-related changes in rat striatal D2 receptor mRNA-containing neurons: a quantitative nonradioactive *in situ* hybridization study. *J Neurosci*. 1995; 15:1735–1740. [PubMed: 7891131]
25. Kreft S, Zajc-Kreft K, Zivin M, Sket D, Grubic Z. Application of the nonradioactive *in situ* hybridization for the localization of acetylcholinesterase mRNA in the central nervous system of the rat; comparison to the radioactive technique. *Pflugers Arch*. 1996; 431:R309–310. [PubMed: 8739388]
26. Hope BT, Michael GT, Knippe KM, Vincent SR. Neuronal NADPH diaphorase is a nitric oxide synthase. *Proc Natl Acad Sci USA*. 1991; 88:2811–2814. [PubMed: 1707173]
27. Smeets WJ, Alonso JR, González A. Distribution of NADPH diaphorase and nitric oxide synthase in relation to catecholaminergic neuronal structures in the brain of the lizard *Gekko gekko*. *J Comp Neurol*. 1997; 377:121–141. [PubMed: 8986877]
28. Sato S, Braham CS, Putnam SK, Hull EM. Neuronal nitric oxide synthase and gonadal steroid interaction in the MPOA of male rats: co-localization and testosterone-induced restoration of copulation and nNOS-immunoreactivity. *Brain Res*. 2005; 1043:205–213. [PubMed: 15862534]
29. Atoji Y, Yamamoto Y, Suzuki Y. Distribution of NADPH diaphorase-containing neurons in the pigeon central nervous system. *J Chem Neuroanat*. 2001; 21:1–22. [PubMed: 11173217]
30. Brüning G, Mayer B. Localization of nitric oxide synthase in the brain of the frog, *Xenopus laevis*. *Brain Res*. 1996; 741:331–343. [PubMed: 9001739]
31. Villani L, Guarnieri T. Ultrastructural localization of NADPH diaphorase in the goldfish brain. *Brain Res*. 1995; 705:332–226. [PubMed: 8821765]
32. Hurtazo HA, Paredes RG, Agmo A. Inactivation of the medial preoptic area/anterior hypothalamus by lidocaine reduces male sexual behavior and sexual incentive motivation in male rats. *Neuroscience*. 2008; 152:331–337. [PubMed: 18262728]
33. Kingston PA, Crews D. Effects of hypothalamic lesions on courtship and copulatory behavior in sexual and unisexual whiptail lizards. *Brain Res*. 1994; 643:349–351. [PubMed: 8032930]
34. Brüning G, Wiese S, Mayer B. Nitric oxide synthase in the brain of the turtle *Pseudemys scripta elegans*. *J Comp Neurol*. 1994; 348:183–206. [PubMed: 7529267]
35. Okamura H, Yokosuka M, McEwen BS, Hayashi S. Colocalization of NADPH diaphorase and estrogen receptor immunoreactivity in the rat ventromedial hypothalamic nucleus: stimulatory effect of estrogen on NADPH diaphorase activity. *Endocrinology*. 1994; 135:1705–1708. [PubMed: 7925135]

36. Mathews D, Greene SB, Hollingsworth EM. VMN lesion deficits in lordosis: partial reversal with pergolide mesylate. *Physiol Behav.* 1983; 31:745–748. [PubMed: 6665063]
37. Kendrick AM, Rand MS, Crews D. Electrolytic lesions to the ventromedial hypothalamus abolish receptivity in female whiptail lizards, *Cnemidophorus uniparens*. *Brain Res.* 1995; 680:226–228. [PubMed: 7663980]
38. Du J, Hull EM. Effects of testosterone on neuronal nitric oxide synthase and tyrosine hydroxylase. *Brain Res.* 1999; 836:90–98. [PubMed: 10415408]
39. Putnam SK, Sato S, Riolo JV, Hull EM. Effects of testosterone metabolites on copulation, medial preoptic dopamine, and NOS-immunoreactivity in castrated male rats. *Horm Behav.* 2005; 47:5, 13–22.
40. Hull EM, Muschamp JW, Sato S. Dopamine and serotonin: influences on male sexual behaviour. *Physiol Behav.* 2004; 83:291–307. [PubMed: 15488546]
41. Witt DM, Young LJ, Crews D. Progesterone and sexual behavior in males. *Psychoneuroendocr.* 1994; 12:553–562.
42. Ishihara T, Orikasa C, Araki T, Sakuma Y. Sex difference in the expression and regulation of nitric oxide synthase gene in the rat preoptic area. *Neurosci Res.* 2002; 43:147–154. [PubMed: 12067750]
43. Chu HP, Sarkar G, Etgen AM. Estradiol and progesterone modulate the nitric oxide/cyclic gmp pathway in the hypothalamus of female rats and in GT1-1 cells. *Endocrine.* 2004; 24:177–184. [PubMed: 15347845]
44. Martini M, Di Sante G, Collado P, Pinos H, Guillaumon A, Panzica GC. Androgen receptors are required for full masculinization of nitric oxide synthase system in rat limbic-hypothalamic region. *Horm Behav.* 2008; 54:557–564. [PubMed: 18582470]
45. Grassman M, Crews D. Progesterone induction of pseudocopulatory behavior and stimulus-response complementarity in an all-female lizard species. *Horm Behav.* 1986; 20:327–335. [PubMed: 3770655]
46. Nelson RJ, Demas GE, Huang PL, Fishman MC, Dawson VL, Dawson TM, Snyder SH. Behavioral abnormalities in male mice lacking neuronal nitric oxide synthase. *Nature.* 1995; 378:383–386. [PubMed: 7477374]
47. Nelson RJ, Trainor BC, Chiavegatto S, Demas GE. Pleiotropic contributions of nitric oxide to aggressive behaviour. *Neurosci Biobehav Rev.* 2006; 30:346–355. [PubMed: 16483891]
48. Chiavegatto S, Dawson VL, Mamounas LA, Koliatsos VE, Dawson TM, Nelson RJ. Brain serotonin dysfunction accounts for aggression in male mice lacking neuronal nitric oxide synthase. *Proc Natl Acad Sci U S A.* 2001; 98:1277–1281. [PubMed: 11158630]
49. Rodríguez-Trelles F, Tarrío R, Ayala FJ. Evolution of cis-regulatory regions versus codifying regions. *Int J Dev Biol.* 2003; 47:665–673. [PubMed: 14756342]
50. Tautz D. Evolution of transcriptional regulation. *Curr Opin Genet Dev.* 2000; 10:575–579. [PubMed: 10980438]

Figure 1. Distribution of NADPH diaphorase histochemistry in the brain of whiptail lizards
Representative sections of the whiptail brain are depicted on the left (A-F) with nNOS-positive cell bodies represented by dots and fibers depicted as grey shading. Micrographs on the right panel indicate NADPH diaphorase staining in the anteroventral periventricular nucleus of the POA (pvPOA, C1) and the ventromedial nucleus of the hypothalamus (VMN, F1). All scale bars are shown at 50µm.

Abbreviations: nucleus accumbens (ACC), Nucleus centralis amygdalae (AMC), Nucleus externus amygdalae (AME), periventricular nucleus of the preoptic region (PvPOA), cortex dorsalis (CXD), cortex lateralis (CXL), cortex medialis (CXM), nucleus dorsalis hypothalami (DH), nucleus dorsolateralis anterior (DL), nucleus dorsomedialis (DM), dorsal ventricular ridge (DVR), lateral forebrain bundle (LFB), lateral hypothalamic area (LHA), lateral POA (LPA), lentiformis thalami pars plicata (LTP), medial preoptic area (mPOA), nucleus sphericus (NS), nucleus sphericus anterior (NSA), nucleus sphericus lateralis (NSL), nucleus sphericus medialis (NSM), optic tract (OT), posterior commissure (PC), nucleus periventricularis hypothalami (PH), nucleus supraopticus (SO), striatum (STR), optic tectum (TECT), ventromedial nucleus of the hypothalamus (VMN).

Figure 2. Regulation of nNOS mRNA expression in the pvPOA by hormone state

(A) P-sensitive males and testosterone implanted *C. inornatus* males, both of which display mounting behaviour, have significantly higher levels of nNOS mRNA in the pvPOA than P-insensitive and castrated (Blank) males who do not mount (Kruskal-Wallis test, $p < 0.0001$, post-hoc Mann-Whitney U test, $p < 0.004$). (B) Postovulatory (PostOv) and ovariectomized + testosterone-implanted *C. uniparens* have higher levels of nNOS mRNA in the pvPOA than preovulatory (PreOv) and ovariectomized (OVX) + blank implanted animals (Kruskal-Wallis test, $p < 0.0001$, post-hoc Mann-Whitney U test, $p < 0.004$). Histograms are means and error bars are at $\pm$ SEM. In the third panel are representative micrographs of PostOv and PreOv *C. uniparens* pvPOA nNOS density. Abbreviations as in Figure 1.

Figure 3. Regulation of nNOS in the pvPOA of *C. uniparens* and colocalization with PR nNOS-positive cells detected by NADPH diaphorase histochemistry (blue staining) were counted in PreOv (A) and PostOv (B) *C. uniparens* females. PostOv animals have significantly more nNOS-positive cells in the pvPOA than PreOv animals (C; Wilcoxon Rank Sum test, $p < 0.0001$). Slides were also processed for PR immunoreactivity (brown staining) and showed most nNOS-positive cells (~90%) also contain PR. Histograms are means and error bars are at $\pm$ SEM. Abbreviations as in Figure 1.

Figure 4. Putative transcription factor binding sites in the rat and whiptail nNOS promoters
The rat nNOS promoter (top) and whiptail nNOS promoter (bottom) show putative progesterone response elements (PRE, black boxes), putative androgen response elements (ARE; white boxes), and putative estrogen response elements (ERE, hatched boxes).

Figure 5. Species differences in nNOS regulation in the pvPOA across the ovarian cycle
 Micrographs of an NADPH diaphorase histochemistry showing nNOS-positive cells within the pvPOA of *C. uniparens* (A) and *C. inornatus* females (B). PostOv *C. uniparens* have significantly more cells within the pvPOA than all other groups suggesting that presence of nNOS facilitates mounting behaviour (Kruskal Wallis test, $p < 0.0001$, post-hoc Mann Whitney U test, $p < 0.002$). Histograms are means and error bars are at $\pm$ SEM. Scale bar shown at 50 μ m. Abbreviations as in Figure 1.