Published in final edited form as:

Microbiol Spectr. 2015 June ; 3(3): . doi:10.1128/microbiolspec.MBP-0006-2014.

Commensal and Pathogenic *Escherichia coli* Metabolism in the Gut

Tyrrell Conway¹ and Paul S. Cohen²

¹Department of Microbiology and Plant Biology, University of Oklahoma, Norman, OK 73019,
Phone: 405-820-7329, tconway@ou.edu

²Department of Cell and Molecular Biology, University of Rhode Island, Kingston, RI 02881,
Phone: 401-874-5920, pco1697u@mail.uri.edu

Introduction

Every mammal on the planet is colonized with *E. coli* (1), as well as cold blooded animals (e.g., fish) at an appropriately warm temperature (2). We estimate there are 10^{21} *E. coli* cells in the human population alone. *E. coli* is frequently the first bacterium to colonize human infants and is a lifelong colonizer of adults (3). *E. coli* is arguably the best understood of all model organisms (4). Yet the essence of how *E. coli* colonizes and/or causes disease is still not completely understood. Certainly, innate immunity, adaptive immunity, and bacterial cell-to-cell communication play important roles in modulating the populations of the 500–1000 different commensal species in the intestine (5–11); however, these topics will not be a focus of this chapter. We have reviewed the mucus layer as habitat for *E. coli* to colonize the intestine, aspects of *E. coli* physiology that enable its success, and the model systems employed for colonization research (12–14). Here, we focus on *E. coli* metabolism in the intestinal mucus layer. We discuss evidence that *E. coli* must obtain nutrients in the mucus layer to colonize, that it resides in the mucus layer as a member of mixed biofilms, and that each *E. coli* strain displays a unique nutritional program in the intestine. We also discuss evidence supporting the “Restaurant” hypothesis for commensal *E. coli* strains, i.e., that they colonize the intestine as sessile members of mixed biofilms obtaining the nutrients they need for growth locally, but compete for nutrients with invading *E. coli* pathogens planktonically.

From ingestion to colonization

When *E. coli* is eliminated by a host animal it is not growing because it cannot grow in the luminal contents of the intestine (15). *E. coli* persists in the environment until its next host consumes viable bacteria in contaminated water or adulterated food. Following ingestion, a stressor faced by *E. coli* is acidity in the stomach, which it survives because stationary phase bacteria induce protective acid resistance systems (16). Extreme acid tolerance makes *E. coli* transmissible by as few as ten bacterial cells (17). Upon reaching the colon, *E. coli* must find the nutrients it needs to exit lag phase and grow from low to high numbers. Failure to transition from lag phase to logarithmic phase will lead to elimination of the invading *E. coli* bacteria (18). Successful colonization of the colon by *E. coli* depends upon competition for nutrients with a dense and diverse microbiota (18), penetration of the mucus layer (19), but not motility (20), and the abilities to avoid host defenses (21, 22), and grow rapidly,

exceeding the turnover rate of the mucus layer (23). *E. coli* resides in mucus until being sloughed into the lumen of the intestine (24, 25), from whence some cells are eliminated in the host feces and the cycle begins again. This circle of colonization and extra-intestinal survival is the reality for commensal and pathogenic *E. coli* alike.

Basic principles of colonization

Colonization is defined as the indefinite persistence of a particular bacterial population without reintroduction of that bacterium. We agree with Rolf Freter, a true pioneer in the field of intestinal colonization, who concluded that although several factors could theoretically contribute to an organism's ability to colonize, competition for nutrients is paramount for success in the intestinal ecosystem (26). According to Freter's nutrient-niche hypothesis, the mammalian intestine is analogous to a chemostat in which several hundreds of species of bacteria are in equilibrium. To co-colonize, each species must use at least one limiting nutrient better than all the other species (18, 27, 28). The nutrient-niche hypothesis further predicts that invading species will have difficulty colonizing a stable ecosystem, such as the healthy intestine. The ability of the microbiota to resist invasion is termed colonization resistance (29), an example of which being that when human volunteers were fed *E. coli* strains isolated from their own feces those *E. coli* failed to colonize (30). Yet, despite colonization resistance, humans are colonized on average with five different *E. coli* strains and there is a continuous succession of strains in individuals (30). This suggests that diversity exists among commensal *E. coli* strains and that different strains may possess different strategies for utilizing growth-limiting nutrients.

If diversity amongst *E. coli* commensal strains plays a role in colonization resistance, then mice pre-colonized with a human *E. coli* commensal strain would resist colonization by the same strain (isogenic challenge strain) because bacteria that consume the nutrients it needs to colonize already occupy its preferred niche. However, if mice pre-colonized with one human *E. coli* commensal strain were subsequently fed a different *E. coli* strain (non-isogenic challenge strain) then if the second strain could occupy a distinct niche in the intestine it would co-colonize with the first strain. The results of such experiments showed that each of several pre-colonized *E. coli* strains nearly eliminated its isogenic challenge strain from the intestine, confirming that colonization resistance can be modeled in mice, but non-isogenic challenge *E. coli* strains grew to higher numbers in the presence of different pre-colonized strains, suggesting that the newly introduced non-isogenic challenge strain either grows faster than the pre-colonized strain on one or more nutrients or uses nutrient(s) not being used by the pre-colonized strain (31).

How might an invading enteric pathogen subvert colonization resistance? According to the nutrient-niche hypothesis, upon reaching the intestine the pathogen would first have to outcompete the resident microbiota for at least one nutrient, allowing it initially to colonize the intestine. However, colonization would not in itself result in pathogenesis if the pathogen must reach the epithelium and either bind to epithelial cells or invade the epithelium. In such instances, the pathogen must presumably penetrate the mucus layer. In a series of groundbreaking studies (32–34), Stecher, Hardt, and colleagues showed that when *Salmonella enterica* serovar Typhimurium induces inflammation in a mouse colitis model,

the composition of the microbiota is changed and its growth is suppressed while serovar Typhimurium growth is enhanced. The authors also showed that serovar Typhimurium is attracted by chemotaxis to galactose-containing nutrients on the mucosal surface (e.g. galactose-containing glycoconjugates and mucin) and, as expected, flagella and motility were required (32). Thus, to quote the authors (34), “Triggering the host’s immune defense can shift the balance between the protective microbiota and the pathogen in favor of the pathogen.”

In streptomycin-treated mice, nutrient consumption by colonized *E. coli* strains can prevent invading *E. coli* strains from colonizing (35). By examining the sugars used by various human commensal *E. coli* strains to colonize, we identified a pair of strains (*E. coli* HS and *E. coli* Nissle 1917) that together use the five sugars previously found to be most important for colonization by the enterohemorrhagic *Escherichia coli* (EHEC) strain EDL933 (O157:H7) (36). When mice were pre-colonized with *E. coli* HS and *E. coli* Nissle 1917, invading *E. coli* EDL933 was eliminated from the intestine (35). Clearly, one therapeutic strategy to prevent pathogenesis would be to outcompete the pathogen for nutrients normally present in the intestine and eliminate it before it can colonize and subsequently cause inflammation (5, 6, 37).

Implicit in the nutrient-niche hypothesis is the idea that different species compete for preferred nutrients from a mixture that is equally available to all species. However, there is growing evidence that at least under some circumstances *E. coli* receives the nutrients it needs through direct interactions with neighboring microbes in the intestinal community. Thus, we take a renewed look at the metabolism of and nutrient flow between members of the intestinal microbiota.

Central metabolism and intestinal colonization

E. coli is a Gram negative, prototrophic, facultative anaerobe with the ability to respire oxygen, use alternative anaerobic electron acceptors, or ferment, depending on electron acceptor availability. Central metabolism in *E. coli* consists of the Embden-Meyerhof-Parnas glycolytic pathway (EMP), the pentose phosphate pathway (PP), the Entner-Doudoroff pathway (ED), the TCA cycle, and diverse fermentation pathways. *E. coli* grows best on sugars, including a wide range of mono- and disaccharides, but it cannot grow on complex polysaccharides because it lacks the necessary hydrolase enzymes (36). *E. coli* also can grow on amino acids and dicarboxylates that feed into the TCA cycle; the metabolism of these nutrients requires gluconeogenesis, the biosynthesis of glucose phosphate to be used as precursors of macromolecules such as LPS and peptidoglycan. Central metabolic pathways in *E. coli* are highly conserved, constituting a significant part of the core *E. coli* genome (38). The role of central metabolism during intestinal colonization has been studied in *E. coli*. The results of these experiments are summarized below (Table 1).

Mutants blocked in glycolysis or the ED pathway, but not the PP pathway, have major colonization defects in competition with their wild type parents (39). Given its role in hexose metabolism, it is expected that glycolysis is important for colonization. Indeed, a *pgi* mutant lacking the key enzyme, phosphoglucose isomerase, of the EMP glycolytic pathway

has a substantial colonization defect when competed against its wild type *E. coli* K-12 parent (Table 1). The role of the EMP pathway goes beyond colonization by *E. coli*. For example, glucose catabolism and glycolysis are known to play a role in intracellular growth of serovar Typhimurium within macrophage vacuoles (40) and proper regulation of glucose catabolism and glycolysis are coupled to virulence factor expression in EHEC (41). A recent study of *Shigella flexneri* revealed similar usage of these central metabolic pathways to support replication within host cells (42). We conclude that glycolysis is important for *E. coli* colonization and other aspects of enteric pathogenesis.

Gluconate was the first nutrient that was shown to be used by *E. coli* to colonize the streptomycin-treated mouse intestine (43). Since gluconate and other sugar acids are primarily catabolized via the ED pathway, it is reasonable to expect that mutants lacking the pathway will be defective in colonization (44). The ED pathway is encoded by the *edd-eda* operon (45). The promoter-proximal *edd* gene encodes 6-phosphogluconate dehydratase, which converts 6-phosphogluconate to 2-keto-3-deoxy-6-phosphogluconate. The *eda* gene encodes 2-keto-3-deoxy-6-phosphogluconate aldolase, which converts 2-keto-3-deoxy-6-phosphogluconate to glyceraldehyde-3-phosphate and pyruvate. *E. coli edd* mutants lacking the ED pathway, but retaining the pentose phosphate (PP) pathway, are poor colonizers of the mouse intestine, suggesting that *E. coli* utilizes the ED pathway for growth in the intestine (43). Other enteric bacteria require the ED pathway. For example, intracellular serovar Typhimurium induces genes of the ED pathway and gluconate catabolism during growth in macrophages (46). Moreover, the ED pathway is induced by *Vibrio cholerae* in vivo and an *edd* mutant failed to colonize the mouse intestine (47).

In contrast to the importance of the ED pathway, an *E. coli gnd* mutant, missing 6-phosphogluconate dehydrogenase and therefore deficient in the oxidative branch of the PP pathway, was as good a mouse intestine colonizer as the wild type (39). It should be noted that *gnd* mutants retain the non-oxidative PP pathway; therefore they retain the ability to make essential precursor metabolites (e.g. ribose-5-phosphate) (48). We conclude that *E. coli* has alternative mechanisms for generating reducing power (NADPH) other than the oxidative PP pathway, but that the ED pathway for sugar acid catabolism is required to colonize efficiently (Table 1).

The role of the TCA cycle in commensal *E. coli* colonization of the intestine and in *E. coli* pathogenesis is poorly studied. It has been reported that an *sdhB* mutant lacking succinate dehydrogenase colonized as well as its wild type parent (39). However, *E. coli* has a second isoform of succinate dehydrogenase: fumarate reductase, which provides redundant enzyme function under some circumstances (49). Indeed, an *E. coli sdhAB frdA* double mutant has a significant colonization defect (Table 1). The role of the TCA cycle in colonization and pathogenesis by other Enterobacteriaceae is better understood, as described immediately below.

A fully functional TCA cycle is required for virulence of *Salmonella enterica* serovar Typhimurium via oral infection of BALB/c mice, i.e., a *sucCD* mutant, which prevents the conversion of succinyl coenzyme A to succinate, was attenuated. Also, an *sdhCDA* mutant, which blocks the conversion of succinate to fumarate, was attenuated, whereas both an *aspA*

mutant and an *frdABC* mutant, deficient in the ability to run the reductive branch of the TCA cycle, were fully virulent (50). Moreover, although it appears that serovar Typhimurium replenishes TCA cycle intermediates from substrates present in mouse tissues, fatty acid degradation and the glyoxylate bypass are not required, since a *fadD*, *fadF*, and *aceA* mutants were all fully virulent during acute infection (50–52). Interestingly, it appears that the TCA cycle is required for virulence of *Edwardsiella ictaluri* in catfish fingerlings (53) and that the glyoxylate bypass is required for serovar Typhimurium persistent infection of mice (51).

The fact that *E. coli* depends on the TCA cycle for colonization implies that gluconeogenesis also is important. Using mutants that are unable to synthesize glucose from fatty acids, acetate, and TCA cycle intermediates because they are blocked in converting pyruvate to phosphoenolpyruvate (*ppsA pckA*), a critical step in gluconeogenesis, it was shown that neither the commensal *E. coli* K-12 strain MG1655 nor EHEC use gluconeogenesis for growth in the streptomycin-treated mouse intestine when each is the only *E. coli* strain fed to mice (54). However, *E. coli* Nissle 1917, the probiotic strain does use gluconeogenesis to colonize (Schinner, et al, unpublished). In addition, while *E. coli* EDL933 did not use gluconeogenic nutrients when it was the only *E. coli* strain in the mouse intestine, it used metabolic flexibility to switch to gluconeogenic nutrients when in competition in the intestine with either *E. coli* MG1655 (54) or *E. coli* Nissle 1917 (Schinner, et al, unpublished). These findings are of extreme interest in view of a recent report showing that *E. coli* EDL933 activates expression of virulence factor genes only under gluconeogenic conditions (41).

Catabolic pathway diversity in *E. coli*

The substrate range of *E. coli* is limited to mono-saccharides, disaccharides, a small number of larger sugars, some polyols, and sugar acids (55). Amino acids and carboxylates also are consumed (55, 56). The corresponding catabolic pathways feed these substrates into central metabolism. While the genes encoding central metabolism in *E. coli* fall within the highly conserved core genome (38), there is predicted to be some variation between strains with respect to the catabolic pathways that feed various substrates into central metabolism, as indicated by genome based metabolic modeling (57). For example, pathogenic *E. coli* strains are predicted to grow on sucrose while commensals are not. In contrast, commensals are predicted to grow on galactonate while pathogens are not. However, most of the substrates predicted by modeling to be used differentially by different *E. coli* strains are not known to be present in the intestine (57). *E. coli* EDL933, the prototypical EHEC strain, is able to grow on sucrose, whereas most commensal strains do not because they lack the *sac* genes, and some strains are missing genes within the N-acetylgalactosamine operon and are thus unable to grow on this substrate (36). Despite the modest differences between strains regarding their substrate range, in laboratory cultures containing a mixture of thirteen different sugars known to be present in mucus polysaccharides, *E. coli* EDL933 and *E. coli* MG1655 each use the sugars in the same order (36). However, although *E. coli* strains have nearly identical catabolic potential, they vary significantly in the sugars that support their colonization (35, 36, 39, 43, 54, 58, 59).

Nutrient availability in the intestine

Fluorescent in situ hybridization microscopy (FISH) of thin sections of the cecum of streptomycin-treated mice show that colonized *E. coli* are surrounded by other members of the microbiota within the mucus layer (19, 54, 60). Indeed, *E. coli* grows well in vitro on cecal mucus, but fails to grow in intestinal contents (15, 19). When a transposon insertion mutant library was screened for poor growth on mucus agar plates, a *waaQ* mutant of *E. coli* K-12 was isolated that also was defective in lipopolysaccharide biosynthesis, sensitive to detergents, clumped in vitro in broth culture, and failed to colonize streptomycin-treated mice (19). While the *waaQ* strain initially (during the first 24 h) grew from low to high numbers in the intestine, it rapidly declined in fecal plate counts and was undetectable by day 7 of the experiment. FISH showed that the *waaQ* mutant formed clumps in the cecal mucus layer of streptomycin-treated mice at 24 h post-feeding, leading to the conclusion that failure to penetrate mucus and grow as dispersed cells within the mucus layer prevented it from colonizing (19).

The sources of nutrients that support intestinal colonization by *E. coli* are shed epithelial cells, dietary fiber, and mucosal polysaccharides (12–14). Most of the amino acids are available in the cecum, as growth of *E. coli* in mucus results in repression of the majority of genes involved in amino acid biosynthesis (39). In rat, mouse, and human, colonic mucus is organized by Muc2, the major glycoprotein, which is a high molecular weight gel-forming glycoprotein containing L-fucose, D-galactose, D-mannose, N-acetyl-D-glucosamine, N-acetyl-D-galactosamine, and N-acetylneuraminic acid (61). Mucin is 80% polysaccharide and 20% protein and is highly viscous (62). In addition to mucin, the mucus layer contains a number of smaller glycoproteins, proteins, glycolipids, and lipids (61–64). There are two mucus layers, a loosely adherent suction-removable layer closest to the lumen of the intestine and an adherent layer firmly attached to the mucosa (61, 65, 66). In the rat colon, the thickness of the adherent layer is about 100 μm and that of the loose layer about 700 μm (61). In the mouse colon, the thickness of the adherent layer is about 50 μm and that of the loose layer about 100 μm (66). The mucus layer itself is in a dynamic state, constantly being synthesized and secreted by the mucin-secreting, specialized goblet cells and degraded to a large extent by the indigenous intestinal microbes (67, 68). Degraded mucus components are shed into the intestinal lumen forming a part of the luminal contents that is excreted in the feces (67).

The loosely adherent mucus layer contains large numbers of bacteria in the mouse, but the inner adherent mucus layer is largely devoid of bacteria (69) and is not penetrated by beads the size of bacteria (70), suggesting that the inner mucus layer protects the colonic epithelium from the commensal microbiota. Commensal *E. coli* strains do not attach to intestinal epithelial cells and growth takes place predominantly in the mucus layer (19, 24). The mucus layer of the conventional mouse large intestine turns over about every 2 hours (23). Hence, to maintain a stable population, the bacterial growth rate in mucus must keep pace with the turnover rate of the mucus layer. For example, *E. coli* BJ4 has a generation time of 40–80 minutes in the streptomycin-treated mouse cecum, which is more than fast enough to maintain its population (23).

E. coli cannot degrade oligosaccharides or polysaccharides, except dextrin (36, 71). In the intestine; this is the job of anaerobes. To obtain the mono- and di-saccharides it needs for growth *E. coli* relies on hydrolysis of complex polysaccharides by members of the intestinal community such as *Bacteroides thetaiotaomicron*, a Gram-negative obligate anaerobe and a major member of the human intestinal microbiota (72–77). This symbiotic relationship is illustrated in Figure 1. Complex polysaccharides derived from epithelial cell debris, dietary fiber, or mucus are degraded by extracellular polysaccharide hydrolases secreted by anaerobes. The anaerobes preferentially take up the resultant oligosaccharides, which are further degraded intracellularly to monosaccharides that enter central metabolism. The mono- and di-saccharides that are released by polysaccharide hydrolysis are discarded by the anaerobes and thereby made available to *E. coli* and organisms with similar metabolism.

Evidence for the model shown in Figure 1 is mounting. In response to competition from *Eubacterium rectale*, *B. thetaiotaomicron* upregulates a number of polysaccharide utilization loci that encode a variety of glycosyl hydrolases (78). In contrast, *E. rectale* responds to *B. thetaiotaomicron* by down regulating expression of loci encoding glycan degrading enzymes and up regulating expression of a number of sugar transporters, suggesting that *E. rectale* uses metabolic flexibility to take advantage of the superior ability of *B. thetaiotaomicron* to degrade polysaccharides (78). In a tour de force series of experiments from the Sonnenberg laboratory, it was recently shown in gnotobiotic mice associated with *B. thetaiotaomicron* and *Salmonella* Typhimurium that fucose and sialic acid catabolic gene systems are up regulated in the latter organism and its growth is stimulated (76). Importantly, *Salmonella* mutants lacking the capacity to catabolize either fucose or sialic acid showed substantial fitness defects in streptomycin-treated mice (76). A *B. thetaiotaomicron* mutant lacking a predicted surface-associated sialidase failed to release free sialic acid in mice, whereas the wild type *B. thetaiotaomicron* released free sialic acid and stimulated intestinal growth of *Clostridium difficile* (76). Free sialic acid was very low in conventional mice and greatly elevated in mice that were treated with streptomycin, which is consistent with the idea that the microbiota consumed sialic acid in the conventional animals, but streptomycin treatment removed those members of the microbiota (76). These experiments prove that polysaccharide hydrolysis by anaerobes in the intestine can provide monosaccharides to members of the microbiota that can use them.

Nutrient limitation in the intestine

Competition for limiting resources drives ecosystems (79). In microbial ecosystems, such as the intestine, the microbial community competes for carbon and energy sources and terminal electron acceptors (27). It appears that *E. coli* uses strategies for maximizing its population in a stiffly competitive environment. *E. coli* can simultaneously utilize a mixture of six sugars under nutrient-limiting conditions in chemostats (80). When presented with a mixture of thirteen sugars in vitro, *E. coli* uses up to nine of them at a time (36). Bacteria expand their transcriptome to induce a number of gene systems for carbon source transport and catabolism when growing slowly (81) or when they are nutrient-deprived (82). While there is a fine line between “hunger” and “starvation”, in fact, the two are distinct. Hunger is a state that is defined by physiological and genetic changes that expand the metabolic capacity of the cell (83). To better understand the role of the hunger lifestyle in vivo, the importance

of carbon stores for colonization fitness was examined (59). It is well known that stored glycogen promotes survival during times of carbon limitation. It was found that mutants of commensal and pathogenic *E. coli* that cannot synthesize or degrade glycogen have significant intestinal colonization defects in mice (59). Furthermore, in support of the hypothesis that *E. coli* uses glycogen to withstand hunger, when a constant supply of a readily metabolized carbon source was supplied in the drinking water of the mice, these colonization defects were rescued (59).

E. coli Nissle 1917 is a non-pathogenic strain that has been used since the early 1920's as a probiotic agent to treat gastrointestinal infections in humans, sold under the name "Mutaflor®" (84). Despite having the *E. coli* Nissle 1917 genome sequence to search for clues as to its success as a probiotic agent (85, 86), little attention has been paid to the possibility that it may out-compete pathogens for essential nutrients. When *E. coli* Nissle 1917 is the only *E. coli* strain in the streptomycin-treated mouse intestine, it appears to use arabinose, fucose, galactose, gluconate, mannose, N-acetylgalactosamine, and sialic acid to colonize (Table 2 and (58)). In contrast, *E. coli* Nissle 1917 does not appear to use ribose to colonize (Table 2). Identifying the nutrients used by *E. coli* Nissle 1917 when competing with *E. coli* pathotypes might lead to new approaches to prevent *E. coli* infections.

Metabolic flexibility is also exhibited by uropathogenic *E. coli* (UPEC) strain CFT073, a human urinary tract pathogen, when its nutritional program in the mouse intestine is compared to that in the mouse urinary tract. *E. coli* CFT073 utilizes several sugars simultaneously to colonize the intestine (Table 2 and (35)), but not to infect the urinary tract. Instead, in the urinary tract, transport of peptides and gluconeogenesis are required for maximum *E. coli* CFT073 growth (87). When *E. coli* EDL933 is the only *E. coli* strain in the mouse intestine, it does not use gluconeogenic substrates to colonize, but it switches to gluconeogenic nutrients when competing in the intestine with *E. coli* MG1655 (54). As an additional example of metabolic flexibility in vivo, neither *E. coli* MG1655 nor *E. coli* Nissle 1917 use ribose for growth in the intestine, unless fucose-1-phosphate accumulates in mutants unable to catabolize it further (58). Moreover, it was shown that fucose at a concentration too low to support growth, stimulated the utilization of ribose by the wild-type *E. coli* strains in vitro, suggesting that fucose-1-phosphate plays a role in regulating the use of ribose as a carbon source by *E. coli* MG1655 and *E. coli* Nissle 1917 in the mouse intestine (58). In summary, to colonize successfully, *E. coli* must compete for limiting nutrients, so it uses several sugars at a time, is flexible in its nutrient preference, and relies on glycogen carbon stores in the intestine.

Competition for nutrients in the intestine

E. coli competes for nutrients in the intestine in (at least) three ways. First, it can use nutrients that are available because no other community member has used it (Table 2). Second, it can outcompete other strains for the nutrients it prefers by growing faster on them (88). Third, it can enter into a symbiotic association with the anaerobe(s) that releases its preferred sugar(s) (60, 89). Evidence for each of these mechanisms comes from competitive fitness studies in streptomycin-treated mice. Streptomycin treatment (5g/l in drinking water) perturbs the microbiota by selectively removing the facultative anaerobes with which *E. coli*

competes, making nutrients available and allowing experimentally introduced strains to overcome colonization resistance (27, 90, 91). Colonization resistance in animal models can also be overcome in other ways, e.g., by gnotobiotic mice in which there is no native microbiota (92), by mice that have a contrived microbiota (93) or by mice with intestinal inflammation, which generates nitrate that is respired by *E. coli* (94).

The three ways that *E. coli* can compete with the microbiota for nutrients are described in detail here. First, different *E. coli* strains use different nutrients in vivo, despite using the same nutrients in the same order in vitro (Table 2). Mutants with deletions in genes corresponding to metabolic pathways induced in mucus were constructed by allelic replacement (29) and were tested for their ability to compete with their wild type parent strain when simultaneously fed to mice in low numbers (10^5 CFU/mouse). The data obtained from these studies showed that the human commensals *E. coli* HS, *E. coli* Nissle 1917, *E. coli* MG1655, and the pathogen *E. coli* EDL933 each occupy unique nutritional niches in the mouse intestine (Table 2 and (35, 36)). Of the 12 sugars available in the mucus layer, *E. coli* HS utilizes 6 for colonization: arabinose, galactose, gluconate, N-acetylglucosamine, lactose, and ribose (Table 2). *E. coli* Nissle 1917 uses a different list of 7 carbon sources to support colonization, including arabinose, fucose, galactose, gluconate, N-acetylglucosamine, and N-acetylneuraminate, and mannose (Table 2) and *E. coli* MG1655 utilizes 5 sugars for colonization: arabinose, fucose, gluconate, N-acetylglucosamine, and N-acetylneuraminate (Table 2). Each of these commensals is capable of colonizing mice that are pre-colonized with any one of the others (31), and each strain is capable of utilizing at least one sugar not used by the others in vivo, which suggests that differences in their in vivo sugar preferences allows them to occupy distinct nutrient-defined niches in the intestine. Furthermore, the intestinal niche occupied by pathogenic *E. coli* EDL933 is also unique and is defined by utilization of 7 sugars: arabinose, galactose, hexuronates, mannose, N-acetylglucosamine, ribose, and sucrose (Table 2).

That *E. coli* MG1655 and *E. coli* EDL933 display different nutritional programs in the mouse intestine, e.g. *E. coli* MG1655 uses N-acetylneuraminate but not mannose whereas *E. coli* EDL933 uses mannose but not N-acetylneuraminate (Table 2), was surprising in view of the fact that *E. coli* MG1655 and *E. coli* EDL933 utilize them equally well in vitro and display identical nutritional preferences in vitro, i.e., they use sugars in the same order in vitro as follows: N-acetylglucosamine, gluconate, ribose, sialic acid, mannose, arabinose, maltose, and fucose (36). These findings will be discussed in the context of the “Restaurant” hypothesis below.

Since different commensal *E. coli* strains use different sugars to colonize the intestine, it seemed reasonable that a potential strategy for preventing colonization by the enterohemorrhagic *E. coli* EDL933 would be to pre-colonize mice with a combination of commensal strains that would fill the sugar-defined nutritional niches normally available to the invading pathogen. When mice were pre-colonized for 10 days with either the commensal *E. coli* MG1655, *E. coli* HS, or *E. coli* Nissle 1917 and then fed 10^5 CFU of the pathogenic *E. coli* EDL933, *E. coli* Nissle 1917 limited growth of *E. coli* EDL933 in the intestine (10^3 to 10^4 CFU/gram of feces), whereas *E. coli* MG1655 and *E. coli* HS allowed growth to higher numbers (10^6 to 10^7 CFU/gram of feces). However, when *E. coli* EDL933

was fed to mice previously pre-colonized with the three commensal *E. coli* strains (MG1655, HS, and Nissle 1917), each of which displays a different nutritional program in vivo (Table 2), *E. coli* EDL933 was eliminated from the intestine (31). Therefore, a combination of as few as 3 commensal *E. coli* strains provided a barrier to *E. coli* EDL933 infection.

If the basis for exclusion of *E. coli* EDL933 was because the 3 commensal *E. coli* strains utilize the nutrients needed by *E. coli* EDL933 to compete and colonize, then any *E. coli* commensal strain or combination of strains that effectively catabolizes the sugars used by *E. coli* EDL933 would prevent its colonization. Indeed, when the ability of *E. coli* EDL933 to colonize mice that were pre-colonized with *E. coli* HS and *E. coli* Nissle 1917 was tested, which the data in Table 2 indicate should be equally effective without *E. coli* MG1655 present, *E. coli* EDL933, was indeed eliminated 5 days following association (35). However, the same 2 commensal *E. coli* strains could not prevent colonization of *E. coli* CFT073, a uropathogenic strain, and *E. coli* E2348/69, an enteropathogenic strain (95). Therefore, it is unlikely that any particular commensal strain(s) of *E. coli* will be generally effective as a probiotic to prevent colonization by enteric pathogens. Nevertheless, the data support the hypothesis that nutrient consumption by commensal *E. coli* can limit nutrient availability to pathogens, which in turn points to the potential of probiotics for preventing disease.

A second way of competing for nutrients in the intestine is illustrated by what happens when *E. coli* MG1655 adapts to the mouse intestine. When mice were fed *E. coli* MG1655, non-motile *flhDC* deletion mutants appeared in the feces 3 days post-feeding and reached approximately 90% of the total population by day 15 of the experiment (96). These mutants had a striking colonization advantage over the wild type *E. coli* MG1655 parent strain. The deletions were of varying length and began immediately downstream of the *IS1* element in the *flhDC* promoter region. One such mutant, designated MG1655*, was a better colonizer than its parent, grew in cecal mucus faster than its parent in vitro (90 ± 2 min generation time vs. 105 ± 2 min, $P < 0.001$), and grew 15–30% faster than its parent on a number of sugars present in the mouse intestine (97). The *E. coli flhDC* operon encodes the FlhD₄C₂ regulatory complex, which is the master positive regulator of the more than 40 gene flagella regulon (98). The FlhD₄C₂ complex has also been reported to negatively regulate *E. coli* K-12 genes involved in galactose transport, the Entner-Doudoroff pathway, and the TCA cycle and positively regulate genes involved in ribose transport (88, 99). *E. coli* MG1655 *flhDC* deletion mutants have also been reported to be selected in the intestines of ex-germfree mono-associated mice (100).

Several high-throughput genomic approaches were taken to further characterize *E. coli* MG1655*. Whole-genome pyrosequencing did not reveal any changes on its genome, aside from the deletion at the *flhDC* locus, that could explain the colonization advantage of *E. coli* MG1655* (88). Microarray analysis revealed modest yet significant induction of catabolic gene systems across the genome in both *E. coli* MG1655* and an isogenic *flhD* mutant constructed in the laboratory (88). Catabolome analysis with Biolog GN2 microplates revealed an enhanced ability of both *E. coli* MG1655* and the isogenic *flhD* mutant to oxidize a variety of carbon sources (88). Collectively, the results showed that intestine-adapted *E. coli* MG1655* is more fit than the wild type for intestinal colonization, because

loss of FlhD results in elevated expression of genes involved in carbon and energy metabolism, allowing the mutants to outcompete their wild type parent for the same nutrients. Hence a second strategy for gaining a colonization advantage is to outcompete other members of the microbiota for their preferred nutrients.

There is a third way that *E. coli* competes for nutrients in the intestine. The intestine selects for mutants that gain a colonization advantage by promoting occupation of a distinct niche. The selection of non-motile *E. coli* MG1655 *flhDC* mutants by the streptomycin-treated mouse intestine is easily explained by the nutrient-niche hypothesis, i.e., the mutants grow 15% faster *in vitro* in mouse cecal mucus and 15%–30% faster on several sugars present in cecal mucus than *E. coli* MG1655 (96). In addition to *flhDC* mutants, *E. coli* MG1655 mutants with reduced motility also were selected by adaptation in the streptomycin-treated mouse intestine and these turned out to be *E. coli* MG1655 *envZ* missense mutants (60). *E. coli* MG1655 *envZ* missense mutants have also been reported to be selected in the intestines of ex-germ free mono-associated mice (101). The *envZ* gene encodes a histidine kinase that is a member of the *envZ/ompR* two-component signal transduction system that modulates gene expression in response to osmolarity. The genes regulated include *flhDC*, the porin genes *ompC* and *ompF*, and several other genes encoding outer membrane proteins (102, 103). These *E. coli* MG1655 *envZ* missense mutants produced more phosphorylated OmpR than both *E. coli* MG1655 and the *E. coli* MG1655 *flhDC* deletion mutants and produced more of the outer membrane porin OmpC and less of the outer membrane porin OmpF (89). As a result, the *E. coli* MG1655 *envZ* missense mutants were more resistant to bile salts and colicin V than *E. coli* MG1655 and the *E. coli* MG1655 *flhDC* deletion mutants (60). One of the *E. coli* MG1655 *envZ* missense mutants, which was studied further, contained the *envZ*_{P41L} missense mutation and grew about 15% slower *in vitro* in mouse cecal mucus and on several sugars present in mucus compared to the *flhDC* deletion mutants, yet was as good an intestinal colonizer as the *flhDC* deletion mutants and far better than *E. coli* MG1655 (60). Moreover, *E. coli* MG1655 *envZ*_{P41L} and the *E. coli* MG1655 *flhDC* deletion mutants appeared to colonize equally well in one major intestinal niche, but *E. coli* MG1655 *envZ*_{P41L} appeared to use galactose to colonize a second intestinal niche either not colonized or colonized poorly by the *E. coli* MG1655 *flhDC* deletion mutants. These data are not consistent with the nutrient-niche hypothesis but they are consistent with what we call the “Restaurant” hypothesis, which will be discussed below.

Since *E. coli* MG1655 *envZ*_{P41L} was a far better intestinal colonizer than wild type *E. coli* MG1655, the *envZ*_{P41L} gene was transferred for further study into *E. coli* Nissle 1917, the human probiotic strain used to treat gastrointestinal infections. Like *E. coli* MG1655 *envZ*_{P41L}, *E. coli* Nissle 1917 *envZ*_{P41L} produced more phosphorylated OmpR than its parent and produced more of the outer membrane porin OmpC and less of the outer membrane porin OmpF (89). It also became more resistant to bile salts and colicin V, grew 50% slower *in vitro* in mucus and 15%–30% slower on several sugars present in mucus, yet was a 10-fold better colonizer than *E. coli* Nissle 1917 (89). Furthermore, like *E. coli* MG1655 *envZ*_{P41L}, *E. coli* Nissle 1917 *envZ*_{P41L} appeared to use galactose to colonize a second intestinal niche either not colonized or colonized poorly by wild type *E. coli* Nissle 1917, despite not growing as well as *E. coli* Nissle 1917 on galactose as a sole carbon source (89).

Moreover, despite being a better colonizer, *E. coli* Nissle 1917 *envZ*_{P41L} was not better than its parent at preventing colonization by enterohemorrhagic *E. coli* EDL933 and, in fact, appeared to be worse (89). The data can be explained according to our “Restaurant” hypothesis for commensal *E. coli* strains, i.e., that they colonize the intestine as sessile members of mixed biofilms obtaining the sugars they need for growth locally, but compete for sugars with invading *E. coli* pathogens planktonically as described below.

Biofilms in the intestine

Much attention has been given to the role played by biofilms in bacterial colonization of many environments, but until recently surprisingly little information was available regarding biofilms in the intestine. On the one hand, the transit time of intestinal contents is short compared to the timescale of biofilm development (104), so it was hard to imagine how a stable biofilm might be maintained in the intestine, yet the kinetics of plasmid transfer between *E. coli* strains in the streptomycin-treated mouse intestine suggested that *E. coli* resides in biofilms in vivo (105). Moreover, the mucus layer itself has many of the characteristics of a secreted biofilm matrix. Add to this the concept of bacterial binding to mucus components and it became reasonable to consider the possibility of bacterial biofilms in the GI tract. Indeed, it's been shown that biofilms form in the mucus layers of the large intestines of healthy humans, rats, baboons and mice (106–109) and that mixed biofilms consisting of *Bacteroides*, *Enterobacter*, and *Clostridia* species form rapidly on strands of mucin in mucus introduced into a growing human microbiota contained in a continuous flow culture system constructed to mimic the human intestine (110). Furthermore, it appears that human colonic mucosal biofilms and bacterial communities in feces differ greatly in composition (111) and dysbiosis in the community structure of mucosal biofilms may play an important role in contributing to chronic inflammatory bowel diseases such as ulcerative colitis and Crohn's disease (106, 107). In *Vibrio cholerae*, biofilm formation is important for pathogenesis (112). Intestinal biofilms would provide the habitat for microbe-microbe interactions such as those that are thought to occur between *E. coli* and the polysaccharide degrading anaerobes.

The “Restaurant” hypothesis

As discussed above, commensal strains of *E. coli* appear to reside in mixed biofilms in the large intestines of mice (60, 89). Moreover, commensal and pathogenic strains of *E. coli* use mono- and disaccharides for growth in the intestine (36). However, these sugars are absorbed in the small intestine, whereas dietary fiber reaches the large intestine intact. In contrast to the anaerobes, most *E. coli* strains do not secrete extracellular polysaccharide hydrolases (71, 113) and therefore cannot degrade dietary fiber-derived and mucin-derived oligo- and polysaccharides. Since commensal and pathogenic *E. coli* strains colonize the mouse large intestine by growing in intestinal mucus (43, 114–118), it appears likely that *E. coli* depends on the anaerobes present in mucus that can degrade oligo- and polysaccharides to provide them with the mono- and disaccharides and maltodextrins they need for growth. Indeed, *Salmonella enterica* serovar Typhimurium, which is in the same family as *E. coli* and has very similar metabolism, catabolizes fucose and sialic acid liberated from mucosal polysaccharides by *Bacteroides thetaiotaomicron* (76). It is therefore possible that anaerobes

in the mixed biofilms provide *E. coli* with the sugars it needs for growth locally, rather than from a perfectly mixed pool available to all species, which is an assumption of the nutrient-niche hypothesis. We call the mixed biofilms that feed the *E. coli* strains “Restaurants” and we hypothesize that different commensal *E. coli* strains reside in different “Restaurants” interacting physically and metabolically with different anaerobes. Each restaurant might serve different nutrients, i.e., each commensal *E. coli* strain could be exposed to a different menu, which explains why different *E. coli* strains display different nutritional programs in the mouse intestine despite displaying identical nutritional programs in vitro (35, 36). The restaurant hypothesis can also explain how *E. coli* strains that grow more slowly in mucus and on several sugars found in mucus are better colonizers than their parents as long as they have a higher affinity for biofilm binding sites than their parents. Indeed, the outer membranes of both *E. coli* MG1655 *envZ*_{P41L} and *E. coli* Nissle 1917 *envZ*_{P41L} are very different from those of their parents (60, 89), which could result in increased affinities for mixed biofilms.

The “Restaurant” hypothesis can also explain why despite being a better colonizer than *E. coli* Nissle 1917, *E. coli* Nissle 1917 *envZ*_{P41L} is not better at limiting enterohemorrhagic *E. coli* EDL933 colonization than *E. coli* Nissle 1917 and, in fact, may be worse (89). It is possible when *E. coli* EDL933 invades the mouse intestine it initially grows planktonically in mucus and not in mixed biofilms. If we are correct that *E. coli* Nissle1917 and *E. coli* Nissle1917 *envZ*_{P41L} colonize the mouse intestine by being served specific sugars by the anaerobes in the mixed biofilms they inhabit, then small amounts of these sugars that escape the mixed biofilms might be available to invading *E. coli* EDL933 as well as to the small numbers of planktonic *E. coli* Nissle1917 or *E. coli* Nissle1917 *envZ*_{P41L} that leave the mixed biofilms. Therefore, it may be that both planktonic *E. coli* Nissle 1917 *envZ*_{P41L} and *E. coli* Nissle 1917 compete directly with planktonic *E. coli* EDL933 for the sugars that escape the biofilms or that are produced by small numbers of planktonic members of the microbiota that leave the biofilms. This scenario would allow planktonic *E. coli* EDL933 to grow to the extent allowed by the available concentrations of those sugars in competition with planktonic *E. coli* Nissle 1917 *envZ*_{P41L} or *E. coli* Nissle 1917, which could explain why *E. coli* Nissle 1917, the faster grower in perfectly mixed bacteria-free mucus in vitro, appears to limit *E. coli* EDL933 growth in the intestine to a greater extent than does *E. coli* Nissle 1917 *envZ*_{P41L} (89).

We stress that the granularity of mixed intestinal biofilms and nutrient flow between the microbes that reside within them is not known. These interactions could be so finite as to allow two different *E. coli* strains to interact with the same anaerobe cell and each grow on a different preferred nutrient. If so, it will not be possible to find zones within the biofilm (i.e., restaurants) that contain only a single population of *E. coli* and interacting anaerobe partner.

Conclusions

It is becoming increasingly clear that once *E. coli* strains reach the large intestine, in order to colonize, they must enter the mucus layer and utilize nutrients there for growth. It is also clear that different strains of *E. coli* display different nutritional programs in the intestine. However, it is not known whether a specific *E. coli* commensal strain utilizes the same

nutrients when it is the only *E. coli* strain in the intestine compared to a situation in which it colonizes along with several different commensal *E. coli* strains with which it must compete for nutrients. Metabolic flexibility could be a key requirement for successful colonization of the intestine by several *E. coli* strains simultaneously. However, the “Restaurant” hypothesis explains long term colonization by several established commensal *E. coli* strains without invoking metabolic flexibility, i.e., each commensal *E. coli* strain resides as a sessile member of a mixed biofilm in the intestine and obtains nutrients locally rather than from a perfectly mixed pot of nutrients. However, as described above, it seems likely that when mice colonized long-term with one *E. coli* strain are fed a pathogenic *E. coli* strain, planktonic members of the pre-colonized strain that escape the mixed biofilm compete directly with the invading pathogen for nutrients from the same perfectly mixed pot, according to the Freter nutrient-niche hypothesis. Since it appears likely that a pathogen must be able to grow in the intestine in order to initiate the pathogenic process, we hope that future research will provide a nutritional framework for colonizing humans with a combination of commensal *E. coli* strains or with one commensal *E. coli* strain that has been engineered to be as effective as several strains and can serve as an effective first line of defense against pathogenic *E. coli* intestinal infections.

Acknowledgements

The work carried out in the authors' laboratories was supported by U. S. Public Health Service grants AI48945 and GM095370. The authors wish to acknowledge the contributions of the following people to work in the authors' laboratories. In Rhode Island: Jimmy Adediran, James Allen, Steven Autieri, Swati Banerjee, Megan Banner, Eric Gauger, Jakob Frimodt-Møller, Mathias Jorgensen, Mary Leatham-Jensen, Regino Mercado-Lubo, Regina Miranda, Matthew Mokszycki, Annette Møller, and Silvia Schinner. In Oklahoma: April Anderson, Amanda Ashby, Matthew Caldwell, Amanda Cernosek, Dong-Eun Chang, Fatema Chowdhury, Andrew Fabich, Terri Gibson, Shari Jones, Amanda Laughlin, Rosalie Maltby, Jessica Meador, Darren Smalley, Stephanie Tison, and Don Tucker.

References

1. Finegold, SM.; Sutter, VL.; Mathisen, GE. Normal indigenous intestinal microflora. In: Hentges, DJ., editor. Human intestinal microflora in health and disease. New York, N.Y.: Academic Press, Inc.; 1983. p. 3-31.
2. Huggins C, Rast HV Jr. Incidence of coliform bacteria in the intestinal tract of *Gambusia affinis holbrooki* (Girard) and in their habitat water. J Bacteriol. 1963; 85:489–490. [PubMed: 13955523]
3. Palmer C, Bik EM, Digiulio DB, Relman DA, Brown PO. Development of the human infant intestinal microbiota. PLoS Biol. 2007; 5:e177. [PubMed: 17594176]
4. Riley M, Abe T, Arnaud MB, Berlyn MK, Blattner FR, Chaudhuri RR, Glasner JD, Horiuchi T, Keseler IM, Kosuge T, Mori H, Perna NT, Plunkett G 3rd, Rudd KE, Serres MH, Thomas GH, Thomson NR, Wishart D, Wanner BL. *Escherichia coli* K-12: a cooperatively developed annotation snapshot--2005. Nucleic Acids Res. 2006; 34:1–9. [PubMed: 16397293]
5. Stecher B, Berry D, Loy A. Colonization resistance and microbial ecophysiology: using gnotobiotic mouse models and single-cell technology to explore the intestinal jungle. FEMS Microbiol Rev. 2013; 37:793–829. [PubMed: 23662775]
6. Stecher B, Hardt WD. Mechanisms controlling pathogen colonization of the gut. Curr Opin Microbiol. 2011; 14:82–91. [PubMed: 21036098]
7. Clarke MB, Sperandio V. Events at the host-microbial interface of the gastrointestinal tract III. Cell-to-cell signaling among microbial flora, host, and pathogens: there is a whole lot of talking going on. Am J Physiol Gastrointest Liver Physiol. 2005; 288:G1105–G1109. [PubMed: 15890712]

8. Cole, AM.; Ganz, T. Defensins and other antimicrobial peptides: Innate defense of mucosal surfaces. In: Nataro, JP.; Cohen, PS.; Mobley, HLT.; Weiser, JN., editors. *Colonization of Mucosal Surfaces*. Washington, D. C.: ASM Press; 2005. p. 17-34.
9. Kaper JB, Sperandio V. Bacterial cell-to-cell signaling in the gastrointestinal tract. *Infect Immun*. 2005; 73:3197–3209. [PubMed: 15908344]
10. Pasetti, MF.; Salerno-Goncalves, R.; Sztein, MB. Mechanisms of adaptive immunity that prevent colonization of mucosal surfaces. In: Nataro, JP.; Cohen, PS.; Mobley, HLT.; Weiser, JN., editors. *Colonization of Mucosal Surfaces*. Washington, D. C.: ASM Press; 2005. p. 35-47.
11. Sansonetti PJ. War and peace at mucosal surfaces. *Nat Rev Immunol*. 2004; 4:953–964. [PubMed: 15573130]
12. Conway, T.; Krogfelt, KA.; Cohen, PS. Chapter 8.3.1.2, The Life of Commensal *Escherichia coli* in the Mammalian Intestine. In: Böck, A.; Curtiss, R., III; Kaper, JB.; Karp, PD.; Neidhardt, FC.; Nyström, T.; Slauch, JM.; Squires, CL.; Usery, D., editors. *EcoSal--Escherichia coli and Salmonella: Cellular and Molecular Biology*, 3rd. Online ed.. Washington, D.C.: ASM Press; 2004.
13. Conway, T.; Krogfelt, KA.; Cohen, PS. *Escherichia coli* at the intestinal mucosal surface. In: Brogden, KA.; Minion, FC.; Cornick, N.; Stanton, TB.; Zhang, Q.; Nolan, LK.; Wannemuehler, MJ., editors. *Virulence mechanisms of bacterial pathogens*. 4 ed.. Washington, D.C.: ASM Press; 2007. p. 175-196.
14. Laux, DC.; Cohen, PS.; Conway, T. Role of the mucus layer in bacterial colonization of the intestine. In: Nataro, JP.; Mobley, HLT.; Cohen, PS., editors. *Colonization of Mucosal Surfaces*. Washington, DC: ASM Press; 2005. p. 199-212.
15. Wadolkowski EA, Laux DC, Cohen PS. Colonization of the streptomycin-treated mouse large intestine by a human fecal *Escherichia coli* strain: role of growth in mucus. *Infect Immun*. 1988; 56:1030–1035. [PubMed: 3281898]
16. Foster JW. *Escherichia coli* acid resistance: tales of an amateur acidophile. *Nat Rev Microbiol*. 2004; 2:898–907. [PubMed: 15494746]
17. Lin J, Smith MP, Chapin KC, Baik HS, Bennett GN, Foster JW. Mechanisms of acid resistance in enterohemorrhagic *Escherichia coli* . *Appl Environ Microbiol*. 1996; 62:3094–3100. [PubMed: 8795195]
18. Freter R, Brickner H, Fekete J, Vickerman MM, Carey KE. Survival and implantation of *Escherichia coli* in the intestinal tract. *Infect Immun*. 1983; 39:686–703. [PubMed: 6339389]
19. Moller AK, Leatham MP, Conway T, Nuijten PJ, de Haan LA, Krogfelt KA, Cohen PS. An *Escherichia coli* MG1655 lipopolysaccharide deep-rough core mutant grows and survives in mouse cecal mucus but fails to colonize the mouse large intestine. *Infect Immun*. 2003; 71:2142–2152. [PubMed: 12654836]
20. McCormick BA, Laux DC, Cohen PS. Neither motility nor chemotaxis plays a role in the ability of *Escherichia coli* F-18 to colonize the streptomycin-treated mouse large intestine. *Infect Immun*. 1990; 58:2957–2961. [PubMed: 2201640]
21. McGuckin MA, Linden SK, Sutton P, Florin TH. Mucin dynamics and enteric pathogens. *Nat Rev Microbiol*. 2011; 9:265–278. [PubMed: 21407243]
22. Bergstrom KS, Sham HP, Zarepour M, Vallance BA. Innate host responses to enteric bacterial pathogens: a balancing act between resistance and tolerance. *Cell Microbiol*. 2012; 14:475–484. [PubMed: 22233392]
23. Rang CU, Licht TR, Midtvedt T, Conway PL, Chao L, Krogfelt KA, Cohen PS, Molin S. Estimation of growth rates of *Escherichia coli* BJ4 in streptomycin-treated and previously germfree mice by in situ rRNA hybridization. *Clin Diagn Lab Immunol*. 1999; 6:434–436. [PubMed: 10225851]
24. Poulsen LK, Lan F, Kristensen CS, Hobolth P, Molin S, Krogfelt KA. Spatial distribution of *Escherichia coli* in the mouse large intestine inferred from rRNA in situ hybridization. *Infect Immun*. 1994; 62:5191–5194. [PubMed: 7927805]
25. Poulsen LK, Licht TR, Rang C, Krogfelt KA, Molin S. Physiological state of *Escherichia coli* BJ4 growing in the large intestines of streptomycin-treated mice. *J Bacteriol*. 1995; 177:5840–5845. [PubMed: 7592332]

26. Freter, R. Factors affecting the microecology of the gut. In: Fuller, R., editor. Probiotics, The scientific basis. London: Chapman and Hall; 1992. p. 111-144.
27. Freter, R. Mechanisms that control the microflora in the large intestine. In: Hentges, DJ., editor. Human intestinal microflora in health and disease. New York, N.Y.: Academic Press, Inc.; 1983. p. 33-54.
28. Freter, R. Mechanisms of bacterial colonization of the mucosal surfaces of the gut. In: Roth, JA., editor. Virulence mechanisms of bacterial pathogens. Washington, D.C.: American Society for Microbiology; 1988. p. 45-60.
29. van der Waaij D, Berghuis-de Vries JM, Lekkerkerk L-v. Colonization resistance of the digestive tract in conventional and antibiotic-treated mice. J Hyg (Lond). 1971; 69:405-411. [PubMed: 4999450]
30. Apperloo-Renkema HZ, Van der Waaij BD, Van der Waaij D. Determination of colonization resistance of the digestive tract by biotyping of Enterobacteriaceae. Epidemiol. Infect. 1990; 105:355-361. [PubMed: 2209739]
31. Leatham MP, Banerjee S, Autieri SM, Mercado-Lubo R, Conway T, Cohen PS. Precolonized human commensal *Escherichia coli* strains serve as a barrier to *E. coli* O157:H7 growth in the streptomycin-treated mouse intestine. Infect. Immun. 2009; 77:2876-2886. [PubMed: 19364832]
32. Stecher B, Barthel M, Schlumberger MC, Haberli L, Rabsch W, Kremer M, Hardt WD. Motility allows *S. Typhimurium* to benefit from the mucosal defence. Cell Microbiol. 2008; 10:1166-1180. [PubMed: 18241212]
33. Stecher B, Hapfelmeier S, Muller C, Kremer M, Stallmach T, Hardt WD. Flagella and chemotaxis are required for efficient induction of *Salmonella enterica* serovar Typhimurium colitis in streptomycin-pretreated mice. Infect Immun. 2004; 72:4138-4150. [PubMed: 15213159]
34. Stecher B, Robbiani R, Walker AW, Westendorf AM, Barthel M, Kremer M, Chaffron S, Macpherson AJ, Buer J, Parkhill J, Dougan G, von Mering C, Hardt WD. *Salmonella enterica* serovar typhimurium exploits inflammation to compete with the intestinal microbiota. PLoS Biol. 2007; 5:2177-2189. [PubMed: 17760501]
35. Maltby R, Leatham-Jensen MP, Gibson T, Cohen PS, Conway T. Nutritional basis for colonization resistance by human commensal *Escherichia coli* strains HS and Nissle 1917 against *E. coli* O157:H7 in the mouse intestine. PLoS One. 2013; 8:e53957. [PubMed: 23349773]
36. Fabich AJ, Jones SA, Chowdhury FZ, Cernosek A, Anderson A, Smalley D, McHargue JW, Hightower GA, Smith JT, Autieri SM, Leatham MP, Lins JJ, Allen RL, Laux DC, Cohen PS, Conway T. Comparison of carbon nutrition for pathogenic and commensal *Escherichia coli* strains in the mouse intestine. Infect Immun. 2008; 76:1143-1152. [PubMed: 18180286]
37. Stecher B, Hardt WD. The role of microbiota in infectious disease. Trends Microbiol. 2008; 16:107-114. [PubMed: 18280160]
38. Cook H, Ussery DW. Sigma factors in a thousand *E. coli* genomes. Environ Microbiol. 2013; 15:3121-3129. [PubMed: 23992563]
39. Chang DE, Smalley DJ, Tucker DL, Leatham MP, Norris WE, Stevenson SJ, Anderson AB, Grissom JE, Laux DC, Cohen PS, Conway T. Carbon nutrition of *Escherichia coli* in the mouse intestine. Proc. Natl. Acad. Sci. U. S. A. 2004; 101:7427-7432. [PubMed: 15123798]
40. Bowden SD, Rowley G, Hinton JC, Thompson A. Glucose and glycolysis are required for the successful infection of macrophages and mice by *Salmonella enterica* serovar typhimurium. Infect Immun. 2009; 77:3117-3126. [PubMed: 19380470]
41. Njoroge JW, Nguyen Y, Curtis MM, Moreira CG, Sperandio V. Virulence meets metabolism: Cra and KdpE gene regulation in enterohemorrhagic *Escherichia coli*. MBio. 2012; 3:e00280-e00212. [PubMed: 23073764]
42. Waligora EA, Fisher CR, Hanovice NJ, Rodou A, Wyckoff EE, Payne SM. Role of Intracellular Carbon Metabolism Pathways in *Shigella flexneri* Virulence. Infect Immun. 2014; 82:2746-2755. [PubMed: 24733092]
43. Sweeney NJ, Laux DC, Cohen PS. *Escherichia coli* F-18 and *E. coli* K-12 *eda* mutants do not colonize the streptomycin-treated mouse large intestine. Infect. Immun. 1996; 64:3504-3511. [PubMed: 8751891]

44. Peekhaus N, Conway T. What's for dinner?: Entner-Doudoroff metabolism in *Escherichia coli* . J Bacteriol. 1998; 180:3495–3502. [PubMed: 9657988]
45. Conway T. The Entner-Doudoroff pathway: history, physiology and molecular biology. FEMS Microbiol Rev. 1992; 9:1–27. [PubMed: 1389313]
46. Eriksson S, Lucchini S, Thompson A, Rhen M, Hinton JC. Unravelling the biology of macrophage infection by gene expression profiling of intracellular *Salmonella enterica* . Mol Microbiol. 2003; 47:103–118. [PubMed: 12492857]
47. Patra T, Koley H, Ramamurthy T, Ghose AC, Nandy RK. The Entner-Doudoroff pathway is obligatory for gluconate utilization and contributes to the pathogenicity of *Vibrio cholerae* . J Bacteriol. 2012; 194:3377–3385. [PubMed: 22544275]
48. Zhao J, Baba T, Mori H, Shimizu K. Global metabolic response of *Escherichia coli* to *gnd* or *zwf* gene-knockout, based on ¹³C-labeling experiments and the measurement of enzyme activities. Appl Microbiol Biotechnol. 2004; 64:91–98. [PubMed: 14661115]
49. Steinsiek S, Frixel S, Stagge S, Sumo, Bettenbrock K. Characterization of *E. coli* MG1655 and *frdA* and *sdhC* mutants at various aerobiosis levels. J Biotechnol. 2011; 154:35–45. [PubMed: 21458504]
50. Tchawa Yimga M, Leatham MP, Allen JH, Laux DC, Conway T, Cohen PS. Role of gluconeogenesis and the tricarboxylic acid cycle in the virulence of *Salmonella enterica* serovar Typhimurium in BALB/c mice. Infect Immun. 2006; 74:1130–1140. [PubMed: 16428761]
51. Fang FC, Libby SJ, Castor ME, Fung AM. Isocitrate lyase (AceA) is required for *Salmonella* persistence but not for acute lethal infection in mice. Infect Immun. 2005; 73:2547–2549. [PubMed: 15784602]
52. Spector MP, DiRusso CC, Pallen MJ, Garcia del Portillo F, Dougan G, Finlay BB. The medium-/long-chain fatty acyl-CoA dehydrogenase (*fadF*) gene of *Salmonella typhimurium* is a phase 1 starvation-stress response (SSR) locus. Microbiology. 1999; 145(Pt 1):15–31. [PubMed: 10206693]
53. Dahal N, Abdelhamed H, Lu J, Karsi A, Lawrence ML. Tricarboxylic acid cycle and one-carbon metabolism pathways are important in *Edwardsiella ictaluri* virulence. PLoS One. 2013; 8:e65973. [PubMed: 23762452]
54. Miranda RL, Conway T, Leatham MP, Chang DE, Norris WE, Allen JH, Stevenson SJ, Laux DC, Cohen PS. Glycolytic and gluconeogenic growth of *Escherichia coli* O157:H7 (EDL933) and *E. coli* K-12 (MG1655) in the mouse intestine. Infect. Immun. 2004; 72:1666–1676. [PubMed: 14977974]
55. Lin, ECC. Sugars, polyols, and carboxylates. In: Neidhardt, FC.; Curtiss, R., III; Ingraham, JL.; Lin, ECC.; Low, KB.; Magasanik, B.; Reznikoff, WS.; Riley, M.; Schaechter, M.; Umberger, HE., editors. *Escherichia coli* and *Salmonella*: cellular and molecular biology. 2nd ed.. Washington, D.C.: ASM Press; 1996. p. 307-342.
56. Reitzer, L. Chapter 3.4.7 Catabolism of Amino Acids and Related Compounds. In: Böck, A.; Curtiss, R., III; Kaper, JB.; Karp, PD.; Neidhardt, FC.; Nyström, T.; Slauch, JM.; Squires, CL.; Uesery, D., editors. *EcoSal--Escherichia coli* and *Salmonella*: Cellular and Molecular Biology, 3rd. Online ed.. Washington, D.C.: ASM Press; 2005.
57. Monk JM, Charusanti P, Aziz RK, Lerman JA, Premyodhin N, Orth JD, Feist AM, Palsson BO. Genome-scale metabolic reconstructions of multiple *Escherichia coli* strains highlight strain-specific adaptations to nutritional environments. Proc Natl Acad Sci U S A. 2013; 110:20338–20343. [PubMed: 24277855]
58. Autieri SM, Lins JJ, Leatham MP, Laux DC, Conway T, Cohen PS. L-fucose stimulates utilization of D-ribose by *Escherichia coli* MG1655 *fucAO* and *E. coli* Nissle 1917 *fucAO* mutants in the mouse intestine and in M9 minimal medium. Infect. Immun. 2007; 75:5465–5475. [PubMed: 17709419]
59. Jones SA, Jorgensen M, Chowdhury FZ, Rodgers R, Hartline J, Leatham MP, Struve C, Krogfelt KA, Cohen PS, Conway T. Glycogen and maltose utilization by *Escherichia coli* O157:H7 in the mouse intestine. Infect. Immun. 2008; 76:2531–2540. [PubMed: 18347038]
60. Leatham-Jensen MP, Frimodt-Moller J, Adediran J, Mokszycki ME, Banner ME, Caughron JE, Krogfelt KA, Conway T, Cohen PS. The streptomycin-treated mouse intestine selects *Escherichia*

coli envZ missense mutants that interact with dense and diverse intestinal microbiota. Infect. Immun. 2012; 80:1716–1727. [PubMed: 22392928]

61. Atuma C, Strugala V, Allen A, Holm L. The adherent gastrointestinal mucus gel layer: thickness and physical state *in vivo*. Am J Physiol Gastrointest Liver Physiol. 2001; 280:G922–C929. [PubMed: 11292601]
62. Allen, A. The structure and function of gastrointestinal mucous. In: Boedeker, EC., editor. Attachment of organisms to the gut mucosa. Vol. II. Boca Raton, Fla., Boca Raton, Fla: 1984. p. 3-11.
63. Kim, YS.; Morita, A.; Miura, S.; Siddiqui, B. Structure of glycoconjugates of intestinal mucosal membranes. In: Boedeker, EC., editor. Attachment of organisms to the gut mucosa. Vol. 2. Boca Raton, FL: CRC Press; 1984. p. 99-109.
64. Slomiany, BL.; Slomiany, A. Lipid and mucus secretions of the alimentary tract. In: Boedeker, EC., editor. Attachment of organisms to the gut mucosa. Vol. II. Boca Raton, Fla., Boca Raton, Fla: 1984. p. 24-31.
65. Johansson ME, Gustafsson JK, Holmen-Larsson J, Jabbar KS, Xia L, Xu H, Ghishan FK, Carvalho FA, Gewirtz AT, Sjoval H, Hansson GC. Bacteria penetrate the normally impenetrable inner colon mucus layer in both murine colitis models and patients with ulcerative colitis. Gut. 2014; 63:281–291. [PubMed: 23426893]
66. Johansson ME, Larsson JM, Hansson GC. The two mucus layers of colon are organized by the MUC2 mucin, whereas the outer layer is a legislator of host-microbial interactions. Proc Natl Acad Sci U S A. 2011; 108(Suppl 1):4659–4665. [PubMed: 20615996]
67. Hoskins, LC. Mucin degradation by enteric bacteria: ecological aspects and implications for bacterial attachment to gut mucosa. In: Boedeker, EC., editor. Attachment of organisms to the gut mucosa. Vol. II. Boca Raton, Fla., Boca Raton, Fla: 1984. p. 51-65.
68. Neutra, MR. The mechanism of intestinal mucous secretion. In: Boedeker, EC., editor. Attachment of organisms to the gut mucosa. Vol. II. Boca Raton, Fla., Boca Raton, Fla: 1984. p. 33-41.
69. Johansson ME, Phillipson M, Petersson J, Velcich A, Holm L, Hansson GC. The inner of the two Muc2 mucin-dependent mucus layers in colon is devoid of bacteria. Proc Natl Acad Sci U S A. 2008; 105:15064–15069. [PubMed: 18806221]
70. Holmen Larsson JM, Thomsson KA, Rodriguez-Pineiro AM, Karlsson H, Hansson GC. Studies of mucus in mouse stomach, small intestine, and colon. III. Gastrointestinal Muc5ac and Muc2 mucin O-glycan patterns reveal a regiospecific distribution. Am J Physiol Gastrointest Liver Physiol. 2013; 305:G357–G363. [PubMed: 23832516]
71. Hoskins LC, Agustines M, McKee WB, Boulding ET, Kriaris M, Niedermeyer G. Mucin degradation in human colon ecosystems. Isolation and properties of fecal strains that degrade ABH blood group antigens and oligosaccharides from mucin glycoproteins. J Clin Invest. 1985; 75:944–953. [PubMed: 3920248]
72. Moore WE, Holdeman LV. Human fecal flora: the normal flora of 20 Japanese-Hawaiians. Appl Microbiol. 1974; 27:961–979. [PubMed: 4598229]
73. Comstock LE, Coyne MJ. *Bacteroides thetaiotaomicron*: a dynamic, niche-adapted human symbiont. Bioessays. 2003; 25:926–929. [PubMed: 14505359]
74. Goodman AL, McNulty NP, Zhao Y, Leip D, Mitra RD, Lozupone CA, Knight R, Gordon JI. Identifying genetic determinants needed to establish a human gut symbiont in its habitat. Cell Host Microbe. 2009; 6:279–289. [PubMed: 19748469]
75. Martens EC, Chiang HC, Gordon JI. Mucosal glycan foraging enhances fitness and transmission of a saccharolytic human gut bacterial symbiont. Cell Host Microbe. 2008; 4:447–457. [PubMed: 18996345]
76. Ng KM, Ferreyra JA, Higginbottom SK, Lynch JB, Kashyap PC, Gopinath S, Naidu N, Choudhury B, Weimer BC, Monack DM, Sonnenburg JL. Microbiota-liberated host sugars facilitate post-antibiotic expansion of enteric pathogens. Nature. 2013; 502:96–99. [PubMed: 23995682]
77. Salyers AA, Pajeau M. Competitiveness of different polysaccharide utilization mutants of *Bacteroides thetaiotaomicron* in the intestinal tracts of germfree mice. Appl Environ Microbiol. 1989; 55:2572–2578. [PubMed: 2557798]

78. Mahowald MA, Rey FE, Seedorf H, Turnbaugh PJ, Fulton RS, Wollam A, Shah N, Wang C, Magrini V, Wilson RK, Cantarel BL, Coutinho PM, Henrissat B, Crock LW, Russell A, Verberkmoes NC, Hettich RL, Gordon JI. Characterizing a model human gut microbiota composed of members of its two dominant bacterial phyla. *Proc Natl Acad Sci U S A*. 2009; 106:5859–5864. [PubMed: 19321416]
79. Tilman D. Resource competition and community structure. *Monogr Popul Biol*. 1982; 17:1–296. [PubMed: 7162524]
80. Lendenmann U, Snozzi M, Egli T. Kinetics of the simultaneous utilization of sugar mixtures by *Escherichia coli* in continuous culture. *Appl Environ Microbiol*. 1996; 62:1493–1499. [PubMed: 8633848]
81. Ihssen J, Egli T. Global physiological analysis of carbon- and energy-limited growing *Escherichia coli* confirms a high degree of catabolic flexibility and preparedness for mixed substrate utilization. *Environ Microbiol*. 2005; 7:1568–1581. [PubMed: 16156730]
82. Liu M, Durfee T, Cabrera JE, Zhao K, Jin DJ, Blattner FR. Global transcriptional programs reveal a carbon source foraging strategy by *Escherichia coli*. *J Biol Chem*. 2005; 280:15921–15927. [PubMed: 15705577]
83. Ferenci T. Hungry bacteria—definition and properties of a nutritional state. *Environ Microbiol*. 2001; 3:605–611. [PubMed: 11722540]
84. Sartor RB. Probiotic therapy of intestinal inflammation and infections. *Curr Opin Gastroenterol*. 2005; 21:44–50. [PubMed: 15687884]
85. Grozdanov L, Raasch C, Schulze J, Sonnenborn U, Gottschalk G, Hacker J, Dobrindt U. Analysis of the genome structure of the nonpathogenic probiotic *Escherichia coli* strain Nissle 1917. *J Bacteriol*. 2004; 186:5432–5441. [PubMed: 15292145]
86. Sun J, Gunzer F, Westendorf AM, Buer J, Scharfe M, Jarek M, Gossling F, Blocker H, Zeng AP. Genomic peculiarity of coding sequences and metabolic potential of probiotic *Escherichia coli* strain Nissle 1917 inferred from raw genome data. *J Biotechnol*. 2005; 117:147–161. [PubMed: 15823404]
87. Alteri CJ, Smith SN, Mobley HL. Fitness of *Escherichia coli* during urinary tract infection requires gluconeogenesis and the TCA cycle. *PLoS Pathog*. 2009; 5:e1000448. [PubMed: 19478872]
88. Fabich AJ, Leatham MP, Grissom JE, Wiley G, Lai H, Najar F, Roe BA, Cohen PS, Conway T. Genotype and phenotypes of an intestine-adapted *Escherichia coli* K-12 mutant selected by animal passage for superior colonization. *Infect. Immun*. 2011; 79:2430–2439. [PubMed: 21422176]
89. Adediran J, Leatham-Jensen MP, Mokszycki ME, Frimodt-Moller J, Krogfelt KA, Kazmierczak K, Kenney LJ, Conway T, Cohen PS, McCormick BA. An *Escherichia coli* Nissle 1917 missense mutant colonizes the streptomycin-treated mouse intestine better than the wild type but is not a better probiotic. *Infect Immun*. 2014; 82:670–682. [PubMed: 24478082]
90. Bohnhoff M, Drake BL, Miller CP. Effect of streptomycin on susceptibility of intestinal tract to experimental *Salmonella* infection. *Proc Soc Exp Biol Med*. 1954; 86:132–137. [PubMed: 13177610]
91. Hentges DJ, Que JU, Casey SW, Stein AJ. The influence of streptomycin on colonization in mice. *Microecol. Theor*. 1984; 14:53–62.
92. Sonnenburg JL, Chen CT, Gordon JI. Genomic and metabolic studies of the impact of probiotics on a model gut symbiont and host. *PLoS Biol*. 2006; 4:e413. [PubMed: 17132046]
93. Samuel BS, Gordon JI. A humanized gnotobiotic mouse model of host-archaeal-bacterial mutualism. *Proc Natl Acad Sci U S A*. 2006; 103:10011–10016. [PubMed: 16782812]
94. Winter SE, Winter MG, Xavier MN, Thiennimitr P, Poon V, Keestra AM, Laughlin RC, Gomez G, Wu J, Lawhon SD, Popova IE, Parikh SJ, Adams LG, Tsolis RM, Stewart VJ, Baumler AJ. Host-derived nitrate boosts growth of *E. coli* in the inflamed gut. *Science*. 2013; 339:708–711. [PubMed: 23393266]
95. Meador JP, Caldwell ME, Cohen PS, Conway T. *Escherichia coli* pathotypes occupy distinct niches in the mouse intestine. *Infect Immun*. 2014
96. Gauger EJ, Leatham MP, Mercado-Lubo R, Laux DC, Conway T, Cohen PS. Role of motility and the *flhDC* Operon in *Escherichia coli* MG1655 colonization of the mouse intestine. *Infect Immun*. 2007; 75:3315–3324. [PubMed: 17438023]

97. Leatham MP, Stevenson SJ, Gauger EJ, Krogfelt KA, Lins JJ, Haddock TL, Autieri SM, Conway T, Cohen PS. Mouse intestine selects nonmotile *flhDC* mutants of *Escherichia coli* MG1655 with increased colonizing ability and better utilization of carbon sources. *Infect. Immun.* 2005; 73:8039–8049. [PubMed: 16299298]
98. Bartlett DH, Frantz BB, Matsumura P. Flagellar transcriptional activators FlhB and FlaI: gene sequences and 5' consensus sequences of operons under FlhB and FlaI control. *J Bacteriol.* 1988; 170:1575–1581. [PubMed: 2832369]
99. Prüß BM, Campbell JW, Van Dyk TK, Zhu C, Kogan Y, Matsumura P. FlhD/FlhC is a regulator of anaerobic respiration and the Entner-Doudoroff pathway through induction of the methyl-accepting chemotaxis protein Aer. *J Bacteriol.* 2003; 185:534–543. [PubMed: 12511500]
100. De Paepe M, Gaboriau-Routhiau V, Rainteau D, Rakotobe S, Taddei F, Cerf-Bensussan N. Trade-off between bile resistance and nutritional competence drives *Escherichia coli* diversification in the mouse gut. *PLoS Genet.* 2011; 7:e1002107. [PubMed: 21698140]
101. Giraud A, Arous S, Paepe MD, Gaboriau-Routhiau V, Bambou JC, Rakotobe S, Lindner AB, Taddei F, Cerf-Bensussan N. Dissecting the genetic components of adaptation of *Escherichia coli* to the mouse gut. *PLoS Genet.* 2008; 4:e2. [PubMed: 18193944]
102. Egger LA, Park H, Inouye M. Signal transduction via the histidyl-aspartyl phosphorelay. *Genes Cells.* 1997; 2:167–184. [PubMed: 9189755]
103. Walthers, D.; Go, A.; Kenney, LJ. Regulation of Porin Gene Expression by the Two-Component Regulatory System EnvZ/OmpR. In: Benz, R., editor. *Bacterial and Eukaryotic Porins*. Wiley-VCH Verlag; 2004.
104. Pratt LA, Kolter R. Genetic analysis of *Escherichia coli* biofilm formation: roles of flagella, motility, chemotaxis and type I pili. *Mol Microbiol.* 1998; 30:285–293. [PubMed: 9791174]
105. Licht TR, Christensen BB, Krogfelt KA, Molin S. Plasmid transfer in the animal intestine and other dynamic bacterial populations: the role of community structure and environment. *Microbiology.* 1999; 145(Pt 9):2615–2622. [PubMed: 10517615]
106. Macfarlane S. Microbial biofilm communities in the gastrointestinal tract. *J Clin Gastroenterol.* 2008; 42(Suppl 3)(Pt 1):S142–S143. [PubMed: 18806707]
107. Macfarlane S, Bahrami B, Macfarlane GT. Mucosal biofilm communities in the human intestinal tract. *Adv Appl Microbiol.* 2011; 75:111–143. [PubMed: 21807247]
108. Palestrant D, Holzkecht ZE, Collins BH, Parker W, Miller SE, Bollinger RR. Microbial biofilms in the gut: visualization by electron microscopy and by acridine orange staining. *Ultrastruct Pathol.* 2004; 28:23–27. [PubMed: 14967595]
109. Swidsinski A, Weber J, Loening-Baucke V, Hale LP, Lochs H. Spatial organization and composition of the mucosal flora in patients with inflammatory bowel disease. *J Clin Microbiol.* 2005; 43:3380–3389. [PubMed: 16000463]
110. Macfarlane S, Woodmansey EJ, Macfarlane GT. Colonization of mucin by human intestinal bacteria and establishment of biofilm communities in a two-stage continuous culture system. *Appl Environ Microbiol.* 2005; 71:7483–7492. [PubMed: 16269790]
111. Zoetendal EG, von Wright A, Vilpponen-Salmela T, Ben-Amor K, Akkermans AD, de Vos WM. Mucosa-associated bacteria in the human gastrointestinal tract are uniformly distributed along the colon and differ from the community recovered from feces. *Appl Environ Microbiol.* 2002; 68:3401–3407. [PubMed: 12089021]
112. Fong JC, Syed KA, Klose KE, Yildiz FH. Role of *Vibrio* polysaccharide (vps) genes in VPS production, biofilm formation and *Vibrio cholerae* pathogenesis. *Microbiology.* 2010; 156:2757–2769. [PubMed: 20466768]
113. Henrissat B, Davies G. Structural and sequence-based classification of glycoside hydrolases. *Curr Opin Struct Biol.* 1997; 7:637–644. [PubMed: 9345621]
114. Franklin DP, Laux DC, Williams TJ, Falk MC, Cohen PS. Growth of *Salmonella typhimurium* SL5319 and *Escherichia coli* F-18 in mouse cecal mucus: role of peptides and iron. *FEMS Microbiol Ecol.* 1990; 74:229–240.
115. Licht TR, Tolker-Nielsen T, Holmstrom K, Krogfelt KA, Molin S. Inhibition of *Escherichia coli* precursor-16S rRNA processing by mouse intestinal contents. *Environ Microbiol.* 1999; 1:23–32. [PubMed: 11207715]

116. Newman JV, Kolter R, Laux DC, Cohen PS. Role of *leuX* in *Escherichia coli* colonization of the streptomycin-treated mouse large intestine. *Microb Pathog.* 1994; 17:301–311. [PubMed: 7723657]
117. Sweeney NJ, Klemm P, McCormick BA, Moller-Nielsen E, Utley M, Schembri MA, Laux DC, Cohen PS. The *Escherichia coli* K-12 *gntP* gene allows *E. coli* F-18 to occupy a distinct nutritional niche in the streptomycin-treated mouse large intestine. *Infect Immun.* 1996; 64:3497–3503. [PubMed: 8751890]
118. McCormick BA, Stocker BA, Laux DC, Cohen PS. Roles of motility, chemotaxis, and penetration through and growth in intestinal mucus in the ability of an avirulent strain of *Salmonella typhimurium* to colonize the large intestine of streptomycin-treated mice. *Infect Immun.* 1988; 56:2209–2217. [PubMed: 3044995]

Figure 1.

Table 1
Central metabolism mutants tested for colonization defects in the mouse intestine

Pathway	Gene Defect	MG1655	EDL933
Glyoxylate bypass	<i>aceA</i>	No	Yes
ED	<i>edd</i>	Yes	Yes
glycolysis	<i>pgi</i>	Yes	ND
PPP	<i>gnd</i>	No	ND
gluconeogenesis	<i>ppsA pckA</i>	No	No
TCA cycle	<i>frdA</i>	Yes	Yes
TCA cycle	<i>sdhAB</i>	No	Yes
TCA cycle	<i>frdA sdhAB</i>	Yes	Yes

Results show the difference in population sizes of wild type verses mutant strains at Day 9.
Yes indicates the difference exceeds a 0.8 log10 colonization advantage and students *t* test value $P<0.05$.
ND indicates not determined.

Table 2

Sugar utilization in the intestine by *E. coli* strains.

Sugar Defect	Mutation	MG1655	Nissle	HS	EDL933	UPEC	EPEC
Arabinose	<i>araBAD</i>	Yes	Yes	Yes	Yes	Yes	Yes
Fucose	<i>fucK</i>	No	Yes	No	No	No	No
Galactose	<i>galK</i>	No	Yes	Yes	Yes	Yes	Yes
Gluconate	<i>gntK/ idnK</i>	Yes	Yes	Yes	No	ND	ND
Hexuronates	<i>axaC</i>	No	No	No	Yes	ND	ND
Lactose	<i>lacZ</i>	No	No	Yes	No	ND	ND
Mannose	<i>manA</i>	No	Yes	No	Yes	Yes	Yes
N-acetylglucosamine	<i>nagE</i>	Yes	No	Yes	Yes	Yes	Yes
N-acetylglactosamine	<i>agaWEFA</i>	NA	Yes	ND	No	ND	ND
N-acetylneuraminate	<i>nanAT</i>	Yes	Yes	Yes	No	ND	ND
Ribose	<i>rbsK</i>	No	No	Yes	Yes	Yes	Yes
Sucrose	<i>sacH</i>	NA	NA	NA	Yes	NA	ND

Results show the difference in population sizes of wild type verses mutant strains at Day 9.

Yes indicates the difference exceeds a 0.8 log10 colonization advantage and students *t* test value $P < 0.05$.

NA indicates that the pathway is not intact in this genetic background.

ND indicates not determined.