Published in final edited form as:

J Health Care Poor Underserved. 2009 August ; 20(3): 756–765. doi:10.1353/hpu.0.0189.

Increasing the Reach of HIV Testing to Young Latino MSM: Results of a Pilot Study Integrating Outreach and Services

Jennifer Toller Erausquin, PhD, MPH, Naihua Duan, PhD, Oscar Grusky, PhD, Aimee-Noelle Swanson, PhD, Dustin Kerrone, MS, and Ellen T. Rudy, PhD

JENNIFER TOLLER ERAUSQUIN is a Research Associate at the University of California Los Angeles (UCLA) School of Public Health. NAIHUA DUAN is the Director of the Division of Biostatistics in the Department of Psychiatry at Columbia University Medical Center and a faculty member of the Departments of Psychiatry and Biostatistics at Columbia University. OSCAR GRUSKY is Emeritus Professor of Sociology, Director of the UCLA AIDS Research Training Program, and Co-Director, Administrative Core in the Center for HIV Identification, Prevention, and Treatment Services. AIMEE-NOELLE SWANSON is Project Director for a NIDA P50 Center for Excellence in the UCLA Department of Family Medicine. DUSTIN KERRONE is Program Supervisor of SPOT, an HIV testing satellite clinic of the Los Angeles Gay and Lesbian Center. ELLEN T. RUDY is an Epidemiologist with the Los Angeles County Department of Public Health, Sexually Transmitted Disease Program.

Abstract

Background—In the U.S., HIV infections are increasing among men who have sex with men (MSM), particularly young, racial/ethnic minority MSM.

Objective—To examine the feasibility of increasing HIV testing among young Latino MSM by integrating tailored outreach strategies with testing, counseling, and HIV medical services.

Design—Descriptive study comparing demographic characteristics, behaviors, and HIV test results of clients from the intervention period with clients who tested during other time periods.

Results—Clients in the intervention period were younger and more likely to be Latino than those in other time periods. In addition, clients who received outreach were more likely than those who did not receive outreach to report methamphetamine use, sex with an HIV-positive person, and sex with a sex worker.

Conclusion—Venue-based and selective media outreach, in combination with linking rapid testing to HIV care, may help overcome some of the barriers to testing among high-risk young Latino MSM.

Keywords

HIV/AIDS; men who have sex with men (MSM); Latino; community-based organization

In the United States, men who have sex with men (MSM) continue to be disproportionately affected by HIV/AIDS. These men make up 46% of all new diagnoses of HIV and AIDS, and 63% of new diagnoses among men.^{1,2} Despite declines in risk behavior in the 1980s and 1990s as more MSM used condoms and reduced their number of sexual partners,^{3,4} recent data indicate that HIV infections and AIDS cases are on the rise among this population. From 2001 to 2006, HIV/AIDS diagnoses among MSM increased 8.6% in the 33 states using confidential name-based reporting.² In addition, HIV in MSM is increasingly concentrated among African American and Latino men, and among young men.^{1,2,5–7} Latino and African American MSM make up 55% of the HIV/AIDS infections reported among MSM in the U.S.² Although the estimated annual percentage change in HIV/AIDS cases from 2001–2006 was 0.7 for White MSM, it was 1.9 for both Latino and African American MSM.² For MSM under age 25, the annual percentage change was 12.4, compared with –1.1 for MSM ages 25–44, suggesting a shift in patterns of infection towards younger men.

One important factor contributing to the sexual transmission of HIV among MSM and the increasing prevalence of HIV among young, racial/ethnic minority MSM is lack of awareness of HIV infection.⁸ People who are HIV-positive and unaware of their infection (designated as *HIV-positive/unaware* or *having an unrecognized infection*) are unlikely to change their behaviors in ways that reduce the risk of HIV transmission to uninfected partners. A meta-analysis of risk behavior and awareness of HIV status showed that HIV-positive/unaware men and women are nearly twice as likely as HIV-positive/aware men and women to report unprotected anal or vaginal sex.⁹ Further, HIV-infected/unaware individuals are 3.5 times more likely to transmit HIV to partners than those who are aware of their HIV infection.¹⁰ Among MSM, men with unrecognized HIV infection have 49% greater odds of engaging in unprotected anal intercourse than men who are aware of their infection.¹¹ Multi-city studies of MSM have shown that Latino, African American, and young MSM are most likely to be HIV infected and unaware.^{11–13}

These data on lack of awareness of HIV infection are alarming because the benefits of knowing one's HIV serostatus are well established. HIV-positive individuals live longer and report an increase in their quality of life with early knowledge of the disease and treatment with antiretroviral medications.^{14,15} Awareness of HIV serostatus can also reduce risk behaviors and thereby decrease HIV transmission.^{16,17} In addition, HIV medical treatment decreases HIV viral load and thus infectivity to others.^{18,19}

A number of programs and policies have attempted to increase HIV testing and awareness of infection, but few have effectively targeted young, racial/ethnic minority MSM. In 2003 the Centers for Disease Control and Prevention (CDC) announced new strategies to promote HIV testing, including incorporating HIV testing into routine medical care, expanded use of rapid testing, and increased opportunities for testing outside of medical settings.^{20,21} In many urban areas where ethnic populations are concentrated, additional strategies are necessary to reach young and at-risk MSM, who frequently do not use a regular source of health care and are reluctant to test for HIV in clinics or mobile vans in neighborhoods where their peers and family may see them.^{22,23}

Young Latino MSM, the focus of this study, frequently face barriers to HIV testing due to socioeconomic status, ethnicity, and sexual behavior.^{7,24,25} Socioeconomic status may decrease access to medical care, including testing for HIV. Experiences or fear of homophobia and racism may increase social isolation and poor self-esteem, which in turn reduce HIV testing and protective behaviors.^{22,24} In addition, young Latino MSM may be further marginalized due to their age. Young MSM are often concerned with concealing their sexual behavior and may be less likely to seek out information, condoms, or HIV testing, particularly if such services are not located in settings where youth can be reached easily.^{23,26}

In an effort to increase HIV testing among young Latino MSM at risk for HIV infection, the Los Angeles Gay and Lesbian Center collaborated with a team of academic researchers to develop and implement an innovative pilot intervention. This study's unique contribution is that it integrates tailored outreach strategies with testing, counseling, and HIV medical services.

Methods

Setting

The Los Angeles Gay and Lesbian Center (LAGLC) is an established community-based organization that provides a broad range of health and social services for the lesbian, gay, bisexual, and transgender community. Since the beginning of the HIV epidemic, LAGLC has worked to promote care of HIV-infected persons, and has been a leader in community HIV testing and prevention efforts. The LAGLC is situated within Los Angeles County, where 43% of new AIDS cases are among Latinos,²⁷ and 70% of Latino AIDS cases are among men who have sex with men.²⁸ In recognition of the need to improve awareness of HIV serostatus among young and Latino MSM, LAGLC staff contacted local university researchers for assistance in developing and implementing a pilot intervention.

Project design

In the formative stage of this project, focus group interviews were conducted with young Latino MSM. The focus group data indicated that Latino MSM at risk for HIV may be marginalized by their ethnicity, sexual behavior, and lack of economic and social-environmental resources. Many participants mentioned reasons for avoiding HIV testing such as fear of a positive HIV test result and not knowing where to go for HIV care. The focus group data suggested that a constellation of factors limited this population's access to health services and their knowledge of where to seek services that are gay-friendly, youth-friendly, and culturally sensitive. It was learned that many young Latino MSM are unable to access HIV testing services, a circumstance contributing to the disproportionate number of HIV-positive young Latino MSM who are unaware of their HIV serostatus. The pilot intervention was designed to address this cycle of vulnerability by increasing awareness of free testing services, providing incentives for getting test results, and improving access to treatment by integrating HIV medical care with testing.

Targeted outreach was provided to young Latino MSM. Trained staff and volunteers recruited participants at Latino-oriented gay club and event nights in the Los Angeles and West Hollywood area. From August through October 2004, trained outreach volunteers that shared characteristics with the target population (young, gay/bisexual, Latino) distributed bilingual outreach cards to encourage young Latino MSM to test for HIV at LAGLC's Service, Prevention, Outreach, Treatment (SPOT) center in West Hollywood. Potential clients were informed that the testing location was within walking distance, that rapid HIV test results would be available in about 20 minutes, and that outreach cards could be traded for a movie pass at the time of testing. In addition to the venue-based outreach, testing information was advertised on two Internet sites and in three widely available gay/bisexual-oriented magazines. The Internet postings included a printable outreach coupon and the magazine advertisements had a coupon that could be exchanged at the time of testing for a movie pass.

The second component of the pilot intervention involved HIV counseling and testing, and early entry into medical care for individuals who tested HIV-positive. Free HIV counseling and testing was provided at the SPOT. The SPOT is a storefront clinic located in the heart of West Hollywood, close to clubs and bars that attract a large gay and bisexual crowd. The clinic has evening hours and, during the intervention period, offered rapid HIV tests (finger prick blood tests with OraQuick®) that allowed clients to receive their results in 20 minutes. All individuals with preliminary positive test results had their preliminary positive test results confirmed with Western blot and were linked to HIV medical care within 72 hours. The HIV medical care was offered at the LAGLC's HIV care clinic or at another clinic convenient for the patient.

Data collection and statistical analysis

The SPOT center gathers standard demographic and risk assessment information from all clients as part of their HIV testing and counseling. This process remained unchanged during the intervention period. A trained staff counselor gathered this information using a California standardized risk assessment form, on which client's age, ethnicity, HIV risk behaviors, reason for testing, previous HIV tests, sexually transmitted disease (STD) history, and the result of the current HIV test are entered. During the intervention period, additional information was gathered regarding the presence of an outreach card or coupon, and the outreach source code identifying the place where the card or coupon originated.

Data from the intervention period (August–October 2004) were entered into a database that also included data from MSM testing for HIV during two comparison periods: May–July 2004 and August–October 2003. These comparison groups were selected in order to reduce the likelihood of mistakenly attributing temporal or seasonal trends to an intervention effect. Statistical analyses were conducted using Stata 9.2/SE.²⁹ First, data from the intervention period was compared with data from the two comparison periods, using one-way Analysis of Variance (ANOVA) for continuous variables (e.g., age, number of sexual partners) and chi-squared analysis or Fisher's exact test for categorical variables (e.g., Latino ethnicity, HIV test result). Next, data from the intervention period was further analyzed, for

differences between MSM who had received outreach and those who had not. One-way ANOVA, chi-squared analysis, and Fisher's exact tests were used, as appropriate.

Results

Table 1 shows descriptive demographic and behavioral characteristics, as well as HIV test results, for the intervention period and two comparison periods. Results are limited to male HIV testing clients age 25 and under who reported sexual activity with a male. During the intervention period, 95 young MSM tested for HIV. They had an average age of 22 years, and reported an average of 10 partners (range: 1–100) in the prior two years or since their last negative HIV test. One-third of testing clients in the intervention period self-reported Hispanic/Latino ethnicity. Clients reported a variety of risk behaviors, including methamphetamine use (16.8%); sometimes not using or never using a condom for anal sex with a male (69.4%); having sex with an HIV-positive individual (16.1%), a sex worker (7.6%), or an injection drug user (11.0%); and exchanging sex for money or drugs (10.5%). Five clients (5.8%) tested positive during the intervention period. Tests of comparison across the intervention period, Summer 2004, and Fall 2003 indicated that MSM clients testing in the intervention period were significantly younger [$F(2, 233) = 3.13, p=.045$] and more likely than clients in Summer 2004 to report being Latino ($\chi^2(2) = 8.83, p=.012$).

Data from the intervention period, stratified by outreach, are presented in Table 2. About half of young MSM testing clients during the intervention period were outreach participants, as evidenced by presenting an outreach card or coupon. The cards and coupons indicated that 78% [36/46] of outreach recipients received in-person outreach; the remainder saw Internet postings, magazine advertisements, or both [results not reported in table]. Outreach recipients significantly differed from clients who did not receive outreach on several key characteristics. Outreach recipients were younger [$F(1, 93) = 27.08, p=.015$] and were more likely than clients who did not receive outreach to be Latino ($\chi^2(1) = 4.64, p=.031$). Outreach recipients also were more likely to report a number of HIV risk behaviors. Strikingly, while 6.1% of other clients reported methamphetamine use, 28.3% of outreach recipients reported using this drug ($\chi^2(1) = 8.30, p=.004$). Outreach recipients were also more than twice as likely to report having had sex with an HIV-positive person (23.9% compared with 8.5%; $\chi^2(1) = 4.08, p=.043$) and more than six times as likely to report having sex with a sex worker (13.0% compared with 2.2%; $\chi^2(1) = 3.87, p=.049$). The proportion of clients testing positive did not differ significantly by outreach. All clients testing positive during the intervention period, regardless of receipt of outreach, were linked to HIV medical care, that is, they received an appointment with an HIV specialist health care provider. Information on how many actually showed up for their appointment was unavailable.

Discussion

The purpose of this study was to describe the effects of a pilot intervention on reaching high-risk, young, and Latino MSM with HIV testing. The intervention, which combined venue-based outreach and targeted media advertisements with testing, counseling, and HIV care, was successful in motivating the target population to test for HIV. The intervention

resulted in more testing among young Latino clients. In addition, the outreach strategies were successful in that a higher proportion of outreach recipients (than of clients who did not receive outreach) was Latino. Furthermore, compared with HIV testing clients who did not receive outreach, more outreach recipients reported behaviors that put them at risk for HIV infection. Despite a relatively low prevalence of HIV infection across all study groups, HIV risk behaviors tended to be higher among outreach recipients. A higher proportion of outreach recipients reported HIV risk behaviors, including methamphetamine use, having sex with an HIV-positive person, and having sex with a sex worker. These findings suggest that the intervention was able to motivate high-risk, young, and Latino MSM to test for HIV.

This study demonstrates that it is feasible to create an effective, integrated outreach and service intervention for young Latino MSM at risk for HIV infection. The results are consistent with other studies of MSM in the United States, which show that Latino MSM have high rates of unprotected anal intercourse and other sexual risk behaviors.^{7,8,13,30–33} Collaboration between a community-based organization and university researchers facilitated use of novel strategies for motivating at-risk young men to test for HIV. Importantly, the intervention design and its implementation accounted for the characteristics and needs of the target population. Expansion of late-night hours for HIV testing and making rapid tests available seemed to be effective strategies. In addition, targeted outreach to clients through in-person, venue-based recruitment and advertisements on selected Internet sites and in magazines appeared successful. Future intervention studies targeting young Latino MSM should assess additional, sustainable community-based strategies for reaching this population with HIV testing and linking those who test positive to HIV medical care.

The results of this study are also relevant in light of recent efforts to expand HIV testing in the U.S. In 2006, the CDC released new recommendations for testing, which emphasized that not only should testing in health care settings be routine, but it should be offered on an opt-out, rather than opt-in, basis.³⁴ These recommendations have yet to be implemented by most health care providers, and many young adults, particularly men, do not seek primary or preventive care. In order to increase HIV testing among groups such as young and racial/ethnic minority men, prevention providers may need to tailor their efforts based on the targets' age and culture.^{11,22,35–38} Innovative strategies will be needed to reach at-risk men, and to help men overcome barriers to testing, such as fears about testing and about follow-up care. The results of this pilot intervention suggest that venue-based and selective media outreach, in combination with linking rapid testing to HIV care, may help overcome some of these barriers.

Acknowledgments

The authors are grateful to the staff, volunteers, and clients of the Los Angeles Gay and Lesbian Center for their participation in this project. The authors received support for this project from the following sources: (1) UCLA School of Nursing Center for Vulnerable Populations Research, "An integrated outreach and service intervention for young Latino MSM" (O. Grusky, PI). The Center is supported by NINR P30 NR00504 (D. Koniak-Griffin, PI). (2) UCLA Center for HIV Identification, Prevention, and Treatment Services, "Evaluation of an integrated outreach and services intervention" (O. Grusky, PI; J.T. Erausquin, Co-PI). The Center is supported by National Institute of Mental Health P30 58107 (M.J. Rotheram-Boris, PI). (3) National Institute of Mental Health T32-MH19127, "Service systems for persons living with HIV/AIDS" (O. Grusky, PI).

Notes

1. Centers for Disease Control and Prevention. HIV/AIDS surveillance report, 2006; vol. 18. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2008. p. 1-55. Available at: <http://www.cdc.gov/hiv/topics/surveillance/resources/reports/>.
2. Centers for Disease Control and Prevention. Trends in HIV/AIDS diagnoses among men who have sex with men—33 states, 2001–2006. MMWR Morb Mortal Wkly Rep. 2008 Jun 27; 57(25):681–686. [PubMed: 18583954]
3. Catania JA, Coates TJ, Stall R, et al. Changes in condom use among homosexual men in San Francisco. Health Psychol. 1991; 10(3):190–199. [PubMed: 1879391]
4. McKusick L, Coates TJ, Morin SF, et al. Longitudinal predictors of reductions in unprotected anal intercourse among gay men in San Francisco: the AIDS Behavioral Research Project. Am J Public Health. 1990 Aug; 80(8):978–983. [PubMed: 2368862]
5. Brooks R, Rotheram-Borus MJ, Bing EG, et al. HIV and AIDS among men of color who have sex with men and men of color who have sex with men and women: an epidemiological profile. AIDS Educ Prev. 2003 Feb; 15(1 Suppl A):1–6. [PubMed: 12630595]
6. Blair JM, Fleming PL, Karon JM. Trends in AIDS incidence and survival among racial/ ethnic minority men who have sex with men, United States, 1990–1999. J Acquir Immune Defic Syndr. 2002 Nov 1; 31(3):339–347. [PubMed: 12439211]
7. Harawa NT, Greenland S, Bingham TA, et al. Associations of race/ethnicity with HIV prevalence and HIV-related behaviors among young men who have sex with men in 7 urban centers in the United States. J Acquir Immune Defic Syndr. 2004 Apr 15; 35(5):526–536. [PubMed: 15021318]
8. Marks G, Millet GA, Bingham T, et al. Understanding differences in HIV sexual transmission among Latino and Black men who have sex with men: the Brothers y Hermanos Study. AIDS Behav. 2008 Aug 28. [Epub ahead of print.].
9. Marks G, Crepaz N, Senterfitt JW, et al. Meta-analysis of high-risk sexual behavior in persons aware and unaware they are infected with HIV in the United States: implications for HIV prevention programs. J Acquir Immune Defic Syndr. 2005 Aug 1; 39(4):446–453. [PubMed: 16010168]
10. Marks G, Crepaz N, Janssen RS. Estimating sexual transmission of HIV from persons aware and unaware that they are infected with the virus in the USA. AIDS. 2006 Jun 26; 20(10):1447–1450. [PubMed: 16791020]
11. MacKellar DA, Valleroy LA, Secura GM, et al. Perceptions of lifetime risk and actual risk for acquiring HIV among young men who have sex with men. AIDS Behav. 2007 Mar; 11(2):263–270. [PubMed: 16791527]
12. MacKellar DA, Valleroy LA, Secura GM, et al. Unrecognized HIV infection, risk behaviors, and perceptions of risk among young men who have sex with men: opportunities for advancing HIV prevention in the third decade of HIV/AIDS. J Acquir Immune Defic Syndr. 2005 Apr 15; 38(5):603–614. [PubMed: 15793373]
13. Sifakis F, Flynn CP, Metsch L, et al. HIV prevalence, unrecognized infection, and HIV testing among men who have sex with men—five U.S. cities, June 2004–April 2005. MMWR Morb Mortal Wkly Rep. 2005 Jun 24; 54(24):597–601. [PubMed: 15973239]
14. Murri R, Fantoni M, Del Borgo C, et al. Determinants of health-related quality of life in HIV-infected patients. AIDS Care. 2003 Aug; 15(4):581–590. [PubMed: 14509872]
15. Mocroft A, Ledergerber B, Katlama C, et al. Decline in the AIDS and death rates in the EuroSIDA study: an observational study. Lancet. 2003 Jul 5; 362(9377):22–29. [PubMed: 12853195]
16. Gardner LI, Metsch LR, Anderson-Mahoney P, et al. Efficacy of a brief case management intervention to link recently diagnosed HIV-infected persons to care. AIDS. 2005 Mar 4; 19(4):423–431. [PubMed: 15750396]
17. Weinhardt LS, Carey MP, Johnson BT, et al. Effects of HIV counseling and testing on sexual risk behavior: a meta-analytic review of published research, 1985–1997. Am J Public Health. 1999 Sep; 89:1397–1405. [PubMed: 10474559]

18. Quinn TC, Wawer MJ, Sewankambo N, et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. Rakai Project Study Group. *N Engl J Med*. 2000 Mar 30; 342(13): 921–929. [PubMed: 10738050]
19. Castilla J, Del Romero J, Hernando V, et al. Effectiveness of highly active antiretroviral therapy in reducing heterosexual transmission of HIV. *J Acquir Immune Defic Syndr*. 2005 Sep 1; 40(1):96–101. [PubMed: 16123689]
20. Centers for Disease Control and Prevention. Advancing HIV prevention: interim technical guidance for selected interventions. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2003.
21. Centers for Disease Control and Prevention. Advancing HIV prevention: new strategies for a changing epidemic—United States, 2003. *MMWR Morb Mortal Wkly Rep*. 2003 Apr 18; 52:329–332. [PubMed: 12733863]
22. Brooks RA, Etzel MA, Hinojos E, et al. Preventing HIV among Latino and African American gay and bisexual men in a context of HIV-related stigma, discrimination, and homophobia: perspectives of providers. *AIDS Patient Care STDS*. 2005 Nov; 19(11):737–744. [PubMed: 16283834]
23. Bell DN, Martinez J, Botwinick G, et al. Case finding for HIV-positive youth: a special type of hidden population. *J Adolesc Health*. 2003 Aug; 33(2 Suppl):10–22. [PubMed: 12888283]
24. Diaz RM, Ayala G, Bein E, et al. The impact of homophobia, poverty, and racism on the mental health of gay and bisexual Latino men: findings from 3 U.S. cities. *Am J Public Health*. 2001 Jun; 91:927–932. [PubMed: 11392936]
25. Marin BV. HIV prevention in the Hispanic community: sex, culture, and empowerment. *J Transcult Nurs*. 2003 Jul; 14(3):186–192. [PubMed: 12861921]
26. Rangel MC, Gavin L, Reed C, et al. Epidemiology of HIV and AIDS among adolescents and young adults in the United States. *J Adolesc Health*. 2006; 39(2):156–163. [PubMed: 16857526]
27. Los Angeles County Department of Health Services. An epidemiologic profile of HIV and AIDS in Los Angeles County 2004. Los Angeles, CA: United Way of Greater Los Angeles; 2005.
28. Los Angeles County Department of Health Services. HIV/AIDS semi-annual surveillance summary June 2008. Los Angeles, CA: Los Angeles County Department of Health Services, HIV Epidemiology Program; 2008.
29. StataCorp.. Stata Statistical Software: Release 9.2. College Station, TX: StataCorp LP; 2006.
30. Remien RH, Wagner G, Dolezal C, et al. Factors associated with HIV sexual risk behavior in male couples of mixed HIV status. *J Psychol Human Sex*. 2001; 13:31–48.
31. Chen SY, Gibson S, Katz MH, et al. Continuing increases in sexual risk behavior and sexually transmitted diseases among men who have sex with men: San Francisco, Calif., 1999–2001, USA. *Am J Public Health*. 2002 Sep; 92(9):1387–1388. [PubMed: 12197957]
32. Jaffe HW, Valdiserri RO, De Cock KM. The reemerging HIV/AIDS epidemic in men who have sex with men. *JAMA*. 2007 Nov 28; 298(20):2412–2414. [PubMed: 18042919]
33. van Kesteren NM, Hospers HJ, Kok G. Sexual risk behavior among HIV-positive men who have sex with men: a literature review. *Patient Educ Couns*. 2007 Jan; 65(1):5–20. Epub 2006 Nov 13. [PubMed: 17098392]
34. Branson BM, Handsfield HH, Lampe MA, et al. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR Recomm Rep*. 2006 Sep 22; 55(RR-14):1–17. [PubMed: 16988643]
35. Irwin KL, Valdiserri RO, Holmberg SD. The acceptability of voluntary HIV antibody testing in the United States: a decade of lessons learned. *AIDS*. 1996 Dec; 10(14):1707–1717. [PubMed: 8970692]
36. Stein JA, Nyamathi A. Gender differences in behavioural and psychosocial predictors of HIV testing and return for test results in a high-risk population. *AIDS Care*. 2000 Jun; 12(3):343–356. [PubMed: 10928212]
37. Spielberg F, Branson BM, Goldbaum GM, et al. Overcoming barriers to HIV testing: preferences for new strategies among clients of a needle exchange, a sexually transmitted disease clinic, and sex venues for men who have sex with men. *J Acquir Immune Defic Syndr*. 2003 Mar 1; 32(3): 318–327. [PubMed: 12626893]

38. Spielberg F, Branson BM, Goldbaum GM, et al. Choosing HIV counseling and testing strategies for outreach settings: a randomized trial. *J Acquir Immune Defic Syndr*. 2005 Mar 1; 38(3):348–355. [PubMed: 15735456]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Demographic characteristics, behavioral characteristics, and HIV test results of male clients age 25 years or younger over three periods of time

Characteristic	Fall 2003 (n=86)	Summer 2004 (n=97)	Intervention period, Fall 2004 (n=95)
Demographic characteristics			
Age *	22.2 ± 2.4	22.5 ± 1.8	21.8 ± 2.2
Number of sexual partners	6.1 ± 6.9	9.2 ± 14.4	10.2 ± 16.0
Latino ethnicity (%) *	36.4	16.8	33.0
Behavioral characteristics			
Methamphetamine use	11.4	8.3	16.8
Anal sex without a condom (%)	53.9	61.1	69.4
Sex with an HIV-positive person (%)	20.5	12.4	16.1
Sex with a sex worker (%)	—	6.2	7.6
Sex with an injection drug user (%)	6.8	6.2	11.0
Sex in exchange for money or drugs (%)	—	6.2	10.5
Test result			
HIV positive (%)	4.6	6.2	5.8

*
p<.05

Table 2

Demographic characteristics, behavioral characteristics, and HIV test results of male clients age 25 years or younger during the intervention period, by outreach status

Characteristic	Outreach recipients (n=46)	Clients who did not receive outreach (n=9)
Demographic characteristics		
Age *	21.2 ± 2.1	22.3 ± 2.1
Number of sexual partners	11.2 ± 16.4	9.2 ± 15.9
Latino ethnicity (%) *	44.2	22.9
Behavioral characteristics		
Methamphetamine use (%) *	28.3	6.1
Anal sex without a condom (%)	72.5	66.7
Sex with an HIV-positive person (%) *	23.9	8.5
Sex with a sex worker (%) *	13.0	2.2
Sex with an injection drug user (%)	15.2	6.7
Sex in exchange for money or drugs (%)	13.0	8.2
Test result		
HIV positive (%)	5.0	6.5

*
p<.05