Hodgkin disease: Hodgkin and Reed–Sternberg cells picked from histological sections show clonal immunoglobulin gene rearrangements and appear to be derived from B cells at various stages of development

RALF KÜPPERS*†, KLAUS RAJEWSKY*, MIN ZHAO‡, GÜNTHER SIMONS‡, RALF LAUMANN*, ROBERT FISCHER‡, AND MARTIN-LEO HANSMANN‡

*Institute for Genetics, and †Department of Pathology, University of Cologne, 50931 Cologne, Germany

Contributed by Klaus Rajewsky, July 20, 1994

ABSTRACT Hodgkin disease (HD) is characterized by a small number of putative malignant cells [Hodgkin and Reed–Sternberg (HRS) cells] among a background of lymphocytes and histiocytes. The lineage of HRS cells is still elusive and a clonal origin of these rare cells has not formally been demonstrated. We isolated HRS cells by micromanipulation from histological sections of three cases of Hodgkin lymphoma (each representing a distinct subtype of the disease) and analyzed individual cells for immunoglobulin variable (V) gene rearrangements by PCR. In each of the three cases a single heavy-chain V (V_H) (and in one case, in addition, a κ light-chain) gene rearrangement was amplified from the HRS cells, identifying these cells as members of a single clone. A potentially functional V_H rearrangement was obtained from a case of nodular sclerosis HD. Somatic mutations and intraclonal diversity in the V_H genes indicate a germinal center B-cell origin of the HRS cells in a case of lymphocyte-predominant HD, whereas in a case of mixed-cellularity HD the sequence analysis revealed only nonfunctional V gene rearrangements, suggesting a pre-B-cell origin. This indicates that HRS cells can originate from B-lineage cells at various stages of development.

Hodgkin disease (HD), one of the most common malignant lymphomas, is characterized by a small number of putative malignant mononuclear Hodgkin and multinucleated Reed–Sternberg cells (HRS cells; usually <1% of the tumor) amidst a major population of lymphocytes, plasma cells, histiocytes, neutrophils, eosinophils, and stromal cells. Four subtypes of HD have been described: lymphocyte predominant (LP), nodular sclerosis (NS), mixed cellularity (MC), and lymphocyte depletion (LD) (1). This subclassification is based on the morphology of the HRS cells as well as on differences of the cellular background. HRS cells in the NS, LD, and MC subtypes of HD are characterized by expression of the CD30 and CD15 antigens, whereas HRS cells in the LP subtype often lack these antigens but express B-cell antigens such as CD20 (2).

Although HD has been known for a long time, the lineage derivation of the putative malignant HRS cells is still a matter of controversy. Immunohistochemical studies detected B and/or T-cell or monocyte markers on HRS cells (2), and a recent analysis of gene expression in HRS cells at the single-cell level demonstrated the coexpression of genes characteristic of several hematopoietic lineages (3). The low frequency of HRS cells in the tumor and the difficulty of isolating these cells make molecular biological studies difficult to interpret. Southern blot analysis for immunoglobulin (Ig) or T-cell antigen receptor (TCR) gene rearrangements in

HD tumor tissue (4) does not allow one to determine whether clonal gene rearrangements detected in HD biopsies are derived from the HRS or from other cells.

The detection of Ig and/or TCR gene rearrangements in 11 of 12 HD-derived cell lines thought to originate from HRS cells (5–7) suggests that HRS cells may be of B- and/or T-lineage origin, but the limited success in establishing cell lines from HD precludes generalization of those observations.

A monoclonal origin of HRS cells in at least some instances is suggested by karyotype analysis (8, 9) and by the finding of monoclonal Epstein–Barr virus DNA in some HD tissues in which HRS cells harbor the virus (10, 11). To confirm the monoclonality of HRS cells and to analyze these cells for a possible B-lineage origin we isolated single HRS cells from histological sections by micromanipulation and analyzed them for immunoglobulin heavy-chain and κ light-chain variable (V_H and V_κ) gene rearrangements by polymerase chain reaction (PCR) (12). The amplification of rearranged Ig genes allows one to unambiguously determine the clonality of the cells (13, 14). In addition, the sequence determination of $V_H D_H J_H$ and $V_\kappa J_\kappa$ gene rearrangements obtained by PCR gives a detailed picture of Ig gene rearrangements in HRS cells and discriminates $V_H D_H J_H$ from $D_H J_H$ rearrangements.[§] Such a discrimination is important, since cross-lineage rearrangements of Ig heavy-chain genes have been described in non-B-cell leukemias (15) but appear to be restricted to $D_H J_H$ joints (16–20). Therefore, the present approach of amplifying $V_H D_H J_H$ rearrangements tests HRS cells for a possible B-lineage origin.

MATERIALS AND METHODS

Tissues and Clinical Data. The patients were a 39-year-old man with LP HD at relapse, a 73-year-old man with NS HD, and a 26-year-old man with MC HD. Inguinal lymph nodes were available from the former two patients and a cervical lymph node from the latter.

Immunostaining and Micromanipulation. Eight- to 10- μ m-thick frozen sections were stained with either anti-CD20 (L26), anti-CD3 (OKT3), or anti-CD30 (BerH2) monoclonal antibodies (21). Stained cells were mobilized and aspirated with the help of a micropipette fixed to a hydraulic micromanipulator (12). Multiple cells were picked from each section.

Abbreviations: HD, Hodgkin disease; HRS, Hodgkin and Reed–Sternberg; MC, mixed cellularity; LP, lymphocyte predominant; NS, nodular sclerosis; LD, lymphocyte depletion; CDR, complementarity-determining region; TCR, T-cell antigen receptor.

[†]To whom reprint requests should be addressed at: Institute for Genetics, Weyertal 121, 50931 Cologne, Germany.

[§]Sequences repeatedly obtained from HRS cells have been deposited in the GenBank data base (accession nos. X77415–X77421).

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

Single-Cell PCR. Rearranged V_H - and V_K -region genes were amplified from single cells by using V-gene family-specific primers and J_H and J_K primers as previously described (12). Both strands of the PCR products were directly sequenced.

RESULTS

Immunohistochemical Analysis. Single HRS cells were micro-manipulated from frozen sections stained with anti-CD30 antibodies for HRS cells in a case of NS and a case of MC type (Fig. 1) and anti-CD20 antibodies for HRS cells in a case of LP HD. HRS cells of the cases of NS and MC type were positive for CD30 and CD15 and negative for CD3 and CD20, whereas in the case of LP HD the HRS cells were CD3, CD15, and CD30 negative and CD20 positive (data not shown).

Validity of the Experimental Approach. In all three cases a repeat experiment was carried out, and in four of the six experiments T and B cells were picked on the same day as HRS cells and analyzed together with the HRS cells in a blind fashion. For a total of 50 T cells only one rearranged V gene was obtained, whereas 50 B cells gave rise to 17 PCR products, 13 of which were sequenced (Table 1). These 13 V-gene sequences were all unique and clonally unrelated to rearrangements amplified from HRS cells.

Clonal Heavy-Chain Gene Rearrangements in a Case of NS HD. From 30 single HRS cells of the NS case 12 PCR products indicating a heavy-chain rearrangement were obtained, 11 with a V_{H3} family-specific primer and one with a V_{H4} primer (Table 1). Sequencing of 8 of these products revealed an identical $V_H D_H J_H$ sequence for 7 cells. (In this

Table 1. Summary of single-cell analyses of three cases of Hodgkin lymphoma for Ig V gene rearrangements

Case	Exp.	Cells positive	PCR products		Rearrangements	
			Total	Sequenced	Repeated	Unique
HRS cells						
NS	1	7/20	9	5 [†]	4 V _H 3+	1 V _H 3-
	2*	3/10	3	3	3 V _H 3+	
MC	1	12/20	30	18 [‡]	4 V _H 1-	1 V _H 1+
					6 V _H 3-	1 V _κ 4-
					3 V _κ 1-	1 V _H 3+
					2 V _κ 4-	
	2*	3/5	7	4 [‡]	1 V _H 3-	1 V _κ 4+
LP	1*	3/11	3	3	3 V _H 5-	
	2*	4/17	6	6	3 V _H 5-	
					3 V _H 4+	
Controls						
NS	2	2/10 B	2	2		1 V _H 1?
		0/10 T				1 V _H 4+
MC	2	1/10 B	1	1		1 V _H 1+
		0/10 T				
LP	1	2/10 B	6	3		1 V _κ 2+
		0/10 T				2 V _H 3+
	2	6/20 B	8	7		Multiple [§]
		1/20 T			1	1

For analysis of HRS cells, two independent experiments were carried out for each case. Some experiments (*) were performed as double-blind experiments meaning that HRS cells as well as control (B and T) cells isolated on the same day were analyzed coded in a blinded fashion. The numbers of PCR products obtained and sequenced are indicated. The sequences repeatedly obtained from HRS cells were deposited in the EMBL/GenBank data library under accession nos. X77415–X77421. In each of the three cases the same repeated rearrangements were obtained in the first and second experiments. It is indicated whether the rearrangements are non-functional (–) or potentially functional (+). The functionality of the rearrangement marked by the question mark is unknown due to a sequence ambiguity in complementarity-determining region (CDR) III. For several PCR products no sequences were obtained because PCR products were lost upon purification.

[†]Three of the four PCR products that were not sequenced represent V_{H3} rearrangements.

[‡]Among 15 PCR products that were not sequenced, 8 V_{H1} family rearrangements with a deletion of about 50 bp and 5 V gene rearrangements belonging to the V_{H3} , V_{K1} , and V_{K4} families were obtained.

[§]Clonally unrelated V_H and V_K gene rearrangements of various V gene families (2 V_{H3} , 1 V_{H4} , 1 V_{K1} , 2 V_{K2} , and 1 V_{K4}) were obtained.

and the other two cases studied, sequences were not obtained for some PCR products, because DNA was lost upon purification; see Table 1.) This rearrangement is potentially functional, because the correct reading frame for translation of the Ig gene was preserved in the $V_H D_H J_H$ rearrangement process. A nonfunctional (out-of-frame) V_{H3} rearrangement and a V_{H4} PCR product were each obtained only once and could potentially represent cellular or other contamination.

Clonal Nonfunctional Heavy- and Light-Chain Gene Rearrangements in a Case of MC HD. Twenty-five HRS cells were analyzed for the case of MC HD, with 15 of these cells giving rise to a total of 37 PCR products, 22 of which were sequenced (Table 1). Two nonfunctional V_H and two nonfunctional V_K rearrangements were obtained repeatedly, with five cells harboring two to four of these rearrangements. The V_{H3} rearrangement, amplified from seven cells, uses a V_{H3} gene with 98% homology to the V_{H3} germline gene DP42 and is nonfunctional because of a stop codon in CDR III. The V_{H1} rearrangement, obtained from four cells, uses a V_{H1} gene with 97% homology to the V_{H1} germline gene DP15 and has a 47-bp deletion of codons 56–71 which leads to loss of the

FIG. 1. (Upper) Frozen section of a lymph node with a binucleated CD30-positive Reed-Sternberg cell in the middle of the picture. ($\times 240$.) (Lower) Frozen section after picking, showing a hole where the Reed-Sternberg cell was located, with a small rim of immunostained cell membrane and cytoplasm.

For three cells (two of which had also given rise to a V_H5 PCR product) a potentially functional V_H4 rearrangement was obtained. The V_H4 gene in this structure has 88% sequence homology to the DP64 germline gene (Fig. 3b) and contains a 6-bp insertion between codons 49 and 50. One of the three sequences (LPVH4.3; Fig. 3b) differs in two positions from the other two sequences. To test whether the

FIG. 2. V gene segments derived from single HRS cells of a case of MC HD, compared with the most homologous germline genes. Codons are numbered according to Kabat *et al.* (22). Corresponding amino acids are shown above each codon; asterisks indicate stop codons. CDRs I-III of the V-region gene are indicated. PCR products of single HRS cells with a V_H1 rearrangement (MCVH1) are compared with germline V_H1 gene DP15 (23), rearranged D_H gene DFS-1 (24), and the J_H6 gene (25). The V_H1 rearrangement harbors a deletion of 47 bp from codons 56-71. PCR products of four cells showed an identical sequence. A sequence stretch of six nucleotides homologous to the conserved heptamer recombination signal sequence for Ig genes close to the 5' end of the deletion is underlined. At the 3' end of the deletion a tetranucleotide motif (CCAG; see text) (26) is underlined.

HD Can Result from Malignant Transformation of B-Lineage Cells at Various Stages of Development. For the case of MC HD two nonfunctional V_H and V_K gene rearrangements were obtained from the HRS cells (Table 1; Fig. 2). Whereas two nonfunctional V_K rearrangements might occur in λ light-chain-expressing B cells (30), the identification of a human B-cell clone with two nonfunctional V_H rearrangements has to our knowledge not been described before. Gene targeting experiments in mice have shown that B-cell progenitors unable to express membrane-bound μ heavy chains are arrested in development and presumably die still within the bone marrow (31, 32). In recent work a cellular compartment

LP HD thus appears related to follicular lymphoma, for which a similar cellular composition and ongoing somatic mutation of V-region genes are main characteristics (41).

Concluding Remarks. The present results demonstrate the clonal expansion of HRS cells in each of the three patients analyzed. The most straightforward interpretation of this result is that in each case the disease resulted from the transformation of a single HRS cell. On the other hand, heterogeneity of HRS cells within a patient has been observed in the past at the level of immunophenotype or pattern of gene expression (3, 42). By using specific Ig gene rearrangements as markers, it should now be possible to decide whether such heterogeneity reflects intra- or interclonal cellular diversity. Indeed, the main strength of the present approach for future work lies in the possibility to use HRS cell-specific V-region gene rearrangements as markers in the search for members of the same clone in the various lymphocyte compartments in the body. Perhaps this approach

will allow us to trace HRS cells back to their progenitors and also to follow disease progression in the patient.

While the present results demonstrate that HRS cells can be derived from the B-cell lineage, they do not exclude the possibility of other lineage origins in other cases of HD. This is also indicated by a recent study of single HRS cells from 13 patients, isolated from cell suspensions (43). However, the failure to obtain $V_H D_H J_H$ PCR products from the HRS cells in the latter analysis could at least in part be due to technical matters. A primer mix hybridizing near to the 5' end of the J_H segment was used which may fail to amplify V_H rearrangements using short J_H segments, and a consensus oligonucleotide used as hybridization probe to detect $V_H D_H J_H$ PCR products harbors two or more mismatches to several V_H germline genes. In fact, two of the five V_H rearrangements detected in our analysis may have been undetectable with the probe used by Roth *et al.* (43). In addition, unequivocal identification of HRS cells is more difficult in suspension than on tissue sections.

T-cell-specific surface antigens have been demonstrated on HRS cells in cases of NS, MC, and LD HD, in the absence of B-cell markers (2). In addition, TCR gene rearrangements are present in some HD-derived cell lines (5). These data indicate that HRS cells may also originate from the T-cell lineage. An analysis of TCR gene rearrangements at the level of single HRS cells is needed to further investigate this problem.

We thank Andrea Klöckner for technical help and Ulf Klein and Diane Mathis for helpful discussions. This work was supported by the Deutsche Forschungsgemeinschaft (Di184, Di1284/1-5, and SFB 243), the Dr. Mildred Scheel Stiftung, the Deutsche Krebshilfe, the Fazit Foundation, and the Land Nordrhein Westfalen.

- Lukes, R. J., Butler, J. J. & Hicks, E. B. (1966) *Cancer* **19**, 317-344.
- Drexler, H. G. (1992) *Leuk. Lymph.* **8**, 283-313.
- Trümper, L. H., Brady, G., Bagg, A., Gray, D., Loke, S. L., Griesser, H., Wagman, R., Braziel, R., Gascoyne, R. D., Vicini, S., Cossman, J. & Mak, T. (1993) *Blood* **81**, 3097-3115.
- Weiss, L. M. & Chang, K. L. (1992) *Semin. Diagn. Pathol.* **9**, 272-278.
- Diehl, V., von Kalle, C., Fonatsch, C., Tesch, H., Jücker, M. & Schaadt, M. (1990) *Semin. Oncol.* **17**, 660-672.
- Kanzaki, T., Kubonishi, I., Eguchi, T., Yano, S., Sonobe, H., Ohyashiki, J. H., Ohyashiki, K., Toyama, K., Ohtsuki, Y. & Miyoshi, I. (1992) *Cancer* **69**, 1034-1041.
- Bargou, R. C., Mapara, M. Y., Zugck, C., Daniel, P. T., Pawlita, M., Döhner, H. & Dörken, B. (1993) *J. Exp. Med.* **177**, 1257-1268.
- Teerenhovi, L., Lindholm, C., Pakkala, A., Franssila, K., Stein, H. & Knuutila, S. (1988) *Can. Genet. Cytogenet.* **34**, 305-311.
- Poppema, S., Kaleta, J. & Hepperle, B. (1992) *J. Natl. Cancer Inst.* **84**, 1789-1793.
- Weiss, L. M., Movahed, L. A., Warnke, R. A. & Sklar, J. (1989) *N. Engl. J. Med.* **320**, 502-506.
- Anagnostopoulos, I., Herbst, H., Niedobitek, G. & Stein, H. (1989) *Blood* **74**, 810-816.
- Küppers, R., Zhao, M., Hansmann, M.-L. & Rajewsky, K. (1993) *EMBO J.* **12**, 4955-4967.
- Tonegawa, S. (1983) *Nature (London)* **302**, 575-581.
- Sanz, I. (1991) *J. Immunol.* **147**, 1720-1729.
- Waldmann, T. A. (1987) *Adv. Immunol.* **40**, 247-321.
- Mizutani, S., Ford, A. M., Wiedemann, L. M., Chan, L. C., Furley, A. J. W., Greaves, M. F. & Molgaard, H. V. (1986) *EMBO J.* **5**, 3467-3473.
- Trainor, K. J., Brisco, M. J., Wan, J. H., Neoh, S., Grist, S. & Morley, A. A. (1991) *Blood* **78**, 192-196.
- Segal, G. H., Wittwer, C. T., Fishleder, A. J., Stoler, M. H., Tubbs, R. R. & Kjeldsberg, C. R. (1992) *Am. J. Pathol.* **141**, 1291-1297.
- Ingrihami, G., Szabolcs, M. J., Yee, H. T., Corradini, P., Cesarman, E. & Knowles, D. M. (1993) *Lab. Invest.* **68**, 746-757.
- Reed, T. J., Reid, A., Wallberg, K., O'Leary, T. J. & Frizzera, G. (1993) *Diagn. Mol. Pathol.* **2**, 42-49.
- Guesdon, J.-L., Ternynck, T. & Avrameas, S. (1979) *J. Histochem. Cytochem.* **27**, 1131-1139.
- Kabat, E. A., Wu, T. T., Reid-Miller, M., Perry, H. M. & Gottesman, K. S. (1987) *Sequences of Proteins of Immunological Interest* (U.S. Govt. Printing Office, Bethesda, MD).
- Tomlinson, I. M., Walter, G., Marks, J. D., Llewellyn, M. B. & Winter, G. (1992) *J. Mol. Biol.* **227**, 776-798.
- Pascual, V., Victor, K., Lelsz, D., Spellerberg, M. B., Hamblin, T. J., Thompson, K. M., Randen, I., Natvig, J., Capra, J. D. & Stevenson, F. K. (1991) *J. Immunol.* **146**, 4385-4391.
- Yamada, M., Wasserman, R., Reichard, B. A., Shane, S., Caton, A. J. & Rovera, G. (1991) *J. Exp. Med.* **173**, 395-407.
- Chou, C. L. & Morrison, S. L. (1993) *J. Immunol.* **150**, 5350-5360.
- Kitchingman, G. R. (1993) *Blood* **81**, 775-782.
- Lewis, S. M. & Hesse, J. E. (1991) *EMBO J.* **10**, 3631-3639.
- Sanz, I., Kelly, P., Williams, C., Scholl, S., Tucker, P. & Capra, J. D. (1989) *EMBO J.* **8**, 3741-3748.
- Graninger, W. B., Goldman, P. L., Morton, C. C., O'Brien, S. J. & Korsmeyer, S. J. (1988) *J. Exp. Med.* **167**, 488-501.
- Kitamura, D., Roes, J., Kühn, R. & Rajewsky, K. (1991) *Nature (London)* **350**, 423-426.
- Ehlich, A., Schaal, S., Gu, H., Kitamura, D., Müller, W. & Rajewsky, K. (1993) *Cell* **72**, 695-704.
- Ehlich, A., Martin, V., Müller, W. & Rajewsky, K. (1994) *Curr. Biol.* **4**, 573-583.
- Jacob, J., Kelsoe, G., Rajewsky, K. & Weiss, U. (1991) *Nature (London)* **354**, 389-392.
- Berek, C., Berger, A. & Apel, M. (1991) *Cell* **67**, 1121-1129.
- Poppema, S. (1992) *Int. Rev. Exp. Pathol.* **33**, 53-79.
- Alavaikko, M. J., Hansmann, M.-L., Nebendahl, C., Parwaresch, M. R. & Lennert, K. (1991) *Am. J. Clin. Pathol.* **95**, 194-200.
- Stein, H., Hansmann, M.-L., Lennert, K., Brandtzaeg, P., Gatter, K. C. & Mason, D. Y. (1986) *Am. J. Clin. Pathol.* **83**, 292-297.
- Kuzu, I., Delsol, G., Jones, M., Gatter, K. C. & Mason, D. Y. (1993) *Histopathology* **22**, 141-144.
- Hell, K., Pringle, J. H., Hansmann, M.-L., Lorenzen, J., Colloby, P., Lauder, I. & Fischer, R. (1993) *J. Pathol.* **171**, 137-143.
- Levy, R., Levy, S., Cleary, M. L., Carroll, W., Kon, S., Bird, J. & Sklar, J. (1987) *Immunol. Rev.* **96**, 43-58.
- Chu, W.-S., Abbondanzo, S. L. & Frizzera, G. (1992) *Am. J. Pathol.* **141**, 11-17.
- Roth, J., Daus, H., Trümper, L., Gause, A., Salamon-Looijen, M. & Pfreundschuh, M. (1994) *Int. J. Cancer* **57**, 799-804.