Published in final edited form as:

Addict Behav. 2015 November ; 50: 70–73. doi:10.1016/j.addbeh.2015.06.011.

The Interactive Effect of Paternal Problem Drinking and Maternal Problem Drinking on Adolescent Internalizing Problems

Christine McCauley Ohannessian, Ph.D.^{a,b}

Children's Center for Community Research, Connecticut Children's Medical Center, University of Connecticut School of Medicine

Abstract

Introduction—This study examined the effects of both paternal problem drinking and maternal problem drinking on adolescent internalizing problems (depression and anxiety symptomatology).

Methods—Surveys were administered to 566 10th and 11th grade students from the Mid-Atlantic region of the U.S. in the spring of 2007 and again in the spring of 2008.

Results—Although significant main effects were not observed, significant interactions were found between paternal problem drinking and maternal problem drinking for internalizing problems, especially for boys. In general, these interactions indicated that when paternal problem drinking was high, depression symptomatology and anxiety symptomatology were lower if maternal problem drinking was low.

Conclusions—Findings from this study highlight the need to consider both paternal and maternal problem drinking when examining the effects that parental problem drinking may have on adolescent adjustment.

1. Introduction

Approximately 15% of children in the U.S. live with at least one adult diagnosed with current alcohol abuse or dependence and roughly 30% have been exposed to an alcoholic parent (Grant, 2000). These children are genetically (Cook, 2001; Foroud, Edenberg, & Crabbe, 2010; Palmera, McGeary, & Franco et al., 2012) and environmentally (McGue,

Correspondence: Please direct all inquiries relating to this article to Christine McCauley Ohannessian, Connecticut Children's Medical Center, 282 Washington Street, Hartford, CT 06106. Phone: (860) 837-5326. COhannessian@connecticutchildrens.org or ohannessian@uchc.edu.

^aChildren's Center for Community Research, Connecticut Children's Medical Center, 282 Washington Street, Hartford, CT 06106. Phone: (860) 837-5326. COhannessian@connecticutchildrens.org or ohannessian@uchc.edu.

^bDepartments of Pediatrics and Psychiatry, University of Connecticut School of Medicine, 263 Farmington Avenue, Farmington, CT 06032. Ohannessian@uchc.edu.

Contributors

Christine McCauley Ohannessian was the sole contributor for this manuscript. She conceptualized the study, conducted the data analysis, and wrote the manuscript.

Conflict of Interest

No conflicts of interest are present in the conduct and reporting of this research.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Iacono, Burt, & Elkins, 2004; Palmera et al., 2012) predisposed for the development of substance use and related psychological problems (King & Chassin, 2008). Increased levels of conflict and stress within alcoholic families (Barnow, Schuckit, Lucht, John, & Freyberger, 2002) places children of alcoholic parents (COAs) at an elevated risk for the development of psychological problems as well (Grekin, Brennan, & Hammen, 2005; Griffin, Amodeo, Fassler, Ellis, & Clay, 2005). Indeed, research has shown that COAs are more likely to experience internalizing problems than are children of non-alcoholic parents (non-COAs) (Hussong, Flora, Curran, Chassin, & Sucker, 2008; Ohannessian, 2009; Park & Schepp, 2014).

A major limitation of the literature on alcoholic families is the failure to differentiate between the effects of paternal and maternal alcoholism on COA adjustment. The majority of studies conducted within the field have focused on paternal alcoholism (Leonard & Eiden, 2007). Although most of the available research has indicated that paternal alcoholism has more detrimental effects on COAs (Chassin, Curran, Hussong, & Colder, 1997; Loukas et al., 2001; Ohannessian & Hesselbrock, 1994; Ohannessian et al., 2005), a few studies have found both paternal and maternal alcoholism to predict problematic substance use in COAs (Slutske et al., 2008; Walden et al., 2007). However, most research to date has relied on relatively small samples of alcoholic mothers and clinical indicators of parental alcoholism. Moreover, these studies primarily have focused on substance use outcomes. The differential effects of paternal and maternal alcoholism on other indicators of adjustment (e.g., internalizing problems) need to be systematically addressed. Finally, studies that have examined maternal alcoholism have focused on main effects of parental alcoholism, neglecting to consider the interactive effects of paternal and maternal alcoholism on COA adjustment.

Given these limitations, the goal of the present study was to examine the main and interactive effects that paternal and maternal alcoholism may have on adolescent psychological adjustment (as indicated by internalizing problems). Of note, parental alcoholism was assessed as a continuous variable to capture the broad range of variability likely to be found in a community sample (as such, "parental alcoholism" is referred to as "parental problem drinking" from this point forward). The assessment of parental alcoholism as a continuous variable is consistent with research (Keller, Cummings, Davies, & Mitchell, 2008) that suggests that the severity of parental drinking should be taken into account when examining COA adjustment.

In sum, the following research questions were examined in a diverse sample of adolescents from the U.S. (a) Does paternal problem drinking predict adolescent internalizing symptomatology? (b) Does maternal problem drinking predict adolescent internalizing symptomatology? (c) Do paternal problem drinking and maternal problem drinking *interactively* predict adolescent internalizing symptomatology? (d) Do these relations differ by the gender of the adolescent?

2. Methods

2.1. Participants

All of the adolescents were involved in a larger research project (Ohannessian, 2009). During the spring of 2007 (Time 1), 10th and 11th grade students attending public high schools in the Mid-Atlantic region of the U.S. (Delaware, Maryland, and Pennsylvania) were invited to participate in the study. These participants were invited to participate again during the spring of 2008 (Time 2). The sample included 566 15–18 year-old girls (51%) and boys from seven public high schools. The mean age of the adolescents was 16.04 (SD=.73) at Time 1. Sixty-five percent of the adolescents were Caucasian, 12% were African-American, 12% were Hispanic, and 4% were Asian (the remainder described themselves as “other”). Most of the mothers (96%) and fathers (97%) had graduated from high school. Some of the parents (35% of mothers and 32% of fathers) also had completed four years of college. Only adolescents who lived with both of their biological parents were included in the present study.

2.2. Measures

2.2.1. Parental problem drinking—The 9-item version of the Short Michigan Alcoholism Screening Test (F-SMAST/M-SMAST) (Crews & Sher, 1992) was used to measure alcohol problems in fathers and mothers. A representative SMAST item is “Has your father/mother ever neglected his/her obligations, family, or work for two or more days in a row because he/she was drinking?” The SMAST items were summed to create separate total scores for fathers ($\alpha=.86$) and mothers ($\alpha=.80$). The SMAST has been shown to have strong psychometric properties (Crews & Sher, 1992).

2.2.2. Adolescent depression—The Center for Epidemiological Studies Depression Scale for Children (CES-DC; Weissman, Orvaschel, & Padian, 1980) was used to assess adolescent depressive symptomatology. Participants were asked to respond to the CES-DC items (e.g., “I felt sad”) in respect to how they felt or acted during the past week. The 20 CES-DC items were summed to create a total depression symptomatology score ($\alpha=.90$). The CES-DC has been shown to be a reliable and valid measure of depressive symptomatology (Ohannessian, Lerner, Lerner, & von Eye, 1999).

2.2.3. Adolescent anxiety—The Screen for Child Anxiety Related Disorders (SCARED) (Birmaher, Khetarpal, Cully, Brent, & McKenzie, 1995) was used to measure adolescent anxiety symptomatology. The 41 SCARED items were completed in relation to the past three months. A sample item is “I get really frightened for no reason at all.” The following SCARED scales were included in this study: Panic disorder ($\alpha=.87$), separation anxiety disorder ($\alpha=.73$), social anxiety disorder ($\alpha=.84$), and generalized anxiety disorder ($\alpha=.87$). Prior research has shown that the SCARED has good psychometric properties (Birmaher, Khetarpal, Cully, Brent, & McKenzie, 2003).

2.3. Procedures

The study protocol was approved by the University of Delaware’s IRB. Before data collection, parents were mailed a consent form describing the study. Parents were instructed

to contact the study staff via phone, e-mail, or mail if they did not want their adolescent to participate. Adolescents who had parental consent and provided assent were given a self-report survey by trained research staff (all of whom were certified with human subjects training). The survey took approximately 40 minutes to complete. After completing the survey, the adolescents were given a movie pass.

2.4. Analysis plan

Hierarchical regression models were conducted to examine the main and interactive effects of paternal and maternal problem drinking (assessed at Time 1) on adolescent internalizing symptomatology (assessed at Time 2). Given that depression and anxiety have been shown to differ by gender during adolescence, with girls at a greater risk for both (Graber & Sontag, 2009; Negri & Susman, 2011; Ohannessian et al., 1999), all models were conducted by gender. For both boys and girls, five models were conducted – one for each of the internalizing problems. Step 1 (the main effects model) included paternal problem drinking, maternal problem drinking, and the baseline indicator of internalizing symptomatology. Step 2 (the interaction effects model) included these variables as well as the two-way interaction between paternal problem drinking and maternal problem drinking. Prior to creating the interaction terms, the parental problem drinking variables were centered to aid in the interpretation of the regression weights and to avoid multicollinearity problems (Jaccard & Turrisi, 2003). For the sake of parsimony, the interaction effects models are reported in the results section.

3. Results

3.1. Results for boys

All of the models were significant for boys [$F(4,135)=14.49, p<.001$; $F(4,137)=14.20, p<.001$; $F(4,142)=13.25, p<.001$; $F(4,142)=25.11, p<.001$; $F(4,142)=36.36, p<.001$; for depression, panic disorder, separation anxiety disorder, social anxiety disorder, and generalized anxiety disorder, respectively]. In these models, none of the main effects were significant. However, for boys, significant interactions were found between paternal problem drinking and maternal problem drinking for depression ($\beta=.18, p<.05$), panic disorder ($\beta=.19, p<.05$), and separation anxiety disorder ($\beta=.22, p<.05$). Similar trends emerged for the interactions between paternal problem drinking and maternal problem drinking for social anxiety disorder ($\beta=.12, p<.10$) and generalized anxiety disorder ($\beta=.11, p<.10$). These interactions indicated that when paternal problem drinking was high, depression symptomatology and anxiety symptomatology were lower if maternal problem drinking was low (see Figures 1–2).

3.2. Results for girls

All of the models were significant for girls as well [$F(4,174)=14.65, p<.001$; $F(4,169)=13.87, p<.001$; $F(4,180)=25.06, p<.001$; $F(4,181)=32.50, p<.001$; $F(4,174)=26.16, p<.001$; for depression, panic disorder, separation anxiety disorder, social anxiety disorder, and generalized anxiety disorder, respectively]. None of the main effects were significant. However, similar to boys, a significant interaction was observed between paternal problem drinking and maternal problem drinking for separation anxiety disorder symptomatology for

girls ($\beta=14, p<.05$). When paternal problem drinking was high, separation anxiety symptomatology was lower when maternal problem drinking was low. Of note, separation anxiety symptomatology was highest when both paternal problem drinking and maternal problem drinking were low.

4. Discussion

This study examined the main and interactive effects of paternal problem drinking and maternal problem drinking on adolescent internalizing symptomatology. Consistent with the literature (Hussong et al., 2008; Ohannessian, 2009; Park & Schepp, 2014), parental problem drinking was associated with internalizing problems during adolescence. However, significant main effects for paternal problem drinking and maternal problem drinking were not observed in any of the models for boys or girls. Nonetheless, significant interactions were found between paternal problem drinking and maternal problem drinking for depression and anxiety symptomatology, especially for boys. In general, these interactions indicated that when paternal problem drinking was high, depression symptomatology and anxiety symptomatology were lower if maternal problem drinking was low.

Only one significant finding diverged from this pattern - low paternal and maternal problem drinking were associated with the highest levels of separation anxiety for girls. It may be that separation anxiety depends on the degree of attachment an adolescent has to his/her parents. Indeed, separation anxiety is defined as “developmentally inappropriate and excessive anxiety concerning separation from home or from those to whom the individual is attached” (American Psychiatric Association, 1994). Of note, adolescent COAs report more parental hostility, less parental attachment and warmth, more adolescent-parent conflict, and fewer interactions with their parents in comparison to non-COAs (Barnow et al., 2002). Because of the negative behaviors associated with parental problem drinking (e.g., neglect, abuse, rejection) (MacPherson, Stewart, & McWilliams, 2001), youth with a substance abusing parent simply may be less attached to that parent and therefore, may experience less separation anxiety in relation to that parent. Simply put, some youth may need to feel emotionally close to and attached to a parent for the development of separation anxiety disorder. This may be especially the case for girls given that they are more relationship oriented than are boys during adolescence (Gilligan, 1983).

4.1. Limitations

Results from this study extend the literature; nevertheless, limitations should be noted. Data in the present study were based on adolescents' self-reports. Although self-report measures have been found to yield valid and reliable data (LaForge, Bosari, & Baer, 2005), it would be useful for future investigations to validate findings from this study with other types of methodology. In addition, all of the participants resided in the Mid-Atlantic region of the United States. As such, the findings may not be generalized to adolescents living outside of this area.

4.2. Contributions and conclusions

Despite the limitations, the present study extends the literature by examining both the main *and* interactive effects of paternal *and* maternal problem drinking on adolescent internalizing behaviors. Notably, the large, diverse sample was drawn from the community and followed over time. Moreover, parental problem drinking was assessed continuously in order to capture the wide variation present within the community. These strengths further validate the results, underscoring the need to take *both* parents into account when examining the effects that parental problem drinking may have on adolescent adjustment.

Acknowledgments

Appreciation is extended to the schools and students who participated in the study. Special thanks go to members of the AAP staff, especially Lisa Fong, Kelly Cheeseman, Alyson Cavanaugh, Sara Bergamo, Juliet Bradley, Ashley Malooly, Ashley Ings, Jessica Schulz, Laura Finan, Kaitlin Flannery, and Magdalena Owczarska. This research was supported by NIH grant number K01-AA015059.

Role of Funding Sources

This study was funded by National Institute on Alcohol and Alcoholism (NIAAA) Grant K01-AA015059 awarded to Christine McCauley Ohannessian. NIAAA did not have a role in the study design, collection, analysis, or interpretation of data, writing the manuscript, or the decision to submit the manuscript for publication.

References

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4. Washington, DC: American Psychiatric Association; 1994.
- Barnow S, Schuckit MA, Lucht M, John U, Freyberger HJ. The importance of a positive family history of alcoholism, parental rejection and emotional warmth, behavioral problems and peer substance use for alcohol problems in teenagers: a path analysis. *Journal of Studies on Alcohol and Drugs*. 2002; 63(3):305.
- Birmaher, B.; Khetarpal, S.; Cully, M.; Brent, DA.; McKenzie, S. Screen for child anxiety related disorders (SCARED). Pittsburgh, PA: Western Psychiatric Institute and Clinic, University of Pittsburgh; 1995.
- Birmaher, B.; Khetarpal, S.; Cully, M.; Brent, DA.; McKenzie, S. Screen for child anxiety related disorders (SCARED) – parent form and child form (8 years and older). In: VandeCreek, L.; Jackson, TL., editors. *Innovations in clinical practice: Focus on children & adolescents*. Sarasota, FL: Professional Resource Press; 2003. p. 99-104.
- Chassin, L.; Curran, PJ.; Hussong, AM.; Colder, CR. The relation of parent alcoholism to adolescent substance use: A longitudinal follow-up study. In: Marlatt, GA.; VandenBos, GR., editors. *Addictive behaviors: Readings on etiology, prevention, and treatment*. Washington, DC: American Psychological Association; 1997. p. 509-533.
- Cook, CC. Genetic predisposition to alcohol dependence and problems. In: Heather, N.; Peters, TJ.; Stockwell, T., editors. *International Handbook of Alcohol Problems and Dependence*. New York: John Wiley & Sons; 2001. p. 257-279.
- Crews TM, Sher KJ. Using adapted short MASTs for assessing parental alcoholism: Reliability and validity. *Alcoholism: Clinical and Experimental Research*. 1992; 16(3):576–584.
- Foroud T, Edenberg HJ, Crabbe JC. Genetic Research: Who Is At Risk for Alcoholism? *Alcohol Research & Health*. 2010; 33(1–2):64. [PubMed: 23579937]
- Gilligan, C. *In a different voice: Psychological Theory and Women's Development*. Cambridge, MA: Harvard; 1983.
- Graber, JA.; Sontag, LM. Internalizing problems during adolescence. In: Lerner, RM.; Steinberg, L., editors. *Handbook of adolescent psychology, Vol 1: Individual bases of adolescent development*. 3. Hoboken, NJ, US: John Wiley & Sons Inc; 2009. p. 642-682.

- Grant BF. Estimates of US children exposed to alcohol abuse and dependence in the family. *American journal of public health*. 2000; 90(1):112. [PubMed: 10630147]
- Grekin ER, Brennan PA, Hammen C. Parental alcohol use disorders and child delinquency: the mediating effects of executive functioning and chronic family stress. *Journal of Studies on Alcohol and Drugs*. 2005; 66(1):14.
- Griffin ML, Amodeo M, Fassler I, Ellis MA, Clay C. Mediating Factors for the Long-term Effects of Parental Alcoholism in Women: The Contribution of Other Childhood Stresses and Resources. *The American Journal on Addictions*. 2005; 14(1):18–34. [PubMed: 15804874]
- Hussong AM, Flora DB, Curran PJ, Chassin LA, Zucker RA. Defining risk heterogeneity for internalizing symptoms among children of alcoholic parents. *Development and psychopathology*. 2008; 20(01):165–193. [PubMed: 18211733]
- Jaccard, J.; Turrisi, R. *Interaction effects in multiple regression*. 2. Thousand Oaks, CA: Sage Publications; 2003.
- Keller PS, Cummings EM, Davies PT, Mitchell PM. Longitudinal relations between parental drinking problems, family functioning, and child adjustment. *Development and Psychopathology*. 2008; 20(1):195. [PubMed: 18211734]
- King KM, Chassin L. Adolescent stressors, psychopathology, and young adult substance dependence: A prospective study. *Journal of Studies on Alcohol and Drugs*. 2008; 69(5):629. [PubMed: 18781237]
- Laforge RG, Borsari B, Baer JS. The utility of collateral informant assessment in college alcohol research: results from a longitudinal prevention trial. *Journal of Studies on Alcohol*. 2005; 66(4):479–487. [PubMed: 16240555]
- Leonard KE, Eiden RD. Marital and family processes in the context of alcohol use and alcohol disorders. *Annual Review of Clinical Psychology*. 2007; 3:285.
- Loukas A, Fitzgerald HE, Zucker RA, von Eye A. Parental alcoholism and co-occurring antisocial behavior: Prospective relationships to externalizing behavior problems in their young sons. *Journal of Abnormal Child Psychology*. 2001; 29(2):91–106. [PubMed: 11321632]
- MacPherson PS, Stewart SH, McWilliams LA. Parental problem drinking and anxiety disorder symptoms in adult offspring: examining the mediating role of anxiety sensitivity components. *Addictive Behaviors*. 2001; 26(6):917–934. [PubMed: 11768552]
- McGue M, Iacono WG, Burt SA, Elkins I. Identifying shared environmental contributions to early substance use: The respective roles of peers and parents. *Journal of Abnormal Psychology*. 2004; 113:440–450. [PubMed: 15311989]
- Minuchin, P. Looking toward the horizon: Present and future in the study of family systems. In: McHale, JP.; Grolnick, WS., editors. *Retrospect and prospect in the psychological study of families*. Mahway, NJ: Lawrence Erlbaum Associates Publishers; 2002. p. 59–278.
- Negriff S, Susman EJ. Pubertal timing, depression, and externalizing problems: A framework, review, and examination of gender differences. *Journal of Research on Adolescence*. 2011; 21(3):717–746.
- Ohannessian CM. Media use and adolescent psychological adjustment: An examination of gender differences. *Journal of child and family studies*. 2009; 18(5):582–593. [PubMed: 21359124]
- Ohannessian CM, Hesselbrock VM, Tennen H, Affleck G. Hassles and uplifts and generalized outcome expectancies as moderators on the relation between a family history of alcoholism and drinking behaviors. *Journal of Studies on Alcohol and Drugs*. 1994; 55(6):754.
- Ohannessian CM, Hesselbrock VM, Kramer J, Kuperman S, Bucholz KK, Schuckit MA, Nurnberger JI. The Relationship Between Parental Psychopathology and Adolescent Psychopathology An Examination of Gender Patterns. *Journal of Emotional and Behavioral Disorders*. 2005; 13(2):67–76.
- Ohannessian CM, Lerner RM, Lerner JV, von Eye A. Does self-competence predict gender differences in adolescent depression and anxiety? *Journal of adolescence*. 1999; 22(3):397–411. [PubMed: 10462430]
- Palmer RHC, McGeary JE, Francazio S, Raphael BJ, Lander AD, Heath AC, Knopik VS. The genetics of alcohol dependence: advancing towards systems-based approaches. *Drug and alcohol dependence*. 2012; 125(3):179–191. [PubMed: 22854292]

- Park S, Schepp KG. A systematic review of research on children of alcoholics: Their inherent resilience and vulnerability. *Journal of Child and Family Studies*. 2014:1–10.
- Slutske WS, D'Onofrio BM, Turkheimer E, Emery RE, Harden KP, Heath AC, Martin NG. Searching for an environmental effect of parental alcoholism on offspring alcohol use disorder: a genetically informed study of children of alcoholics. *Journal of Abnormal Psychology*. 2008; 117(3):534. [PubMed: 18729607]
- U.S. Census Bureau. US Census Bureau: State and county QuickFacts 2008. 2008. Retrieved from <http://quickfacts.census.gov/qfd/states/10/10003.html>
- Walden B, Iacono WG, McGue M. Trajectories of change in adolescent substance use and symptomatology: Impact of paternal and maternal substance use disorders. *Psychology of Addictive Behaviors*. 2007; 21(1):35. [PubMed: 17385953]
- Weissman MM, Orvaschel H, Padian N. Children's symptom and social functioning self-report scales comparison of mothers' and children's reports. *The Journal of Nervous and Mental Disease*. 1980; 168(12):736–740. [PubMed: 7452212]

Highlights

- Paternal and maternal problem drinking interactively predicted depression for boys.
- Paternal and maternal problem drinking interactively predicted anxiety for boys and girls.
- Results underscore the need to consider both paternal and maternal problem drinking.

Figure 1.
Depression Symptomatology for Boys

Figure 2.
Panic Disorder Symptomatology for Boys