Published in final edited form as:

European J Org Chem. 2015 March ; 2015(7): 1602–1605. doi:10.1002/ejoc.201403615.

Alkyl Sulfinates: Formal Nucleophiles for Synthesizing TosMIC Analogs

J. Armando Lujan-Montelongo^{[a],[b]}, Angel Ojeda Estevez^[b], and Fraser F. Fleming^[b]

J. Armando Lujan-Montelongo: jalujanm@cinvestav.mx; Fraser F. Fleming: flemingf@duq.edu

^[a]Departamento de Química, Centro de Investigación y Estudios Avanzados del IPN, Av. Instituto Politecnico Nacional 2508, San Pedro Zacatenco, 07360 Ciudad de México, Distrito Federal, Mexico, <http://www.quimica.cinvestav.mx/Directorio/Investigadores/DrLujanMontelongoJesus.aspx>

^[b]Department of Chemistry and Biochemistry Duquesne University 600 Forbes Ave, Pittsburgh, PA 15202, USA, <http://www.scienceresearch.duq.edu/chem/chemfac/ffleming/>

Abstract

Alkyl sulfinates function as formal nucleophiles in Mannich-type reactions to give sulfonyl formamides, which are readily dehydrated to the corresponding sulfonylmethyl isonitriles. The efficient, two-step synthesis provides a general route to sulfonylmethyl isonitriles from readily available methyl sulfinates or thiols. Mechanistic analysis reveals that the unusual nucleophilicity of the alkyl sulfinates arises from the in situ release of sulfinic acids.

Keywords

Sulfinates; Isonitriles; Umpolung; Isocyanides; Nucleophilic addition; Mannich reaction

Introduction

Sulfonylmethyl isonitriles such as TosMIC [(tolylsulfonyl)methyl isocyanide; Scheme 1; compound **3**, $R^1 = p\text{Tol}$], are extremely valuable precursors to multifarious isonitriles,^[1] heterocycles,^[2] and N-heterocyclic carbene complexes.^[3] Traditionally, metal sulfinates **1** have featured prominently in the stoichiometric^[4] and metal-catalyzed^[5] syntheses of sulfonylmethyl isonitriles and sulfones.^[6] The versatility of metal sulfinates **1** stems from the potent nucleophilicity^[7] of the central sulfur atom, whose reaction with electrophiles directly generates sulfones (Scheme 1; **1** → **2**). The interception of metal sulfinates with iminium ions forges aminomethyl sulfones^[8] **2**, which are readily dehydrated to give versatile sulfonyl isonitriles **3**.

Esterification of metal sulfinates **1** inverts the reactivity at the central sulfur atom to create alkyl sulfinates **4**, which are potent electrophiles with reactivities comparable to those of the

corresponding sulfonyl chlorides^[9] (Scheme 1). Organometallic addition to alkyl sulfonates provides a valuable route to sulfoxides **5**, particularly chiral sulfoxides.^[10]

Although alkyl sulfonates are electrophilic, the presence of the two lone pairs of electrons on the adjacent oxygen atom increases the nucleophilicity of the central sulfur atom.^[11] The potential nucleophilicity of alkyl sulfonates make them an attractive replacement for metal sulfonates **1**, which suffer from three significant disadvantages: few metal sulfonates are commercially available, many show modest solubility, and often metal sulfonates are not particularly stable.^[12] Alkyl sulfonates, in contrast, have good stability, excellent solubility, and are rapidly synthesized from commercially available disulfides or thiols.^[13] Using alkyl sulfonates as formal nucleophiles in Mannich reactions addresses the longstanding challenge of synthesizing structurally diverse sulfonylmethyl isonitriles (Scheme 1; **4** → **2** → **3**).

Results and Discussion

In exploratory experiments, commercially sourced methyl benzenesulfonate (**4a**) was employed as a prototype in a Mannich reaction with formamide, formic acid, and paraformaldehyde (Scheme 2). In initial reaction optimization, microwave heating was used to facilitate precise temperature control, and formamide **2a** was formed at temperatures above 50 °C. Further experimentation identified 90–110 °C as optimal; at lower temperatures the reaction rate was slow, whereas higher temperatures resulted in lower yields. Solvent screening revealed an essential role for toluene, which presumably reflects the unusual solvation; the bi-phasic mixture^[14] formed at room temperature coalesces into a single phase on heating. After 2–3 h, formamide **2a** was obtained in high yield. The crude formamide (i.e., **2a**) was formed cleanly and efficiently, and, because of the strong absorption during silica gel chromatography,^[15] was dehydrated without prior purification to directly give isocyanide **3a** (72% yield over two steps).

The generality of the two-step Mannich–dehydration sequence was ascertained by converting a series of methyl sulfonates into the corresponding sulfonylmethyl isonitriles (i.e., **3**; Table 1). Varying the electronic nature of the sulfonate had a minimal impact on the reaction efficiency (Table 1; compare Entries 1–3, 4–7, 8–10, and 11–13). The efficiency is sensitive to steric compression adjacent to the sulfonate, with *p*-tolylsulfonate **4b** reacting significantly more efficiently than *o*-tolylsulfonate **4c** (Table 1; Entries 1 and 2). Similarly, *o*-phenoxysulfonate **4j** reacted more efficiently than *o*-methoxysulfonate **4i** (Table 1; Entries 8 and 9).^[16] In general, the isolated yields of aliphatic sulfonylmethyl isonitriles **3l–3n** were lower, which reflects an instability of the isonitriles to purification and storage.^[15] With the exception of **3l**, all the isonitriles had minimal odor.

In most instances, the reaction was equally efficient with microwave or conventional heating. Comparative microwave and conventional heating gave **4h** in yields of 89 and 87%, respectively. However, the reactions that generated **4m** and **4n** with microwave heating were sluggish, whereas conventional heating significantly improved the conversion. Although speculative, the pressure increase that accompanies microwave irradiation, and not conventional heating, may be responsible for the difference in efficiency.

Insight into the mechanism was obtained by comparing the relative conversion rates of a series of alkyl toluene-sulfonates in the Mannich reaction (Figure 1). Increasing the steric demand of the alkyl substituent from methyl to ethyl and isopropyl decreased the conversion rate. A similar rate correlation occurs in the NBS-promoted racemization of alkyl sulfonates, where methyl arylsulfonates racemize ten times faster than sterically more demanding isopropyl sulfonates.^[11b]

The dependence of the conversion rate on alkyl sulfonate steric demand is consistent with at least two different mechanisms: nucleophilic attack from the sulfonate followed by alkyl cleavage of a sulfurane intermediate^[17] (Scheme 3; **4o** → **7o** → **2o**); or alkyl cleavage followed by nucleophilic attack from a sulfonic acid (**4o** → **9o** → **10o** → **2o**). Phenethyl sulfonate **4o**^[18] was selected as a probe to differentiate between these two mechanisms, because the substantial molecular weight of the phenethyl substituent facilitates tracking the fate of the alkyl fragment. NMR spectroscopic analysis of the crude reaction mixture showed equal amounts of formate **8** and formamide **2o**,^[19] which is consistent with an alkylation of formic acid by sulfonium salt **7o**, but does not distinguish between the two mechanisms. However, a control experiment in which **4o** was treated with formic acid in formamide *without* paraformaldehyde gave formate **8**, which implies that alkyl cleavage occurs before sulfonate attack. The mechanistic experiments suggest that the reaction proceeds by sulfonate protonation (**4o** → **9o**)^[20] and formate alkylation by the sulfonium salt^[21] to generate a sulfonic acid **10o**^[22] as the true nucleophilic species, and that this then intercepts the iminium ion (i.e., **6**).^[23]

Insight into the reaction mechanism stimulated a second approach to sterically congested sulfonylmethyl isonitriles. The new strategy harnessed the greater nucleophilicity of sulfides to address the challenge of preparing isonitriles with sterically demanding alkyl substituents (Table 2).^[24] Subjecting dimethoxythiophenol **11p** to the standard Mannich conditions gave the corresponding formamide (i.e., **12p**), which was sequentially oxidized with *m*CPBA (3-chloroperbenzoic acid) and dehydrated to give **3p** (Table 2; Entry 1). Analogous three-step sequences with the 2,6-di-substituted thiol **11q**, 1,2-substituted naphthalene **11r**, and even adamantane **11t** formed the corresponding isonitriles (Table 2; Entries 2 and 3). None of these isonitriles (i.e., **3p–3t**) were accessible by the methyl sulfonate method. Subjecting menthol-derived thiol **11s** to the Mannich–oxidation–dehydration sequence efficiently gave isocyanide **3s**, thus demonstrating the viability of preparing chiral, secondary sulfonylmethyl isonitriles (Table 2; Entry 4).

Conclusions

A diverse array of sulfonylmethyl isonitriles were easily prepared in Mannich-type condensations with alkyl sulfonates or thiols. The strategy features a formal polarity reversal of alkyl sulfonates through in situ cleavage to nucleophilic sulfonic acids. The resulting formamides are readily dehydrated to provide an efficient synthesis of sulfonylmethyl isonitriles. This method demonstrates that alkyl sulfonates are a practical alternative to metal sulfonates in providing access to a diverse range of sulfonylmethyl isonitriles.

Experimental Section

Representative Experimental Procedure

The methyl sulfinate, paraformaldehyde (5 equiv.), formamide (6 equiv.), formic acid (5 equiv.), and toluene (5 equiv.) were added sequentially to a Biotage[®] microwave vial. The vial was capped and purged with N₂, and then it was irradiated at 100 °C. In cases where the internal pressure rose above 20 psi, the vial was vented after 30 min, and then the heating was continued. After 3 h, the contents were poured onto an ice/water mixture, and the resulting mixture was extracted with EtOAc (4×). The combined organic layers were washed with brine, and dried (Na₂SO₄). Some formamides solidified upon cooling, and could be recrystallized from benzene/pentane. All form-amides were sufficiently pure to be used directly in the dehydration reaction (after complete removal of volatiles).

A round-bottomed flask containing the crude formamide was purged with N₂ (3×), and then a 2:1 mixture of THF/MeCN (1.5 M) was added to dissolve the formamide. The flask was cooled to –10 °C, *i*Pr₂NH was added dropwise (9.3 equiv.), and POCl₃ (3.3 equiv.) was added dropwise at a sufficiently slow rate to keep the temperature below 5 °C. After 1 h, the mixture was poured onto a 50:50 mixture of ice/NaHCO₃ (satd. aq.). The resulting mixture was extracted with CH₂Cl₂ (4×), and the combined organic layers were washed with brine and dried (Na₂SO₄). The crude products were prepurified by passing through a short silica gel plug, eluting with hexanes/diethyl ether (70:30), and then purified by radial chromatography. Complete experimental details are provided in the Supporting Information.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support for this research from the NIH (2R15AI051352-04), CONACYT, and release time from the National Science Foundation (NSF) (IRD) is gratefully acknowledged. The opinions expressed in this manuscript are those of the authors and do not necessarily reflect the views of the NSF or the NIH.

References

1. Chakrabarty S, Choudhary S, Doshi A, Liu F-Q, Mohan R, Ravindra MP, Shah D, Yang X, Fleming FF. *Adv Synth Catal.* 2014; 356:2135–2196. [PubMed: 25484847]
2. a) Suginome M, Ito Y. *Sci Synth.* 2012; 44:445–531. b) Van Leusen D, Van Leusen AM. *Org React.* 2001; 57:417–666.
3. a) Manzano R, Rominger F, Hashmi ASK. *Organometallics.* 2013; 32:2199–2203. b) Blanco Jaimes MC, Böhling CRN, Serrano-Becerra JM, Hashmi ASK. *Angew Chem Int Ed.* 2013; 52:7963–7966. *Angew Chem.* 2013; 125:8121–8124. c) Hashmi ASK, Yu Y, Rominger F. *Organometallics.* 2012; 31:895–904. d) Hashmi ASK, Riedel D, Rudolph M, Rominger F, Oeser T. *Chem Eur J.* 2012; 18:3827–3830. [PubMed: 22378616] e) Hashmi ASK, Lothschütz C, Graf K, Häffner T, Schuster A, Rominger F. *Adv Synth Catal.* 2011; 353:1407–1412. f) Hashmi ASK, Lothschütz C, Böhling C, Hengst T, Hubbert C, Rominger F. *Adv Synth Catal.* 2010; 352:3001–3012.
4. For recent examples, see: Xiao F, Chen H, Xie H, Chen S, Yang L, Deng G-J. *Org Lett.* 2014; 16:50–53. [PubMed: 24328422] Liang S, Zhang R-Y, Xi L-Y, Chen S-Y, Yu X-Q. *J Org Chem.* 2013; 78:11874–11880. [PubMed: 24251813] Orgies DH, Forgione P. *Synlett.* 2013; 24:1715–

1721. Umierski N, Manolikakes G. *Org Lett.* 2013; 15:4972–4975. [PubMed: 24047409] Umierski N, Manolikakes G. *Org Lett.* 2013; 15:188–191. [PubMed: 23252771]
5. For recent examples, see: Li H-S, Liu G. *J Org Chem.* 2014; 79:509–516. [PubMed: 24354639] Deeming AS, Russell CJ, Hennessy AJ, Willis MC. *Org Lett.* 2014; 16:150–153. [PubMed: 24308313] Lin X, Wang G, Li H, Huang Y, He W, Ye D, Huang K-W, Yuan Y, Weng Z. *Tetrahedron.* 2013; 69:2628–2632. Chawla R, Singh AK, Yadav LDS. *Tetrahedron.* 2013; 69:1720–1724.
6. Sulfonylmethyl isonitriles are typically synthesized by a multistep sequence involving sulfinate displacement, oxidation, and dehydration: Hundscheld FJA, Tandon VK, Rouwette PHFM, van Leusen AM. *Tetrahedron.* 1987; 43:5073–5088.
7. Baidya M, Kobayashi S, Mayr H. *J Am Chem Soc.* 2010; 132:4796–4805. [PubMed: 20225879]
8. a) Buschmann, H.; Puetz, C. WO 2003068204 A1. 2003. b) Barendse, NCME. US 4922016A. 1990.
9. García Ruano JL, Parra A, Yuste F, Mastranzo VM. *Synthesis.* 2008:311–312.
10. O'Mahony GE, Kelly P, Lawrence SE, Maguire AR. AR-KIVOC (Gainesville, FL, US). 2011:1–110.
11. a) Kice JL, Cleveland JP. *J Am Chem Soc.* 1973; 95:104–109. b) Drabowicz J. *Phosphorus Sulfur Relat Elem.* 1987; 31:123–131.
12. Kamiyama T, Enomoto S, Inoue M. *Chem Pharm Bull.* 1988; 36:2652–2653.
13. Brownbridge P, Jowett IC. *Synthesis.* 1988:252–254.
14. The use of formamide alone resulted in a severe increase of temperature and pressure in the reaction vessel, triggering the microwave safety sensor to halt further heating.
15. a) Kotha S, Sreenivasachary N. *Bioorg Med Chem Lett.* 1998; 8:257–260. [PubMed: 9871665] b) Kotha S, Brahmachary E, Sreenivasachary N. *Tetrahedron Lett.* 1998; 39:4095–4098.
16. Although the A value is higher for a phenoxy group than for a methoxy group, the edge-on steric demand of rotationally restricted phenyl groups significantly lowers the steric demand: Eliel EL, Wilen SH, Mander LN. *Stereochemistry of Organic Compounds.* Wiley New York 1994:597–606. 698. and particularly
17. For a related mechanism, see: Wenschuh E, Fahsl R, Hoehne R. *Synthesis.* 1976:829–830.
18. Attempts to hydrolyze methyl benzenesulfinate (**4a**) to the corresponding sulfinic acid gave a mixture of sulfinic and sulfonic acids.
19. Purification gave sulfone **2o** in 40% yield, and formate **8** in 22% yield. Presumably, the lower isolated yield of formate **8** is due to easy hydrolysis during silica gel chromatography.
20. Sulfonates derived from benzylic alcohols can ionize and recombine to directly form sulfones, but this mechanism appears to be unlikely, because most of the sulfonates (i.e. **4**) do not contain stabilizing benzylic or allylic substituents: Drabowicz J, Bujnicki B, Biscarini P, Mikolajczyk M. *Tetrahedron: Asymmetry.* 1999; 10:3177–3187.
21. Direct hydrolysis of the sulfinate by adventitious water provides a conceivable, but unlikely, alternative route to sulfinic acid **10**. Sulfinate hydrolysis would concomitantly release phenylethanol, but the presence of this compound in the crude reaction mixture could not be definitively excluded. Independently subjecting phenylethanol to the reaction conditions did cause esterification to give formate **8** (21% yield), but the reaction was incomplete, so unreacted phenylethanol should be detected in the crude reaction mixture if a hydrolysis mechanism was operating.
22. For a closely related mechanism, see: Li H-H, Dong D-J, Jin Y-H, Tian SK. *J Org Chem.* 2009; 74:9501–9504. [PubMed: 19911839]
23. A mechanistically distinct cleavage–recombination mechanism is possible for the conversion of allylic and benzylic sulfinate esters to sulfones. [22] For selected examples, see: Douglass IB, Farah BS. *J Org Chem.* 1958; 23:805–807. Cope AC, Morrison DE, Field L. *J Am Chem Soc.* 1950; 72:59–67. c) ref. [6] and references cited therein.
24. van Leusen AM, Boerma GJM, Helhholdt RB, Siderius H, Strating J. *Tetrahedron Lett.* 1972; 13:2367–2368.

Figure 1.
Influence of steric demand of the R group of $p\text{TolSO}_2\text{R}$ on conversion.

Alkyl sulfinates: electrophiles

Alkyl sulfinates: **formal nucleophiles**

Scheme 1.
Contrasting reactivity of sulfinate anions and esters.

Scheme 2.
Methyl sulfinate Mannich–dehydration sequence.

Scheme 3.
Mechanistic pathways for the sulfinate Mannich reaction.

Table 1

Conversion of methyl sulfonates to sulfonyl isonitriles.

$ \begin{array}{c} \text{R}^1\text{-SO}_2\text{OMe} \xrightarrow[\text{HCO}_2\text{H, } -(\text{CH}_2\text{O})_n]{\text{H}_2\text{NCHO}} \text{R}^1\text{-SO}_2\text{OCH}_2\text{NHC(=O)H} \xrightarrow[\text{iPr}_2\text{NH}]{\text{POCl}_3} \text{R}^1\text{-SO}_2\text{OCH}_2\text{NC} \\ \mathbf{4} \qquad\qquad\qquad \mathbf{2} \qquad\qquad\qquad \mathbf{3} \end{array} $			
Entry	Sulfonate	Isonitrile	Yield ^[d]
1			71% ^[b]
	4b	3b	
2			57% ^[b]
	4c	3c	
3			70% ^[a]
	4d	3d	
4			81% ^[b]
	4e	3e	
5			71% ^[b]
	4f	3f	
6			57% ^{[a][c]}
	4g	3g	
7			72% ^{[a][c]}
	4h	3h	
8			60% ^[b]
	4i	3i

Entry	Sulfinate	Isonitrile	Yield ^[d]
9			72% ^[a]
	4j	3j	
10			45% ^[a]
	4k	3k	
11			45% ^[b]
	4l	3l	
12			51% ^[a]
	4m	3m	
13			57% ^[a]
	4n	3n	

^[a] Reaction was performed by heating with an oil bath.

^[b] Reaction was performed in a microwave reactor.

^[c] Et₃N was used instead of *i*Pr₂NH.

^[d] Yield over two steps after purification.

Table 2

Three-step sulfide to isonitrile synthesis.

Reaction scheme showing the three-step synthesis of isonitriles from thiols:

$$\text{R}^1\text{-SH} \xrightarrow[\text{-(CH}_2\text{O)}_n]{\text{H}_2\text{NCHO, HCO}_2\text{H}} \text{R}^1\text{-S-CH}_2\text{-NH-C(=O)-H} \xrightarrow[\text{2. POCl}_3, \text{ } i\text{Pr}_2\text{NH}]{\text{1. } m\text{-CPBA}} \text{R}^1\text{-S(=O)}_2\text{-CH}_2\text{-NC}$$

Entry	Thiol	Isonitrile	Yield ^[d]
1 ^[a]	 11p	 3p	40%
2 ^[a,c]	 11q	 3q	46%
3 ^[b]	 11r	 3r	56%
4 ^[a]	 11s	 3s	55%
5 ^[a]	 11t	 3t	32%

^[a] Heating was performed with a conventional oil bath.^[b] Microwave heating was used.^[c] Et₃N was used instead of *i*Pr₂NH.^[d] Yields over three steps after purification.